

## ELEPHANTS, WOODLANDS AND BIODIVERSITY IN SOUTHERN AFRICA

By: Cumming, D.H.M., Fenton, M.B., Rautenbach, I.L., Taylor, R.D., Cumming, G.S., Cumming, M.S., Dunlop, J.M., Ford, G.S., Hovorka, M.D., Johnston, D.S., Kalcounis, M.C., Mahlangu, Z., and C.V. Portfors

Cumming, D.H.M., Fenton, M.B., Rautenbach, I.L., Taylor, R.D., Cumming, G.S., Cumming, M.S., Dunlop, J.M., Ford, G.S., Hovorka, M.D., Johnston, D.S., [Kalcounis, M.C.](#), Mahlangu, Z., and C.V. Portfors. 1997. Elephants, woodlands and biodiversity in miombo woodland in southern Africa . *South African Journal of Science* 93: 231-236.

Made available courtesy of F R D: <http://www.sajs.co.za/>

*Please note that copyright was held by the FRD, now the National Research Foundation, at the time that issue was published but now resides with the ASSAf.*

**\*\*\*Note: Figures may be missing from this format of the document**

### **TABLE 1 CAN BE FOUND AT THE END OF THE ARTICLE**

When elephant densities exceed approximately 0.5 per km<sup>2</sup>, savanna woodlands are generally converted to shrub-lands or grasslands. The impact of such elephant-mediated habitat change on biodiversity in African game reserves has seldom been measured. We examined species richness of woody plants, birds, bats, mantises and ants in reserves where elephants had destroyed the miombo woodland and in adjacent but intact miombo woodlands outside the reserves. Species richness of woodland birds and ants was significantly lower where elephants had removed the tree canopy. Our findings may have important policy implications for conserving biodiversity in many African reserves in the face of rapidly growing elephant populations (approximately 5% per annum). The problem is further compounded by international public pressures against reducing elephant densities within game reserves while, outside these protected areas, savanna woodlands and their associated faunas are being lost to agriculture. Where then will refugia for habitat-sensitive species exist if not within the region's largest protected areas?

Preserving large populations of elephants while maintaining bio-diversity in national parks and protected areas in East and southern Africa is becoming increasingly problematic. In southern Africa human and elephant populations are growing at rates of about 3% and 5% per annum, respectively[1-5] and in some areas wildland is being converted to subsistence agriculture at similar rates.[6] The results are further reductions in elephant range, increased density of elephants within protected areas[7,8] and human expansion into marginal lands.[1] Together these processes are leading to the deforestation of large areas of savanna woodland of high biodiversity but low agricultural productivity.[9-11] Besides affecting biodiversity, deforestation, particularly of upland woodlands, is likely to affect seasonal patterns of water storage, discharge and stream flow from otherwise protected landscapes.[12,13]

Contrary to popular perceptions, elephant numbers were not always high in East and southern Africa and high densities such as presently occur in many national parks were seldom encountered by early explorers. In southern Africa elephant numbers reached very low levels by the 1880s due to hunting and by 1900 their extinction south of the Zambezi was considered likely.[14,15] Elephant numbers at the turn of the century in Zimbabwe, for example, were less than 5 000 and the human population was below 500 000 compared to present numbers of more than 60 000 and 11 million, respectively. [16,17]

Large generalist herbivores in the absence of their predators can have devastating effects on biodiversity; introduced domesticated species, such as goats on oceanic islands, have provided spectacular examples.[18] The African elephant is an example of a large, generalist herbivore which has often transformed natural habitats in game reserves[19-32] where it has been protected from human predation. In many East African parks and reserves widespread elephant impacts were greatly reduced during the 1970s and 1980s by illegal hunting for ivory.[33,34] In southern Africa, where elephant populations are still growing,[4,5,35] attempts by wildlife management authorities to mitigate effects on woodlands by culling elephants have been resisted by public pressure groups both within Africa and beyond.[36-41] Ecological counter-arguments to culling have also been advanced[42-45] but elephant impacts on biodiversity, as opposed to their influence on vegetation structure and scenery, have seldom been considered in the debate because of the absence of appropriate studies. Recent exceptions are the studies of elephant impacts on plant species diversity of succulent thicket in the Eastern Cape of South Africa[46] and effects on bird diversity in northern Botswana.[47]

Miombo woodlands (species-rich, broad-leaved, deciduous woodlands dominated by trees of the genus *Brachystegia* and *Julbernardia*) prevail over much of southern central Africa from Tanzania south to Zimbabwe and from Angola across to Mozambique.[48] Throughout much of this region woodland has been cleared for cultivation, and in parts of the Zambezi valley in Zimbabwe this has occurred at the rate of 4% per annum over the last three decades. [6]High elephant densities, combined with fire, have had a major effect on formerly pristine miombo woodlands in many protected areas of the region and particularly in Zimbabwe, where veterinary control fences have confined expanding elephant populations to protected areas. Although elephant impact on miombo woodland structure and woody species is well known, [23-25] the effect of high elephant densities on other taxa has been neglected. We therefore compared woodland structure and biodiversity in elephant-affected miombo woodland (within protected areas) with biodiversity in equivalent, undisturbed, or intact habitats outside protected areas.

### *Study areas and methods*

We worked in miombo woodlands in the southern escarpment of the Zambezi valley in Zimbabwe (Fig. 1). Elephants were seldom present in the intact woodlands (e.g. Fig. 1b) until recently, whereas elephant densities in the affected woodlands (e.g. Fig. 1c) had exceeded 1 elephant per km<sup>2</sup> since the mid-1970s (Department of National Parks and Wild Life Management, unpub. censuses). In the Mana area (Fig. 2) intact and impacted woodlands were separated by a game fence, while in the the Kanyati/Matusadona area they were separated by the Sanyati Gorge, which is impassable to elephants. The game fence on the southern boundary of the Mana Pools National Park and the Hurungwe Safari Area was erected in 1968, as part of

tsetse control operations in the region, to prevent the southward movement of elephant and other game into the adjacent Hurungwe Communal Lands.

In the Mana area on each of nine nights between 15 and 24 November 1994, we sampled an intact woodland site north of the fence and an impacted woodland site south of the fence to provide nine paired samples. In the Kanyati woodlands six intact sites were sampled over three nights (26-28Nov.), followed by samples from six impacted sites, also over three nights, in the Matusadona woodlands (30 Nov.-2 Dec.). The distance between the Kanyati and Matusadona areas (approximately 40 km) precluded simultaneous sampling of intact and impacted sites. Our sample sites were near the crests of ridges in undulating terrain and were in similar geological formations and soil types. On each sampling night two teams simultaneously sampled bats and insects at two sites and the next morning at each site we listed birds and sampled vegetation.

Woody vegetation was sampled in two ways. At each site all species, in three size classes (small shrubs <1 m; shrubs 1-3 m; trees >3 m), were recorded within an area of about 1 ha to provide a measure of species richness. The number of woody plants by species in each of the three size classes was then recorded in a 50-m-long belt-transect orientated north-south or east-west, whichever maximised overall site variability. Transect width was adjusted between 2.5 and 7.5 m to include at least 15 individuals in each size class. Overall, we identified approximately 50-100 individual plants in each transect and measured their basal diameters.[49] Basal diameters were measured to accommodate the prevalence of multi-stemmed and coppicing plants.

At each site we listed all the bird species encountered by one observer in a 45-min sampling period starting at either 05:00 or 07:00, with the starting time alternating on consecutive days between disturbed and undisturbed woodlands. Sampling time did not significantly affect the number of species of birds observed ( $\chi^2 = 3.47$ ; d.f. = 1;  $P > 0.05$ ). Birds, identified visually and by their songs, were recorded while walking within a radius of approximately 250 m of the previous evening's bat and insect sampling station.

We sampled flying bats using macro-mist nets[50] erected at dusk and set open from 18:30 to 23:30 when they were monitored continuously. For each captured bat, we recorded the time of capture, species and sex.

Insect samples were collected from two light traps at each sampling site: a 12-V white fluorescent neon tube suspended above a dish of soapy water and turned on from 19:00 to 20:30; and a 12-V ultraviolet neon tube suspended under a white sheet spread over a tripod of 1-m-long poles. The UV light was turned on for the first 15 min of each 30-min period (starting at 18:30). During the last five minutes insects were collected off the sheet into a wide-necked killing jar. Insects from the UV samples were sorted by size class and order. Formicids and mantids from both sampling systems were separated by species. We lacked the resources to separate other insect taxa to species level.

## *Results*

Woodland structure differed markedly between impacted and intact woodlands (Table 1). Tree cover and the density of large trees (basal diameter >15 cm) was markedly lower in affected

sites. Differences in mean overall tree density were not as high because some sites, particularly in the Matusadona, had high densities of small, regenerating, *Brachystegia boehmii* trees. Associated with reduced tree cover in impacted woodlands was a nearly fourfold increase in shrub cover and a twofold increase in small-shrub cover (Table 1). Neither the mean number nor the overall number of woody plant species found in intact and impacted woodland differed significantly but the number of woody species in the >3-m height class, namely trees, was markedly lower in the affected woodland (Table 1).

Significantly fewer bird species were found in impacted woodlands and, as expected, the greatest contrast in species richness was found in woodland species where species richness was significantly lower in impacted woodland (Table 1). We recorded a total of 121 species of birds with a total of 99 species in intact woodland and 73 in impacted woodland. The loss of woodland species in affected areas was not compensated for by an increase in non-woodland species (Table 1). Among the arboreal passer-inc bird species present in the intact woodland but missing from the impacted woodlands were four miombo woodland endemics (Miombo Rock Thrush, *Monticola angolensis*; Spotted Creeper, *Salpornis spilonotus*; Mashona Hyliota, *Hyliota australis*; Miombo double collared Sunbird, *Nectarinia manoensis*; Cabanis's Bunting, *Ernberiza cabanisi*) and several species which, although occurring in other habitats, might be described as miombo woodland specialists in the sense that their distribution is largely confined to this habitat type.[51,52]

We captured a total of 344 bats and, because all of the adult females were lactating or post-lactating, the populations appear to be resident and breeding. Bat species richness in upland areas was low with 11 species being caught over 15 nights of trapping, while the greatest number of species caught at any one site was eight. The number of bat species caught in intact and impacted woodlands did not differ significantly (Table 1). Fruit bats (*Epomophorus* species) were caught only in intact woodland while seven species of vespertilionids were caught in both intact and impacted sites.

We consistently caught more species of ants by night in the intact woodland in the Maria area and the same pattern prevailed for mantises; these patterns were less marked in the Kanyati and Matusadona sites (Table 1). We found and heard cicadas only in the intact woodlands and seven specimens from two genera (*Lacetes* and *Oxypleura*) were captured at the light traps.

### *Discussion*

Our results show that where elephants have existed at high densities (>0.5 per km<sup>2</sup>) for more than a decade (Cumming and Taylor, unpublished census data) the structure of woodlands is changed markedly and the diversity of canopy trees and of associated bird and insect faunas may be reduced. Most woody species were still present in the shrub layer and the potential for regeneration of the woodland, albeit via a thicket phase, was still present. Bird species diversity is known to be correlated with foliage height diversity in woodlands [53,54] so that the greatly reduced vertical and horizontal heterogeneity in the elephant-affected woodlands would account for the lower species richness of birds in these woodlands.

While frequent hot late dry-season bums can open up miombo woodlands[55] in the absence of elephants, it is the interaction of elephants and fire that is of prime importance in the context of

our study areas. Since the onset of tsetse control operations in the mid-1960s, intact miombo woodlands in the Mana and Kanyati areas were subjected to early burns and elephants were removed. Within the national parks (impacted areas) confined elephant populations increased and unsustainable rates of tree felling were evident by the late 1960s and early 1970s (personal observation and blanket aerial photography). Attempts in the early 1970s to protect woodlands from fire only served to increase the risk and frequency of late hot burns[56] because wild fires could seldom be controlled. An early-burning policy was introduced to pre-empt the risk of hot, late season burns but was not consistently implemented in the Mana area. In the Matusadona highlands early burning combined with a reduction in elephant numbers facilitated the regeneration of *Brachystegia boehmii* in some areas (R.D. Taylor, unpublished data). A similar reduction in elephant numbers but without an effective early burning regime in the Mana Pools National Park inhibited regeneration. The persistence of large tracts of intact woodlands south of the Mana fence, where fire was not controlled but from which elephants were excluded, is consistent with the conclusion that high elephant densities were primarily responsible for transforming closed canopy miombo woodlands to open bushlands or grasslands.

Local loss of biodiversity under high elephant densities has important implications for the role of such game reserves in conserving biodiversity locally, nationally and regionally. Elephant numbers in southern Africa presently exceed 170 000 with more than eight populations of over 5 000 animals living in reserves of more than 5 000 km<sup>2</sup>. [57] With a probability of >99% that an elephant population of 2 500 held at 50% of carrying capacity (i.e. at a density of 0.5 elephant per km<sup>2</sup> in a 5 000 km<sup>2</sup> reserve) will be genetically and demographically secure for 1000 years, [58] there is no compelling conservation argument to increase elephant numbers and densities in reserves within the region. Lower elephant densities (i.e. at <0.5 elephant per km<sup>2</sup>) may, on the other hand, conserve higher levels of plant and animal biodiversity, safeguard habitats and reduce the risk of local population collapses of elephants or other species, or both. Examples include the initial collapse of elephant and black rhino populations in Tsavo National Park [59-61] and the extinction of bushbuck and lesser kudu from Amboseli National Park as a result of elephant-induced habitat change. [62]

The intermediate disturbance hypothesis [63,64] provides an appropriate theoretical framework within which to explore the effects of elephant disturbance on biodiversity and particularly their effects on spatial and temporal heterogeneity of habitats but this work has scarcely started. [65-67] The links between overall elephant population density (as used here), temporal and spatial shifts in local population densities and heterogeneity in woodland habitats need further examination. It is already clear that localised woodland damage and loss of certain tree species will occur even at elephant densities of <0.2 elephant per km<sup>2</sup> in a wide range of woodlands. [5,35] At persistent densities >0.5-1 elephant per km<sup>2</sup> large areas of several types of woodland have been reduced to scrub with an associated loss in spatial heterogeneity. [19-29] In Zimbabwe, localised culling in such circumstances reduced elephant densities and resulted in regeneration and a return of some spatial heterogeneity (Cumming and Taylor, pers. obs.). At the other end of the spectrum 'old growth' woodlands may well support a characteristic fauna and flora which needs to be conserved but this question has not surfaced in the elephant/ woodland debate.

An alternative thesis holds that eruptions of elephant populations, woodland clearance and elephant die-offs are 'natural' and necessary for protected areas to maintain habitat resilience, stability and biodiversity in the long term (e.g. ref. 42). This view assumes, inter alia, that such cycles occurred naturally in the past, were sufficiently localised to avoid regional mass extinctions, and that refugia existed (and will exist in the future) from which species can re-invade habitats destroyed by elephants. However, none of these assumptions holds for our study area. The Zambezi Valley, Great Zimbabwe in the Save River catchment, and Mapungubwe south of the Limpopo were an integral part of the ancient trade in gold and ivory which was centred on the East African coast.[68-70] Ivory was harvested and traded in southeast Africa at surprisingly consistent levels over at least four centuries before the over-exploitation of African elephants occurred in the late 19th century.[71] The writings of early hunters and explorers provide no evidence that densities of 1-3 elephant per km<sup>2</sup> were prevalent in East and southern Africa and we are not aware of any evidence to support the view that elephants existed at high densities over extensive areas in prehistoric times. It is thus very likely that intact miombo woodlands persisted over large areas for at least the last 500-1000 years, and, given current levels of endemicity, did so for very much longer.

Presently about 88% (6 800 km<sup>2</sup>) of the miombo woodland in protected areas of the Zambezi Valley in Zimbabwe has been heavily impacted by elephant with spatial heterogeneity largely obliterated. Only the Duma Safari Area (945 km<sup>2</sup>) remains relatively intact (Cumming and Taylor, pers. obs.). With woodlands outside protected areas being converted to agriculture at about 4% per annum,[6] the opportunity for species re-invasions to occur from intact refugia within Zimbabwe will soon be lost. A woodland regeneration time of several decades further reduces the chances of re-invasion.

The practical problems of reducing elephant densities in large populations such as now occur in many game reserves, and other woodland habitats, in southern Africa are daunting. To reduce densities to <0.5 elephant per km<sup>2</sup> within Zimbabwe's protected areas, it would now be necessary to remove more than 35 000 elephants. One costly option is to move herds into new reserves or open new areas into which elephants can disperse. Both have been used in southern Africa but the scope for further translocations within the region is limited and the gains short-lived because confined, protected elephant populations soon reach levels at which they start to impact on woodlands. Culling, an option successfully used in the past, is now inhibiting because the CITES ban on trade in elephant products precludes the sale of ivory and hides and thus the opportunity to recover the costs of culling elephants. Animal welfare groups have campaigned against the use of culling, arguing instead for the use of contraceptives. For elephants this alternative has yet to be developed and tested.

Our results and this overview of the elephant problem highlight the potential risks to biodiversity conservation in African savanna game reserves of maintaining high elephant densities. The generality of our specific findings for miombo needs to be critically examined for other savanna woodland types in the region and further research on this question is urgently needed if the current impasse between protecting elephants and conserving woodland biodiversity is to be resolved rationally. Equally important is the observation that while miombo woodlands and their associated biodiversity are at risk within game reserves, woodlands outside of protected areas are also under siege from agricultural development despite the obvious long-term, adverse

consequences for people and biodiversity of subsidising agricultural development on marginal soils.

We thank the Director, Department of National Parks and Wildlife Management, and the Research Council of Zimbabwe for permission to carry out this work in Zimbabwe. The research was supported by grants in aid to I.L.R. from the Anglo American and De Beers Chairman's Fund and the Foundation for Research Development (South Africa), to M.B.F. from the National Science and Engineering Research Council of Canada and by WWF -- World Wide Fund for Nature. Martin Villet, Rhodes University, kindly assisted in the identification of ants and cicadas. We are also grateful to Colin Craig, Peter Frost, Norman Owen-Smith and an anonymous referee for valuable and constructive criticism of earlier drafts of this paper, to Charles Mackie, who piloted the aircraft for the aerial photograph of the study area, and to Judy Murray of Action Magazine for producing a composite digital image for Fig. 1.

### *References*

1. World Resources 1990-91 (1990). Oxford University Press, New York,
2. Calef G.W. (1988). Maximum rate of increase in the African elephant. *Afr. J. Ecol.* 26, 323-327.
3. Cumming D.H.M. (1980). The management of elephant and other large mammals in Zimbabwe. In *Problems in Management of Locally Abundant Wild Mammals*, eds. R.A. Jewell, S. Holt, and D. Hart, pp. 91-118. Academic Press, New York.
4. Hall-Martin A.J. (1992). Distribution and status of African elephant *Loxodonta africana* in South Africa, 1652-1992. *Koedoe* 35, 65-88.
5. Spinage C.A. (1990). Botswana's problem elephants. *Pachyderm* 13, 14-19.
6. Cumming D.H.M. (1994). WWF/EC study of land-use trends, natural resource use and agro-ecosystem sustainability in the Zambezi Valley of Zimbabwe. In *Proceedings of a Workshop to Co-ordinate Studies of Land-use change in the Zambezi Valley of Zimbabwe*, eds. D.H.M. Cumming and I.M. Daniels, pp. 30-44. Regional Tsetse and Trypanosomiasis Control Programme and WWF, Harare.
7. Hanks J. (1979). *A Struggle for Survival: The Elephant Problem*. Struik, Cape Town.
8. Owen-Smith N. (1988) *Megaherbivores: The Influence of Very Large Body Size on Ecology*. Cambridge University Press, Cambridge.
9. Huston M. (1993). Biological diversity, soils and economics. *Science* 262,1676-1680.
10. Huston M. (1994). Saving the planet. *Bull. Ecol. Soc. Am.* 76, 97-99.
11. Huntley B.J. (1982). Southern African savannas. In *Ecology of Tropical Savannas*, eds B.J. Huntley and B.H. Walker, pp. 101-119. Springer-Verlag, Berlin.
12. Hough J. (1986). Management alternatives for increasing dry season base flow in the miombo woodlands of southern Africa. *Ambio* 15, 341-346.
13. Pereira H.C. (1973), *Land Use and Water Resources*. Cambridge University Press, Cambridge.
14. Bryden H.A. (1903). The decline and fall of the elephant in southern Africa. *Fortnightly Rev.* 79, 100-108.
15. Skinner J.D. and Smithers R.H.N. (1990). *The Mammals of the Southern African Sub-region*. University of Pretoria, Pretoria.

16. Cumming D.H.M. (1991). Wildlife products and the market place: A view from southern Africa. In *Wildlife Production: Conservation and Sustainable Development*, eds. L.A. Renecker and R.J. Hudson, pp. 11-25, AFES misc. pub., University of Alaska Fairbanks, Fairbanks, Alaska.
17. Murphree M.W. and Cumming D.H.M. (1993). Savanna land use: Policy and practice in Zimbabwe. In *The World's Savannas: Economic Driving Forces, Ecological Constraints and Policy Options for Sustainable Land Use*, eds. M.D. Young and O.T. Solbrig, pp. 139-178, UNESCO and Parthenon, Paris.
18. Pimm S.L. (1982). *Food Webs*. Chapman and Hall, London.
19. Buechner H.K. and Dawkins H.C. (1961). Vegetation change induced by elephants and fire in Murchison Falls National Park. *Ecology* 42, 752-766.
20. Glover R (1963). The elephant problem at Tsavo. *E. Afr. Wildl. J.* 1, 30-39.
21. Van Wyk P. and Fairall N. (1969). The influence of the African elephant on the vegetation of the Kruger National Park. *Koedoe* 9, 57-95.
22. Laws R.M. (1970). Elephants as agents of habitat and landscape change in East Africa. *Oikos* 21, 1-15,
23. Anderson G.D. and Walker B.H. (1974). Vegetation composition and elephant damage in the Sengwa Wildlife Research Area, Rhodesia. *J. sth Afr. Wildl. Mgmt Ass.* 4, 1-14
24. Thomson P.J. (1975). The role of elephants, fire and other agents in the decline of a *Brachystegia boehmii* woodland. *J. sth Afr. Wildl. Mgmt Ass.* 5, 11-18.
25. Guy P.R. (1981). Changes in the biomass and productivity of woodlands in the Sengwa Wildlife Research Area, Zimbabwe. *J. appl. Ecol.* 18,507-519
26. Cumming D.H.M. (1982). The influence of large herbivores on savanna structure in Africa. in *Ecology of Tropical Savannas*, eds B.J. Huntley and B.H. Walker, pp. 217-245. Springer-Verlag, Berlin.
27. Barnes R.F.W. (1983). Effects of elephant browsing on woodlands in a Tanzanian National Park -- measurements, models and management. *J. appl. Ecol.* 20, 521-540.
28. McShane T.O. (1987). Elephant-fire relationships in Combretum/Terminalia woodland in south-west Niger. *Afr. J. Ecol.* 25, 79-94.
29. Lewis D.M. (1991). Observations on tree growth, woodland structure and elephant damage on *Colophospermum mopane* in Luangwa Valley, Zambia. *Afr. J. Ecol.* 29, 207-221.
30. Stuart-Hill G.C. (1992). Effects of elephants and goats on the Kaffrarian succulent thicket of the Eastern Cape, South Africa. *J. appl. Ecol.* 29, 699-710.
31. Ben-Shahar R. (1993). Patterns of elephant damage to vegetation in northern Botswana. *Biol. Conserv.* 65,249-256.
32. Swanepoel C.M. (1993). Baobab damage in Mana Pools National Park, Zimbabwe. *Afr. J. Ecol.* 31,220-225.
33. Cumming D.H.M. and Jackson R (1984). The status and conservation of Africa's elephants and rhinos. *Proceedings of the joint meeting of IUCN/SSC African Elephant and African Rhino Specialist Groups, Hwange Safari Lodge, August 1980.* IUCN, Gland.
33. Cumming D.H.M., du Toit R.F. and Stuart S.N. (eds) (1990). *African Elephants and Rhinos. Status Survey and Conservation Action Plan.* IUCN/SSC African Elephant and Rhino Specialist Group. IUCN, Gland.
34. Martin R.B., Craig C.G. and Booth V.R. (eds) (1989). *Elephant management in Zimbabwe.* Dept. National Parks and Wild Life Management, Harare.



35. Hawthorne R (1995). Time June 5, 84.
36. Anon. (1996). Cash if you don't cull. Newsweek February 26, 4.
37. Anon. (1996). An elephantine problem. The Economist March 30, 85.
38. Anon. (1996). Killing to be kind. The Economist April 20, 86-87.
39. Anon. (1996). Earthday reconsidered. The Wall Street Journal Europe April 24.
40. Trevor S. (1992). Elephant as architect. BBC Wildlife 10(9), 50-54.
41. Caughley G. (1976). The elephant problem -- an alternative hypothesis. E. Afr. Wildl. J. 14, 265-283.
42. Caughley G. (1980). Overpopulation. In Problems in Management of Locally Abundant Wild Mammals, eds P.A. Jewell, S. Holt and D. Hart, pp. 7-19. Academic Press, New York.
43. Walker B.H., Emslie R.N., Owen-Smith R.N. and Scholes R.J. (1987). To cull or not to cull: lessons from a southern African drought. J. appl. Ecol. 24,381-401.
44. Walker B.H. (1989). Diversity and stability in ecosystem conservation. In Conservation for the Twenty-first Century, eds D. Western and M Pearl, pp. 121-130. Oxford University Press, Oxford.
45. Moolman H.J. and Cowling R.M. (1994). The impact of elephant and goats on the endemic flora of South African succulent thicket. Biol. Conserv. 68, 53-61.
46. Herremans M. (1995). Effects of woodland modification by African elephant *Loxodonta africans* on bird diversity in northern Botswana. Ecography 18, 440-454.
47. White F. (1983). The Vegetation of Africa. UNESCO, Paris.
48. Mueller-Dombois D. and Ellenberg H. (1974). Aims and Methods of Vegetation Ecology. John Wiley, New York.
49. Rautenbach I.L. (1985). A new technique for the efficient use of macro-mist nets. Koedoe 28, 81-86.
50. Stuart-Irwin M. (1964). Report of the Mafungabusi Expedition, Rhodesia Schools Exploration Society. Bulawayo, Rhodesia.
51. MacLean G.L. (1984). Roberts' Birds of Southern Africa, 5th edn. John Voelker Bird Book Fund, Cape Town.
52. MacArthur R.H. (1964). Environmental factors affecting bird species diversity. Am. Nat. 98, 387-397
53. Rosenzweig M.L. (1995). Species Diversity in Time and Space. Cambridge University Press, Cambridge.
54. Trapnell C.G. (1959). Ecological results of woodland burning experiments in Northern Rhodesia. J. Ecol. 47,129-168.
55. Starfield A.M., Cumming D.H.M., Taylor R.D. and Quadling M.S. (1993). A frame-based paradigm for dynamic ecosystem models. AI Applications 7, 1-13.
56. Said M.Y., Chunge R.N., Craig C.G., Thouless C.R., Barnes R.F.W. and Dublin H.T. (1995). African Elephant Database: 1995. Occasional Paper of the IUCN Species Survival Commission No. 11. IUCN, Gland.
57. Armbruster P. and Lande R. (1993). A population viability analysis for African elephant (*Loxodonta africans*): How big should reserves be? Conserv. Biol. 7, 602-610.
58. Botkin D.B. (1990). Discordant Harmonies. Oxford University Press, New York.
59. Goddard J. (1970). Food preferences of black rhinoceros in the Tsavo National Park. E. Afr. Wildl. J. 8, 145-161.
60. Martin E.B. and Martin C.B. (1982). Run Rhino Run. Chato and Windus, London.

61. Western D. and Gichohi H. (1993). Segregation effects and the impoverishment of savanna parks: the case for ecosystem viability analysis. *Afr. J. Ecol.* 31, 269-281.
62. Connell J.H. (1978). Diversity in tropical forests and coral reefs. *Science* 199, 1302-1310.
63. Petraitis P.S., Latham R.E. and Niesenbaum R.A. (1989). The maintenance of species diversity by disturbance. *Quart. Rev. Biol.* 64, 393-418.
64. Owen-Smith N. (1987). Pleistocene extinctions: the pivotal role of megaherbivores. *Paleobiology* 13, 351-362.
65. Owen-Smith N. (1989). Megafaunal extinctions: The conservation message from 11 000 years B.R *Conserv. Biol.* 3, 405-412.
66. Western D. (1989). The ecological role of elephants in Africa. *Pachyderm* 12, 42.
67. Summers R. (1963). *Zimbabwe: A Rhodesian mystery*. Nelson, Johannesburg.
68. Alpers E.A. (1975). *Ivory and Slaves in East Central Africa*. Heinemann, London.
69. Campbell A. (1990). History of elephants in Botswana. In *The Future of Botswana's Elephants*, ed. P. Hancock, pp. 5-15. Proceedings of Kala-hari Conservation Society Symposium, Gaborone.
70. Spinage C.A. (1973). A review of ivory exploitation trends in Africa. *E. Afr. Wildl. J.* 11, 281-289.

Table 1. Comparison of woody plant density and cover and species richness of woody plants, birds, bats, ants and mantises in intact and elephant-impacted woodland sites. (A) Tree and shrub density (density of woody plants in three height classes and large trees where basal diameter (BD) > 15 cm); (B) woody plant cover reflected by basal area; (C) species richness of woody vegetation, birds, bats, ants and mantises. (Means +/- standard error of mean, n = 15 intact and 15 impacted sites (see text) except for birds where n = 14; probabilities for single-sided, 2-sample t-test; ns indicates not significant at 5% level.)

Legend for Chart:

- A - Group or taxon
- B - Intact woodland (mean +/- s.e.m.)
- C - Impacted woodland (mean +/- s.e.m.)
- D - Probability
- E - Direction

A	B	C
	D	E

A) Density woody plants (plants ha<sup>-1</sup>)

Trees (>3 m)	408 +/- 39.9	245 +/- 70.9
--------------	--------------	--------------

<0.05[\*]      arrow down

Trees with BD > 15 cm    193.8 +/- 19.9    56 +/- 22.0  
<0.001[\*\*\*]      arrow down

Shrubs (1-3 m)      257.8 +/- 44.9    1412.0 +/- 209.5  
<0.001[\*\*\*]      arrow up

Small shrubs (<1 m)    3482.7 +/- 391.9    6029.3 +/-724.4  
<0.01[\*\*]      arrow up

B) Woody plant cover  
(basal area m<sup>2</sup>  
ha<sup>-1</sup>)

Trees (height >3 m)    9.5 +/- 1.2      3.5 +/- 1.2  
<0.001[\*\*\*]      down arrow

Shrubs (height 1-3 m)    2.5 +/- 0.7      12.4 +/- 1.8  
<0.001[\*\*\*]      up arrow

Small shrubs (height <1 m)    7.1 +/- 0.9      15.4 +/- 2.0  
<0.001[\*\*\*]      up arrow

C) Species richness  
(no. species per site)

Trees                  25.5 +/- 1.0      11.2 +/- 0.8  
<0.001[\*\*\*]      arrow up

Shrubs                  35.6 +/- 1.8      35.5 +/- 2.8  
>0.5 ns              left and right arrow

Total woody plants    41.1 +/- 2.3      29.5 +/- 3.8  
>0.2 ns              down arrow

Woodland birds        16.3 +/- 2.0      10.2 +/- 0.5  
<0.001[\*\*\*]      down arrow

Other birds            5.7 +/- 0.5      8.7 +/- 1.1  
>0.2 ns              up arrow

Total birds            22.0 +/- 2.3      18.8 +/- 1.3  
<0.01[\*\*]          arrow down

Bats                    3.5 +/- 0.6      3.3 +/- 0.8

	>0.05 ns	arrow down
Ants	9.2 +/- 0.7 <0.05[*]	7.3 +/- 0.5 arrow down
Mantises	3.0 +/- 0.3 >0.09 ns	1.8 +/- 0.5 arrow down
Total animals		
(birds, bats, ants & mantises)	37.7 +/- 3.2 <0.01[**]	31.3 +/- 2.0 arrow down

PHOTO (COLOR): Fig. 1. Elephant impact on woodland. (a) Oblique aerial view of the southern boundary of Mana Pools National Park/Hurungwe Safari Area, showing elephant-affected woodland to the right (north) of the tsetse fence and intact woodland in the adjacent communal land to the south. Note clearance for agriculture in the middle distance (May 1996). (b) Ground view of intact miombo woodland south of the fence in Communal Land (November 1994). (c) Ground view of elephant and fire-affected woodland to the north of the fence inside game reserve (November 1994). (Photographs by David Cumming)

MAP: Fig. 2. Study area on the Zambezi Valley escarpment in Zimbabwe showing protected areas (NP, national park; SA, safari area) and sites sampled (1) on the southern boundary of Mana Pools National Park and Hurungwe Safari Area, the Kanyati Communal Land and the Matusadona National Park. Intact miombo woodlands within the protected areas occur in the southern part of Charara (south of former tsetse fence) and in the Doma Safari Area.