

Elevated atmospheric carbon dioxide and ozone alter forest insect abundance and community composition

MICHAEL L. HILLSTROM and RICHARD L. LINDROTH Department of Entomology, University of Wisconsin, Madison, WI, USA

Abstract. 1. Human-induced climate changes threaten the health of forest ecosystems. In particular, carbon dioxide (CO₂) and tropospheric ozone (O₃) will likely have significant but opposing impacts on forests and their associated insect communities. Compared with other animal groups, insect communities are expected to be especially sensitive to changes in global climate.

2. This study examined the effects of elevated CO₂ and O₃ (eCO₂ and eO₃) individually and in combination on the abundance, diversity and composition of forest insect communities. Insects were sampled using yellow pan traps in an aggrading aspen-birch forest at the Aspen Free Air CO₂ Enrichment (FACE) site in northern Wisconsin, USA. We trapped for 24 h every 10–15 days throughout the summers (June to September) of 2000–2003.

3. We examined 47 415 insects from 4 orders and 83 families. Elevated CO₂ reduced abundance of phloem-feeding herbivores and increased abundance of chewing herbivores, although results were not statistically significant. Enriched CO₂ increased numbers of some parasitoids. The effects of eO₃ on insect abundance were generally opposite those of eCO₂. No significant differences in arthropod family richness were found among treatments. However, eCO₂, eO₃, or both significantly affected insect community composition in all years.

4. Carbon dioxide and tropospheric ozone have the potential to alter significantly forest insect communities. Feeding guild may strongly influence insect response to environmental change and may provide the best opportunity to generalise for conservation efforts. Because insect communities influence forest health and ecosystem services, continued research on their response to global change is critically important to forest management and conservation.

Key words. Biodiversity, carbon dioxide, climate change, FACE, forest, insect community, NMDS, pan trap, ozone.

Introduction

As the evidence for human-induced climate change mounts, conservation of natural communities has become a scientific, social and political priority. Ecosystems face not only increases in carbon dioxide levels, temperature, and altered precipitation, but also increased habitat fragmentation and loss, making adaptation increasingly difficult (Lovejoy & Hannah, 2005). Responses of organisms to global change will be species-specific and occur at different rates, potentially disaggregating communities and causing alteration or loss of ecosystem services critical to human existence (Lovejoy & Hannah, 2005).

Correspondence: M. L. Hillstrom, Department of Entomology, University of Wisconsin, 237 Russell Laboratories, 1630 Linden Drive, Madison, WI 53706-1598, USA. E-mail: hillstro@entomology.wisc.edu

Biodiversity will play a key role in how ecosystems respond to global change (Hansen & Dale, 2001) as loss of species can lead to reduced ecosystem function (Cardinale *et al.*, 2006). Much of the planet's biodiversity resides in two groups of organisms: plants and insects. Forests are of particular interest as they cover one third of Earth's land mass (Ozanne *et al.*, 2003), host millions of species and are expected to be especially sensitive to changes in the global environment (Fowler *et al.*, 1999). Forest insects play key ecological roles, influencing forest productivity, species composition, energy flow, and nutrient cycling (Mattson & Addy, 1975; Schowalter *et al.*, 1986; Veblen *et al.*, 1991). Insects are also important herbivores, pollinators, predators and parasitoids in forests (Altermatt, 2003). Thus, changes in insect abundance, diversity or community composition have the potential to alter forest ecosystems and the services they provide.

Two gases in particular, carbon dioxide (CO₂) and tropospheric ozone (O₃), will likely have significant but opposing impacts on forest ecosystems (Dickson *et al.*, 1998; Saxe *et al.*, 1998). Since the beginning of the industrial revolution, CO₂ levels have increased from 280 to 380 ppm (IPCC, 2007). Over this same time period, ambient tropospheric O₃ levels have risen to over 40 ppb (from ~10 ppb in 1850) in many regions of the world and are projected to reach 70 ppb (mean monthly 24-h ozone concentrations) during the summer in the USA, Europe and East Asia by 2100 (Sitch *et al.*, 2007). Elevated CO₂ and eO₃ have been shown to alter tree growth and chemical characteristics (reviewed for Aspen Free Air CO₂ Enrichment (FACE) in Lindroth *et al.*, 2001; Karnosky *et al.*, 2003). Numerous studies have demonstrated changes in insect performance at eCO₂ and eO₃ (Bezemer & Jones, 1998; Coviella & Trumble, 1999), but more recent work suggests that population and community responses may differ from individual responses (Percy *et al.*, 2002; Awmack *et al.*, 2004). Likewise, the artificial conditions of many of the early studies confound population and community level predictions because natural communities experience many indirect biotic and abiotic interactions and feedbacks that cannot be accounted for in studies with simplified environments (Bailey & Whitham, 2002). Recent work at FACE facilities has provided some evidence for the effects of CO₂ and O₃ on natural insect communities, but no study to date has examined the effects of CO₂ and O₃ simultaneously on above-ground forest insect communities. Multifactor, ecologically realistic studies of environmental change are necessary for conservation strategies to be developed.

The effects of eCO₂ or eO₃ on insect communities are likely to be indirect. For example, effects on herbivorous insect populations will depend on the magnitude of change in host plant quality, natural enemy impact and changes in the physical environment (tree architecture, etc.). Elevated CO₂ and eO₃ may change natural enemy populations via shifts in the diversity, abundance and quality of prey or changes in host-finding mechanisms. Natural enemies may be particularly sensitive to eCO₂ or eO₃ through changes in searching behaviour (Gate *et al.*, 1995; Vuorinen *et al.*, 2004; Pinto *et al.*, 2007) or the behaviour of their prey (Mondor *et al.*, 2004).

This study was designed to examine the effects of eCO₂ and eO₃, individually and in combination, on the abundance, biodiversity and composition of forest insect communities. Insects were sampled using yellow pan traps in planted aspen-birch stands at the Aspen FACE site in northern Wisconsin. Yellow pan traps are known to preferentially attract predators and parasitoids (Duelli *et al.*, 1999), two of the most under-represented groups in climate change research. We recognise that pan traps sample only a subset of the insect community but chose them because they allow for easy collection of large quantities of insects across a broad range of families.

Methods

Aspen FACE

The Aspen FACE site is a 32-ha experimental station located near Rhinelander, Wisconsin. The site consists of 12 rings each

30 m in diameter planted with trees spaced 1 m apart. Sampling occurred in the interplanted trembling aspen (*Populus tremuloides* Michx.) and paper birch (*Betula papyrifera* Marsh.) section of each ring. The aspen-birch community is one of the most widespread communities in northern forests and contains the primary pulpwood species of the Great Lakes region (Piva, 2006). Trees were planted as seedlings in 1997 and fumigation began in 1998 (Dickson *et al.*, 2000; Karnosky *et al.*, 2003).

Aspen FACE is set up as a randomised complete block design with three blocked sets of four treatment rings: control (ambient CO₂, ambient O₃), elevated CO₂ (+CO₂; 560 μM M⁻¹ CO₂), elevated O₃ (+O₃; 1.5 × ambient) and elevated CO₂ and O₃ (+CO₂+O₃). Elevated CO₂ and O₃ levels represent concentrations predicted for 2060. Ambient O₃ concentration is monitored daily and the elevated rings are fumigated based on a diurnal concentration profile approximately 1.5 times that of ambient levels (Dickson *et al.*, 2000).

Insect sampling

The intent of this study was to sample a large number of insects across a broad range of families. We choose pan traps to accomplish this task although we recognise the limitations of this methodology. Pan traps do not attract all families of insects equally but are particularly good at attracting predators and parasitoids (Duelli *et al.*, 1999; Triplehorn & Johnson, 2005), the intended focus of this study. Pan traps are also passive traps so location of the trap will affect the insects captured. To account for this sampling effect, we always placed the traps in the same location in each ring and in similar surrounding vegetation. It is also possible that some percentage of the insects collected were only flying through the rings and were unaffected by the treatment. We accounted for this by analyzing only the 15 most abundant families (each accounting for greater than one percent of total abundance) in both the community and individual analyses. Each of the 15 families had over 500 individuals collected over the 4 years so they are unlikely to be merely passing through. Although pan traps have limitations, we accounted for these concerns in both sampling technique and data analysis.

We placed one yellow pan trap (22 cm in diameter) on the ground in the aspen-birch section of each FACE ring for 24-h collections during the summers of 2000–2003. Sampling occurred approximately every 10 days from June to September. Traps contained 0.5% scentless detergent solution 1 cm deep and were placed at fixed locations each time to preclude spatial differences in catches among sample dates. Insects were removed from the traps by filtering the detergent solution through a fine strainer. Samples were stored in 70% ethyl alcohol. Insects were identified under a dissecting microscope to order and family. Identifications and feeding guild assignment were made using Triplehorn & Johnson (2005).

Insect community analysis

Arthropod abundance was analysed two ways. First, total abundance of individuals across all insect families collected

Table 1. Average (\pm SE) total number of insects collected per pan trap per year, by treatment, for the 15 most abundant families of arthropods. Statistics represent the results of two-way crossed repeated measures ANOVA tests (df 1,8). Significant *P*-values (< 0.05) are indicated in bold.

Feeding guild/family	Control	+CO ₂	+O ₃	+CO ₂ +O ₃	CO ₂		O ₃		CO ₂ ×O ₃	
					<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
Herbivores – phloem feeders										
Aphididae	51 ± 15	58 ± 15	83 ± 15	56 ± 15	1.1	0.334	0.6	0.449	6.6	0.033
Cicadellidae	57 ± 10	32 ± 10	78 ± 10	56 ± 10	3.8	0.087	3.8	0.087	0.6	0.463
Miridae	10 ± 3	9 ± 3	24 ± 3	13 ± 3	4.1	0.077	7.8	0.024	2.1	0.185
Herbivores – chewers										
Curculionidae	14 ± 2	16 ± 2	9 ± 2	13 ± 2	3.8	0.087	4.7	0.062	0.4	0.553
Anthomyiidae	101 ± 23	139 ± 23	76 ± 23	141 ± 23	5.1	0.054	0.3	0.612	0.3	0.576
Parasitoids										
Braconidae	51 ± 5	49 ± 5	32 ± 5	35 ± 5	0.2	0.649	10.6	0.011	0	0.961
Chalcidoidea	97 ± 6	93 ± 6	76 ± 6	65 ± 6	1.6	0.248	15	0.005	0.1	0.765
Figitidae	23 ± 6	52 ± 6	12 ± 6	19 ± 6	9.3	0.016	13.5	0.006	3.4	0.102
Ichneumonidae	99 ± 15	122 ± 15	55 ± 15	75 ± 15	6	0.040	14	0.006	0	0.921
Predators										
Dolichopodidae	41 ± 5	33 ± 5	35 ± 5	28 ± 5	2	0.191	1.1	0.320	0	0.945
Saprophages										
Drosophilidae	16 ± 4	20 ± 4	9 ± 4	12 ± 4	0.9	0.376	4	0.079	0	0.874
Phoridae	191 ± 27	143 ± 27	90 ± 27	141 ± 27	0	0.955	3.6	0.100	3.3	0.108
Sciaridae	160 ± 30	162 ± 30	141 ± 30	94 ± 30	0.6	0.455	2.4	0.159	1.1	0.332
Herbivores/Predators										
Formicidae	31 ± 6	12 ± 6	30 ± 6	21 ± 6	7.3	0.027	1.3	0.292	0.9	0.362
Predators/Saprophages										
Sarcophagidae	43 ± 8	42 ± 8	48 ± 8	30 ± 8	5.1	0.054	0.2	0.708	2.1	0.183
Total	1094 ± 90	1069 ± 90	903 ± 90	887 ± 90	0.1	0.807	5.3	0.050	0	0.956

(= total arthropod abundance) was evaluated. Second, the 15 most abundant families were examined individually in more detail (Table 1). All data were pooled across an equal number of sample dates for each of the 4 years of this study. The data were normalised using a square root or log transformation until the Shapiro–Wilks *W* test statistic exceeded 0.85. Treatment effects were analysed using two-way crossed repeated measures ANOVA with block, CO₂ (= eCO₂) and O₃ (= eO₃) as the main effects, year as the repeated measures term and year by main effect interactions as the subplot terms. Year is considered a repeated measure because traps were located in the same spot for each of the 4 years. We also included the interaction of the main effects (CO₂ × O₃). CO₂ main effects (eCO₂) represent differences between plots with enriched CO₂ (+CO₂ and +CO₂+O₃) and plots with ambient CO₂ (control and +O₃). Similarly, O₃ main effects (eO₃) represent differences between plots with enriched O₃ (+O₃ and +CO₂+O₃) and plots with ambient O₃ (control and +CO₂). *P*-values are considered significant at $\alpha = 0.05$ for individual family tests. We did not employ Bonferroni corrections because these individual family tests are almost certainly not independent of one another due to direct and indirect biotic interactions among the insects and host plants. If the tests are not independent, then Bonferroni correction is excessively conservative leading to high Type II error rates (Gotelli & Ellison, 2004; Moran, 2003). Nevertheless, we interpret the results with caution.

We analysed arthropod family richness using sample-based rarefaction curves produced with the program ESTIMATES 7.5.0 (Colwell, 2006). Rarefaction curves account for differential

sampling effort whereas other commonly used alpha diversity measures (e.g. Shannon–Wiener index, Simpson's index) do not, producing potentially misleading results (Colwell *et al.*, 2004; Buddle *et al.*, 2005). Curves were re-scaled to individuals to account for differing densities of individuals in samples (Colwell *et al.*, 2004). For each year, samples were pooled across all sample dates and Mao Tau species richness and 95% confidence intervals were calculated for each treatment as suggested in Colwell *et al.* (2004). Treatment differences were tested using a two-way crossed ANOVA for each year. Treatments were compared at an abundance shared by all treatments on the asymptotic part of the rarefaction curves, where species richness had stabilised. All data met normality and equal variance assumptions.

We assessed arthropod community composition (which accounts for abundance, richness and family identity) using non-metric multidimensional scaling (NMDS) (Shepard, 1962; Kruskal, 1964). NMDS is commonly used in arthropod community analysis (e.g. Wimp *et al.*, 2005) for several reasons: (i) it makes few assumptions about the nature of the data (e.g. it does not require a Gaussian distribution), (ii) it relies on only the rank order of similarities rather than their actual values (results are easy to understand), and (iii) it provides robust results (Clark & Warwick, 2001; McCune & Grace, 2002). We pooled data across sample dates within each year and analysed years separately using PRIMER 6. Pooled raw abundance data for each year were square root transformed and all families that did not comprise at least 1% of total abundance were excluded (Clark & Warwick, 2001). We then produced a dissimilarity matrix using the Bray–Curtis

dissimilarity coefficient (Faith *et al.*, 1987). NMDS was run and stress levels were evaluated to select the best representation of the data by assessing how well the distances between samples were maintained from the dissimilarity matrix to the ordination. A two-dimensional solution was selected because it maintained a consistently low stress (< 0.2) across multiple runs. We tested for differences in community composition using a two-way crossed ANOVA on the NMDS scores. NMDS scores represent the relative similarity of each of the 12 FACE rings to each other (points in ordination plots that are near one another are more similar than those that are farther apart). Because NMDS plots are based only on the relative similarities of the specific data points, axes have no intrinsic meaning.

We used indicator analysis (Dufrene & Legendre, 1997) in PC-ORD5 to examine whether particular families were significant indicators of control, +CO₂, +O₃, or +CO₂+O₃ treatments. This analysis is specific to treatments because we could not test both CO₂ and O₃ main effects simultaneously as we did for our abundance and community composition data. Indicator analysis accounts for each family's fidelity to a treatment as well as its abundance within that treatment. Values range from zero (no indication) to 100 (perfect indication) (McCune & Grace, 2002). Perfect indication for this data set would mean a family was found in only one treatment. Results for data pooled across all 4 years are presented.

Results

In total, we examined 47 415 insects from 4 orders and 83 families. Over half of the specimens collected were Diptera (54.8%), followed by Hymenoptera (28.4%), Hemiptera (13.9%) and Coleoptera (2.9%). The 15 most abundant families consisted of six Diptera, five Hymenoptera, three Hemiptera and one Coleoptera.

Arthropod abundance

Changes in total arthropod abundance were determined by changes in individual family abundance. Elevated CO₂ had no effect on total abundance. Elevated O₃, alternatively, reduced total abundance by 17% compared with ambient O₃ conditions, largely as a result of the negative effects on parasitoids (Table 1). Total abundance also varied by collection year, with the highest abundances in 2000 and 2003 and much lower abundance in 2001 [raw abundance (no. of individuals) 2000: 15 442; 2001: 6923; 2002: 9793; 2003: 15 257] ($F_{3,24} = 37.33$, $P < 0.001$).

Analysis of individual family abundance reveals that eCO₂ and eO₃ had significant effects on numbers of some insect families (Table 1). Elevated CO₂ substantially decreased the abundance of Formicidae (46%) compared with ambient CO₂. In contrast, eCO₂ increased numbers of Figitidae (103%) and Ichneumonidae (28%) parasitoids. Trends for decreased numbers of Cicadellidae (35%), Miridae (35%) and Sarcophagidae (21%) and increased numbers of Curculionidae (26%) and Anthomyiidae (58%) were apparent but these results were not statistically significant. Overall, the trend under eCO₂ was for the abundance of sucking

Table 2. Mean (± 1 SE) number of insect families collected per pan trap per year.

Year	Control	+CO ₂	+O ₃	+CO ₂ +O ₃
2000	38 \pm 2	34 \pm 1	35 \pm 1	37 \pm 1
2001	35 \pm 2	34 \pm 3	35 \pm 1	34 \pm 2
2002	34 \pm 1	35 \pm 4	39 \pm 2	34 \pm 1
2003	40 \pm 1	37 \pm 4	41 \pm 2	38 \pm 2

herbivores and families using resources similarly to sucking herbivores (e.g. honeydew and nectaries) to decrease whereas parasitoids tended to increase. Relative changes in abundance due to eCO₂ were generally consistent across years.

The effect of eO₃ on insect abundance was generally opposite that of eCO₂. Elevated O₃ had a strong negative effect on parasitoid abundances, reducing Braconidae by 33%, Chalcidoidea by 26%, Figitidae by 59% and Ichneumonidae by 41% (Table 1). A pronounced O₃ \times year interaction occurred, however, for Figitidae ($F_{3,24} = 4.23$, $P = 0.034$). From 2000–2002, Figitidae abundance was decreased 22–59% but in 2003, numbers increased 66%. Elevated O₃ also decreased Curculionidae (27%) and Drosophilidae (42%) but these changes were not statistically significant. In contrast to the largely negative effects of eO₃ on other families, Miridae abundance increased 95%. Cicadellidae abundance increased 51% but again the change was not significant. Generally, eO₃ had the strongest impact on parasitoids (negative) but also influenced some sucking herbivores (positive). Families other than Figitidae showed more consistent relative changes in abundance due to eO₃ across years.

The effects of CO₂ and O₃ on insect abundance were independent of one another. The single exception was for the family Aphididae, for which eCO₂ increased abundance under aO₃, but decreased abundance under eO₃ (Table 1).

Arthropod family richness

We found no significant differences in arthropod family richness among fumigation treatments for any year (Table 2). FACE rings contained 34–41 families across CO₂ and O₃ treatments and years.

Arthropod community composition

Arthropod community composition shifted in response to CO₂ and O₃ treatments in all 4 years of this study. Elevated CO₂ affected composition in 2001, 2002 and 2003, whereas eO₃ affected composition in 2000 and 2003 (Fig. 1). Overall model P -values were significant in 2001 ($F_{3,8} = 4.00$, $P = 0.050$) and 2003 ($F_{3,8} = 12.30$, $P = 0.002$) but not significant in 2000 ($F_{3,8} = 3.30$, $P = 0.080$) and 2002 ($F_{3,8} = 2.70$, $P = 0.119$). CO₂ and O₃ affected the insect community independently (no significant CO₂ \times O₃ interactions). Thus, although family richness was not altered by eCO₂ or eO₃, shifts in insect abundance due the treatments led to changes in community composition that varied among years.




Fig. 1. 2000–2003 non-metric multidimensional scaling (NMDS) ordinations of community composition. Differences among treatments were tested using two-way crossed ANOVA on NMDS axis 1. A CO_2 main effect tests filled versus unfilled symbols, whereas an O_3 main effect tests circles versus triangles.

(a) 2000
(b) 2001
(c) 2002
(d) 2003

○ Control ● $+\text{CO}_2$ △ $+\text{O}_3$ ▲ $+\text{CO}_2+\text{O}_3$

Table 3. Indicator analysis (Dufrene & Legendre, 1997) across 2000–2003 for the 15 most abundant families. Indicator values can range from 0–100, with 100 signifying complete fidelity to the indicated treatment.

Feeding guild/family	Indicator value	Indicated treatment	<i>P</i>
Herbivores – phloem feeders			
Aphididae	26.5	$+\text{O}_3$	0.082
Cicadellidae	28.7	$+\text{O}_3$	0.025
Miridae	23.3	$+\text{O}_3$	0.001
Herbivores – chewers			
Curculionidae	15	$+\text{CO}_2$	0.447
Anthomyiidae	29.7	$+\text{CO}_2+\text{O}_3$	0.025
Parasitoids			
Braconidae	25	Control	0.1
Chalcidoidea	27.6	Control	0.085
Figitidae	30.3	$+\text{CO}_2$	0.001
Ichneumonidae	33.2	$+\text{CO}_2$	0.001
Predators			
Dolicopodidae	20.7	Control	0.315
Saprophages			
Drosophilidae	14	$+\text{CO}_2$	0.163
Phoridae	32.2	Control	0.002
Sciaridae	28	$+\text{CO}_2$	0.122
Herbivores/Predators			
Formicidae	25.2	$+\text{O}_3$	0.014
Predators/Saprophages			
Sarcophagidae	18.1	$+\text{O}_3$	0.344

Community composition indicator analysis

Indicator analysis designated 7 of the 15 most abundant families as significant indicators of particular treatments (Table 3). The $+\text{CO}_2$ treatment was indicated by two families, Figitidae and Ichneumonidae, in agreement with their increased abundance at $e\text{CO}_2$. Increased abundance at $e\text{O}_3$, similarly, led to three

families that indicated $+\text{O}_3$ treatments: Cicadellidae, Miridae and Formicidae. Aphididae also tended to increase when exposed to $e\text{O}_3$ but the trend was not significant. Current atmospheric concentrations of CO_2 and O_3 (control) allowed for the greatest abundances of Phoridae. The combined elevated treatment ($+\text{CO}_2+\text{O}_3$), in contrast, was indicated by Anthomyiidae. Anthomyiidae may have indicated $+\text{CO}_2+\text{O}_3$ due to $e\text{CO}_2$ significantly increasing Anthomyiidae abundance, but doing so slightly more in the $+\text{CO}_2+\text{O}_3$ treatment than in the $+\text{CO}_2$ treatment. In general, indicator analysis results reflect the changes in family abundances.

Discussion

Insect communities in the second half of the 21st century will likely be significantly different from present forest insect communities due to elevated concentrations of CO_2 and O_3 . Our results suggest that $e\text{CO}_2$ may favour some parasitoid families but limit populations of sucking insects. In contrast, forests exposed to $e\text{O}_3$ may have significantly fewer insects, especially parasitoids, although phloem-feeding insects may benefit. CO_2 and O_3 effects may be expected to influence families/feeding guilds differently, leading to altered community composition. Insects exposed to both $e\text{CO}_2$ and $e\text{O}_3$ appear to respond to each gas independently. Therefore, offsetting effects of $e\text{CO}_2$ and $e\text{O}_3$ may occur for insect families affected positively by one gas and negatively by the other. The lack of interactions suggests that, at least for some global change factors, accurate predictions may be drawn from studies manipulating single variables.

Abundance

Total insect abundance did not change under $e\text{CO}_2$ but was reduced under $e\text{O}_3$. Similarly, total insect abundance was unaffected

by eCO₂ at the Oak Ridge FACE site (Sanders *et al.*, 2004) and at SoyFACE (minus a large effect on Japanese beetles for one sample date; Hamilton *et al.*, 2005). Total insect abundance at SoyFACE was also unaffected by eO₃ (Hamilton *et al.*, 2005). Our eO₃ results likely differ from those of Hamilton *et al.*, (2005) because they sampled only herbivores and our major reduction in abundance was due to the strongly reduced abundance of parasitoids. Changes in total insect abundance appear to depend on the families or feeding guilds sampled.

Abundances of four of the five herbivore families we examined exhibited strong mean responses to eCO₂, eO₃ or both, but low replication precluded statistical significance. Changes in abundance due to eCO₂ and eO₃ were dependent on herbivore feeding guild (chewing vs. phloem feeding) as suggested in reviews by Bezemer & Jones (1998) and Koricheva *et al.* (1998). We found that chewing herbivores increased in abundance under eCO₂ and decreased in abundance under eO₃, while phloem-feeders showed the converse. Bezemer & Jones (1998) and Stiling & Cornelissen (2007) suggested the opposite pattern for eCO₂. Our results may differ from their predictions because we sampled families with feeding habits different from those of previous work, as well as families that were not previously examined. For example, species of the two chewing herbivore families we trapped, Curculionidae and Anthomyiidae, feed on roots below ground as larvae. Indeed, our results match those of Altermatt (2003) for weevil abundance. We also recognise that pan traps are not an effective means of capturing many families of herbivorous insects, which limits our conclusions. Broader sampling of many herbivore families will be necessary to determine if our trends hold across multiple feeding guilds.

Global change is predicted to have larger impacts at higher trophic levels (Voight *et al.*, 2003) and our data support that perspective. Parasitoid abundance increased under eCO₂ for two of the four families we examined and decreased under eO₃ for all four families. Previous work suggested that eCO₂ and eO₃ can increase or decrease parasitoid abundance, respectively, for short periods, but have no effect on parasitoid abundance over an entire season (Percy *et al.*, 2002; Awmack *et al.*, 2004). Furthermore, predation and parasitism rates under eCO₂ have been found to increase (Stiling *et al.*, 1999, 2003), decrease (Gate *et al.*, 1995) or not change (Stacy & Fellowes, 2002). While increased predation at eCO₂ could benefit pest control, the particularly strong negative impacts of O₃ on parasitoids could have major negative implications for pest control. Reduced ability to find prey, or the host plants of prey, is one possible explanation for the substantial reduction of parasitoids at eO₃. A 1995 study by Gate *et al.* found that eO₃ interferes with wasp searching behaviour. Furthermore, recent work suggests that eCO₂ and eO₃ inhibit the ability of wasps to detect damaged plants (Vuorinen *et al.*, 2004; Pinto *et al.*, 2007). Parasitoids could also be influenced by changes in herbivore behaviour at eCO₂ and eO₃ (Mondor *et al.*, 2004).

Saprophagous insects showed the fewest significant responses to the treatments, despite previous work at Aspen FACE suggesting strong influences of eCO₂ and eO₃ on litter microarthropods (Loranger *et al.*, 2004). Our results likely differ because Loranger *et al.* (2004) focused on Collembola which feed on bacteria whereas we examined saprophagous species feeding on decaying

organic material or fungi. In agreement with our results, Sanders *et al.* (2004) found no effect of CO₂ on detritivore abundance.

Insect species richness

Neither the CO₂ nor O₃ treatments affected species richness. Similarly, insect species richness was unaffected by both CO₂ in a sweetgum understorey at the Oak Ridge National Laboratory FACE site (Sanders *et al.*, 2004) and a natural O₃ pollution gradient in California forests (Jones & Paine, 2006). In contrast, eCO₂ did decrease insect species richness on beech and oak trees, although sample sizes were small (Altermatt, 2003). Together, these studies suggest that eCO₂ and eO₃ will have negligible impacts on species richness via direct or plant-mediated effects, at least over the short term. The effects of CO₂ and O₃ may, however, magnify over longer timescales. The strong negative effect of O₃ on parasitoids could increase extinction risk and reduce diversity over time. CO₂ and O₃ may also alter diversity by changing insect species dominance (e.g. making the most abundant species more abundant) in the community (Jones & Paine, 2006). Finally, the effects of CO₂ and O₃ may be mediated more strongly via their function as greenhouse gases. Decreases in insect diversity over the past three decades appear to be linked to elevated temperatures (Lovejoy & Hannah, 2005).

We acknowledge that the taxonomic level of family may be too coarse to evaluate fully the richness for this system. However, species level analyses have also revealed no change in richness in response to eCO₂ and eO₃ (Sanders *et al.*, 2004; Jones & Paine, 2006).

Arthropod community composition

Few studies have examined the effects of eCO₂ or eO₃ on insect community composition. We found that eCO₂ altered forest insect communities in 3 of 4 years, while eO₃ modified communities in 2 of 4 years. Previous work at another forest FACE site showed that eCO₂ had no effect on understorey arthropod community composition (Sanders *et al.*, 2004). Elevated CO₂ has, however, been shown to cause the most abundant herbivores to become more abundant on forest trees, (Altermatt, 2003) an effect that could alter community composition. We found increased abundance of the most abundant insects in some years but not in others. Insect communities could also be modified by changes in the ratio of chewing to sucking herbivores. Jones & Paine (2006) found that insect communities exposed to elevated O₃ concentrations may shift from phloem-feeding to chewing-dominated herbivore communities. Our results suggest the opposite effect, with phloem-feeders benefiting from eO₃ environments.

Mechanisms and implications

Changes in insect abundance and community composition at eCO₂ and eO₃ are likely due to a combination of bottom-up, top-down and abiotic effects. Meta-analysis has clearly shown

that both plant quality and herbivore performance are altered by CO₂ and O₃ (Zvereva & Kozlov, 2006; Valkama *et al.*, 2007). Altered herbivore quality may in turn influence predator fitness (Ode, 2006), altering top-down effects on the herbivore community. Insect communities also likely differ due to eCO₂- and eO₃-mediated alterations in forest environment. For example, forest stand development in eCO₂ rings reached canopy closure over the 4 years of this study, which may be responsible for the strengthening eCO₂ effect. The forest environment along with insect populations and plant quality are also affected directly and indirectly by major weather patterns (e.g. temperature, precipitation) each year.

Predicting abiotic conditions of the future is critical to conservation efforts. New regional climates, in combination with individualistic species responses, may mean disassembly of current communities and establishment of 'no-analogue' communities (Williams & Jackson, 2007). Under environmental conditions of the future, many plants and insects will likely be forced to adapt, move to new areas, or go extinct. Plants will respond directly and indirectly to changes in the abiotic environment, whereas insects will respond to both changes in plant community composition and climate (Schaffers *et al.*, 2008). Additionally, insects may respond to new conditions with altered feeding preferences (Goverde & Erhardt, 2003; Agrell *et al.*, 2006) and food consumption (Hamilton *et al.*, 2004; Hamilton *et al.*, 2005; Knepp *et al.*, 2005). The consequences of potential 'no-analogue' communities for ecosystem function remains to be determined.

Conclusions

Elevated CO₂ and eO₃ altered insect abundance and community composition but not family level richness. Feeding guild may strongly influence insect response to environmental change and may provide the best opportunity to generalise for conservation efforts. Because insects play key roles in forest ecosystems, changes in insect abundance, diversity or community composition have the potential to alter forest ecosystems and the services they provide. Continued multifactor research and increased public awareness will be essential to forest management and conservation efforts.

Acknowledgments

Aspen FACE is principally supported by the Office of Science (BER), US Department of Energy, Grant No. DE-FG02-95ER62125 to Michigan Technological University, and Contract No. DE-AC02-98CH10886 to Brookhaven National Laboratory, the US Forest Service Northern Global Change Program and North Central Research Station, Michigan Technological University, and Natural Resources Canada – Canadian Forest Service. We thank C. Gratton and T. Meehan for statistical advice, C. Awmack, E. Mondor and M. Tremblay for help collecting pan traps, and D. Young and S. Krauth for help with insect identification. This work was supported by National Science Foundation grant DEB-0129123, US Department of Energy (Office of Science, BER) grant DE-FG02-06ER64232, and University of Wisconsin Hatch grant WIS04898 to R.L. Lindroth.

References

- Agrell, J., Anderson, P., Oleszek, W., Stochmal, A. & Agrell, C. (2006) Elevated CO₂ levels and herbivore damage alter host plant preferences. *Oikos*, **112**, 63–72.
- Altermatt, F. (2003) Potential negative effects of atmospheric CO₂-enrichment on insect communities in the canopy of a mature deciduous forest in Switzerland. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft Bulletin de la Societe Entomologique Suisse*, **76**, 191–199.
- Awmack, C.S., Harrington, R. & Lindroth, R.L. (2004) Individual performance does not predict aphid population responses to elevated atmospheric CO₂ or O₃. *Global Change Biology*, **10**, 1414–1423.
- Bailey, J.K. & Whitham, T.G. (2002) Interactions among fire, aspen, and elk affect insect diversity: reversal of a community response. *Ecology*, **83**, 1701–1712.
- Bezemer, T.M. & Jones, T.H. (1998) Plant-insect herbivore interactions in elevated atmospheric CO₂: quantitative analyses and guild effects. *Oikos*, **82**, 212–222.
- Buddle, C.M., Beguin, J., Bolduc, E., Mercado, A., Sackett, T.E., Selby, R.D., Varady-Szabo, H. & Zeran, R.M. (2005) The importance and use of taxon sampling curves for comparative biodiversity research with forest arthropod assemblages. *Canadian Entomologist*, **137**, 120–127.
- Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M. & Jouseau, C. (2006) Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, **443**, 989–992.
- Clark, K.R. & Warwick, R.M. (2001) *Change in Marine Communities: An Approach to Statistical Analysis and Interpretation*, 2nd edn. PRIMER-E, Plymouth, UK.
- Colwell, R.K. (2006) *ESTIMATES*. Available from URL: <http://viceroy.eeb.uconn.edu/estimates>
- Colwell, R.K., Mao, C.X. & Chang, J. (2004) Interpolating, extrapolating, and comparing incidence based species accumulation curves. *Ecology*, **85**, 2717–2727.
- Coviella, C. & Trumble, J.T. (1999) Effects of elevated atmospheric CO₂ on insect-plant interactions. *Conservation Biology*, **13**, 700–712.
- Dickson, R.E., Coleman, M.D., Riemenschneider, D.E., Isebrands, J.G., Hogan, G.D., & Karnosky, D.F. (1998) Growth of five hybrid poplar genotypes exposed to interacting elevated CO₂ and O₃. *Canadian Journal of Forest Research*, **28**, 1706–1716.
- Dickson, R.E., Lewin, K.F., Isebrands, J.G., Coleman, M.D., Heilman, W.E., Riemenschneider, D.E., Sober, J., Host, G.E., Zak, D.R., Hendry, G.R., Pregitzer, K.S. & Karnosky, D.F. (2000) *Forest Atmosphere Carbon Transfer Storage-II (FACTS II) – The Aspen Free-Air CO₂ and O₃ Enrichment (FACE) Project: an Overview*, pp. 68. General Technical Report NC-214. USDA Forest Service, North Central Research Station, Rhinelander, Wisconsin.
- Duelli, P., Obrist, M.K. & Schmatz, D.R. (1999) Biodiversity evaluation in agricultural landscapes: above-ground insects. *Agriculture, Ecosystems and Environment*, **74**, 33–64.
- Dufrene, M. & Legendre, P. (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs*, **67**, 345–366.
- Faith, D.P., Minchin, P.R. & Belbin, L. (1987) Compositional dissimilarity as a robust measure of ecological distance. *Vegetatio*, **69**, 57–68.
- Fowler, D., Cape, J.N., Coyle, M., Flechard, C., Kuylenstierna, J., Hicks, K., Derwent, D., Johnson, C. & Stevenson, D. (1999) The global exposure of forests to air pollutants. *Water, Air and Soil Pollution*, **116**, 5–32.
- Gate, I.M., McNeill, S. & Ashmore, M.R. (1995) Effects of air pollution on the searching behavior of an insect parasitoid. *Water, Air and Soil Pollution*, **85**, 1425–1430.

- Gotelli, N.J. & Ellison, A.M. (2004) *A Primer of Ecological Statistics*. Sinauer & Associates, Inc. Sunderland, Massachusetts.
- Goverde, M. & Erhardt, A. (2003) Effects of elevated CO₂ on development and larval food-plant preference in the butterfly *Coenonympha pamphilus* (Lepidoptera, Satyridae). *Global Change Biology*, **9**, 74–83.
- Hamilton, J.G., Dermody, O., Aldea, M., Zangerl, A.R., Rogers, A., Berenbaum, M.R. & DeLucia, E.H. (2005) Anthropogenic changes in tropospheric composition increase susceptibility of soybean to insect herbivory. *Environmental Entomology*, **34**, 479–485.
- Hamilton, J.G., Zangerl, A.R., Berenbaum, M.R., Phippen, J., Aldea, M. & DeLucia, E.H. (2004) Insect herbivory in an intact forest understory under experimental CO₂ enrichment. *Oecologia*, **138**, 566–573.
- Hansen, A. & Dale, V. (2001) Biodiversity in US forests under global climate change. *Ecosystems*, **4**, 161–163.
- Intergovernmental Panel on Climate Change (2007) *Climate Change 2007: The Physical Scientific Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK.
- Jones, M.E. & Paine, T.D. (2006) Detecting changes in insect herbivore communities along a pollution gradient. *Environmental Pollution*, **143**, 377–387.
- Karnosky, D.F., Zak, D.R., Pregitzer, K.S., Awmack, C.S., Bockheim, J.G., Dickson, R.E., Hendrey, G.R., Host, G.E., King, J.S., Kopper, B.J., Kruger, E.L., Kubiske, M.E., Lindroth, R.L., Mattson, W.J., McDonald, E.P., Noormets, A., Oksanen, E., Parsons, W.F.J., Percy, K.E., Podila, G.K., Riemenschneider, D.E., Sharma, P., Thakur, R.C., Sober, A., Sober, J., Jones, W.S., Anttonen, S., Vapaavuori, E., Mankovska, B., Heilman, W.E. & Isebrands, J.G. (2003) Tropospheric O₃ moderates responses of temperate hardwood forests to elevated CO₂: a synthesis of molecular to ecosystem results from the Aspen FACE project. *Functional Ecology*, **17**, 289–304.
- Knepp, R.G., Hamilton, J.G., Mohan, J.E., Zangerl, A.R., Berenbaum, M.R. & DeLucia, E.H. (2005) Elevated CO₂ reduces leaf damage by insect herbivores in a forest community. *New Phytologist*, **167**, 207–218.
- Koricheva, J., Larsson, S., Haukioja, E. & Keinänen, M. (1998) Insect performance on experimentally stressed woody plants: a meta-analysis. *Annual Review of Entomology*, **43**, 195–216.
- Kruskal, J.B. (1964) Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*, **29**, 1–26.
- Lindroth, R.L., Kopper, B.J., Parsons, W.F.J., Bockheim, J.G., Karnosky, D.F., Hendry, G.R., Pregitzer, K.S., Isebrands, J.G. & Sober, J. (2001) Consequences of elevated carbon dioxide and ozone for foliar chemical composition and dynamics in trembling aspen (*Populus tremuloides*) and paper birch (*Betula papyrifera*). *Environmental Pollution*, **115**, 395–404.
- Loranger, G.I., Pregitzer, K.S. & King, J.S. (2004) Elevated CO₂ and O₃ concentrations differentially affect selected groups of fauna in temperate forest soils. *Soil Biology & Biochemistry*, **36**, 1521–1524.
- Lovejoy, T.E. & Hannah, L. (2005) *Climate Change and Biodiversity*. Yale University Press, New Haven & London.
- Mattson, W.J. & Addy, N.D. (1975) Phytophagous insects as regulators of forest primary production. *Science*, **190**, 515–522.
- McCune, B. & Grace, J.B. (2002) *Analysis of Ecological Communities*. MjM Software Design, Gleneden Beach, Oregon.
- Mondor, E.B., Tremblay, M.N., Awmack, C.S. & Lindroth, R.L. (2004) Divergent pheromone-mediated insect behavior under global atmospheric change. *Global Change Biology*, **10**, 1820–1824.
- Moran, M.D. (2003) Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos*, **100**, 403–405.
- Ode, P.J. (2006) Plant chemistry and natural enemy fitness: effects on herbivore and natural enemy interactions. *Annual Review of Entomology*, **51**, 163–185.
- Ozanne, C.M.P., Anhof, D., Boulter, S.L., Keller, M., Kitching, R.L., Korner, C., Meinzer, F.C., Mitchell, A.W., Nakashizuka, T., Silva Diaz, P.L., Stork, N.E., Wright, S.J. & Yoshimura, M. (2003) Biodiversity meets the atmosphere: a global view of forest canopies. *Science*, **301**, 183–186.
- Percy, K.E., Awmack, C.S., Lindroth, R.L., Kubiske, M.E., Kopper, B.J., Isebrands, J.G., Pregitzer, K.S., Hendrey, G.R., Dickson, R.E., Zak, D.R., Oksanen, E., Söber, J., Harrington, R. & Karnosky, D.F. (2002) Altered performance of forest pests under atmospheres enriched by CO₂ and O₃. *Nature*, **420**, 403–407.
- Pinto, D.M., Nerg, A. & Holopainen, J.K. (2007) The role of ozone-reactive compounds, terpenes, and green leaf volatiles (GLV's), in the orientation of *Cotesia plutellae*. *Journal of Chemical Ecology*, **33**, 2218–2228.
- Piva, R.J. (2006) *Pulpwood Production in the North-Central Region, 2004*. United States Department of Agriculture Forest Service North Central Research Station. Resource Bulletin NC-265. St. Paul, Minnesota.
- Sanders, N.J., Belote, R.T. & Weltzin, J.F. (2004) Multitrophic effects of elevated CO₂ on understory plant and arthropod communities. *Environmental Entomology*, **33**, 1609–1616.
- Saxe, H., Ellsworth, D.S. & Heath, J. (1998) Tree and forest functioning in an enriched CO₂ atmosphere. *New Phytologist*, **139**, 395–436.
- Schaffers, A.P., Raemakers, I.P., Sykora, K.V. & ter Braak, C.J.F. (2008) Arthropod assemblages are best predicted by plant species composition. *Ecology*, **89**, 782–794.
- Schowalter, T.D., Hargrove, W.W. & Crossley, D.A. Jr. (1986) Herbivory in forested ecosystems. *Annual Review of Entomology*, **31**, 177–196.
- Shepard, R.N. (1962) The analysis of proximities: multidimensional scaling with an unknown distance function. *Psychometrika*, **27**, 125–139.
- Sitch, S., Cox, P.M., Collins, W.J. & Huntingford, C. (2007) Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. *Nature*, **448**, 791–795.
- Stacy, D.A. & Fellowes, M.D.E. (2002) Influence of elevated CO₂ on interspecific interactions at higher trophic levels. *Global Change Biology*, **8**, 668–678.
- Stiling, P. & Cornelissen, T. (2007) How does elevated carbon dioxide (CO₂) affect plant-herbivore interactions? A field experiment and meta-analysis of CO₂-mediated changes on plant chemistry and herbivore performance. *Global Change Biology*, **13**, 1–20.
- Stiling, P., Moon, D.C., Hunter, M.D., Colson, J., Rossi, A.M., Hymus, G.J. & Drake, B. (2003) Elevated CO₂ lowers relative and absolute herbivore density across all species of scrub-oak forest. *Oecologia*, **134**, 82–87.
- Stiling, P., Rossi, A.M., Hungate, B., Dukstra, P., Hinkle, C.R., Knott III, W.M. & Drake, B. (1999) Decreased leaf-miner abundance in elevated CO₂: reduced leaf quality and increased parasitoid attack. *Ecological Applications*, **9**, 240–244.
- Triplehorn, C.A. & Johnson, N.F. (2005) *Borror and DeLong's Introduction to the Study of Insects*, 7th edn. Thomson Learning, Belmont, California.
- Valkama, E., Koricheva, J. & Oksanen, E. (2007) Effects of elevated O₃, alone and in combination with elevated CO₂, on tree leaf chemistry and insect herbivore performance: a meta-analysis. *Global Change Biology*, **13**, 184–201.
- Veblen, T.T., Hadley, K.S., Reid, M.S. & Rebertus, A.J. (1991) The response of subalpine forests to spruce beetle outbreak in Colorado. *Ecology*, **72**, 213–231.
- Voight, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bahrmann, R., Fabian, B., Heinrich, W., Kohler, G., Lichter, D., Marstaller, R. & Sander, F.W. (2003) Trophic levels are differentially sensitive to climate. *Ecology*, **84**, 2444–2453.
- Vuorinen, T., Nerg, A., Ibrahim, M.A., Reddy, G.V.P. & Holopainen, J. (2004) Emission of *Plutella xylostella*-induced compounds from

- cabbages grown at elevated CO₂ and orientation behavior of the natural enemies. *Plant Physiology*, **135**, 1984–1992.
- Williams, J.W. & Jackson, S.T. (2007) Novel climates, no-analog communities, and ecological surprises. *Frontiers in Ecology and the Environment*, **5**, 475–482.
- Wimp, G.W., Martinsen, G.D., Floate, K.D., Bangert, R.K. & Whitham, T.G. (2005) Plant genetic determinants of arthropod community structure and diversity. *Evolution*, **59**, 61–69.
- Zvereva, E.L. & Kozlov, M.V. (2006) Consequences of simultaneous elevation of carbon dioxide and temperature for plant-herbivore interactions: a metaanalysis. *Global Change Biology*, **12**, 27–41.

Accepted 9 September 2008

Editor: Simon Leather

Associate editor: Martin Bezemer