



**HAL**  
open science

## Elevated expression of complement C4 in the mouse prefrontal cortex causes schizophrenia-associated phenotypes

Mélanie Druart, Marika Nosten-Bertrand, Stefanie Poll, Sophie Crux, Felix Nebeling, Célia Delhaye, Yaëlle Dubois, Manuel Mittag, Marion Leboyer, Ryad Tamouza, et al.

### ► To cite this version:

Mélanie Druart, Marika Nosten-Bertrand, Stefanie Poll, Sophie Crux, Felix Nebeling, et al.. Elevated expression of complement C4 in the mouse prefrontal cortex causes schizophrenia-associated phenotypes. *Molecular Psychiatry*, Nature Publishing Group, 2021, 10.1038/s41380-021-01081-6 . hal-03203920

**HAL Id: hal-03203920**

**<https://hal.sorbonne-universite.fr/hal-03203920>**

Submitted on 21 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Elevated expression of complement C4 in the mouse prefrontal cortex**  
2 **causes schizophrenia-associated phenotypes**

3  
4 **Running title:** Elevated C4 causes schizophrenia-like phenotypes

5  
6 Mélanie Druart<sup>1,2,3</sup>, Marika Nosten-Bertrand<sup>1,2,3\*</sup>, Stefanie Poll<sup>4\*</sup>, Sophie Crux<sup>4\*</sup>, Felix  
7 Nebeling<sup>4\*</sup>, Célia Delhaye<sup>1,2,3</sup>, Yaëlle Dubois<sup>1,2,3</sup>, Manuel Mittag<sup>4</sup>, Marion Leboyer<sup>5,6</sup>, Ryad  
8 Tamouza<sup>5,6</sup>, Martin Fuhrmann<sup>4</sup>, Corentin Le Magueresse<sup>1,2,3#</sup>

9  
10 <sup>1</sup>INSERM UMR-S 1270, 75005 Paris, France

11 <sup>2</sup>Sorbonne Université, 75005 Paris, France

12 <sup>3</sup>Institut du Fer à Moulin, 17 rue du Fer à Moulin, 75005 Paris, France

13 <sup>4</sup>Neuroimmunology and Imaging Group, German Center for Neurodegenerative Diseases,  
14 53175 Bonn, Germany

15 <sup>5</sup>INSERM U955, Neuro-Psychiatrie Translationnelle, Université Paris-Est, Créteil, France.

16 <sup>6</sup>AP-HP, DMU IMPACT, Département Médical Universitaire de Psychiatrie, Hôpitaux  
17 Universitaires Henri Mondor, Créteil, France.

18  
19 \* Equal contribution

20 # Correspondence: corentin.le-magueresse@inserm.fr

21  
22  
23  
24  
25 **Abstract**

26 Accumulating evidence supports immune involvement in the pathogenesis of  
27 schizophrenia, a severe psychiatric disorder. In particular, high expression variants of *C4*, a  
28 gene of the innate immune complement system, were shown to confer susceptibility to  
29 schizophrenia. However, how elevated C4 expression may impact brain circuits remains  
30 largely unknown. We used *in utero* electroporation to overexpress C4 in the mouse prefrontal  
31 cortex (PFC). We found reduced glutamatergic input to pyramidal cells of juvenile and adult,  
32 but not of newborn C4-overexpressing (C4-OE) mice, together with decreased spine density,  
33 which mirrors spine loss observed in the schizophrenic cortex. Using time-lapse two-photon  
34 imaging *in vivo*, we observed that these deficits were associated with decreased dendritic

35 spine gain and elimination in juvenile C4-OE mice, which may reflect poor formation and/or  
36 stabilization of immature spines. In juvenile and adult C4-OE mice we found evidence for  
37 NMDA receptor hypofunction, another schizophrenia-associated phenotype, and synaptic  
38 accumulation of calcium-permeable AMPA receptors. Alterations in cortical GABAergic  
39 networks have been repeatedly associated with schizophrenia. We found that functional  
40 GABAergic transmission was reduced in C4-OE mice, in line with diminished GABA release  
41 probability from parvalbumin interneurons, lower GAD67 expression and decreased intrinsic  
42 excitability in parvalbumin interneurons. These cellular abnormalities were associated with  
43 working memory impairment. Our results substantiate the causal relationship between an  
44 immunogenetic risk factor and several distinct cortical endophenotypes of schizophrenia, and  
45 shed light on the underlying cellular mechanisms.

46

## 47 **Introduction**

48 Genome-wide association studies (GWAS) identified genes involved in immune and  
49 in autoimmune processes that are associated with Schizophrenia (SZ)<sup>1-4</sup>. The major GWAS  
50 locus in SZ lies within the major histocompatibility complex (MHC) locus on chromosome  
51 6<sup>4</sup>. This includes the *C4* gene which consists of *C4A* and *C4B* gene copy number variations  
52 and insertion/deletion of the human endogenous retrovirus K (HERVK). A detailed analysis  
53 of this locus showed a strong relationship between the genetic diversity of the C4 region and  
54 SZ<sup>5</sup>. In particular, the presence of several copies of *C4A* with HERVK insertions correlated  
55 with higher *C4A* expression and increased risk of SZ. The complement pathway is a central  
56 determinant of synaptic pruning, a developmental process in which the synaptic connections  
57 between neurons are refined in the brain from childhood until early adulthood<sup>6</sup>. Thus, the  
58 identification of *C4A* as a risk factor in SZ established a link between complement-mediated  
59 synaptic pruning and dendritic spine loss in the cortex of schizophrenic patients, a  
60 characteristic feature of the disease which had long been hypothesized to reflect excessive  
61 synapse refinement in adolescence<sup>7,8</sup>.

62 In mice, the *C4b* gene presents homology with both the *C4A* and *C4B* genetic motifs  
63 of the human *C4* gene locus. Connectivity between the retina and the visual thalamus is  
64 impaired in C4 knock-out mice<sup>5</sup>, indicating that the mouse C4 protein participates in synaptic  
65 pruning. It has been hypothesized that high C4 expression may result in increased synaptic  
66 pruning in upper layers of the neocortex<sup>5</sup>. Indeed *C4A* and *C4B* gene copy numbers positively  
67 correlated with neuropil contraction, which is often considered as a mesoscale correlate of

68 developmental synapse elimination<sup>9</sup>. *C4A* gene copy numbers are associated with increased  
69 neuronal complement deposition and microglial synapse uptake in co-cultures of human  
70 iPSC-derived neurons and microglial cells<sup>10</sup>. Furthermore, it was shown that elevated C4  
71 expression in the mouse prefrontal cortex causes dendritic spine loss, increased engulfment of  
72 synaptic material by microglial cells and impaired social behaviors<sup>11</sup>. A recently published  
73 study also showed that mice overexpressing human C4A display reduced cortical synapse  
74 density, increased microglial engulfment of synapses and altered social behavior, anxiety  
75 levels and spatial short-term memory<sup>12</sup>.

76 In addition to spine loss, several neural endophenotypes have been identified in the  
77 cortex of schizophrenic patients. Decreased NMDA receptor (NMDAR)-mediated  
78 neurotransmission in SZ has been inferred from at least four lines of evidence<sup>13</sup>: i) NMDAR  
79 antagonists induce schizophrenic-like behaviors; ii) the expression of NMDAR subunits is  
80 reduced in post-mortem brain tissue from schizophrenic patients iii) auto-antibodies against  
81 NMDA receptors cause clinical symptoms of SZ; iv) *GRIN2A*, an NMDAR subunit-encoding  
82 gene, has been associated with SZ in recent GWAS. Moreover, multiple deficits in  
83 GABAergic circuits have been identified in SZ. Abnormalities in GABAergic networks  
84 include reduced GABA concentration in the brain of schizophrenic patients, decreased  
85 expression of GAD67, an enzyme responsible for GABA synthesis, and abnormalities at  
86 parvalbumin-expressing (PV) neuron-pyramidal cell synapses<sup>14</sup>. What causes these distinct  
87 alterations, and how they relate to each other, is poorly understood. The relationship between  
88 C4 and SZ-associated cortical endophenotypes such as reduced NMDA receptor-mediated  
89 transmission, loss of GAD67 expression and altered GABAergic synapses remains unknown.

90 Here we examined the consequences of complement C4 overexpression in layer II/III  
91 pyramidal cells of the mouse prefrontal cortex on neuronal circuit development and function.  
92 C4 overexpression recapitulated several cortical phenotypes that have been associated SZ,  
93 including decreased spine number and reduced NMDA receptor function in pyramidal cells.  
94 Intriguingly, dendritic spine loss was accompanied by decreased spine turnover in juvenile  
95 mice. We also report several abnormalities in GABAergic circuits in C4-OE mice which are  
96 consistent with the alterations of GABAergic markers seen in the brain of SZ patients.  
97 Finally, our results reveal that the selective overexpression of C4 in PFC layer II/III is  
98 sufficient to impair working memory, which is reminiscent of SZ-associated cognitive  
99 deficits.

100

101 **Methods**

102

103 A brief summary of experimental procedures is provided here with additional details available  
104 in the Supplemental Information accompanying this article. Additionally, Supplementary  
105 Tables S1-4 provide an overview of the number of recordings, animals and statistics in the  
106 different experiments.

107

108 **Animals**

109 All animal experiments were performed according to guidelines of the European Community  
110 for the use of animals in research and were approved by the local ethical committees (C2EA-  
111 05 in France; LANUV NRW in Germany). All experiments were performed on mice that had  
112 been electroporated at embryonic stage E15 and were on a C57Bl/6J background.

113

114 ***In utero* electroporation**

115 Pregnant mice at E15 were anesthetized by isoflurane 4% and were injected with Flunixin  
116 (0.05 mg/kg body weight) before surgery. The surgery was performed under isoflurane 2% on  
117 a heated blanket and breathing was carefully monitored during all the procedure.

118 Between 0.5  $\mu$ l to 1 $\mu$ l of a solution containing pCAG-C4HA vector (0.8  $\mu$ g/ $\mu$ l) together with  
119 pCAG-TdTomato vector (0.4  $\mu$ g/ $\mu$ l) (molar ratio 4:1) in NaCl 0.9% were injected into the  
120 bilateral ventricles. In the control condition, a solution containing pCAG-TdTomato (0.8  
121  $\mu$ g/ $\mu$ l) alone was injected. Electrical pulses (50V; 50 ms) were applied five times at intervals  
122 of 950 ms with an electroporator (see Supplementary Information).

123

124 ***In vivo* two-photon imaging**

125 Two-photon imaging was performed right after surgery at a Zeiss 7 multiphoton microscope  
126 with a Zeiss 20 x objective (NA = 1). TdTomato fluorescence was excited at 980 nm using an  
127 InSight X3 tunable laser (Spectra-Physics) and collected through a 617/73 band-pass filter on  
128 a highly sensitive non-descanned detector. After the first imaging session, the mice woke up  
129 and were placed in a cage with their littermates for 4-5 h (P15-18). For the second imaging  
130 session the mice were again anesthetized with isoflurane (initially 3%, then 1-1.5% for  
131 anesthesia maintenance) and imaged to retrieve the same dendritic structures recorded in the  
132 first imaging session (see Supplementary Information).

133

134 **Histology**

135 Mice aged P30 were anesthetized with pentobarbital (150 mg/kg) and intracardially perfused  
136 with PBS followed by 4% buffered formaldehyde (Histofix, Roti). Free-floating sections were  
137 processed for immunohistochemistry as described in the Supplementary Information. Images  
138 were acquired using a confocal microscope (SP5, Leica) and analyzed according to  
139 procedures described in the Supplementary Information.

140

141 **Acute slice preparation, *ex vivo* electrophysiology and optogenetics**

142 250  $\mu\text{m}$ -thick slices were prepared from brains of control and C4-OE mice. For patch-clamp  
143 recordings, slices were transferred to the recording chamber where they were continuously  
144 superfused with ACSF (30-32°C) buffered by continuous bubbling with 95% O<sub>2</sub>-5% CO<sub>2</sub>.

145 Whole-cell current and voltage-clamp recordings were performed in PFC layer II/III.  
146 Stimulus delivery and data acquisition were performed using Patchmaster software  
147 (Multichannel Systems). Signals were acquired with an EPC10-usb amplifier (Multichannel  
148 Systems), sampled at 20 kHz and filtered at 4 kHz. For optogenetic experiments, pulses of  
149 blue light (wavelength 470 nm) were emitted by a light emitting diode (CoolLED). Offline  
150 analysis was performed using Clampfit (Molecular Devices) and Igor Pro (Wavemetrics). For  
151 the study of miniature post synaptic currents, recordings were filtered offline at 2 kHz and  
152 analyzed using MiniAnalysis (Synaptosoft). See Supplementary Information.

153

154 **Behavior**

155 Tests for locomotor activity (actimeter, Open Field), and working memory (learning of a Non-  
156 Matching-To-Sample task, Odor Span Test, Spatial Span Test) were performed and analyzed  
157 as described in the Supplementary Information.

158

159 **Statistical analysis**

160 Data are presented as mean  $\pm$  SEM. Statistical analyses were performed using Prism  
161 (Graphpad). The normality of data distribution was tested using Shapiro-Wilk's test. Unpaired  
162 two-tailed T-tests (for normally distributed datasets) or Mann-Whitney tests (for non-  
163 normally distributed datasets) were used for comparisons between two groups. For multiple  
164 comparisons we used two-way ANOVA followed by Sidak's test. Values of  $P < 0.05$  were  
165 considered statistically significant.  $P$  values are reported as follows: \*  $P < 0.05$ ; \*\*  $P < 0.01$ ;  
166 \*\*\*  $P < 0.001$ ; \*\*\*\*  $P < 0.0001$ .

167

168 **Results**

169

170 **Generation of C4-overexpressing (C4-OE) mice using *in utero* electroporation**

171 Using *in utero* electroporation in C57Bl/6J mice we selectively overexpressed the C4-encoding  
172 *C4b* gene, the mouse homolog of the human *C4A* and *C4B* genes, in layer II/III pyramidal cells of  
173 the prefrontal cortex (PFC). Targeted PFC areas included the cingulate, prelimbic and infralimbic  
174 cortices (Supplementary figure S1A). To identify C4-OE neurons using immunohistochemistry,  
175 an HA tag was added at the N-terminal end of the *C4b* overexpression construct before the signal  
176 peptide. In addition, we co-electroporated the *C4b* construct with a tdTomato-expressing  
177 construct, in order to enable the identification of C4-expressing, tdTomato-positive (tdTom+)  
178 neurons in acute brain slices (Supplementary figure S1B-C). To ensure that most tdTom+ neurons  
179 indeed overexpress C4, we co-electroporated the *C4b* construct and the tdTomato construct with a  
180 4:1 ratio. Immunohistochemistry for HA and tdTomato confirmed that  $98.7 \pm 1.8$  % of tdTom+  
181 neurons also expressed C4 (Supplementary figure S1D-E). Control mice were electroporated with  
182 the tdTomato construct only. Due to the selectivity of the *in utero* electroporation procedure,  
183 which exclusively targets dividing progenitors of the ventricular zone, C4 overexpression was  
184 restricted to pyramidal cells. Interneurons or glial cells did not express the electroporated  
185 plasmids. Accordingly, all tdTomato-expressing cells had the characteristic morphology of  
186 cortical pyramidal cells, including a prominent apical dendrite projecting towards the pial surface.  
187 In immunoblots on lysates from the electroporated area, the HA tag was only detected at the  
188 molecular weight of the full-length form of C4 before its proteolytic cleavage (prepro-C4, ~ 200  
189 kDa) in the C4-OE condition, suggesting that the HA tag was correctly cleaved together with the  
190 signal peptide (Supplementary figure S1F-G). To quantify C4 overexpression in the  
191 electroporated area using western blot, we first evaluated several anti-C4 antibodies and found  
192 that an anti-C4 antibody raised against a motif of the  $\alpha$ -chain of human C4 reliably detected a  
193 band of the expected molecular weight in cortical tissue from control and C4-OE mice  
194 (Supplementary figure S1G). Using this antibody we found that C4 protein expression was  
195 increased approximately 1.38-fold in P30 mice in the C4-OE condition (Supplementary figure  
196 S1H). We observed no neuronal mispositioning in C4-OE mice, suggesting the absence of major  
197 migration defects (Supplementary figure S2).

198

199

200 **Loss of excitatory synapses in C4-OE neurons from juvenile and adult, but not neonatal**  
201 **mice**

202 We first evaluated passive and active cellular properties in tdTom+ layer II/III pyramidal cells  
203 using whole-cell patch-clamp *ex vivo*. The firing properties, membrane resistance, cell capacitance  
204 and resting membrane potential were similar in control and C4-OE neurons (Supplementary  
205 figure S3; Supplementary table S1). We next examined excitatory synaptic transmission at three  
206 developmental stages. The frequency of miniature excitatory postsynaptic currents (mEPSCs) in  
207 control tdTom+ cells markedly increased from early postnatal development (age P9-P10) to the  
208 juvenile stage (P25-P30), consistent with increased synapse generation, before decreasing from  
209 the juvenile stage to adulthood (P150-P220), in line with synapse elimination in the maturing  
210 cortex (Figure 1A-B; Supplementary table S1). The paired-pulse ratio of evoked EPSCs, a  
211 measure which is sensitive to alterations in presynaptic release probability, did not change  
212 significantly in control tdTom+ cells throughout development (Figure 1C-D; Supplementary table  
213 S1). Interestingly, mEPSC frequency was similar in control and C4-OE mice at P9-P10, but was  
214 significantly lower in C4-OE mice in juvenile and adult mice (Figure 1B; two-way ANOVA,  $P <$   
215  $0.0001$ ; juvenile: Sidak's post-hoc test  $P < 0.0001$ ; adulthood: Sidak's post-hoc test,  $P < 0.05$ ;  
216 Supplementary table S1). The paired-pulse ratio of evoked EPSCs did not differ between C4-OE  
217 and control mice in any age group (Figure 1D). We also observed a decrease in mEPSC amplitude  
218 from early development to the juvenile stage, but this was comparable in control and C4-OE mice  
219 (Figure 1B; Supplementary table S1).

220

221 **C4 overexpression results in diminished formation and elimination of spines in juvenile**  
222 **PFC**

223 To establish whether decreased functional excitatory input in C4-OE pyramidal cells  
224 correlates with reduced spine density, we imaged dendritic spines of tdTom+ pyramidal cells  
225 in layer I/II of the prefrontal cortex using *in vivo* 2-photon imaging in juvenile (P30) mice  
226 (Figure 1E). We found a substantial spine density decrease in C4-OE neurons (Figure 1F-G;  
227 T-test,  $P = 0.0023$ ; Supplementary table S2). These results suggested that the reduction in  
228 spine density in C4-OE neurons is due to imbalanced spine dynamics (gain and loss)  
229 occurring prior to P30. Therefore, we performed dynamic imaging studies in mice aged P15-  
230 P18, which corresponds to a period of heightened structural synaptic plasticity<sup>15-17</sup>. Here, we  
231 imaged dendrites twice at a 4-hour interval (Figure 2A) to study fast spine kinetics at this  
232 stage of active synaptic reorganization. Similar to the P30 cohort, C4-OE neurons in P15-P18  
233 mice displayed a robust decrease in spine density (Figure 2B-C; Mann Whitney Test,  $P <$



234 0.0001; Supplementary table S2). While the overall spine density within the experimental  
235 groups did not change over the course of 4 h (Figure 2D-E; table S2), C4 overexpression had  
236 a strong effect on the structural plasticity of single spines; first, C4-OE neurons formed fewer  
237 stable spines (Figure 2F; T-test,  $P = 0.0007$ ; Supplementary table S2). Second, C4-OE  
238 neurons showed decreased numbers of both gained and lost spines (Figure 2G-H; gained: T-  
239 test,  $P = 0.0018$ ; lost: T-test,  $P < 0.0001$ ; also see Figure 2F; Supplementary table S2)  
240 resulting in an overall significantly reduced turnover of dendritic spines (Figure 2I; T-test,  $P =$   
241  $0.0039$ ; Supplementary table S2). In summary we show here that C4 overexpression results in  
242 diminished formation/stabilization and elimination of spines in juvenile PFC *in vivo*. To study  
243 dendritic spines with improved spatial resolution, we imaged fixed post-mortem samples from  
244 P15-P18 control and C4-OE mice using confocal microscopy with an Airyscan detector, and  
245 obtained morphological reconstructions of dendritic spines using the Imaris software. These  
246 data confirmed the significant decrease in spine density in C4-OE neurons (Supplementary  
247 figure S4A-B; T-test,  $P = 0.023$ ). We plotted key morphological parameters (dendritic spine  
248 length, maximum head diameter, ratio of mean head to neck diameters) in three dimensions.  
249 The 3D plot revealed differences in the distributions of the morphological parameters  
250 (Supplementary figure S4C). A cluster analysis<sup>18</sup> ('Agglomerative Hierarchical Clustering'  
251 analysis based on Euclidian distance in the parameter space, see Supplementary Materials and  
252 methods) revealed at least three distinct populations of spines, which resemble the different  
253 categories commonly used in the literature to classify dendritic spines ('small'  $\cong$  cluster 1,  
254 'thin'  $\cong$  cluster 2 and 'mushroom-like'  $\cong$  cluster 3; Supplementary figure S4D-F). The  
255 distribution of spines in the three clusters was similar in the control and C4-OE conditions  
256 (Supplementary figure S4E-F).

257

### 258 **NMDAR hypofunction in C4-OE pyramidal cells**

259 To investigate NMDAR-mediated transmission in the cortex of C4-OE mice, we measured the  
260 AMPA/NMDA ratio of evoked EPSCs in tdTom+ cells. The AMPA/NMDA ratio was  
261 comparable in control and C4-OE neonates, but was significantly increased in juvenile and  
262 adult C4-OE mice (Figure 3A-B; Two-way ANOVA,  $P < 0.0001$ ; juvenile and adult mice:  
263 Sidak's test,  $P < 0.001$ ; Supplementary table S1). The amplitude of mEPSCs, which is  
264 determined by the number of AMPAR per synapse, was not altered by C4 overexpression at  
265 any age examined (Figure 1B; Supplementary table S1). Furthermore, the decay time constant  
266 of NMDAR-mediated currents, which depends on the subunit composition of NMDA  
267 receptors, was similar in control and C4-OE mice of the same age group (Figure 3C;

268 Supplementary table S1). Therefore the increased AMPA/NMDA ratio in C4-OE neurons  
269 results from reduced NMDAR-mediated currents. Increased AMPA/NMDA ratio has been  
270 repeatedly associated with altered synaptic expression of the AMPAR GluA2 subunit<sup>19, 20</sup> that  
271 controls the calcium permeability of AMPARs<sup>21, 22</sup>. Therefore we assessed the rectification of  
272 synaptic AMPAR-mediated currents which is sensitive to the presence of the GluA2 subunit.  
273 We found that the rectification index of evoked AMPAR currents was significantly decreased  
274 in juvenile and adult C4-OE mice, but not in pups (Figure 3D-E; two-way ANOVA,  $P <$   
275  $0.0001$ ; juvenile: Sidak's test,  $P < 0.001$ ; adulthood: Sidak's test,  $P < 0.05$ ; Supplementary  
276 table S1). This suggests that synaptic AMPARs in C4-OE neurons show reduced GluA2  
277 subunit content, thus becoming more calcium-permeable, concomitantly with NMDAR  
278 hypofunction.

279

#### 280 **C4 overexpression results in reduced GABAergic transmission**

281 Abnormalities in GABAergic networks constitute another hallmark of the schizophrenic  
282 cortex. They include synaptic abnormalities as well as decreased expression of GAD67 in  
283 parvalbumin-expressing (PV) interneurons. We examined spontaneous inhibitory synaptic  
284 transmission in tdTom+ pyramidal cells from juvenile mice using a high chloride intracellular  
285 solution to maximize the signal/noise ratio. Both the frequency and amplitude of miniature  
286 inhibitory postsynaptic currents (mIPSCs) were significantly decreased in C4-OE neurons  
287 (Figure 4A-B; frequency: Mann-Whitney test,  $P = 0.0004$ ; amplitude: T-test,  $P = 0.0003$ ;  
288 Supplementary table S1). Reduced mIPSC frequency may result from decreased release  
289 probability from GABAergic terminals in the maternal immune activation (MIA) model of  
290 SZ<sup>23</sup> which has been associated with increased expression of complement C1q and C4<sup>24, 25</sup>.  
291 We investigated GABA release probability using a similar protocol<sup>23</sup>. IPSCs were evoked by  
292 four consecutive electrical stimuli delivered at 20 Hz while holding voltage-clamped tdTom+  
293 pyramidal cells at a membrane potential of +10 mV (Figure 4C). Short-term synaptic  
294 depression, which is sensitive to changes in release probability, showed a non-significant  
295 trend towards reduction in C4-OE neurons (Figure 4D; RM two-way ANOVA,  $P = 0.089$ ;  
296 Supplementary table S1). In the MIA model, altered release probability from PV neurons, but  
297 not from other types of interneurons, has been inferred from optogenetic studies of short-term  
298 plasticity<sup>23</sup>. We reasoned that selectively activating the PV neuron-pyramidal cells synapse in  
299 C4-OE mice might allow to uncover significant differences in short-term depression. To test  
300 this hypothesis, we expressed Channelrhodopsin2 (ChR2) fused to YFP in PV-expressing  
301 interneurons using timed matings between lox-STOP-lox ChR2 mice and PV-Cre mice

302 (Supplementary figure S5A). Pregnant females were electroporated at embryonic stage E15  
303 either with a tdTomato-encoding plasmid (control group) or with tdTomato + C4 (C4-OE  
304 condition). To verify the specificity of ChR2 expression, we patched-clamped ChR2-YFP+  
305 cells in the control group and recorded their firing pattern in response to a 500 ms pulse of  
306 blue light (470 nm). All tested ChR2-YFP cells exhibited the characteristic fast-spiking firing  
307 pattern of PV interneurons (n = 3 cells; Supplementary figure S5B). Light-evoked PSCs in  
308 pyramidal cells were blocked by GABA<sub>A</sub>zine, confirming their GABAergic nature  
309 (Supplementary figure S5C-D). Next we assessed short-term depression of PV neuron IPSCs  
310 in pyramidal cells using four consecutive blue light pulses (0.5 ms) delivered at 20 Hz (Figure  
311 4E). We found that PV neuron-pyramidal cell synapses in C4-OE mice displayed significantly  
312 reduced short-term depression, consistent with decreased GABA release probability (Figure  
313 4F; RM two-way ANOVA, main effect,  $P = 0.014$ ; Supplementary table S1). Thus, similar to  
314 the MIA model, decreased release probability from PV neurons may contribute to diminish  
315 GABAergic transmission onto pyramidal cells in C4-OE mice.

316

### 317 **C4 overexpression in pyramidal cells causes reduced input resistance, decreased** 318 **excitability, and lower GAD67 expression in PV neurons**

319 The availability of presynaptic GABA is one of the factors determining release probability.  
320 GAD67 is an enzyme responsible for basal cortical GABA synthesis<sup>26</sup>, and reduced  
321 expression of GAD67, especially in PV cells, has been found in post-mortem cortical tissue  
322 from schizophrenic patients in multiple studies<sup>14, 27</sup>. In mouse models, reducing GAD67  
323 expression in PV neurons decreases both the frequency of spontaneous inhibitory currents<sup>28</sup>  
324 and the amplitude of evoked GABAergic currents in pyramidal cells<sup>29</sup>. Therefore we  
325 measured GAD67 expression in PV interneurons located in the vicinity of control or C4-OE  
326 tdTom+ pyramidal cells. As GAD67 is expressed in both presynaptic terminals and in the cell  
327 soma, we quantified GAD67 expression in the clearly delineated cell soma (Figure 4G)<sup>23</sup>.  
328 GAD67 expression in PV cells was significantly lower in C4-OE mice than in control mice  
329 (Mean intensity compared to control, T-test,  $P = 0.0094$ ; Figure 4H; Supplementary table S2).  
330 These results suggest that reduced GABA synthesis may contribute to the deficit in  
331 GABAergic transmission at PV neuron – pyramidal cell synapses in C4-OE mice. It has been  
332 proposed that reduced expression of activity-dependent biomarkers, including GAD67, in PV  
333 neurons, may result from decreased PV neuron activity in the schizophrenic cortex<sup>14</sup>. We  
334 investigated the number and distribution of PV neurons in the cortex of C4-OE juvenile mice  
335 and found that they were not altered (Supplementary figure S6; Supplementary table S2). We

336 then electroporated C4 + tdTomato or tdTomato alone in PV-Cre::RCE mice which express  
337 EGFP selectively in PV neurons. We patch-clamped EGFP+ PV neurons located in the  
338 vicinity of tdTom+ pyramidal cells, in control and C4-OE mice aged P25-P30 (Figure 4I).  
339 The intrinsic excitability of PV neurons, i.e. their propensity to fire action potentials when  
340 subjected to an input current in the presence of drugs blocking synaptic transmission, was  
341 significantly diminished in the C4-OE condition (Figure 4J; RM two-way ANOVA, main  
342 effect  $P = 0.013$ ; Supplementary table S3). This was associated with lower input resistance of  
343 PV cells (Figure 4K; Mann-Whitney test,  $P = 0.0024$ ; Supplementary table S3), in the  
344 absence of significant changes in cell capacitance or resting membrane potential  
345 (Supplementary figure S7A-B; Supplementary table S3).

346

#### 347 **C4 overexpression results in increased excitatory input on PV neurons**

348 Since synaptic inputs also contribute to neuronal excitability, we recorded miniature  
349 excitatory and inhibitory inputs (mEPSCs and mIPSCs) in EGFP+ PV neurons. We observed  
350 a significant increase in mEPSC frequency, and a trend towards increased mEPSC amplitude,  
351 in the C4-OE condition (Supplementary figure S7C-D; frequency: Mann-Whitney test,  $P =$   
352  $0.018$ ; amplitude: Mann-Whitney test,  $P = 0.069$ ; Supplementary table S3). The paired-pulse  
353 ratio of evoked EPSCs in PV neurons remained unchanged (Supplementary figure S7E-F;  
354 table S3). Neither the frequency nor the amplitude of mIPSCs was changed following C4  
355 overexpression in pyramidal cells (Supplementary figure S7G-H; Supplementary table S3).  
356 Thus, in contrast to C4-OE pyramidal cells, PV neurons exhibit increased excitatory input in  
357 C4-OE mice.

358

#### 359 **Working memory deficits in C4-OE mice**

360 Finally we investigated whether the changes that we observed at the cellular level may result  
361 in long-lasting behavioral alterations in C4-OE mice. In SZ, deficits in working memory are  
362 central to cognitive impairments and appear to result, at least in part, from abnormalities in  
363 PFC function<sup>30, 31</sup>. Therefore we asked whether increased C4 expression in the PFC alters  
364 working memory. Since classical working memory tests rely on locomotion, we first verified  
365 that control and C4-OE mice showed comparable spontaneous locomotor activity in an  
366 actimeter (Figure 5A; Supplementary table S4). Spontaneous activity of C4-OE mice also did  
367 not differ from control mice in the open field assay (Figure 5B; Supplementary table S4). We  
368 then used two different tests of working memory, namely the odor span test and the spatial  
369 span test. C4-OE mice performed equally well as control mice during the acquisition of the

370 non-matching to sample task, in which mice learned to associate the presence of a new odor  
371 and food reward (Figure 5C-E; Supplementary table S4). In the subsequent span tests,  
372 however, when mice had to remember an increasing number of odors or positions to obtain  
373 the reward, C4-OE mice exhibited reduced working memory capacity compared to control  
374 mice (Figure 5F-I; odor span: Mann-Whitney test,  $P = 0.0022$ ; spatial span: Mann-Whitney  
375 test,  $P = 0.045$ ; Supplementary table S4). Taken together, these results indicate that working  
376 memory is impaired in C4-OE mice, in line with neuronal abnormalities in PFC networks.

377

## 378 **Discussion**

379 GWAS have provided unprecedented identification of predisposing genetic risk factors in SZ.  
380 High expression variants of the C4 gene contribute to the strong association between SZ and a  
381 genomic region within the MHC locus on Chromosome 6 in GWAS<sup>4, 5</sup>. Here we show that  
382 overexpressing C4 in the mouse PFC recapitulates several cortical phenotypes associated with  
383 SZ, including synapse loss, decreased NMDAR-mediated neurotransmission, impaired  
384 function of GABAergic circuits, and deficits in working memory (Supplementary figure S8).

385

### 386 **Decreased spine turnover in C4-OE mice**

387 Elevated expression of C4 in neurons enhances microglia-mediated synaptic engulfment<sup>10-12</sup>.  
388 Our *in vivo* imaging results clearly show an overall decrease in gained and lost dendritic  
389 spines in juvenile mice as a consequence of C4 overexpression. This apparent discrepancy  
390 may reflect an underappreciated role of complement in regulating spine formation or  
391 immature spine stabilization. The decrease in spine turnover observed in our model could be  
392 due to excessive microglial phagocytosis of the synapses in formation or just formed,  
393 preventing their stabilization. The disappearance of these spines in less than 4 hours, the time  
394 window used for our time-lapse 2-photon microscopy acquisitions, would thus result in  
395 decreased spine gain.

396

### 397 **Time course of synaptic deficits in C4-OE mice**

398 It is noteworthy that the decrease in excitatory input to C4-OE neurons is not present at the  
399 early stages of postnatal development, but only appears between P10 and P25, a period of  
400 strong synaptic remodeling in the cortex. This may be relevant for the pathophysiology of SZ,  
401 a disorder in which even the earliest prodromal symptoms occur at a relatively late stage of  
402 postnatal development. The reduction in NMDAR-mediated transmission is concomitant with  
403 the loss of functional excitatory synapses in C4-OE mice, suggesting that C4 overexpression

404 may selectively impair the formation or maintenance of “weak” (high NMDAR, low  
405 AMPAR) synapses in juvenile mice. Another study reported that synaptic deficits were  
406 accompanied by a decrease in cellular capacitance in C4-OE pyramidal cells<sup>11</sup>. We found that  
407 cell capacitance was unchanged following C4-OE. While we cannot exclude that different  
408 experimental settings account for this difference, the absence of significant changes in  
409 capacitance may be in line with the unchanged morphology of C4-OE pyramidal neurons<sup>11</sup>.  
410 We show that synaptic deficits persist into adulthood in C4-OE mice. Our  
411 immunohistochemistry results indicate that C4 is still overexpressed in the PFC of adult C4-  
412 OE mice, suggesting that increased C4 expression may persistently affect spine turnover in  
413 C4-OE mice. However, since synaptic refinement is particularly pronounced in the  
414 developing cortex<sup>16, 17</sup>, it is likely that elevated expression of C4 predominantly affects circuit  
415 remodeling in juvenile mice.

416

#### 417 **GABAergic network abnormalities**

418 While another study did not detect significant changes in the inhibitory input to C4-OE  
419 pyramidal cells<sup>11</sup>, we observed a significant reduction in mIPSC frequency and amplitude.  
420 This discrepancy may be due to the higher driving force for chloride ions at hyperpolarized  
421 membrane potentials, and hence improved signal/noise ratio of IPSC recordings, in our  
422 experimental conditions of high intracellular chloride. We also show that C4-OE in pyramidal  
423 cells decreases GABA release from PV neurons onto pyramidal cells, consistent with  
424 decreased GAD67 expression in PV neurons and reduced PV neuron excitability. The cause  
425 of the well-substantiated decrease in GAD67 cortical expression in SZ is unclear. It has been  
426 hypothesized that lower activity of layer II/III pyramidal cells, possibly due to their reduced  
427 number of dendritic spines, could result in fewer excitatory inputs onto PV cells (for instance  
428 through lower neuronal activity-regulated pentraxin secretion from pyramidal cells), leading  
429 to reduced PV neuron excitation and hence decreased activity-dependent GAD67  
430 expression<sup>14, 32</sup>. In C4-OE mice, reduced GAD67 expression in PV neurons may result from  
431 altered intrinsic properties of PV neurons as a consequence of homeostatic adaptation to  
432 reduced network activity, rather than from decreased synaptic excitatory input. Indeed we  
433 show that PV neuron excitability is impaired in C4-OE mice, while increased mEPSC  
434 frequency in PV neurons indicates that PV neurons are spared from excitatory synapse loss  
435 associated with C4 overexpression in pyramidal cells.

436

437

438

439 **Linking neural dysfunctions and cognitive impairment in C4-OE mice**

440 Cognitive dysfunction is a central feature of SZ that is not treated by current  
441 pharmacotherapy, and determines the functional outcome of the illness<sup>33</sup>. Impaired working  
442 memory has been proposed to be central in the cognitive impairments associated with SZ<sup>30,31</sup>.  
443 Spine loss, reduced NMDAR transmission and PV neuron dysfunction in the PFC may all  
444 contribute to impair working memory. For example, decreased cortical connectivity in SZ has  
445 repeatedly been associated with working memory deficits<sup>34</sup>. Decreased synapse density in  
446 layer II/III may particularly affect connectivity, since layer II/III pyramidal cells make major  
447 contributions to cortico-cortical connections, including callosal connections, and therefore  
448 integrate information across cortical areas and hemispheres. Furthermore, blocking NMDA,  
449 but not AMPA, receptors during a working memory task abolishes persistent activity in  
450 prefrontal neurons<sup>35</sup>. Finally, PV neurons contribute to the mechanisms underlying working  
451 memory<sup>36</sup>, in line with the well-established role of PV neurons in the generation of fast,  
452 gamma frequency oscillations in neocortex<sup>37, 38</sup>. Our results suggest that these distinct  
453 alterations occur together during development as a consequence of elevated C4 expression,  
454 making it difficult to attribute cognitive dysfunction to a single one of them. The combination  
455 of reduced connectivity, NMDAR hypofunction and altered PV neuron function might  
456 collectively modify the dynamics of specific attractors, which are thought to be the substrate  
457 for persistent neural activity underlying short-term memory<sup>39,40</sup>.

458

459 **Conclusion**

460 Decreased spine density, reduced NMDAR function, and alterations in GABAergic networks  
461 have been associated with SZ. We show that elevated C4 expression in PFC pyramidal cells  
462 results in functional alterations that are consistent with these SZ-associated phenotypes, and  
463 causes working memory impairment, a characteristic feature of cognitive deficits in SZ. In  
464 particular, we propose that the persistent decrease in NMDAR-mediated neurotransmission in  
465 C4-OE mice might be the consequence of impaired stabilization of immature synapses. This  
466 could provide a conceptual framework to relate the hypotheses of excessive synaptic pruning  
467 and NMDAR hypofunction in SZ.

468

469 **Conflict of interest**

470 The authors declare no conflict of interest.

471

472 **Acknowledgements**

473 MD was the recipient of fellowships from Sorbonne University and from FRM/Venite  
474 Cantemus (FDT20190400802). The project was supported by funds from the Investissements  
475 d'Avenir program under reference ANR-11-IDEX-0004-02 to CLM and RT, the Emergence  
476 program of Sorbonne University to CLM, and from an ERANET Neuron Grant to RT (ANR-  
477 18-0008-01), MF (BMBF 01EW1905) and CLM (ANR-18-0008-02). We thank Drs L  
478 Maroteaux, P Gaspar and J-C Poncer for critically reading the manuscript.

479

480 **References**

- 481 1. Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF *et al.* Common  
482 polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 2009;  
483 **460**(7256): 748-752.  
484
- 485 2. Shi J, Levinson DF, Duan J, Sanders AR, Zheng Y, Pe'er I *et al.* Common variants on  
486 chromosome 6p22.1 are associated with schizophrenia. *Nature* 2009; **460**(7256): 753-757.  
487
- 488 3. Stefansson H, Ophoff RA, Steinberg S, Andreassen OA, Cichon S, Rujescu D *et al.* Common  
489 variants conferring risk of schizophrenia. *Nature* 2009; **460**(7256): 744-747.  
490
- 491 4. Ripke S, Neale BM, Corvin A, al. e. Biological insights from 108 schizophrenia-associated  
492 genetic loci. *Nature* 2014; **511**(7510): 421-427.  
493
- 494 5. Sekar A, Bialas AR, de Rivera H, Davis A, Hammond TR, Kamitaki N *et al.* Schizophrenia risk  
495 from complex variation of complement component 4. *Nature* 2016; **530**(7589): 177-183.  
496
- 497 6. Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N *et al.* The classical  
498 complement cascade mediates CNS synapse elimination. *Cell* 2007; **131**(6): 1164-1178.  
499
- 500 7. Glantz LA, Lewis DA. Decreased dendritic spine density on prefrontal cortical pyramidal  
501 neurons in schizophrenia. *Arch Gen Psychiatry* 2000; **57**(1): 65-73.  
502
- 503 8. Feinberg I. Schizophrenia: caused by a fault in programmed synaptic elimination during  
504 adolescence? *Journal of psychiatric research* 1982; **17**(4): 319-334.  
505
- 506 9. Prasad KM, Chowdari KV, D'Aiuto LA, Iyengar S, Stanley JA, Nimgaonkar VL. Neuropil  
507 contraction in relation to Complement C4 gene copy numbers in independent cohorts of  
508 adolescent-onset and young adult-onset schizophrenia patients-a pilot study. *Translational*  
509 *psychiatry* 2018; **8**(1): 134.  
510
- 511 10. Sellgren CM, Gracias J, Watmuff B, Biag JD, Thanos JM, Whittredge PB *et al.* Increased  
512 synapse elimination by microglia in schizophrenia patient-derived models of synaptic  
513 pruning. *Nat Neurosci* 2019; **22**(3): 374-385.  
514



- 515 11. Comer AL, Jinadasa T, Sriram B, Phadke RA, Kretsge LN, Nguyen TPH *et al.* Increased  
516 expression of schizophrenia-associated gene C4 leads to hypoconnectivity of prefrontal  
517 cortex and reduced social interaction. *PLoS Biol* 2020; **18**(1): e3000604.  
518
- 519 12. Yilmaz M, Yalcin E, Presumey J, Aw E, Ma M, Whelan CW *et al.* Overexpression of  
520 schizophrenia susceptibility factor human complement C4A promotes excessive synaptic loss  
521 and behavioral changes in mice. *Nat Neurosci* 2020.  
522
- 523 13. Balu DT. The NMDA Receptor and Schizophrenia: From Pathophysiology to Treatment. *Adv*  
524 *Pharmacol* 2016; **76**: 351-382.  
525
- 526 14. Dienel SJ, Lewis DA. Alterations in cortical interneurons and cognitive function in  
527 schizophrenia. *Neurobiol Dis* 2019; **131**: 104208.  
528
- 529 15. Coiro P, Padmashri R, Suresh A, Spartz E, Pendyala G, Chou S *et al.* Impaired synaptic  
530 development in a maternal immune activation mouse model of neurodevelopmental  
531 disorders. *Brain Behav Immun* 2015; **50**: 249-258.  
532
- 533 16. Zuo Y, Lin A, Chang P, Gan WB. Development of long-term dendritic spine stability in diverse  
534 regions of cerebral cortex. *Neuron* 2005; **46**(2): 181-189.  
535
- 536 17. Holtmaat AJ, Trachtenberg JT, Wilbrecht L, Shepherd GM, Zhang X, Knott GW *et al.* Transient  
537 and persistent dendritic spines in the neocortex in vivo. *Neuron* 2005; **45**(2): 279-291.  
538
- 539 18. Pfeiffer T, Poll S, Bancelin S, Angibaud J, Inavalli VK, Keppler K *et al.* Chronic 2P-STED imaging  
540 reveals high turnover of dendritic spines in the hippocampus in vivo. *eLife* 2018; **7**.  
541
- 542 19. Bellone C, Luscher C. Cocaine triggered AMPA receptor redistribution is reversed in vivo by  
543 mGluR-dependent long-term depression. *Nat Neurosci* 2006; **9**(5): 636-641.  
544
- 545 20. Wang HX, Gao WJ. Development of calcium-permeable AMPA receptors and their correlation  
546 with NMDA receptors in fast-spiking interneurons of rat prefrontal cortex. *J Physiol* 2010;  
547 **588**(Pt 15): 2823-2838.  
548
- 549 21. Hollmann M, Hartley M, Heinemann S. Ca<sup>2+</sup> permeability of KA-AMPA-gated glutamate  
550 receptor channels depends on subunit composition. *Science* 1991; **252**(5007): 851-853.  
551
- 552 22. Geiger JR, Melcher T, Koh DS, Sakmann B, Seeburg PH, Jonas P *et al.* Relative abundance of  
553 subunit mRNAs determines gating and Ca<sup>2+</sup> permeability of AMPA receptors in principal  
554 neurons and interneurons in rat CNS. *Neuron* 1995; **15**(1): 193-204.  
555
- 556 23. Canetta S, Bolkan S, Padilla-Coreano N, Song LJ, Sahn R, Harrison NL *et al.* Maternal immune  
557 activation leads to selective functional deficits in offspring parvalbumin interneurons. *Mol*  
558 *Psychiatry* 2016; **21**(7): 956-968.  
559
- 560 24. Duchatel RJ, Meehan CL, Harms LR, Michie PT, Bigland MJ, Smith DW *et al.* Increased  
561 complement component 4 (C4) gene expression in the cingulate cortex of rats exposed to  
562 late gestation immune activation. *Schizophr Res* 2018; **199**: 442-444.  
563
- 564 25. Han M, Zhang JC, Hashimoto K. Increased Levels of C1q in the Prefrontal Cortex of Adult  
565 Offspring after Maternal Immune Activation: Prevention by 7,8-Dihydroxyflavone. *Clinical*

566 *psychopharmacology and neuroscience : the official scientific journal of the Korean College of*  
567 *Neuropsychopharmacology* 2017; **15**(1): 64-67.  
568

569 26. Asada H, Kawamura Y, Maruyama K, Kume H, Ding RG, Kanbara N *et al.* Cleft palate and  
570 decreased brain gamma-aminobutyric acid in mice lacking the 67-kDa isoform of glutamic  
571 acid decarboxylase. *Proc Natl Acad Sci U S A* 1997; **94**(12): 6496-6499.  
572

573 27. Hashimoto T, Volk DW, Eggan SM, Mirnics K, Pierri JN, Sun Z *et al.* Gene expression deficits in  
574 a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J*  
575 *Neurosci* 2003; **23**(15): 6315-6326.  
576

577 28. Brown JA, Ramikie TS, Schmidt MJ, Baldi R, Garbett K, Everheart MG *et al.* Inhibition of  
578 parvalbumin-expressing interneurons results in complex behavioral changes. *Mol Psychiatry*  
579 2015; **20**(12): 1499-1507.  
580

581 29. Fujihara K, Miwa H, Kakizaki T, Kaneko R, Mikuni M, Tanahira C *et al.* Glutamate  
582 Decarboxylase 67 Deficiency in a Subset of GABAergic Neurons Induces Schizophrenia-  
583 Related Phenotypes. *Neuropsychopharmacology* 2015; **40**(10): 2475-2486.  
584

585 30. Silver H, Feldman P, Bilker W, Gur RC. Working memory deficit as a core neuropsychological  
586 dysfunction in schizophrenia. *Am J Psychiatry* 2003; **160**(10): 1809-1816.  
587

588 31. Silver H, Feldman P. Evidence for sustained attention and working memory in schizophrenia  
589 sharing a common mechanism. *The Journal of neuropsychiatry and clinical neurosciences*  
590 2005; **17**(3): 391-398.  
591

592 32. Hoftman GD, Datta D, Lewis DA. Layer 3 Excitatory and Inhibitory Circuitry in the Prefrontal  
593 Cortex: Developmental Trajectories and Alterations in Schizophrenia. *Biol Psychiatry* 2017;  
594 **81**(10): 862-873.  
595

596 33. Kahn RS, Keefe RS. Schizophrenia is a cognitive illness: time for a change in focus. *JAMA*  
597 *psychiatry* 2013; **70**(10): 1107-1112.  
598

599 34. Cao H, Dixson L, Meyer-Lindenberg A, Tost H. Functional connectivity measures as  
600 schizophrenia intermediate phenotypes: advances, limitations, and future directions. *Curr*  
601 *Opin Neurobiol* 2016; **36**: 7-14.  
602

603 35. Wang M, Yang Y, Wang CJ, Gamo NJ, Jin LE, Mazer JA *et al.* NMDA receptors subserve  
604 persistent neuronal firing during working memory in dorsolateral prefrontal cortex. *Neuron*  
605 2013; **77**(4): 736-749.  
606

607 36. Rossi MA, Hayrapetyan VY, Maimon B, Mak K, Je HS, Yin HH. Prefrontal cortical mechanisms  
608 underlying delayed alternation in mice. *J Neurophysiol* 2012; **108**(4): 1211-1222.  
609

610 37. Cardin JA, Carlen M, Meletis K, Knoblich U, Zhang F, Deisseroth K *et al.* Driving fast-spiking  
611 cells induces gamma rhythm and controls sensory responses. *Nature* 2009; **459**(7247): 663-  
612 667.  
613

614 38. Uhlhaas PJ, Singer W. Abnormal neural oscillations and synchrony in schizophrenia. *Nat Rev*  
615 *Neurosci* 2010; **11**(2): 100-113.  
616

- 617 39. Brody CD, Romo R, Kepecs A. Basic mechanisms for graded persistent activity: discrete  
618 attractors, continuous attractors, and dynamic representations. *Curr Opin Neurobiol* 2003;  
619 **13**(2): 204-211.  
620
- 621 40. Inagaki HK, Fontolan L, Romani S, Svoboda K. Discrete attractor dynamics underlies  
622 persistent activity in the frontal cortex. *Nature* 2019; **566**(7743): 212-217.  
623  
624

## 625 **Figure legends**

### 626 **Figure 1. Reduced excitatory input onto C4-OE pyramidal cells**

627 **(A)** Sample traces of mEPSCs in control and C4-OE pyramidal cells at three developmental  
628 stages; scale bars: 20pA/100ms.

629 **(B)** mEPSC frequency was affected by C4-OE in juvenile and adult mice, but not in pups  
630 (two-way ANOVA, main effect, \*\*\*\*  $P < 0.0001$ ; Sidak's post-hoc test #####  $P < 0.0001$ , #  $P$   
631  $< 0.05$ , n.s. not significant). mEPSC amplitude was not significantly affected (control: early  
632 postnatal,  $n = 21$  cells from 3 mice; juvenile,  $n = 16$  cells from 5 mice; adult,  $n = 20$  cells from  
633 4 mice; C4-OE: early postnatal,  $n = 21$  cells from 3 mice; juvenile,  $n = 16$  cells from 5 mice;  
634 adult,  $n = 20$  cells from 4 mice).

635 **(C)** Average traces of eEPSCs evoked by paired-pulse stimulation with 50 ms interpulse  
636 interval, recorded from control and C4-OE pyramidal cells at three developmental stages;  
637 scale bars: 50 pA/50 ms.

638 **(D)** Similar paired-pulse ratio in control and C4-OE pyramidal cells (n.s., not significant.  
639 control: early postnatal,  $n = 16$  cells from 3 mice; juvenile,  $n = 17$  cells from 3 mice; adult,  $n$   
640  $= 19$  cells from 5 mice; C4-OE: early postnatal,  $n = 21$  cells from 3 mice; juvenile,  $n = 21$   
641 cells from 3 mice; adult,  $n = 17$  cells from 4 mice). All data are represented as mean  $\pm$  SEM.  
642 Each open circle represents an individual cell.

643 **(E)** Left, schematic drawing illustrating two-photon *in vivo* imaging in PFC at P30. Right,  
644 cranial window.

645 **(F)** Examples of apical dendrites from electroporated pyramidal cells in control and C4-OE  
646 mice aged P30.

647 **(G)** C4-OE decreased dendritic spine density in electroporated pyramidal cells from P30  
648 mice; scale bar: 2  $\mu\text{m}$ . (T-test, \*\*  $P < 0.023$ ; control:  $n = 10$  mice; C4-OE:  $n = 7$  mice). All  
649 data are represented as mean  $\pm$  SEM. Each open circle represents an individual mouse.

650

651 **Figure 2. Decreased spine turnover in young (P15-P18) C4-OE mice**

652 (A) Experimental timeline of *in vivo* two-photon imaging to investigate spine formation and  
653 elimination in electroporated pyramidal neurons. Spines already present at  $t_0$  are in blue;  
654 spines that were stable, gained or lost at  $t_0 + 4h$  are in blue, green and magenta, respectively.

655 (B) Examples of apical dendrites from control and C4-OE electroporated pyramidal neurons  
656 imaged at  $t_0$ ; scale bar: 2  $\mu m$ .

657 (C) Spine density at  $t_0$  in tdTom+ pyramidal neurons (t-test, \*\*\*\*  $P < 0.0001$ ; control: n = 5  
658 mice; C4-OE: n = 7 mice).

659 (D) Example images showing spine turnover from  $t_0$  to  $t_0 + 4h$  in control and C4-OE  
660 pyramidal cells; scale bar: 2  $\mu m$ .

661 (E) Evolution of the spine density between  $t_0$  and  $t_0 + 4h$  in control and control and C4-OE  
662 pyramidal cells (control: n = 5 mice; C4-OE: n = 7 mice).

663 (F-H) Stable, gained and lost spine density in control and C4-OE neurons. C4-OE caused a  
664 significant reduction in the number of both gained and lost spines (stable spine: T-test, \*\*\*  $P$   
665 = 0.0007; gained spines: t-test, \*\*  $P = 0.0018$ ; lost spines: T-test, \*\*\*\*  $P < 0.0001$ ; control: n  
666 = 5 mice; C4-OE: n = 7 mice).

667 (I) Spine turnover in control and C4-OE dendrites. The turnover rate is decreased in C4-OE  
668 dendrites (Mann-Whitney test, \*\*  $P = 0.0039$ ; control: n = 5 mice; C4-OE: n = 7 mice).

669 All data are represented as mean  $\pm$  SEM. Each open circle represents an individual mouse.

670

671 **Figure 3. Altered AMPA/NMDA ratio and AMPAR rectification index in C4-OE mice**

672 (A) Sample traces of evoked synaptic AMPA and NMDA currents recorded at -65 mV and  
673 +55 mV respectively in the presence of GABAazine at 3 developmental stages; The NMDAR  
674 component was measured 50 ms after stimulation (arrows). The AMPAR component was  
675 measured as the peak of the evoked EPSC at -65mV. Scale bars: 50pA/50ms.

676 (B-C) AMPA/NMDA ratio and NMDA decay time throughout development in control and  
677 C4-OE pyramidal cells. The AMPA/NMDA ratio was increased in C4-OE neurons from  
678 juvenile and adult mice, but not from neonates. The decay time of NMDARs was not affected  
679 by C4-OE at any stage of development, indicating that C4-OE does not alter NMDAR subunit

680 composition. This together with unchanged mEPSC amplitude in C4-OE mice (Fig.2B)  
681 indicates that the increased AMPA/NMDA ratio reflects reduced NMDAR-mediated  
682 excitatory transmission in C4-OE pyramidal neurons (two-way ANOVA, main effect, \*\*\*\*  $P$   
683  $< 0.0001$ , Sidak's post-hoc test ###  $P < 0.001$ ; n.s., not significant. Control: early postnatal,  $n$   
684  $= 17$  cells from 3 mice; juvenile,  $n = 23$  cells from 4 mice; adult,  $n = 17$  cells from 4 mice;  
685 C4-OE: early postnatal,  $n = 21$  cells from 3 mice; juvenile,  $n = 20$  cells from 5 mice; adult,  $n$   
686  $= 16$  cells from 4 mice).

687 **(D)** Sample traces of evoked AMPA-mediated currents recorded at  $-65\text{mV}$  and  $+55\text{mV}$  at 3  
688 developmental stages; scale bars:  $50\text{ pA}/50\text{ ms}$ .

689 **(E)** The rectification index of AMPR-mediated currents was decreased in pyramidal cells  
690 from juvenile and adult C4-OE mice, suggesting the insertion of GluA2 subunit-lacking  
691 AMPAR at excitatory synapses (two-way ANOVA, \*\*\*\*  $P < 0.0001$ , Sidak's post-hoc test ##  
692  $P < 0.01$ , #  $P < 0.05$ . Control: early postnatal,  $n = 20$  cells from 3 mice; juvenile,  $n = 21$  cells  
693 from 3 mice; adult,  $n = 17$  cells from 4 mice; C4-OE: early postnatal,  $n = 20$  cells from 3  
694 mice; juvenile,  $n = 16$  cells from 3 mice; adult,  $n = 24$  cells from 3 mice). All data are  
695 represented as mean  $\pm$  SEM. Each open circle represents an individual cell.

696

#### 697 **Figure 4. Alterations of GABAergic networks in C4-OE mice**

698 **(A)** Sample traces of mIPSCs recordings at a holding potential of  $-65\text{ mV}$  in control and C4-  
699 OE pyramidal cells from juvenile mice (P25-P30); scale bars:  $20\text{pA}/100\text{ms}$ .

700 **(B)** Both mIPSC frequency and amplitude were decreased in C4-OE pyramidal cells  
701 (frequency: Mann-Whitney test \*\*\*  $P = 0.0004$ ; amplitude: t-Test \*\*\*  $P = 0.0003$ . Control:  $n$   
702  $= 16$  cells from 4 mice; C4-OE:  $n = 16$  cells from 4 mice).

703 **(C)** Schematic drawing illustrating the electrical stimulation of GABAergic release onto  
704 electroporated pyramidal cells (left) and example traces of 4 consecutive eIPSCs evoked with  
705  $50\text{ms}$  interpulse intervals, in control and C4-OE pyramidal cells (right); scale bars:  $50\text{ pA}/50$   
706  $\text{ms}$ .

707 **(D)** Ratios of the second, third and fourth eIPSC peak amplitude to the first eIPSC peak  
708 amplitude. Paired-pulse depression exhibited a trend towards being decreased in C4-OE  
709 neurons (two-way RM ANOVA, main effect,  $P = 0.089$ . Control:  $n = 29$  cells from 3 mice;  
710 C4-OE:  $n = 37$  cells from 3 mice).

711 (E) Schematic drawing illustrating the optogenetic activation of PV neuron and resulting  
712 IPSC recorded in electroporated pyramidal cells (left); and example traces of 4 consecutive  
713 light-evoked IPSCs with 50 ms interpulse intervals in control and C4-OE pyramidal cells  
714 (right); scale bars: 50 pA/50 ms.

715 (F) Ratios of the second, third and fourth light-evoked IPSC peak amplitude to the first IPSC  
716 peak amplitude. The increased ratio in C4-OE pyramidal cells indicates reduced presynaptic  
717 GABA release at PV neuron-pyramidal cell synapses (two-way RM ANOVA, main effect, \*  
718  $P = 0.014$ , Sidak's post-hoc test #  $P < 0.05$ . Control:  $n = 43$  cells from 4 mice; C4-OE:  $n = 30$   
719 cells from 3 mice). All data are represented as mean  $\pm$  SEM. Each open circle represents an  
720 individual cell.

721 (G) Example of immunostainings for tdTomato/PV and GAD67 on PFC slices from control  
722 and C4-OE mice. Arrows indicate GAD67-expressing PV cells; scale bar: 20 $\mu$ m.

723 (H) Mean intensity  $\pm$  SEM of GAD67 expression in PV+ cells. GAD67 expression in PV  
724 cells was decreased in C4-OE mice (T-test; \*\*  $P = 0.0094$ ; control:  $n = 28$  mice; C4-OE: 33  
725 mice). Each open circles represents an individual mouse.

726 (I) Schematic drawing depicting the patch-clamp recording of a PV interneuron located in the  
727 vicinity of tdTomato pyramidal cells (left) and sample spike trains in PV interneurons evoked  
728 by a 250 pA somatic current injection in control and C4-OE mice in the presence of synaptic  
729 blockers (right); scale bars: 20 mV/100 ms.

730 (J) PV interneurons exhibited reduced intrinsic excitability in C4-OE mice (two-way RM  
731 ANOVA, main effect, \*  $P = 0.013$ . Control:  $n = 20$  cells from 3 mice; C4-OE:  $n = 21$  cells  
732 from 4 mice).

733 (K) The input resistance of PV neurons was significantly decreased in C4-OE mice (Mann-  
734 Whitney test, \*\*  $P = 0.0024$ . Control:  $n = 20$  cells from 3 mice; C4-OE:  $n = 21$  cells from 4  
735 mice). Each open circle represents an individual cell.

736

### 737 **Figure 5. Impaired working memory in C4-OE mice**

738 (A) Locomotor activity measured at weaning in an actimeter, in control ( $n = 32$  mice) and  
739 C4-OE male mice ( $n = 21$  mice). Each dot represents spontaneous activity averaged over a  
740 period of 5 minutes.

741 **(B)** Locomotor activity measured one week after weaning in the open field, in control (n = 20  
742 mice) and C4-OE mice (n = 14 mice).

743 **(C)** Time line of working memory tests. Animals were first trained to dig into scented sawdust  
744 to find a reward pellet (shaping) before learning a non-matching to sample (NMTS) task,  
745 followed by 4 days of odour span test, and 4 days of spatial span test.

746 **(D)** NMTS test procedure. Mice were exposed to pairs of odors and learned to associate a new  
747 odor with a reward.

748 **(E)** Performance in the NMTS odor test was similar in control and C4-OE mice (Control: n =  
749 15 mice; C4-OE: 14 mice).

750 **(F)** Mice were exposed to an increasing number of odors in the odor span test.


751 **(G)** Working memory capacity in the odor span test is impaired in C4-OE mice (Control: n =  
752 15 mice; C4-OE: 14 mice; Mann-Whitney test, \*\*  $P = 0.0022$ ).

753 **(H)** Mice were exposed to an increasing number of positions in the spatial span test.










754 **(I)** Working memory capacity in the spatial span test is also affected in C4-OE mice (Control:  
755 n = 15 mice; C4-OE: 14 mice; Mann-Whitney test, \*  $P = 0.0448$ ).






756 All data are represented as mean  $\pm$  SEM. Each open circle represents an individual mouse.


757














**A****B****C****D****E****F****G****H****I**





**A****B****C****D****E****F****G****H****I**

## Supplementary Information

### KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<b>Antibodies</b>		
Rat monoclonal anti-HA	Sigma	Cat# 11867431001; RRID: AB_390919
Rabbit anti-Parvalbumin	Swant	Cat# PV27; RRID: AB_2631173
Mouse monoclonal anti-Glutamate Decarboxylase 67	Millipore-Merk	Cat# G5419; RRID: AB_261978
Goat anti-rat Alexa Fluor 488	Invitrogen	Cat# A-11006; RRID: AB_2534074
Donkey anti-rabbit Cy3	Jackson ImmunoResearch	Cat# 711-165-152; RRID: AB_2307443
Goat anti-mouse Alexa Fluor 647	Jackson ImmunoResearch	Cat# 115-605-003; RRID: AB_2338902
Chicken anti-GAPDH	Millipore-Merk	Cat# AB2302; RRID: AB_10615768
Rabbit anti-C4 alpha Chain	Proteintech	Cat# 22233-1-AP; RRID: AB_2879042
Goat anti-chicken IR700	Advansta	Cat# R-05057-250; RRID: N/A
Goat anti-rat IR800	Advansta	Cat# R-05064-250; RRID: N/A
Goat anti-rabbit IR700	Invitrogen	Cat# 35568; RRID: AB_2633284
<b>DNA and Plasmids</b>		
C4b mouse sequence	NCBI Resources	NM_009780.2
pCAG-HAC4b	This paper	N/A
pCAG-TdTomato	This paper	N/A
<b>Experimental Models: Organisms/Strains</b>		
Mouse: C57Bl6/J	Janvier Labs	C57BL/6JRj
Mouse: PV-Cre (B6;129P2-Pvalb <sup>tm1(cre)Arbr/J</sup> )	The Jackson Laboratory	JAX: 017320
Mouse: ChR2-YFP-loxP (B6;129S-Gt(ROSA) <sup>26Sortm32(CAG-COP4*H134R/EYFP)Hze/J</sup> )	The Jackson Laboratory	JAX: 012569
Mouse: RCE-loxP (Gt(ROSA) <sup>26Sortm1.1(CAG-EGFP)Fsh/Mmjax</sup> )	The Jackson Laboratory	JAX : 032037
<b>Software and Algorithms</b>		
GraphPad Prism	<a href="https://www.graphpad.com/">https://www.graphpad.com/</a>	RRID: SCR_002798
ImageJ	<a href="https://imagej.nih.gov/ij/">https://imagej.nih.gov/ij/</a>	RRID: SCR_003070
MiniAnalysis	<a href="http://www.synaptosoft.com/MiniAnalysis/">http://www.synaptosoft.com/MiniAnalysis/</a>	RRID: SCR_002184
IgorPro	<a href="https://www.wavemetrics.com/">https://www.wavemetrics.com/</a>	RRID: SCR_000325
PatchMaster	<a href="http://www.heka.com/products/products_main.html#soft_pm">http://www.heka.com/products/products_main.html#soft_pm</a>	RRID:SCR_000034
NeuroMatic	<a href="http://www.neuromatic.thinkrandom.com/">http://www.neuromatic.thinkrandom.com/</a>	RRID:SCR_004186
Imetronic	<a href="http://www.imetronic.com/">http://www.imetronic.com/</a>	N/A
Viewpoint	<a href="https://viewpoint.fr/en/p/software/videotrack">https://viewpoint.fr/en/p/software/videotrack</a>	N/A
Image Studio Lite	<a href="https://www.licor.com/bio/image-studio-lite/">https://www.licor.com/bio/image-studio-lite/</a>	RRID: SCR_013715

## Materials and methods

### Animals

All animal experiments were performed according to guidelines of the European Community for the use of animals in research and were approved by the local ethical committees (C2EA-05 in France; LANUV NRW in Germany). All experiments were performed on mice that had been electroporated at embryonic stage E15. Mice were housed in their home cages and kept under a 12-h light/dark cycle at  $22 \pm 2^\circ\text{C}$ . Food and water were available ad libitum. For optogenetics experiments, we used electroporated transgenic  $\text{Pvalb}^{\text{tm1(cre)Arbr}} (+/-)::\text{ChR2(H134R)-YFP(lox/-)}$  mice obtained after crossing homozygous  $\text{Pvalb}^{\text{tm1(cre)Arbr}}$  (JAX n°017320) females expressing Cre recombinase under the control of the Pvalb promoter (kindly provided by Dr JC Poncez, Institut du Fer à Moulin, Paris) and homozygous lox-STOP-lox ChR2-YFP males (JAX n°012569) (kindly provided by Dr MC Angulo, Institute of Psychiatry and Neuroscience, Paris). To identify and patch-clamp PV interneurons ex vivo, we used PV-Cre mice crossed with RCE:LoxP reporter mice (JAX n°010701) expressing enhanced green fluorescent protein (EGFP) under the control of the endogenous Gt(ROSA)26Sor promoter/enhancer regions and the CAG promoter. All mice were on a C57Bl/6J background. Females and males were used indifferently for electrophysiology, immunohistochemistry and imaging experiments. Only males were used for behavioral experiments.

### DNA plasmid constructs

pCAG-HAC4 expression vector design was based on pCAG-HT2B-HA. The pCAG-HT2B-HA was a gift from I Moutkine (Fer à Moulin Institute, Paris). The HT2B gene was removed and replaced by the full coding sequence for murine C4b (NM\_009780.2). The cDNA of C4b was obtained by reverse transcription PCR of total RNA from whole brain of adult 129S1/SvImJ mice. An HA tag was introduced at the N-terminus of the C4b sequence to allow the identification of the overexpressed C4 peptide. pCAG-tdTomato expression vector design was based on pCAG-mGFP which was a gift from Dr. C. Cepko <sup>1</sup> (Addgene plasmid # 14757). mGFP was replaced by tdTomato in the pCAG backbone. Subcloning of C4b and tdTomato sequences were done using InFusin (Clontech) and DNA were purified with Nucleospin kit (Macherey-Nagel). All constructs were verified by sequencing the entire coding region of the inserts.

### ***In utero* electroporation**

Pregnant mice at E15 were anesthetized by isoflurane 4% and were injected with Flunixin (0.05mg/kg body weight) before surgery. The surgery was performed under isoflurane 2% on a heated blanket and breathing was carefully monitored during all the procedure.

A midline laparotomy was performed and the uterine horns were exposed and moistened with sterile NaCl 0.9%. Between 0.5µl to 1µl of a solution containing pCAG-C4HA vector (0.8 µg/µl) together with pCAG-TdTomato vector (0.4 µg/µl) (molar ratio 4:1) in NaCl 0.9% were injected into the bilateral ventricles with a glass micropipette made from a microcapillary tube (#5-000-1001-X10, Drummond Scientific). In the control condition, a solution containing pCAG-TdTomato (0.8 µg/µl) alone was injected. The injected plasmid solution contained Fast Green (0.1%) to monitor the injection. The embryo's head in the uterus was held between the tweezers-type electrode consisting of two disc electrodes of 5 mm diameter (CUY650P5, Nepagene). The cathode was placed just above the olfactory bulbs and the anode was placed at the back of the head below the cerebellum. Electrical pulses (50V; 50 ms) were applied five times at intervals of 950 ms with an electroporator (CUY21 Edit, Nepagene). The uterine horn was placed back into the abdominal cavity. The abdominal cavity was filled up with sterile NaCl 0.9% and the abdominal muscle and skin were sutured separately. After recovery, pregnant mice were returned to their home cage and sutures were examined every day until birth. Correct tdTomato expression was first checked through the intact skin shortly after birth (age P1-P2) and confirmed *post mortem* on brain slices from each experimental animal. Mice without tdTomato expression in the prefrontal cortex (PFC) were excluded from the analysis. Control and C4-OE electroporations were carried out in parallel (either on the same day or during the same week) in order to minimize experimental biases. The survival rate of electroporated control and C4-OE mice was comparable.

### **Cranial window surgery**

At postnatal day 15 to 18 (P15-18) (and P30 respectively) a cranial window surgery was performed as previously described<sup>2-4</sup>. Mice were anesthetized with an intraperitoneal injection of ketamin/xylazin (0.13/0.01 mg/g body weight). For analgesia an additional subcutaneous injection of buprenorphin (0.05 mg/kg) was given. The anesthetic profile was deepened and maintained by inhalation of isoflurane (0.2-0.4% in 0.2 l/min oxygen). After the withdrawal reflex of the hind limb had subsided, the surgery begun. The eyes were covered with eye ointment (Bepanthen) to prevent drying and the mouse was placed on a heating pad

to maintain body temperature. Subsequently, mice were fixed to a stereotactic frame and the skin of the head was disinfected with 70% ethanol. The skin above the skull was removed using surgical scissors. To assure correct placement of the cranial window, the expression of the fluorophore was located using a UV light source. Next, the skull bone was dried and a 3 to 4-mm-diameter circular piece of bone was removed (3 mm at P15-P18, 4 mm at P30) over the prefrontal cortex using a dental drill. The dura mater was carefully removed and a circular coverslip ( $\varnothing$  3mm at P15-P18,  $\varnothing$  4 mm at P30) was inserted into the craniotomy and glued to the bone. The window was placed unilaterally above the electroporated hemisphere with the following coordinates describing the center of the circular glass (anteroposterior: +2mm, mediolateral: + or - 2mm (P30) and anteroposterior: +1.5 mm, mediolateral: + or - 1.5mm (P15-18). To allow stable and repetitive fixation of the mouse under the microscope, a small metal bar was attached next to the cortical window using UV-curable dental adhesive.

### **Two-photon *in vivo* imaging and analysis**

Two-photon imaging was performed right after surgery at a Zeiss 7 multiphoton microscope with a Zeiss 20 x objective (NA = 1). The anesthesia was maintained with 1-1.5% inhalation of isoflurane while the mouse was head fixed to a custom-made imaging frame. 2P imaging was performed spanning a depth of 20-200  $\mu\text{m}$  from the brain surface. This allowed us to image the full length of apical dendrites of Layer II/III pyramidal neurons. TdTomato fluorescence was excited at 980 nm using an InSight X3 tunable laser (Spectra-Physics) and collected through a 617/73 band-pass filter on a highly sensitive non-descanned detector. First, a 600  $\mu\text{m}$  (x) x 600  $\mu\text{m}$  (y) x 400  $\mu\text{m}$  (z) overview z-stack (4  $\mu\text{m}$  z-spacing) was acquired to retrieve the region of interest over multiple sessions. Then z-stacks of single dendrites (1  $\mu\text{m}$  z-spacing) with a pixel size of 0.089  $\mu\text{m}$  were acquired. After the first imaging session, the mice woke up and were placed in a cage with their littermates for 4-5 h (P15-18). For the second imaging session the mice were again anesthetized with isoflurane (initially 3%, then 1-1.5% for anesthesia maintenance) and imaged to retrieve the same dendritic structures recorded in the first imaging session.

Datasets were blinded prior to analysis by an experimenter not involved in the image analysis. In each animal 4-19 dendrites of 30-120  $\mu\text{m}$  length were analyzed. Spines that extended out laterally from the dendritic shaft were counted manually by scrolling through the z-stack, as previously described<sup>5</sup>. Spine density was calculated by the number of spines divided by the



length of the respective dendrite. Turnover of dendritic spines was calculated as follows:  $\text{turnover [\%]} = 100 * ((\#gained + \#lost) / (\#T1 + \#T2))$ . Here, #gained refers to the number of gained and #lost to the number of lost spines, respectively and #T1 and #T2 to the number of all spines at time-points T1 and T2, respectively. T1 and T2 refer to 0h and 4h.

## **Confocal imaging of dendritic spines and analysis**

### *Tissue preparation and immunochemistry*

All mice were injected with a lethal dose of pentobarbital (200 mg/kg) and perfused transcardially with saline solution followed by 4% (w/v) paraformaldehyde in phosphate buffered saline. Brains were removed, post-fixed in 4% paraformaldehyde for 20 hours, and then sectioned at 50  $\mu\text{m}$  in the coronal plane using a vibratome (VT1200, Leica). Sections were mounted (Dako Fluorescence mounting medium) directly on coverslips for confocal imaging.

### *Fixed tissue imaging*

Images of fixed brain slices were acquired on a commercial confocal microscope equipped with an Airyscan detector (Zeiss LSM800, Carl Zeiss Microscopy GmbH, Jena, Germany), using a 63X/1.4NA oil objective. Excitation of tdTomato was achieved by a 561nm laser and emission was collected using a 565-700 nm bandpass filter. Image stacks of dendrite fragments of pyramidal neurons were acquired with a pixel size of 0.049  $\mu\text{m}$ , a z-step size of 0.200  $\mu\text{m}$  and at a scan speed of 2.55  $\mu\text{s}/\text{pixel}$ . Images acquired with the Airyscan detector were processed by the ZEN software internal algorithm that reassigns the collected pixels to their correct positions to achieve a super-resolution image; for this a super-resolution threshold of 2 was used.

### *Reconstruction of dendritic fragments*

Image stacks were 3D reconstructed in IMARIS x64 8.1.2 (Bitplane AG, Zurich). Basal and apical dendrites and their associated spines were reconstructed semi-automatically using the Filament Tracer module. Spines that extended radially from the dendrite in three dimensions were drawn manually by turning the 3D visualized dendritic fragment.

Filament parameters, that is small diameter = 0.08  $\mu\text{m}$ , large diameter = 0.80  $\mu\text{m}$  and contrast threshold = 0.10 were defined empirically by measuring a random set of acquired images and comparing the rendered filament with the preprocessed image. Spine density per dendritic fragment and morphological parameters of spines (ratio of mean head to neck diameter,

maximum head diameter and spine length) were measured, exported for further analysis (Cluster analysis) and plotted in a 3D graph.

### *Cluster analysis*

The cluster analysis was performed as previously described<sup>6</sup>. The spine parameters, ratio of the mean head to neck diameters ( $\text{ØHead}/\text{ØNeck}$ ), the maximum head diameter ( $\text{Ømax Head}$ ) and spine length were taken into account. Agglomerative Hierarchical Clustering Analysis (AHC) was performed using Python's scikit-learn toolbox. Before cluster analysis, the parameters were centered and scaled to unit variance using the standard scaler from Python's scikit-learn toolbox. AHC dissimilarity level was calculated based on Euclidean distance. Agglomeration was performed using Ward's method. By fitting the hierarchical clustering to the population of all analyzed spines we identified three clusters.

## **Histology**

### *Immunohistochemistry*

Mice aged P30 and P180 were anesthetized with pentobarbital (150 mg/kg) and intracardially perfused with PBS followed by 4% buffered formaldehyde (Histofix, Roti). The brains were removed and fixed overnight in 4% Histofix. 40  $\mu\text{m}$  sagittal sections were cut using a vibratome (VT1000S, Leica). Free-floating sections were permeabilized and blocked in 0.2% Triton-PBS, 3% BSA-PBS for 60 min before incubation with the first antibody in 0.2% Triton, 3% BSA-PBS overnight at 4°C. After three washes in PBS, sections were incubated with the secondary antibody in PBS for 2h, washed again in PBS, and mounted using Fluoromount G (Invitrogen). Primary antibodies were: rat anti-HA (1:300, Sigma), rabbit anti-Parvalbumin (PV27, 1:1000, Swant), mouse anti-Glutamate Decarboxylase 67 (GAD67, clone 1G10.2, 1:500, Millipore-Merk). Secondary antibodies were: goat anti-rat Alexa Fluor 488 (1:400, Invitrogen), donkey anti-rabbit Cy3 (1:1000, Jackson Immuno Research), goat anti-mouse Alexa Fluor 647 (1:1000, Jackson Immuno Research). Hoechst 33342 was used for nuclear staining. Images were acquired using a confocal microscope (SP5, Leica).

### *PV cell number and distribution*

The distribution of PV neurons in different PFC layers was quantified using maximal intensity projections of 3D confocal stacks. For each mouse, 4 PFC sagittal sections were imaged using a confocal microscope (Leica SP5, 40X objective). Layers were identified based on Hoechst nuclear staining and delineated using ImageJ's ROI manager software. The

distribution of PV neurons in different PFC layers was quantified using maximal intensity projections of 3D confocal stacks. The density of PV cells (number of PV cells per surface unit) in each layer was obtained by averaging the density across the 4 slices. During image acquisitions and quantifications, the investigator was blind to the condition (control or C4-OE).

#### *GAD67 expression*

GAD67 expression levels in PV cells were measured by analyzing confocal images (Leica SP5, 40X objective) using ImageJ. For each animal, 3 to 4 PFC sagittal slices were acquired. At least 80 PV neurons per mouse were analyzed. A single confocal section located 10  $\mu\text{m}$  below the surface of the slice was acquired in each slice. The same acquisition parameters were used for each slice and were set such that the brightness of the GAD67 immunostaining was below saturation values. The brightness of the PV immunostaining was set above saturation level, allowing to use PV as a marker to delineate the shape of PV neurons. The density of GAD67 immunostaining within PV cells was then automatically calculated using Image J. Several experiments were performed with a minimum of 6 mice per condition. As the fluorescence intensity is only comparable for samples processed simultaneously, the results are presented as a percentage of the mouse controls performed at the same time. During image acquisitions and quantifications, the investigator was blind to the condition (control or C4-OE).

#### **Acute slice preparation**

250  $\mu\text{m}$ -thick coronal slices were prepared from brains of control and C4-OE mice. Mice were deeply anesthetized with sodium pentobarbital (150 mg/kg) and perfused transcardially with ice-cold (0–4°C) oxygenated (95% O<sub>2</sub>-5% CO<sub>2</sub>) solution containing (in mM): 110 Choline chloride, 2.5 KCl, 25 glucose, 25 NaHCO<sub>3</sub>, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 0.5 CaCl<sub>2</sub>, 7 MgCl<sub>2</sub>, 11.6 L-ascorbic acid and 3.1 sodium pyruvate. The brain was extracted and acute sagittal slices were cut in the same ice-cold solution using a vibroslicer (HM 650 V, Microm), then stored in artificial cerebrospinal fluid (ACSF) containing the following (in mM): 125 NaCl, 2.5 KCl, 25 glucose, 25 NaHCO<sub>3</sub>, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 2 CaCl<sub>2</sub>, and 1 MgCl<sub>2</sub>, continuously bubbled with 95% O<sub>2</sub>-5% CO<sub>2</sub>. Slices were incubated in ACSF at 32°C for 20 minutes and then at room temperature (20-25°C). For patch-clamp recordings, slices were transferred to the recording chamber where they were continuously superfused with ACSF (30-32°C) buffered by continuous bubbling with 95% O<sub>2</sub>-5% CO<sub>2</sub>.

## **Electrophysiology and optogenetics**

Whole-cell current and voltage-clamp recordings were performed in PFC layer II/III. Recorded pyramidal cells all expressed tdTomato. Recorded PV interneurons were in the vicinity of electroporated tdTomato+ pyramidal cells and were visually identified based on the expression of EGFP (in PV-Cre::RCE mice) or YFP (in PV-Cre::lox-STOP-lox ChR2-YFP mice). Stimulus delivery and data acquisition were performed using Patchmaster software (Multichannel Systems). Signals were acquired with an EPC10-usb amplifier (Multichannel Systems), sampled at 20 kHz and filtered at 4 kHz. Offline analysis was performed using Clampfit (Molecular Devices) and Igor Pro (Wavemetrics). For the study of miniature post synaptic currents, recordings were filtered offline at 2 kHz and analyzed using MiniAnalysis (Synaptosoft).

Patch-clamp pipettes (3–6 Mohm resistance) were prepared from borosilicate glass (BF150-86-10; Harvard Apparatus) using a DMZ pipette puller (Zeitz). Current-clamp experiments were performed using the following intracellular solution (in mM): 105 K-gluconate, 10 HEPES, 10 phosphocreatine-Na, 0.3 Na<sub>3</sub>GTP, 4 MgATP, 30 KCl (pH 7.25, adjusted with KOH). For voltage-clamp experiments the following intracellular solution was used (in mM): 120 Cs-methane sulfonate, 10 CsCl, 10 Hepes, 10 Phosphocreatine, 0.2 EGTA, 8 NaCl, 2 ATP-Mg, 3 QX 314 (pH 7.25, adjusted with CsOH). To record miniature and evoked inhibitory postsynaptic currents (mIPSCs and eIPSCs), we used an intracellular solution containing (in mM): 60 Cs methane sulfonate, 70 CsCl, 10 Hepes, 10 Phosphocreatine, 0.2 EGTA, 8 NaCl, 2 ATP-Mg (pH 7.25, adjusted with CsOH). Liquid junction potentials (-14.9 mV for Cs-based intracellular solutions and -5mV for the K<sup>+</sup>-based intracellular solution) were left uncorrected. The cell membrane resistance, membrane capacitance and series resistance were calculated in the seal-test configuration from the decay and steady-state of a current generated in response to a 10 mV hyperpolarizing voltage step. The series resistance was left uncompensated.

### *Excitability*

Pyramidal cell and PV interneuron intrinsic properties and excitability were recorded in current-clamp in presence of the following synaptic blockers: SR95531 hydrobromide (Gabazine, 10  $\mu$ M, Hello Bio), 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, 10  $\mu$ M, Hello Bio) and D-2-amino-5-phosphonopentanoic acid (D-APV, 50  $\mu$ M, Hello Bio). Spike frequency-depolarization curves were generated by injecting series of 500 ms depolarizing current steps increasing by 50 pA. The firing pattern of patched-clamped YFP<sup>+</sup> cells in the

cortex of PV-Cre::lox-STOP-lox Chr2-YFP mice was monitored using 500 ms pulse of blue light emitted by a light emitting diode (CoolLED).

#### *Miniature EPSCs and evoked EPSCs*

Miniature excitatory post synaptic currents (mEPSCs) were recorded at a holding potential of -65 mV in the presence of tetrodotoxin (TTX, 0.5  $\mu$ M, Hello Bio) and Gabazine (10  $\mu$ M). At least 100s were analysed for each recording. Evoked excitatory post synaptic currents (eEPSCs) were recorded at a holding potential of -65 mV and triggered at a frequency of 0.1 Hz in the presence of GABAzine (10  $\mu$ M) with a monopolar electrode connected to a constant current stimulator (Iso-Flex, A.M.P.I.), positioned in the vicinity of the recorded neuron. At least 12 consecutive sweeps were acquired for each cell. The paired-pulse ratio (PPR) was calculated as the ratio of the peak amplitude of the averaged current response evoked by the second stimulation to the peak amplitude of the averaged current response evoked by the first stimulation.

#### *Miniature IPSCs and evoked IPSCs*

mIPSCs were recorded at a holding potential of -65 mV in the presence of TTX (0.5  $\mu$ M, Hello Bio), CNQX, (10  $\mu$ M) and D-APV (50  $\mu$ M). At least 100 s were analysed for each recording. Evoked IPSCs were recorded at a holding potential of +10mV to avoid unclamped action potentials commonly associated with evoked IPSCs measured at hyperpolarized potentials in our high-chloride recording conditions, in the presence of CNQX (10  $\mu$ M) and APV (50  $\mu$ M). At least 25 consecutive sweeps were acquired and averaged for each cell. For electrically evoked IPSC (eIPSCs), four stimulations at 20 Hz were applied with a monopolar electrode connected to a constant current stimulator positioned in layer 3, 100-150  $\mu$ m away from of the recorded neuron. For light-evoked IPSCs, the recorded pyramidal cell was placed in the center of the field of view. The response to four 0.5-ms pulses of blue light (470nm) at 20 Hz emitted by a light emitting diode (CoolLED) was recorded. The paired-pulse ratio of eIPSCs was calculated as the ratio of the peak amplitude of the averaged current response evoked by the second, third and fourth stimulations to the peak amplitude of the averaged current response evoked by the first stimulation.

#### *AMPA/NMDA ratio and rectification index of AMPAR-mediated currents*

To measure the AMPA/NMDA ratio and the rectification index, EPSC were evoked with a monopolar electrode positioned in layer 3, 100-150  $\mu$ m away from the recorded neuron. At

least 12 consecutive sweeps were acquired and averaged at each holding potential. The AMPA/NMDA ratio was recorded in presence of GABA<sub>A</sub>zine (10  $\mu$ M). The AMPA current ( $I_{\text{AMPA}}$ ) was measured as the peak amplitude of the current response at a holding potential of -65mV. The NMDA current ( $I_{\text{NMDA}}$ ) was measured at a holding potential of +55mV, 50ms after stimulation, when AMPAR-mediated current have receded due to the fast deactivation kinetics of AMPARs. The ratio is calculated as  $I_{\text{AMPA}}/I_{\text{NMDA}}$ .

To study the rectification AMPA-mediated currents, pyramidal cells were recorded in the presence of GABA<sub>A</sub>zine (10  $\mu$ M) and D-APV (50  $\mu$ M). Spermine (10  $\mu$ M) was added to the intracellular solution. The rectification index was defined as the peak amplitude of synaptic currents measured at a holding potential of +55 mV ( $\approx$  +40 mV after correcting for junction potential) multiplied by 2 and divided by the peak amplitude of synaptic currents measured at a holding potential of -65mV ( $\approx$  -80 mV after correcting for junction potential).

## **Behavior**

### *Actimeter*

We used actimeter racks of 8 circular corridors (Imetronic, <http://www.imetronic.com/devices/circular-corridor/>) equipped with infrared sensors to detect and automatically record locomotion. Animals were introduced individually in corridors and their spontaneous, novelty-driven activity was recorded for a total of 60 minutes. Activity, measured as photocell beam breaks, was averaged over periods of 5 minutes. Control and C4-OE male mice were tested at weaning (age P30).

### *Open Field*

The test took place in a white open-field apparatus (ViewPoint, Lyon, France) under high illumination ( $\sim$ 100 lux). Animal were individually placed in the corner of the open-field and were allowed to explore for 9 min. A video tracking system, which included a computer-linked overhead camera, was used to monitor general locomotor activity (<https://viewpoint.fr/en/p/software/videotrack>, ViewPoint, Lyon, France). Control and C4-OE male mice were tested a week after weaning (age P35 – P39).

### *Odor Span and Spatial Span Test*

Working memory capacity was tested using a protocol adapted from a previous study<sup>7</sup>. All scented mixtures were prepared with 100 g of sawdust, 18 ground powdered pellets

(Chocapic, Nestle) to mask the scent of the reward pellet and 3 g of a ground powder odor (Savory, Cinnamon, Cumin, Nutmeg, Mint, Fennel, Four spices, Herbs of Provence, Coriander, Basil, Coffee, Ginger). During shaping and testing, adult (4- to 6-month-old) male mice were maintained at 80-85% of their body weight. During the shaping phase, mice were trained to dig into a bowl of unscented sawdust to retrieve the hidden reward pellet. Once all animals had learned the procedure, the non-matching to sample (NMTS) task started for 5 consecutive days. Each day, the mouse was placed on a square platform and required to dig into the first bowl (odor 1) to retrieve the reward and to remember the scent of that bowl. Upon consumption of the reward pellet, a second baited bowl (odor 2) was added. The first bowl (no longer baited) and the new bowl were randomly distributed on the platform. Following retrieval of the pellet, the first bowl was removed and a novel baited bowl (odor 3) randomly placed. This process was repeated until each individual mouse had been exposed to 11 pairs of scented bowls or stopped after a maximum of 12 minutes when the mouse did not perform the task, and the number of correct pairs (direct digging into the baited bowl) was recorded. The order of the odors was randomized across days.

Following the NMTS, animals were then tested 4 consecutive days for the odor span task, using the same procedure except that, following reward retrieval, the no longer baited bowls were not removed so that mice had to remember an increasing numbers of odors (span). The span length was calculated as the number of correct responses with more than one bowl on the platform prior to the first mistake. Thus, using 12 odors, the maximum span length is 11. The span score was given by the first mistake, but each mouse was allowed to continue the task until the 12 odors had been sampled or for a maximum of 12 minutes. The span length was the best score obtained by each tested mouse over a period of 4 consecutive days. As for the NMTS task, the order of the odors was randomized across days.

The same mice were subsequently tested for 4 consecutive days in the spatial span task. They had to remember an increasing numbers of locations, following a procedure similar to the odor span test, except that bowls of unscented sawdust were identified according to their spatial location on the platform. The span length was the best score obtained by each tested mouse over a period of 4 consecutive days.

The investigator was blind to the condition (control or C4-OE) during behavioral tests and analysis.

## Immunoblotting

Mice were euthanized under anesthesia and brains were removed from the skull. TdTomato-expressing areas of the prefrontal cortex were dissected under a stereomicroscope. Extracts were lysed in RIPA buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 10 µl/ml NP-40, 5 µg/ml sodium deoxycholate, 1 µg/ml SDS) containing Complete protease inhibitors (Roche). 10 brain extracts in each condition were pooled and equal amounts of proteins (60µg) were loaded on NuPAGE 8–12% Bis-Tris gels (Thermo Fisher Scientific) and transferred to 0.45 µm nitrocellulose membrane in 25 mM Tris–HCl, pH 7.4, 192 mM glycine and 200 ml/l ethanol for 2h at 4°C. Membranes were blocked with 50 g/l non-fat dry milk in Tris-buffered saline (TBS; 0.25 M Tris/0.5 M NaCl, pH 7.5) for 1 h at room temperature and then incubated with primary antibodies in the same buffer containing Tween 0.1% (TBST) for at least 2h. After three washes in TBST, the membranes were incubated for 1h with IRDye-conjugated secondary antibodies, and imaged using Odyssey Imaging System (LI-COR Biosciences). Quantifications were performed using Image Studio Lite software (LI-COR Biosciences). C4 levels were quantified by normalizing to the loading control, GAPDH. Primary antibodies were: rat anti-HA (1:500, Sigma), rabbit anti-C4α (1:2000, Proteintech), chicken anti-GAPDH (1:2000, Millipore). Secondary antibody were: goat anti-rat IR800 (1:10000, Advansta), goat anti-rabbit IR700 (1:10000, Invitrogen) and goat anti-chicken IR700 (1:10000, Advansta).

## Statistical analysis

Data are presented as mean ± SEM. Statistical analyses were performed using Prism (Graphpad). The normality of data distribution was tested using Shapiro-Wilk's test. Unpaired two-tailed T-tests (for normally distributed datasets) or Mann–Whitney tests (for non-normally distributed datasets) were used for comparisons between two groups. For multiple comparisons we used two-way ANOVA followed by Sidak's test. Values of  $P < 0.05$  were considered statistically significant.  $P$  values are reported as follows: \*  $P < 0.05$ ; \*\*  $P < 0.01$ ; \*\*\*  $P < 0.001$ ; \*\*\*\*  $P < 0.0001$ .


## Supplementary references

1. Matsuda T, Cepko CL. Electroporation and RNA interference in the rodent retina in vivo and in vitro. *Proc Natl Acad Sci U S A* 2004; **101**(1): 16-22.
2. Bittner T, Fuhrmann M, Burgold S, Ochs SM, Hoffmann N, Mitteregger G *et al.* Multiple events lead to dendritic spine loss in triple transgenic Alzheimer's disease mice. *PLoS One* 2010; **5**(11): e15477.



3. Fuhrmann M, Bittner T, Jung CK, Burgold S, Page RM, Mitteregger G *et al.* Microglial Cx3cr1 knockout prevents neuron loss in a mouse model of Alzheimer's disease. *Nat Neurosci* 2010; **13**(4): 411-413.
4. Fuhrmann M, Mitteregger G, Kretzschmar H, Herms J. Dendritic pathology in prion disease starts at the synaptic spine. *J Neurosci* 2007; **27**(23): 6224-6233.
5. Gu L, Kleiber S, Schmid L, Nebeling F, Chamoun M, Steffen J *et al.* Long-term in vivo imaging of dendritic spines in the hippocampus reveals structural plasticity. *J Neurosci* 2014; **34**(42): 13948-13953.
6. Pfeiffer T, Poll S, Bancelin S, Angibaud J, Inavalli VK, Keppler K *et al.* Chronic 2P-STED imaging reveals high turnover of dendritic spines in the hippocampus in vivo. *eLife* 2018; **7**.
7. Young JW, Kerr LE, Kelly JS, Marston HM, Spratt C, Finlayson K *et al.* The odour span task: a novel paradigm for assessing working memory in mice. *Neuropharmacology* 2007; **52**(2): 634-645.
8. Gabbott PL, Warner TA, Jays PR, Salway P, Busby SJ. Prefrontal cortex in the rat: projections to subcortical autonomic, motor, and limbic centers. *J Comp Neurol* 2005; **492**(2): 145-177.
9. Schwab M, Agid Y, Glowinski J, Thoenen H. Retrograde axonal transport of 125I-tetanus toxin as a tool for tracing fiber connections in the central nervous system; connections of the rostral part of the rat neostriatum. *Brain Res* 1977; **126**(2): 211-224.
10. Arikuni T, Kubota K. The organization of prefrontocaudate projections and their laminar origin in the macaque monkey: a retrograde study using HRP-gel. *J Comp Neurol* 1986; **244**(4): 492-510.
11. Reiner A, Hart NM, Lei W, Deng Y. Corticostriatal projection neurons - dichotomous types and dichotomous functions. *Front Neuroanat* 2010; **4**: 142.

## Supplementary figures, tables and legends



## Supplementary Figure 1: C4 overexpression in prefrontal cortex using in utero electroporation

(A) Schematic drawing illustrating the in utero electroporation (IUE) method at embryonic day 15 (E15) for specific targeting of layer II/III PFC pyramidal neurons. Targeted PFC areas include the anterior cingulate cortex (Cg1), the prelimbic cortex (PrL) and the infralimbic cortex (II). VZ: ventricular zone; OB: olfactory bulb.

(B) Left panel: Electroporated brain at postnatal day 30 (P30) exhibiting tdTomato expression in PFC; scale bar: 5 mm. Middle panel: Sagittal slice showing the distribution of TdTomato positive cells in PCF layer II/III. Note that the expression of tdTomato in the striatum originates from tdTom+ PFC neurons in layer II-III which are known to send direct axonal projections to the striatum in mammals<sup>8-11</sup>. OB: olfactory bulb; scale bar: 500  $\mu$ m. Right panel: typical morphology of an electroporated pyramidal cell; scale bar: 30  $\mu$ m.

(C) Plasmid constructs used for IUE. Embryonic brains were co-electroporated with pCAG-HA-C4 and pCAG-tdTomato for C4 overexpression (C4-OE), or only with pCAG-tdTomato (control). pCAG-HA-C4 and pCAG-tdTomato were co-electroporated with a 4:1 ratio to ensure that most tdTomato+ cells overexpressed C4.


(D) Immunostaining for TdTomato (red) and HA (green) in PFC layer II/III at P30 (upper panels) and P180 (bottom panels). The three panels on the right are enlargements of the boxed area in the left panel. Note that tdTom+ neurons also expressed HA-C4; scale bar: 100  $\mu$ m; scale bar in magnified image: 20  $\mu$ m.

(E) Quantification of the results obtained from P30 mice as exemplified in (D).

(F) Schematic drawing illustrating the processing of C4: C4 is synthesized as a single chain precursor molecule (prepro-C4) and processed to a three-chain structure ( $\beta$ - $\alpha$ - $\gamma$ ) before secretion. The HA tag was added at the 5' end of the signal peptide-encoding sequence, i.e. before the signal peptide region, in order not to interfere with the folding and processing of C4. The black dot represents the antigenic region recognized by the anti-C4  $\alpha$ -chain antibody used for western blot experiments in (G).


(G) Immunoblot of HA (left) and C4  $\alpha$ -chain (right) in lysates from the electroporated region of control and C4-OE mice aged P30. To compensate for the lack of material and relatively low levels of C4 expression, lysates from the electroporated region were prepared from individual mice and pooled for the western blot analysis (control mice: N = 10 ; C4-OE mice: N = 10). Left panel: HA was only detected at the molecular weight of prepro-C4 (~ 200 kDa) in the C4-OE condition, suggesting that the HA tag was correctly cleaved together with the signal peptide. Right panel: a band was detected at the molecular weight of the C4  $\alpha$ -chain (~ 93 kDa). Data were normalized to GAPDH.

(H) Densitometry quantification of the results illustrated in (G). Data were normalized to GAPDH for each experiment and expressed as a fold change of control mean density. The experiment was repeated four times. Each circle represents the result of an individual experiment. The bar represents the mean of four experiments.



**Supplementary Figure 2: Similar laminar position of electroporated pyramidal cells in control and C4-OE mice**

Left: sagittal PFC sections from control and C4-OE mice; scale bar: 300 $\mu$ m. Right, enlargements of the boxed area in the left panel showing PFC layers. All tdTomato+ cells are located in PFC layer III; scale bar: 100  $\mu$ m; M2: motor cortex; PFC: prefrontal cortex; OB: olfactory bulb.




### Supplementary Figure 3: Passive and active neuronal properties in control and C4-OE mice

(A) Sample spike trains evoked by a 400 pA somatic current injection in a control and in a C4-OE pyramidal cell; scale bars: 20 mV/100 ms.

(B) f-i curves for control (n = 20 cells from 3 mice) and C4-OE neurons (n = 17 cells from 3 mice).

(C) Input resistance (left), Cell capacitance (middle) and resting membrane potential (right) in control (n = 20 cells from 3 mice) and C4-OE neurons (n = 17 cells from 3 mice). Neither firing frequency nor intrinsic neuronal properties differed between control and C4-OE pyramidal cells.

All data are represented as mean  $\pm$  SEM. Each open circle represents an individual neuron.



## Supplementary Figure 4: Cluster analysis of dendritic spines in control and C4-OE mice


(A) 3D reconstruction of dendritic fragments confocally imaged in fixed tissue of control (CTR) and C4-OE animals aged P15-P18.

(B) Spine density measured on reconstructed dendrites; data from  $n = 7$  CTR and  $n=8$  C4-OE mice with in total 49 (CTR) and 66 (C4-OE) dendrites; single dots show individual mice; diagram shows mean  $\pm$  SEM;  $p=0.0229$ , unpaired t-test; statistics were performed using individual mice as the unit of analysis.

(C, D) 3D morphology plots visualizing the spines of CTR and C4-OE mice (C), and their affiliation to identified clusters 1, 2 and 3 (D); plotted are, the ratio of mean head to neck diameters ( $\emptyset_{\text{Head}}/\emptyset_{\text{Neck}}$ ), spine length and maximum head diameter ( $\emptyset_{\text{max Head}}$ ) of in total 8123 (CTR) and 10946 spines (C4-OE).

(E) Quantification of spine proportions within identified clusters, comparing spines of CTR and C4-OE mice.

(F) Table summarizing the morphological parameters utilized for the cluster analysis:  $\emptyset_{\text{Head}}/\emptyset_{\text{Neck}}$ ,  $\emptyset_{\text{max Head}}$  and length of spines, for CTR (grey) and C4-OE animals (red). Data are represented as median and interquartile range (25th–75th percentile).





### Supplementary Figure 5: Light-evoked firing of ChR2-expressing PV neurons

(A) Homozygous PV-Cre::RCE females were crossed with homozygous lox-STOP-lox ChR2-YFP males before electroporation of timed-pregnant females with tdTomato or C4 + tdTomato. Electroporated offspring (age P25-P30) were used for ex vivo patch-clamp and optogenetics.

(B) Left, patch-clamped YFP+ cell (arrow); right, spike train evoked by a 500 ms pulse of blue (470 nm) light. YFP+ cells exhibited the typical fast-spiking pattern of PV neurons ( $n = 3$  neurons).

(C) Light-evoked IPSCs were recorded at a holding potential of +10 mV in electroporated PFC pyramidal cells from control or C4-OE PVCre-RCE::lox-STOP-lox ChR2-YFP mice.

(D) Light-evoked IPSC blocked by the addition of Gabazine (10  $\mu$ M), demonstrating its GABAergic nature; scale bars: 50 pA/50ms.


**A****B**

### Supplementary Figure 6: Laminar distribution of PV neurons in control and C4-OE mice

(A) PV staining in PFC sagittal slices from a control and from a C4-OE mouse; scale bar: 100  $\mu$ m.

(B) The repartition by layer (left) as well as the total number of histologically identified PV (right) was unchanged in C4-OE mice (n = 10 mice) compared to control mice (n = 9 mice) (Two-Way ANOVA,  $P > 0.05$ ). All data are represented as mean  $\pm$  SEM. ns, not significant.





## Supplementary Figure 7: Increased excitatory inputs onto PV neurons in C4-OE mice

(A) Sample traces of mEPSCs recordings in PV interneurons from control and C4-OE juvenile (P25-P30) mice recorded at  $-65$  mV in the presence of TTX and GABazine; scale bars: 20 pA/100ms.

(B) mEPSC frequency, but not IPSC amplitude, was lower in C4-OE mice ( $n = 16$  cells from 6 mice) than in control mice ( $n = 20$  cells from 3 mice; frequency: Mann-Whitney test, \*  $P = 0.0184$ ).


(C) Average of 12 consecutive recordings; EPSCs were evoked by paired-pulse stimulation and recorded in PV interneurons from control and C4-OE mice at a holding potential of  $-65$  mV; scale bars: 50 pA/50 ms.

(D) Summary histogram of the paired-pulse ratio in control and C4-OE (control:  $n = 24$  cells from 4 mice; C4-OE:  $n = 21$  cells from 3 mice).

(E) Sample traces of mIPSCs recorded at a holding potential of  $-65$  mV with a high-chloride intracellular solution in PV interneurons from control and C4-OE juvenile (P25-P30) mice, in the presence of TTX, APV and CNQX; scale bar: 20 pA/100 ms.

(F) Mean frequency (left) and amplitude (right) of mIPSCs in control ( $n = 18$  cells from 4 mice) and C4-OE neurons ( $n = 13$  cells from 4 mice).

All data are represented as mean  $\pm$  SEM. Open circles represent single cells.



**Supplementary Figure 8: Overview diagram summarizing the main findings at different stages of postnatal development**

**Table S1**

	Control	n	N	L	C4-OE	n	N	L	Test	p value	Significant	F statistics (ANOVA)	
<b>Intrinsic properties at P25-P30</b>													
Membrane resistance (MΩ)	91.9 ± 4.97	20	3	2	98.6 ± 7.62	17	3	3	T-test	0.456	ns		
Resting membrane potential (mV)	-70.9 ± 1.16	20	3	2	-71.2 ± 1.17	17	3	3	T-test	0.841	ns		
Membrane capacitance (pF)	327 ± 24.7	20	3	2	315 ± 31.6	17	3	3	T-test	0.773	ns		
Firing frequency		20	3	2		17	3	3	Two-way RM ANOVA	0.304	ns	F <sub>(1,35)</sub> = 1.09	
<b>mEPSC frequency (Hz)</b>										Two-way ANOVA	C4 factor: <0.0001	****	F <sub>(1,105)</sub> = 17.4
											Age factor: <0.0001	****	F <sub>(2,105)</sub> = 54.4
											Interaction: 0.0059	**	F <sub>(2,105)</sub> = 5.4
<i>P9-P10</i>	2.03 ± 0.17	21	3	2	2.02 ± 0.15	18	4	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	10.2 ± 1.31	16	5	4	6.20 ± 0.90	14	6	3	Sidak's post-hoc test		#####		
<i>P150-P220</i>	4.58 ± 0.36	20	6	4	2.59 ± 0.21	22	5	4	Sidak's post-hoc test		#		
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		#####		
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		#####		
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		##		
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		#####		
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		###		
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		ns		
<b>mEPSC amplitude (pA)</b>										Two-way ANOVA	C4 factor: 0.399	ns	F <sub>(1,105)</sub> = 0.716
											Age factor: <0.0001	****	F <sub>(2,105)</sub> = 85.82
											Interaction: 0.477	ns	F <sub>(2,105)</sub> = 0.745
<i>P9-P10</i>	17.2 ± 1.241	21	3	2	17.7 ± 0.855	18	4	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	8.87 ± 0.586	16	5	4	8.10 ± 0.478	14	6	3	Sidak's post-hoc test		ns		
<i>P150-P220</i>	9.927 ± 0.442	20	6	4	8.637 ± 0.350	22	5	4	Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		#####		
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		#####		
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		#####		
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		#####		
<b>Excitatory PPR</b>										Two-way ANOVA	C4 factor: 0.809	ns	F <sub>(1,105)</sub> = 0.033
											Age factor: 0.0011	**	F <sub>(1,105)</sub> = 7.31
											Interaction: 0.806	ns	F <sub>(1,105)</sub> = 0.213
<i>P9-P10</i>	0.916 ± 0.056	16	3	2	0.875 ± 0.031	21	3	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	1.04 ± 0.04	17	3	2	1.04 ± 0.05	21	3	2	Sidak's post-hoc test		ns		

<i>P150-P220</i>	1.06 ± 0.07	19	5	3	1.08 ± 0.04	17	4	3	Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		ns		
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		ns		
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		#		
<b>AMPA/NMDA ratio</b>										Two-way ANOVA	C4 factor: <0.0001	****	F (1, 108) = 18.9
										Age factor: <0.0001	****	F (2, 108) = 40.6	
										Interaction: 0.007	**	F (1, 108) = 5.2	
<i>P9-P10</i>	0.856 ± 0.078	17	3	2	0.844 ± 0.085	21	3	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	0.928 ± 0.028	23	4	2	1.294 ± 0.086	20	5	3	Sidak's post-hoc test		###		
<i>P150-P220</i>	1.317 ± 0.073	17	4	2	1.752 ± 0.085	16	4	3	Sidak's post-hoc test		###		
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		ns		
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		###		
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		###		
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		####		
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		###		
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		####		
<b>NMDA decay time (ms)</b>										Two-way ANOVA	C4 factor: 0.490	ns	F (1, 108) = 0.48
										Age factor: <0.0001	****	F (2, 108) = 73.09	
										Interaction: 0.332	ns	F (1, 108) = 1.12	
<i>P9-P10</i>	0.098 ± 0.008	17	3	2	0.104 ± 0.006	21	3	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	0.056 ± 0.003	23	4	2	0.051 ± 0.002	20	5	3	Sidak's post-hoc test		ns		
<i>P150-P220</i>	0.058 ± 0.003	17	4	2	0.050 ± 0.002	16	4	3	Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		####		
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		####		
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		####		
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns		
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		####		
<b>Rectification Index</b>										Two-way ANOVA	C4 factor: <0.0001	****	F (1, 112) = 18.09
										Age factor: 0.0058	**	F (2, 112) = 5.40	
										Interaction: 0.196	ns	F (2, 112) = 1.65	
<i>P9-P10</i>	1.09 ± 0.06	20	3	2	1.02 ± 0.05	20	3	2	Sidak's post-hoc test		ns		
<i>P25-P30</i>	1.31 ± 0.05	21	3	3	1.06 ± 0.05	16	3	2	Sidak's post-hoc test		##		

<i>P150-P220</i>	1.13 ± 0.03	17	4	2	0.97 ± 0.03	24	3	2	Sidak's post-hoc test		#	
<i>Control: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		##	
<i>Control: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		#	
<i>Control: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		ns	
<i>C4-OE: P9-P10 vs P25-P30</i>									Sidak's post-hoc test		ns	
<i>C4-OE: P25-P30 vs P150-P220</i>									Sidak's post-hoc test		ns	
<i>C4-OE: P9-P10 vs P150-P220</i>									Sidak's post-hoc test		ns	
<b>mIPSC Amplitude at P25-P30 (pA)</b>	17.1 ± 0.98	16	4	2	12.74 ± 0.46	16	6	4	T-test	0.0003	***	
<b>mIPSC Frequency at P25-P30 (Hz)</b>	7.08 ± 0.57	16	4	2	4.46 ± 0.61	16	6	4	Mann-Whitney test	0.0004	***	
<b>Short-term plasticity of evoked IPSCs at P25-P30 (electrical stim)</b>									Two-way RM ANOVA	0.089	ns	F <sub>(1,64)</sub> = 2.988
<i>Pulse II/ pulse I</i>	0.542 ± 0.028	29	3	2	0.610 ± 0.027	37	3	2	Sidak's post-hoc test		ns	
<i>Pulse III/ pulse I</i>	0.437 ± 0.025	29	3	2	0.488 ± 0.024	37	3	2	Sidak's post-hoc test		ns	
<i>Pulse IV/ pulse I</i>	0.397 ± 0.022	29	3	2	0.442 ± 0.021	37	3	2	Sidak's post-hoc test		ns	
<b>Short-term plasticity of evoked IPSCs at P25-P30 (optogenetics)</b>									Two-way RM ANOVA	0.014	*	F <sub>(1,71)</sub> = 6.85
<i>Pulse II/ pulse I</i>	0.541 ± 0.016	43	4	3	0.592 ± 0.021	30	3	2	Sidak's post-hoc test		ns	
<i>Pulse III/ pulse I</i>	0.400 ± 0.013	43	4	3	0.465 ± 0.019	30	3	2	Sidak's post-hoc test		#	
<i>Pulse IV/ pulse I</i>	0.357 ± 0.014	43	4	3	0.411 ± 0.018	30	3	2	Sidak's post-hoc test		ns	

**Table S1. Summary table of patch-clamp results (pyramidal cells)**

N refers to the number of mice, n refers to the number of neuronal recordings, L refers to the number of litters. All data are presented as mean ± SEM. Sidak's test #  $P < 0.05$ ; ##  $P < 0.01$ ; ###  $P < 0.001$ ; ####  $P < 0.0001$ .

**Table S2**

	Control	N	L	C4-OE	N	L	Test	p value	Significant	F statistics (ANOVA)
<b><i>In vivo</i> 2-photon imaging at P30</b>										
Spine density	0.880 ± 0.043	10	3	0.666 ± 0.029	7	3	T-test	0.0023	**	
<b><i>In vivo</i> 2-photon imaging at P15-P18</b>										
Spine density at 0h	1.145 ± 0.028	5	3	0.921 ± 0.021	7	3	Mann-Whitney test	0.0025	**	
Spine density from 0h to 4h		5	3		7	3	Two-way RM ANOVA	C4 factor: < 0.0001	****	F <sub>(1,10)</sub> = 44.8
								Time factor: 0.780	ns	F <sub>(1,10)</sub> = 0.083
								Interaction 0.366	ns	F <sub>(1,10)</sub> = 0.899
Stable spine density	1.044 ± 0.024	5	3	0.883 ± 0.002	7	3	T-test	0.0007	***	
Gained spine density	0.092 ± 0.009	5	3	0.045 ± 0.007	7	3	T-test	0.0018	**	
Lost spine density	0.099 ± 0.006	5	3	0.039 ± 0.006	7	3	T-test	< 0.0001	****	
Turnover rate (%)	8.767 ± 0.545	5	3	5.091 ± 0.728	7	3	T-test	0.0039	**	
<b>Fixed tissue at P15-P18</b>										
Spine density	1.731 ± 0.035	7	3	1.541 ± 0.061	8	3	T-test	0.023	*	
<b>GAD 67 expression</b>										
Mean intensity (% control group)	100 ± 3.36	28	9	86.62 ± 3.59	33	10	T-test	0.0094	**	
<b>PV cell distribution</b>		9	3		10	3	Two-way ANOVA	0.563	ns	F <sub>(1,68)</sub> = 0.338
<i>Layer II/III</i>	21.1 ± 1.4	9	3	20.4 ± 1.9	10	3	Sidak's post-hoc test		ns	
<i>Layer IV</i>	31.1 ± 2.3	9	3	29.5 ± 2.0	10	3	Sidak's post-hoc test		ns	
<i>Layer V</i>	33.8 ± 2.5	9	3	33.4 ± 1.5	10	3	Sidak's post-hoc test		ns	
<i>Layer VI</i>	11.8 ± 1.2	9	3	11.5 ± 1.0	10	3	Sidak's post-hoc test		ns	
<b>PV cell total number</b>	24.4 ± 1.5	9	3	23.7 ± 1.21	10	3	T-test	0.705	ns	

**Table S2. Summary table of imaging results**

N refers to the number of mice, L refers to the number of litters. All data are presented as mean ± SEM.

**Table S3**

	Control	n	N	L	C4-OE	n	N	L	Test	p value	Significant	F statistics (ANOVA)
<b>Intrinsic properties</b>												
Membrane resistance (M $\Omega$ )	129.4 $\pm$ 8.8	20	3	2	98.3 $\pm$ 7.0	21	4	2	Mann-Whitney test	0.0024	**	
Resting membrane potential (mV)	-68.0 $\pm$ 0.8	20	3	2	-68.6 $\pm$ 0.8	21	4	2	T-test	0.881	ns	
Membrane capacitance (pF)	151 $\pm$ 15	20	3	2	124 $\pm$ 14	21	4	2	Mann-Whitney test	0.127	ns	
Firing		20	3	2		21	4	2	Two-way ANOVA	C4 factor: 0.0129	*	F <sub>(1,37)</sub> = 6.82
										Depolarization: <0.0001	****	F <sub>(10,370)</sub> = 371.8
										Interaction: <0.0001	****	F <sub>(10,370)</sub> = 4.17
mEPSC Amplitude (pA)	13.8 $\pm$ 0.7	18	5	3	15.1 $\pm$ 0.6	16	6	3	Mann-Whitney test	0.069	ns	
mEPSC Frequency (Hz)	17.9 $\pm$ 1.4	18	5	3	27.9 $\pm$ 3.1	16	6	3	Mann-Whitney test	0.018	*	
<b>Excitatory PPR</b>	1.04 $\pm$ 0.06	24	4	2	1.12 $\pm$ 0.05	21	3	2	T-test	0.251	ns	
mIPSC Amplitude (pA)	13.9 $\pm$ 0.8	18	4	2	15.4 $\pm$ 1.4	13	4	3	T-test	0.312	ns	
mIPSC Frequency (Hz)	9.25 $\pm$ 0.97	18	4	2	10.8 $\pm$ 1.43	13	4	3	T-test	0.365	ns	

**Table S3. Summary table of patch-clamp results (PV interneurons)**

N refers to the number of mice, n refers to the number of neuronal recordings, L refers to the number of litters. All data are presented as mean  $\pm$  SEM.

**Table S4**

	Control	N	L	C4-OE	N	L	Test	p value	Significant	F statistics (ANOVA)
<b>Locomotion</b>										
Actimeter (repeated measures)		32	12		21	9	Two-way RM ANOVA	C4 factor: 0.765	ns	$F_{(1,51)} = 0.09$
								Time factor: < 0.0001	****	$F_{(11,561)} = 31.82$
								Interaction 0.366	ns	$F_{(11,561)} = 0.584$
Actimeter total activity	384 ± 21	32	12	396 ± 38	21	9	Mann-Whitney test	0.531	ns	
Open field (repeated measures)		20	10		14	8	Two-way RM ANOVA	C4 factor: 0.127	ns	$F_{(1,32)} = 2.458$
								Time factor: < 0.0001	****	$F_{(8,256)} = 4.250$
								Interaction 0.0037	**	$F_{(8,256)} = 2.975$
Open field total activity	2242 ± 121	20	10	1950 ± 140	14	8	Mann-Whitney test	0.189	ns	
<b>Working memory</b>										
Non-matching to sample		15	5		14	6	Two-way RM ANOVA	C4 factor: 0.556	ns	$F_{(1,135)} = 0.348$
								Time factor: < 0.0001	****	$F_{(1,135)} = 51.80$
								Interaction 0.364	ns	$F_{(4,135)} = 0.364$
Odor Span	8.40 ± 0.83	15	5	4.43 ± 0.54	14	6	Mann-Whitney test	0.0022	**	
Spatial Span	6.40 ± 0.73	15	4	4.21 ± 0.68	14	6	Mann-Whitney test	0.045	*	

**Table S4. Summary table of behavioral studies**

N refers to the number of mice, L refers to the number of litters. All data are presented as mean ± SEM.