

Elliptic blowup equations for 6d SCFTs. Part II. Exceptional cases

Jie Gu,^a Albrecht Klemm,^{b,c} Kaiwen Sun^d and Xin Wang^{b,e}

^a*Département de Physique Théorique et Section de Mathématiques, Université de Genève, Genève, CH-1211 Switzerland*

^b*Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

^c*Hausdorff Center for Mathematics, Universität Bonn, D-53115 Bonn, Germany*

^d*Scuola Internazionale Superiore di Studi Avanzati (SISSA), via Bonomea 265, 34136, Trieste, Italy*

^e*Max Planck Institute for Mathematics, Vivatsgasse 7, D-53111 Bonn, Germany*

E-mail: jie.gu@unige.ch, aklemm@th.physik.uni-bonn.de, ksun@sissa.it, wxin@mpim-bonn.mpg.de

ABSTRACT: The building blocks of 6d (1,0) SCFTs include certain rank one theories with gauge group $G = \text{SU}(3), \text{SO}(8), F_4, E_{6,7,8}$. In this paper, we propose a universal recursion formula for the elliptic genera of all such theories. This formula is solved from the elliptic blowup equations introduced in our previous paper. We explicitly compute the elliptic genera and refined BPS invariants, which recover all previous results from topological string theory, modular bootstrap, Hilbert series, 2d quiver gauge theories and 4d $\mathcal{N} = 2$ superconformal H_G theories. We also observe an intriguing relation between the k -string elliptic genus and the Schur indices of rank k H_G SCFTs, as a generalization of Del Zotto-Lockhart's conjecture at the rank one cases. In a subsequent paper, we deal with all other non-Higgsable clusters with matters.

KEYWORDS: Conformal Field Models in String Theory, Field Theories in Higher Dimensions, Solitons Monopoles and Instantons, Topological Strings

ARXIV EPRINT: [1905.00864](https://arxiv.org/abs/1905.00864)

Contents

1	Introduction	1
2	Elliptic non-compact CY 3-folds and generalised blowup equations	7
2.1	Geometry of elliptic fibrations	7
2.2	Semiclassical partition functions	10
2.3	Determination of r fields	13
2.4	De-affinisation	16
3	Elliptic blowup equations	18
3.1	Modularity of elliptic blowup equations	19
3.2	Universality of elliptic blowup equations	20
3.3	Unity blowup equations	22
3.3.1	Recursion formulas for elliptic genera	23
3.3.2	Uniqueness of recursion formulas	26
3.4	Vanishing blowup equations	27
4	Elliptic genera for 6d (1, 0) minimal SCFTs	29
4.1	Universal behaviors of elliptic genera	29
4.1.1	Universal expansion	29
4.1.2	Symmetric product approximation	32
4.1.3	Symmetries	34
4.2	Revisiting $G = \text{SU}(3)$ and $\text{SO}(8)$	35
4.3	$G = F_4$	38
4.4	$G = E_6$	41
4.5	$G = E_7$	45
4.6	$G = E_8$	48
5	On the relation with 4d SCFTs of type $H_G^{(k)}$	51
5.1	Rank k H_G theories	52
5.2	Hall-Littlewood and Schur indices	54
5.3	Rank one: Del Zotto-Lockhart's conjecture	57
5.4	Rank two	60
5.5	Rank three and higher	69
6	Conclusion and outlook	73
A	Lie algebraic convention	76
B	Mirror symmetry for elliptic non-compact Calabi-Yau three-folds	77
C	Geometric data	81

D Useful identities	82
E Relation with modular ansatz	84
F Elliptic genera	87
G Refined BPS invariants	94

1 Introduction

Six is the highest dimension in which representation theory allows for interacting superconformal quantum theories [1]. Limits of non-perturbative string theory compactifications [2] and in particular the decoupling of gravity in F-theory compactifications to 6d provided the first examples [3, 4] and lead recently to a complete classification of geometrically engineered 6d superconformal quantum field theories [5–7]. Such a classification in 6d is highly desirable, as it might lead by further compactifications, to an exhaustive classification of superconformal theories.

The 6d geometry is the one of an — in general desingularised — elliptic fibration with a contractable configuration of desingularised elliptic surfaces fibred over a configuration of curves in the base. In the decoupling limit the volume outside of the configuration of elliptic surfaces is scaled to infinite size, leaving us with an, in general reducible, configuration of complex desingularised elliptic surfaces that can be contracted within a non-compact Calabi-Yau threefold. Because compact components can be contracted such geometries are sometimes called local Calabi-Yau spaces. We will call the above specific ones for short *elliptic non-compact Calabi-Yau geometries* X and describe them in more detail in section 2.1.

The full topological string partition function on these elliptic non-compact CY geometries has received much attention as it contains important information about protected states of the 6d superconformal theories [3, 8]. Solving the topological string partition function on compact Calabi-Yau manifolds is currently an open problem. On non-compact Calabi-Yau spaces with an $U(1)_R$ isometry a refined topological string partition function $Z(\underline{t}, \epsilon_1, \epsilon_2)$, which depends on the Kähler parameters \underline{t} and two Ω background parameters ϵ_1, ϵ_2 is defined as generating function of refined stable pair invariants.¹ The refinement of the stable pair invariants [9, 10] and the relation to the refined BPS invariants $N_{j_l, j_r}^\beta \in \mathbb{N}$ was given in [11, 12]. Here $\beta \in H_2(X, \mathbb{Z})$ is the degree and the half integers (j_l, j_r) label a spin representation in the $SU(2)_l \times SU(2)_r$ little group of the 5d Poincaré group, which can be identified with Lefschetz actions on the moduli space of D2 and D0-branes. On

¹In this section we underline a symbol, if it is a vector. After the introduction section, we drop the underline when there is no risk of confusion.

toric non-compact Calabi-Yau spaces the refined partition function² $Z(\underline{t}, \epsilon_1, \epsilon_2)$ can be efficiently calculated by large N techniques [13],³ torus localisation [11], the integration of the refined holomorphic anomaly equations [14] and a recursive solution of blowup equations [15] generalized from the Göttsche-Nakajima-Yoshioka K-theoretic blowup equations in the context of 5d $\mathcal{N} = 1$ supersymmetric gauge theories [16–18].

The class of elliptic non-compact Calabi-Yau relevant for the $(1, 0)$ 6d SCFT is non-toric, but has a $U(1)_R$ isometry, and can be viewed as the borderline case for calculating $Z(\underline{t}, \epsilon_1, \epsilon_2)$. Since the techniques based on toric localisation and large N expansions fail, two related new methods have been developed. Similar as in heterotic/Type II duality one can calculate⁴ [8] the world-sheet elliptic genus of dual 2d quiver $(0, 4)$ gauge theories with supersymmetric localisation techniques [20, 21] leading to Jeffrey-Kirwan integrals. These elliptic genera $\mathbb{E}_d(\tau, \underline{a}, \underline{m}, \epsilon_1, \epsilon_2)$ transform as a Jacobi form and are identified with the topological string partition function Z_d at different winding d of the base [8] up to certain prefactor. The Kähler parameter τ of the elliptic fibre class becomes the modular parameter while (ϵ_1, ϵ_2) as well as Kähler parameters \underline{a} of the desingularisations and eventual further sections \underline{m} in the elliptic fibration become elliptic parameters. The refined holomorphic anomaly equations and other B-model techniques also apply and lead to a modular bootstrap approach where different winding contributions Z_d are identified with meromorphic Jacobi forms with weight zero and an index, which depends quadratically on the base degree d . The Z_d are so constrained by modularity, the pole — as well as the refined BPS structure of the topological string that they can be completely reconstructed in many examples [22–24].

In the 2d approach one needs for higher d to consider ever more complicated quiver gauge theories, while in the modular approach one has to deal with more and more complicated rings of weak Jacobi forms. For this reason we further develop in this paper the recursive approach based on the elliptic blowup equations [25] for the calculation of $Z(\underline{t}, \epsilon_1, \epsilon_2)$ that is further based on a specialisation of the generalized blowup equation in [15] to the elliptic non-compact Calabi-Yau geometries. The main advantage of this approach is that it needs as input only⁵ the classical topological data of X , i.e. the classical triple intersection numbers as well as the evaluation of the Chern classes on the elements of the Chow group, and yields with a non ambiguous efficient recursive procedure the string partition function iteratively in the base degree d and for each d exact in (ϵ_1, ϵ_2) and all other Kähler parameters.

²The holomorphic all genus partition function $Z(\underline{t}, \lambda) = \exp(\sum_{g=0}^{\infty} \lambda^{2g-2} F_g(\underline{t}))$ containing the information of all genus Gromov-Witten invariants is obtained as specialisation $\epsilon_1 = -\epsilon_2$ and $\lambda^2 = -\epsilon_1 \epsilon_2$.

³Strictly speaking the refined topological vertex applies directly only to geometries which engineer $\mathcal{N} = 2$ gauge theories, as these have the required preferred direction in the torus action. In blow downs and transitions of gauge theories geometries with Chern-Simons terms to geometries which have no immediate gauge theory interpretation, $Z(\underline{t}, \epsilon_1, \epsilon_2)$ for the latter can often be recovered [13].

⁴The more supersymmetric case of the M-strings has been pioneered in [19].

⁵This holds for all non-compact CY 3-folds studied in [15] and elliptic non-compact CY 3-folds studied in [25]. For the elliptic non-compact CY 3-folds associated to exceptional gauge symmetry studied in this paper, we also input Z_0 to the blowup equations, which can be easily calculated from the intersection and the multi-covering of isolated rational curves, see (2.3) below.

Let us first give a short summary of the structure behind the blowup equations in four, five and six dimensions. Non-compact Calabi-Yau spaces with $U(1)_R$ isometry and the (refined) topological string partition function feature prominently in the geometric engineering approach [26] to 5d and 4d supersymmetric gauge theories as $Z(\underline{t}, \epsilon_1, \epsilon_2)$ is related with the K-theoretic extension of Nekrasov’s 4d gauge theory instanton partition [27] on those non-compact Calabi-Yau spaces [16, 17], which do engineer supersymmetric gauge theories. In the geometric engineering approach, given mirror symmetry, it is physically obvious that world-sheet instantons and space time instantons are related. Simply because the former correct the topological string theory or $\mathcal{N} = 2$ supergravity prepotential, while the latter correct the rigid $\mathcal{N} = 2$ or Seiberg-Witten prepotential, which is related to the former in a well defined limit in the B-model, that decouples gravity as described in [26]. If the geometry engineers five dimensional $U(N)$ gauge theory, the full correspondence states that the K-theoretic partition function of the latter is identified with $Z(\underline{t}, \epsilon_1, \epsilon_2)$ and provides an alternative definition [16, 18]. The K-theoretic blowup equation for $U(N)$ theories without or with Chern-Simons terms has been rigorously established in [16] and [17, 18] respectively.

Nakajima and Yoshioka derived the original blowup equations [28] in the context of 4d $\mathcal{N} = 2$ supersymmetric $SU(N)$ framed gauge instanton calculus, by studying invariants on moduli space $\widehat{M}(N, k, n)$ of framed torsion free sheaves (E, Φ) on $\widehat{\mathbb{P}}^2$ — a \mathbb{P}^1 blowup of \mathbb{P}^2 — via the Atiyah-Bott localization formalism w.r.t. an induced toric action $T = \mathbb{C}^* \times \mathbb{C}^* \times \text{Gl}_N$ on $\widehat{M}(N, k, n)$. Here Φ is the framing automorphism, N is the rank of E , $n = \langle c_2(E) - \frac{N-1}{2N}c_1^2(E), [\widehat{\mathbb{P}}^2] \rangle$ and $k = -\langle c_1(E), [\mathbb{P}^1] \rangle$. A general feature of this calculation is that the Euler class of the tangent space of $\widehat{M}(N, k, n)$ at all relevant fix loci is always a product of contributions from two fixed points at the north and the south poles of the exceptional \mathbb{P}^1 , which arise due to the action of the $\mathbb{C}^* \times \mathbb{C}^*$ on $\widehat{\mathbb{P}}^2$ parametrized by ϵ_1 and ϵ_2 . Upon evaluation of the Atiyah-Bott localization formula the two contributions yield — up to calculable factors — a sum of products of the original partition function on \mathbb{P}^2 at shifted $\epsilon_{1,2}$ and Coulomb branch (in type IIA normalisable Kähler) parameters. The partition function on $\widehat{\mathbb{P}}^2$ can be also directly specialised for $k = 0$ to the one on \mathbb{P}^2 . The identification of the two results gives rise to a finite set of equations for the partition function of the 4d supersymmetric theory on the Omega background. A similar mechanism applies to the K-theoretic instanton calculus [16, 18], and leads to blowup equations for the partition function of the 5d SYM on the Omega background. The latter setup is directly relevant to the calculation of $Z(\underline{t}, \epsilon_1, \epsilon_2)$ on non-compact Calabi-Yau engineering supersymmetric gauge theories. The blowup equations can then be reformulated in terms of the geometric data of the non-compact Calabi-Yau X and refined topological string partition function as follows.

Let $C = (C_{ij})$ be the intersection matrix between compact divisor classes $[\mathcal{D}_i]$, $i = 1, \dots, b_4^c$ and compact curve classes $[\Sigma_j]$, $j = 1, \dots, b_2^c$ of X . Then one defines vector

$$\underline{R}_n = C \cdot \underline{n} + \underline{r}/2, \tag{1.1}$$

with $\underline{n} \in \mathbb{Z}^{b_4^c}$ and $\underline{r} \in \mathbb{Z}^{b_4^c}$ which parametrise the shift of the Kähler parameters. With $|\underline{n}| = \sum_{i=1}^{b_4^c} n_i$, the generalized blowup equations can be cast in the following form [15]

(see also section 8 of [23] and [29])

$$\sum_{\underline{n} \in \mathbb{Z}^{b_4^c}} (-1)^{|\underline{n}|} \widehat{Z}(\underline{t} + \epsilon_1 \underline{R}_{\underline{n}}, \epsilon_1, \epsilon_2 - \epsilon_1) \widehat{Z}(\underline{t} + \epsilon_2 \underline{R}_{\underline{n}}, \epsilon_1 - \epsilon_2, \epsilon_2) = \begin{cases} 0, & \underline{r} \in \mathcal{S}_v, \\ \Lambda(\epsilon_1, \epsilon_2, \underline{m}, \underline{r}) Z(\underline{t}, \epsilon_1, \epsilon_2) & \underline{r} \in \mathcal{S}_u. \end{cases} \tag{1.2}$$

Here we have separated the Kähler parameters \underline{m} from the Kähler parameters \underline{t} to denote those curve classes that do not intersect with compact divisors $[\mathcal{D}_k]$, $k = 1, \dots, b_4^c$. These \underline{m} correspond to mass parameters in the gauge theory context, while the other Kähler parameters correspond to Coulomb branch parameters, thus are also called “true” parameters. If a local mirror curve exists, e.g. for non-compact toric Calabi-Yau spaces, the “true” Kähler parameters are mapped to the complex structure parameters of the (hyperelliptic) mirror curve (of genus $g = b_4^c$) and the \underline{m} correspond to the residues of the meromorphic differential λ . The hat over Z means the Kähler moduli in the instanton partition function have already been shifted

$$\widehat{Z}(\underline{t}, \epsilon_1, \epsilon_2) = Z^{\text{cls}}(\underline{t}, \epsilon_1, \epsilon_2) Z^{\text{inst}}(\underline{t} + \pi i \underline{r}, \epsilon_1, \epsilon_2). \tag{1.3}$$

The integral vector \underline{r} , which we call the \underline{r} -field, is consistent with the checkerboard pattern of refined BPS invariants N_{j_l, j_r}^β , in other words, they satisfy

$$2j_l + 2j_r + 1 \equiv \underline{r} \cdot \beta \pmod{2} \tag{1.4}$$

for non-vanishing N_{j_l, j_r}^β . The set of \underline{r} -fields in (1.2) have to be only considered modulo $2C \cdot \underline{n}$, which leaves two classes of finite sets \mathcal{S}_v and \mathcal{S}_u . The \underline{r} -fields in these two sets are called *vanishing* and *unity* \underline{r} -fields. It is important that Λ , whose form is known, depends beside on $\epsilon_{1,2}$ only on \underline{r} -fields in the two classes and the mass parameters \underline{m} .

Given the simple form of (1.2) and the method of proof in the gauge theory context [16, 18], it seems reasonable to conjecture [15, 23, 25] that these equations, called the *generalised blowup equations*, should hold for the refined partition functions $Z(\underline{t}, \epsilon_1, \epsilon_2)$ of all non-compact Calabi-Yau threefolds with a global $U(1)_R$ symmetry so that the refined invariants or equivalently the corresponding BPS index for the space time theory with an Ω background can be defined [11, 12]. At the technical level the precise non-trivial claim is that $\mathcal{S}_v \cup \mathcal{S}_u$ should be non-empty. In addition it was observed in [15] that the classical topological data of X mentioned above and the genus zero sector *determine* $Z(\underline{t}, \epsilon_1, \epsilon_2)$ recursively, and many examples were already checked in great detail. In particular in [25] this approach was used to compute the refined BPS invariants of elliptic non-compact Calabi-Yau geometries associated to minimal 6d (1, 0) SCFTs with gauge group $G = SU(3), SO(8)$. In this paper we extend this approach to the remaining minimal 6d SCFTs with exceptional gauge groups $G = F_4, E_{6,7,8}$ and give a universal description for all minimal building blocks without matter in the classification of 6d (1, 0) SCFTs [5, 6].

The elliptic non-compact Calabi-Yau geometry corresponding to minimal SCFTs with gauge group G and no matter contains base surface $\mathcal{O}(-\mathbf{n}) \rightarrow \mathbb{P}^1$. For $\mathbf{n} = 3, 4, 5, 6, 8, 12$ which are of interest in this paper, the Kodaira singularity of the elliptic fibration gives

the gauge group $G = \text{SU}(3), \text{SO}(8), F_4, E_{6,7,8}$ respectively. We find that for these geometries, the generalized blowup equations can be uniformly written as the following recursive relations of the elliptic genera \mathbb{E}_d of the corresponding 6d SCFT:

$$\begin{aligned}
 & \sum_{\underline{\omega} \in \phi_\lambda(Q^\vee), d_{1,2} \in \mathbb{N}}^{\frac{1}{2} \|\underline{\omega}\|^2 + d_1 + d_2 = d} (-1)^{|\phi_\lambda^{-1}(\underline{\omega})|} \theta_i^{[a]} \left(\mathbf{n}\tau, (\mathbf{n}-2)(\epsilon_1 + \epsilon_2) - \mathbf{n} \left(\left(\frac{1}{2} \|\underline{\omega}\|^2 + d_1 \right) \epsilon_1 + \left(\frac{1}{2} \|\underline{\omega}\|^2 + d_2 \right) \epsilon_2 - \underline{m} \cdot \underline{\omega} \right) \right) \\
 & \quad \times A_\omega(\underline{m}) \mathbb{E}_{d_1}(\tau, \underline{m} - \epsilon_1 \underline{\omega}, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_{d_2}(\tau, \underline{m} - \epsilon_2 \underline{\omega}, \epsilon_1 - \epsilon_2, \epsilon_2) \\
 & = \begin{cases} 0, & d \notin \mathbb{N}, \\ \theta_i^{[a]}(\mathbf{n}\tau, (\mathbf{n}-2)(\epsilon_1 + \epsilon_2)) \cdot \mathbb{E}_d(\tau, \underline{m}, \epsilon_1, \epsilon_2), & d \in \mathbb{N}. \end{cases} \tag{1.5}
 \end{aligned}$$

Here \underline{m} are the Coulomb parameters⁶ associated to gauge group G . The subscript of theta functions i is 4 if \mathbf{n} is odd and 3 if \mathbf{n} is even, and the characteristic $a = k/\mathbf{n} - 1/2$, $k = 0, 1, \dots, \mathbf{n} - 1$. Besides, ϕ_λ is an embedding of the coroot lattice Q^\vee of G into the weight lattice P . Here the \underline{r} -field is implicit in a and ϕ_λ . Since the number of different embeddings is $|P : Q^\vee|$, the total number of non-equivalent blowup equations is $\mathbf{n}|P : Q^\vee|$. The function $A_\omega(\underline{m})$ is composed of θ_1 and η functions, see (3.5) for the definition. Due to the Jacobi form nature of every component of the above equations, we call (1.5) as *elliptic blowup equations*. In fact, they can be regarded as the natural elliptic lift of the K-theoretic blowup equations for 5d gauge theories [16, 30]. Moreover, the unity elliptic blowup equations in (1.5) ultimately lead to a complete solution of the elliptic genera \mathbb{E}_d in terms of an universal recursion formula, as will be shown in (3.32). The blowup equations are not only effective tools to calculate the refined partition functions, but also together with the general constraints from modularity and BPS structure shed some new light on the structure of $Z(\underline{t}, \epsilon_1, \epsilon_2)$. In particular it is possible to derive from the structure of the blowup equations the index and the weight of the Jacobi forms that constitute the building blocks in (1.5). The recursive structure also helps clarify the form of the denominator of elliptic genus in the modular bootstrap approach as discussed in appendix E, and predicts many non-trivial relations among these Jacobi forms, one particular of which is proven in section 3.3.1.

In the program of classifying superconformal field theories in various dimensions the 6d SCFTs play a similar role as 11d $\mathcal{N} = 1$ supergravity or more precisely M-theory play for the classification of supergravity in lower dimensions. For this reason we expect that various limits as well as suitable expansion of the partition function $Z(\underline{t}, \epsilon_1, \epsilon_2)$ of the 6d (1, 0) minimal SCFTs relate to the protected quantities in lower dimensional supersymmetric theories.

Since the elliptic blowup equations determine $Z(\underline{t}, \epsilon_1, \epsilon_2)$ in particular the elliptic genus \mathbb{E}_d completely, we could make many detailed and indeed successful checks on our results. We summarise the current status of the knowledge on the elliptic genera of all 6d (1, 0) minimal SCFTs from various approaches in table 1. For $\mathbf{n} = 5, 6, 8, 12$ which are of main interest in the current paper, our complete recursive solution for the elliptic genera from blowup equations reproduces *all* previous partial results. Since we made many checks of

⁶Do not confuse with \underline{m} in (1.2). The Coulomb parameters are *not* mass parameters.

n	1	2	3	4	6	8	12	5	7
features	E-strings	M-strings	SU(3)	SO(8)	E_6	E_7	E_8	F_4	$E_7 + \frac{1}{2}\mathbf{56}$
2d quiver	\mathbb{E}_k [32, 33]	\mathbb{E}_k [19]	\mathbb{E}_k [34]	\mathbb{E}_k [8]	?				
B-model	low genus [35]	?	genus zero [8]						
modular bootstrap	\mathbb{E}_k [23, 36]		\mathbb{E}_1 [24, 37]	\mathbb{E}_1 with fugacities off [37, 38]					
topological vertex	\mathbb{E}_k [39]	\mathbb{E}_k [19]	$\mathbb{E}_k^{q \rightarrow 0}$ [40]					?	
Hilbert series	-	-	$\mathbb{E}_k^{q \rightarrow 0}$ [41–43]						-
HL index	-	-	?	$\mathbb{E}_k^{q \rightarrow 0}$ [44, 45]				-	
twisted H_G theories	-	-	?	\mathbb{E}_1 [37, 46, 47]	\mathbb{E}_1 [48]	?	-		
domain walls	$\mathbb{E}_{1,2}$ [49] \mathbb{E}_3 [50]	\mathbb{E}_k [49]	-						
5d blowup equations	trivial		$\mathbb{E}_k^{q \rightarrow 0}$ [30]					$\mathbb{E}_k^{q \rightarrow 0}$ [51]	
6d blowup equations	\mathbb{E}_k [15, 23, 52]	\mathbb{E}_k [52]	\mathbb{E}_k [25]	\mathbb{E}_k , current paper				\mathbb{E}_k [51]	

Table 1. Known results on the elliptic genera of 6d minimal (1,0) SCFTs from various approaches. Here $q = Q_\tau = e^{2\pi i \tau}$ is the modular parameter. - means the method does not apply, and ? means possible applicable but results not yet attained. HL means Hall-Littlewood.

the elliptic blowup equations based on extensive calculations, which might yield further insights, we provide the results of these calculations on a webpage [31].

Part of these checks indicated in table 1 are quite obvious as for example the 5d limit gives a good confirmation of our results. Others are highly non-trivial and indicate new exciting connections to the protected quantities in lower dimensional theories. For example one of the most important tools for the analysis of the spectra and phenomena like Seiberg duality in four dimensional SCFTs are the superconformal indices for $\mathcal{N} = 1, 2, 4$ SCFT, which count operators in the chiral rings of these theories. These indices have in turn various limits such as Macdonald indices, Hall-Littlewood indices and Schur indices which are relatively easy to compute. As explained in section 5 the latter two occur in a quite non-trivial manner in the expansion of the elliptic genera that we can efficiently calculate. This surprising relation between elliptic genera and superconformal indices was found for the rank one H_G theories in [37]. We will push the study on such relation for all rank two and even some rank three cases. This not only sheds light on the structure of these objects, but also allows to calculate them efficiently for example in theories with no Lagrangian description in which other methods are quite difficult to carry through.

This paper is organized as follows: in section 2, we review the geometric construction of elliptic non-compact Calabi-Yau threefolds that engineer 6d (1,0) minimal SCFTs, the basic properties of the generalized blowup equations in [15], and the de-affinisation procedure which was essentially already used in [25] to obtain elliptic blowup equations for $G = \text{SU}(3)$ and $\text{SO}(8)$. In section 3, we discuss both unity and vanishing elliptic blowup equations in detail, and derive a universal recursion formula for the elliptic genera of all minimal SCFTs with $G = \text{SU}(3), \text{SO}(8), F_4, E_{6,7,8}$. We also prove two important properties of the elliptic blowup equations, i.e. modularity and universality. In section 4, we explicitly show for each G the one- and two-string elliptic genera computed from our universal recur-

sion formula and also some relevant information for the blowup equations, such as triple intersection numbers and the r -fields. In section 5, we discuss a surprising relation between the elliptic genera of 6d minimal SCFTs and the Hall-Littlewood indices and Schur indices of 4d $\mathcal{N} = 2$ superconformal H_G theories, as was revealed for rank one in [37]. We find analogous relation indeed exist for rank two and higher. Finally, in section 6, we discuss various possible application and future directions. In a series of appendices we explain our convention, some technical details, and collect more results on elliptic genera and refined BPS invariants too lengthy to be put in the main text.

2 Elliptic non-compact CY 3-folds and generalised blowup equations

The *generalised blowup equations* proposed in [15] (see also section 8 of [23] and [29]) generalise the K-theoretic blowup equations of Nakajima and Yoshioka [16, 18] for 5d SYM theories to all non-compact Calabi-Yau geometries that have a $U(1)_R$ isometry which may or may not engineer 5d supersymmetric field theories. In subsection 2.1 we describe the geometric data of the non-compact elliptic Calabi-Yau threefolds associated to the minimal 6d SCFTs with pure gauge bulk theory. With this input and b_i^{ns} calculated in subsection 2.2, the r fields can be determined in subsection 2.3 and the generalised blowup equations can be expanded to extract BPS constraints that allow for solution of refined BPS invariants as in [15].

We then consider the expansion of the partition function in the base degrees and describe how to recast the generalised blowup equations, with some additional input, as functional equations of elliptic genera of the 6d SCFTs. The latter, which we call the *elliptic blowup equations*, will be discussed in full detail in the next section.

2.1 Geometry of elliptic fibrations

In this paper we are specifically interested in the non compact elliptic Calabi-Yau threefolds on which F-theory compactification yields minimal 6d $(1, 0)$ SCFTs with $\mathfrak{n} = 3, 4, 5, 6, 8, 12$ so that the bulk theory has a pure gauge group G .

We discuss some generic features of these Calabi-Yau threefolds, in particular the compact curves and the compact divisors in the Calabi-Yau. The compact curves and compact divisors in these geometries are best illustrated in [53] and summarised in [25]. Let us go over them here quickly. Suppose the gauge group G has rank $\text{rk}(G)$ and its associated Lie algebra is \mathfrak{g} , then there are $b_2^c = \text{rk}(G) + 2$ linearly independent compact curves in the Calabi-Yau threefold.⁷ One of them is the $-\mathfrak{n}$ curve Σ_B in the base, while the remaining $\text{rk}+1$ curves Σ_I ($I = 0, 1, \dots, \text{rk}$) are \mathbb{P}^1 s resulting from resolution of the singular elliptic fiber fibered over Σ_B . These $r + 1$ curves intersect with each other according to the affine Dynkin diagram of \mathfrak{g} , where each node corresponds to a rational curve of self-intersection -2 and two curves intersect with intersection number 1 if the corresponding nodes are linked. We denote the curve corresponding to the affine node by Σ_0 , and it is

⁷From now on, we simply denote $\text{rk}(G)$ as rk to lighten the notation.




Figure 1. Compact curves and compact divisors in the $n = 5$ local elliptic Calabi-Yau from [53]. The indices $(-n, n - 2)$ on each rational section S or $S' = S + kF$ of \mathbb{F}_k are the degrees of its normal bundles $\mathcal{O}(-n) \oplus \mathcal{O}(n - 2)$ in the corresponding direction. \mathbb{F}_6 meets \mathbb{F}_1 in a double section. Globally the fibration has a \mathbb{Z}_2 monodromy encircling Σ_B that corresponds to an outer automorphism of the E_6 Dynkin diagram and folds its sphere tree to an F_4 type sphere tree over Σ_B . Note that the curve Σ_b plays the role of a Mori cone generator and is related to the base curve Σ_B by (4.43).

the only \mathbb{P}^1 which intersects with Σ_B . The linear combination

$$\sum_{I=0}^{\text{rk}} a_I [\Sigma_I] = [\delta] \tag{2.1}$$

with a_I the marks of $\hat{\mathfrak{g}}$ is homologous to the generic elliptic fiber. We denote the complexified Kähler parameters of Σ_B and Σ_I by t_B and t_I respectively.

The $b_4^c = \text{rk} + 1$ vertical compact divisors \mathcal{D}_I for $I = 0, 1, \dots$ are fibrations of Σ_I over Σ_B . They are argued in [53] to be Hirzebruch surfaces of various degrees, and the Σ_I are the \mathbb{P}^1 fibers of these Hirzebruch surfaces. It is then easy to deduce that the $(\text{rk} + 2) \times (\text{rk} + 1)$ matrix C encoding the intersections between Σ_I, Σ_B and \mathcal{D}_I is given by

$$C = \begin{pmatrix} & -\hat{A} \\ -\mathbf{n}, & 0, \dots, 0 \end{pmatrix} \tag{2.2}$$

where \hat{A} is the affine Cartan matrix of G . We illustrate compact curves and compact divisors in the example of the $n = 5$ model with $G = F_4$ in figure 1.

The topological string partition function on elliptic fibrations can be expanded in terms of the base degree d w.r.t. the base curve Σ_B labelled by $Q_B = e^{t_B}$

$$Z = Z^{\text{cls}} Z_0 \left(1 + \sum_{d=1}^{\infty} Z_d Q_B^d \right). \tag{2.3}$$

Z^{cls} comes from the degree zero maps and depends hence on the classical topological data of X . We discuss them in section 2.2. $Z_0(t_I, \epsilon_-, \epsilon_+)$ gets contributions from the rational curves Σ_I in the elliptic fibre that form the affine Dynkin diagram. These can be directly calculated from the geometry reflecting the affine group structure, i.e. the intersection matrix \hat{A} in (2.2) using $N_{0, \frac{1}{2}}^{[\Sigma_I]} = 1, I = 0, \dots, r$ for rational curves [11] as well as the general multi cover formula (3.21), which leads to (2.36). These contribution of isolated rational curves can be also calculated as one loop correction to the gauge coupling [26], which is the reason that Z_0 is sometimes identified as $Z_0 = Z^{1\text{-loop}}$. The coefficients $Z_{d>0}$ in the expansion, on the other hand, encode the BPS invariants that do wrap the base curve Σ_B , and are rather difficult to compute.

At this point a clarification of subtlety is in order. The curve classes Σ_B, Σ_I actually do not give a good basis for computing the BPS invariants, as they are not all Mori cone generators. To remedy this, one should keep Σ_I and replace Σ_B by the \mathbb{P}^1 base of the Hirzebruch surface with the lowest degree in the chain, so that it cannot be expressed as linear combinations of other curves with non-negative coefficients. We will illustrate this point in example section 4.

The topological string partition function is identical with the BPS partition function of the corresponding 6d SCFT in the tensor branch, put on the Omega background $\mathbb{R}^4 \times_{\epsilon_1, \epsilon_2} T^2$. In the latter point of view, it is more natural to use another set of Kähler parameters $t_{\text{ell}}, \tau, m_i$ ($i = 1, \dots, \text{rk}$), which are related to t_B, t_I by

$$t_{\text{ell}} = t_B - \frac{\mathbf{n} - 2}{2} \tau, \quad \tau = \sum_{I=0}^{\text{rk}} a_I t_I, \quad m_i = t_i, \quad i = 1, \dots, \text{rk}. \quad (2.4)$$

t_{ell} is defined such that the coefficients \mathbb{E}_d in the expansion of the BPS partition function in terms of $Q_{\text{ell}} = e^{t_{\text{ell}}}$

$$Z = Z^{\text{cls}} Z^{1\text{-loop}} \left(1 + \sum_{d=1}^{\infty} \mathbb{E}_d Q_{\text{ell}}^d \right) \quad (2.5)$$

are elliptic genera of the self-dual strings present in the 6d SCFT. τ measures the volume of the generic elliptic fiber $[\delta]$, and since δ intersect with no compact divisor, it is a mass parameter of the theory. In addition, it is also identified with the complex structure modulus of the torus T^2 . m_i are now interpreted as the Wilson loops of the vector multiplets in T^2 .

Note that the 6d SCFT can be reduced to a 5d pure SYM with the same gauge group G if we decompactify Σ_0 and send its volume $-t_0$ to infinity. In the resulting 5d theory, the only mass parameter is the instanton counting parameter t_q , and we find, by looking for curve class not intersecting with divisors, that

$$t_q = t_{\text{ell}} - \frac{\mathbf{n} - 2}{2} \tau. \quad (2.6)$$

We finally comment that in light of the correspondence between m_i and nodes in Dynkin diagram of \mathfrak{g} , we can collect the Kähler moduli m_i into a single vector m taking value in the complexified Cartan subalgebra $\mathfrak{h}_{\mathbb{C}} = \mathbb{C}^{\text{rk}}$ with

$$m = \sum_{i=1}^{\text{rk}} m_i \omega_i, \quad m_i = \langle \alpha_i, m \rangle, \quad i = 1, \dots, \text{rk}, \quad (2.7)$$

where ω_i are the fundamental weights, α_i the simple roots of \mathfrak{g} , and \langle, \rangle the natural pairing between $\mathfrak{h}_{\mathbb{C}}$ and $\mathfrak{h}_{\mathbb{C}}^*$. This allows a reformulation of the generalised blowup equations in terms of Lie algebraic data, which we use heavily in the uniform formula (1.5). The convention of Lie algebra we use is given in appendix A.

2.2 Semiclassical partition functions

We summarise the computation of the semiclassical partition functions here. These are the minimal initial data one needs in order to extract refined BPS invariants from the generalised blowup equations. First of all the semiclassical contribution $Z^{\text{cls}}(t, \epsilon_1, \epsilon_2) = \exp(F^{\text{cls}}(t, \epsilon_1, \epsilon_2))$ can be written as

$$\begin{aligned}
 F^{\text{cls}}(t, \epsilon_1, \epsilon_2) &= \frac{1}{\epsilon_1 \epsilon_2} F_{(0,0)}^{\text{cls}}(t) + F_{(1,0)}^{\text{cls}}(t) - \frac{(\epsilon_1 + \epsilon_2)^2}{\epsilon_1 \epsilon_2} F_{(0,1)}^{\text{cls}}(t) \\
 &= \frac{1}{\epsilon_1 \epsilon_2} \left(\frac{1}{6} \sum_{i,j,k=1}^{b_2^c} \kappa_{ijk} t_i t_j t_k \right) + \sum_{i=1}^{b_2^c} b_i^{\text{GV}} t_i - \frac{(\epsilon_1 + \epsilon_2)^2}{(\epsilon_1 \epsilon_2)} \sum_{i=1}^{b_2^c} b_i^{\text{NS}} t_i .
 \end{aligned}
 \tag{2.8}$$

The coefficients κ_{ijk} are the triple intersection numbers of divisors J_i Poincaré dual to the curve classes Σ_i with volumes t_i . b_i^{GV} are intersections of the divisors J_i with the second Chern class of the Calabi-Yau threefold. The coefficients b_i^{NS} , on the other hand, do not have a geometric meaning, they are usually computed by the refined holomorphic anomaly equations [14, 54], which are difficult to apply here. The Nekrasov partition function of a 5d pure SYM also has a semiclassical contribution which takes the same form as (2.8), and the linear coefficients are subject to the relation

$$b_i^{\text{GV}} + b_i^{\text{NS}} = 0 .
 \tag{2.9}$$

In our previous paper [25] we argued that $b_i^{\text{GV}}, b_i^{\text{NS}}$ of the minimal 6d SCFTs with $G = \text{SU}(3), \text{SO}(8)$ can be computed by uplifting the semiclassical Nekrasov partition function of the 5d pure SYM with the same gauge G aided by the nontrivial automorphism of the affine Dynkin diagrams, and we found these coefficients also satisfy the relation (2.9). In the remaining minimal SCFTs with $G = F_4, E_6, E_7, E_8$, not all the affine Dynkin diagrams have a non-trivial automorphism, and the method of uplifting does not always work. Instead we assume (2.9) to be true and only compute b_i^{GV} by geometric means.⁸

To compute κ_{ijk} and b_i^{GV} , we need to embed the Calabi-Yau threefold $X \rightarrow \mathcal{O}_{\mathbb{P}^1}(-n)$ in a compact Calabi-Yau, for instance the elliptic fibration \hat{X} over \mathbb{F}_n with a single section, and first compute these intersection numbers in the compact geometry. The compact Calabi-Yau \hat{X} can be realised as a hypersurface in a toric variety [8]. Let us look at an example in detail. The threefold $\pi : X \rightarrow \mathcal{O}_{\mathbb{P}^1}(-5)$ is the zero loci of the section of the anti-canonical bundle of the toric variety \mathbb{P}_{Δ} , whose toric data is given in table 2 [8].⁹ It has 7 Mori cone generators with charges $l^{(i)} = (l_n^{(i)})$ for $i = 1, \dots, 7$ (We will use the same notation

⁸Up to a irrelevant τ term, the numbers $b_i^{\text{GV}}, b_i^{\text{NS}}$ can be also predicted from blowup equations by requiring the consistency of BPS invariants. For all the minimal SCFTs, these predictions agree with the values computed from the method we will describe later.

⁹See also the geometric description in [55].

D	ν_i^*				$l^{(1)}$	$l^{(2)}$	$l^{(3)}$	$l^{(4)}$	$l^{(5)}$	$l^{(6)}$	$l^{(7)}$	$l_{F_4}^{(0)}$	$l_{F_4}^{(1)}$	$l_{F_4}^{(2)}$	$l_{F_4}^{(3)}$	$l_{F_4}^{(4)}$	$l_{F_4}^{(b)}$
D_0	0	0	0	0	-2	0	0	0	0	0	0	0	0	0	0	-2	0
D_1	-1	0	0	0	0	-2	0	0	0	0	1	1	0	0	0	0	0
D_2	0	-1	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0
D_3	0	1	0	-1	0	3	0	0	0	1	-2	-2	1	0	0	0	0
D_4	1	2	0	-2	2	0	0	0	0	-2	1	1	-2	0	0	2	0
S'	2	3	0	-1	0	1	-2	0	1	0	0	0	0	-2	1	0	0
S''	2	3	0	-2	1	-2	1	-1	0	0	0	0	0	1	-2	1	-1
S'''	2	3	0	-3	-2	0	0	-1	0	1	0	0	1	0	1	-2	-1
K	2	3	0	0	0	0	1	0	-2	0	0	0	0	1	0	0	0
F	2	3	-1	-5	0	0	0	1	0	0	0	0	0	0	0	0	1
S	2	3	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0
F	2	3	1	0	0	0	0	1	0	0	0	0	0	0	0	0	1

Table 2. The toric data of \mathbb{P}_Δ for $G = F_4$.

for both the curves and their toric charges). Note the number of Mori cone generators is $(\text{rk} + 3)$. Among these curves $l^{(4)}, l^{(5)}$ are identified as the (-5) curve (i.e. Σ_B) and the (0) curve in the base \mathbb{F}_5 ,¹⁰ while the other five toric curves combine linearly into Σ_I for $I = 0, 1, \dots, 4$ [8]

$$\Sigma_0 = l^{(3)}, \quad \Sigma_1 = l^{(2)} + l^{(6)} + 2l^{(7)}, \quad \Sigma_2 = l^{(1)}, \quad \Sigma_3 = l^{(6)}, \quad \Sigma_4 = l^{(7)}. \quad (2.10)$$

The toric divisors listed in the first column of table 2 can also be identified. D_0 is associated to the canonical bundle of \mathbb{P}_Δ . When \hat{X} is written in the Weierstrass form, D_1, D_2 correspond to the divisors $x = 0$ and $y = 0$ respectively, while K is the zero section at $x \rightarrow \infty$. F, S', S are respectively the vertical divisors pulled back from the $(0), (-5)$, and the (5) curves in \mathbb{F}_5 in the base. Since the elliptic fibration over $\pi(S') = \Sigma_B$ factorises to an intersecting tree of Hirzebruch surfaces \mathfrak{D}_I for $I = 0, 1, \dots, 4$, S' should actually be identified with \mathfrak{D}_0 . The remaining four divisors D_3, D_4, S'', S''' are identified (up to linear combination) as the exceptional divisors \mathfrak{D}_i for $i = 1, \dots, 4$. Only $\text{rk}(F_4) + 3 = 7$ of these divisors are linearly independent.

Once the toric data of \mathbb{P}_Δ are specified, there are standard techniques in toric geometry to compute the triple intersection numbers $\hat{\kappa}_{ijk}$ of the divisors \hat{J}_i , and the intersection numbers with $c_2(\hat{X})$ are given by [56]¹¹

$$-24\hat{b}_i^{\text{GV}} = \int_{\hat{X}} c_2(\hat{X}) \wedge \hat{J}_i = \frac{1}{2} \sum_{jk} \hat{\kappa}_{ijk} (l_0^{(j)} l_0^{(k)}) - \sum_{n>0} l_n^{(j)} l_n^{(k)}. \quad (2.11)$$

¹⁰Strictly speaking, the (0) curve should be corrected by certain linear combination of other curves. But after decompactifying this curve to arrive at the non-compact threefold X , this difference disappears.

¹¹We use the same notation for the divisor \hat{J}_i and its dual 2-form.

Alternatively, $\hat{\kappa}_{ijk}$ can be computed by the special geometry relation. For a compact Calabi-Yau threefold realised as a hypersurface in a toric variety one can define the deformed fundamental period $\hat{\omega}_0(z; \rho)$ as a holomorphic function of the Batyrev coordinates z_i and compute the homogeneous A- and B-periods (see for instance [57])

$$\begin{aligned}\hat{\Pi}_i^{(1)}(z) &= \partial_{\rho_i} \hat{\omega}_0(z; \rho)|_{\rho_i=0}, \\ \hat{\Pi}_i^{(2)}(z) &= \frac{1}{2} \hat{\kappa}_{ijk} \partial_{\rho_j} \partial_{\rho_k} \hat{\omega}_0(z; \rho)|_{\rho_i=0}.\end{aligned}\tag{2.12}$$

They are interpreted as the masses of the D2-, D4-branes supported on the curves $l^{(i)}$ and the dual divisors \hat{J}_i . The affine A- and B-periods defined by

$$\hat{t}_i = \hat{\Pi}_i^{(1)} / \hat{\omega}_0(z; 0), \quad \hat{F}_i = \hat{\Pi}_i^{(2)} / \hat{\omega}_0(z; 0)\tag{2.13}$$

satisfy the relation

$$\hat{F}_i = \frac{\partial \hat{F}_{(0,0)}}{\partial \hat{t}_i}.\tag{2.14}$$

The existence of the prepotential $\hat{F}_{(0,0)}$ uniquely fixes the coefficients $\hat{\kappa}_{ijk}$.

The non-compact Calabi-Yau X is obtained by decompactifying \hat{X} in the direction of the (0)-curve in the base. In practise, this corresponds to taking the Kähler parameter in the decompactified direction t_{dc} to infinity in A-model or taking the corresponding complex structure parameter z_{dc} to zero in B-model. In A-model, this limit can be understood as taking some of the compact (1, 1) cycles to infinite size, keeping the other (1, 1) cycles finite. The periods of the geometry will be rearranged so that only one A-period and some B-periods go to infinity. In our current case, the B-period $\frac{\partial}{\partial \tau} F_{(0,0)}$ goes to infinity, while the corresponding A-period remains finite, and becomes the elliptic fiber parameter τ . We can then integrate over the new periods to get the triple intersection numbers of the non-compact geometry. However, this method will always have a integration constant term τ^3 unfixed, which is very important for the refined BPS invariants in the τ direction. To determine the τ^3 term, we study the normalization scheme of τ derivative of the genus zero free energy $\frac{\partial}{\partial \tau} F_{(0,0)}$.

In the example of the $G = F_4$ model visited above, this is z_5 associated to the curve $l^{(5)}$. In the limit $z_{dc} \rightarrow 0$ the affine A-period associated to the (0)-curve diverges, while the other $\text{rk} + 2$ affine A-periods remain finite. We can choose a basis of the latter to be¹²

$$(t_I), \quad I = 0, 1, \dots, \text{rk}, B,\tag{2.15}$$

which correspond to the curve classes $\Sigma_0, \dots, \Sigma_r, \Sigma_B$ discussed in the previous section. At the same time, both the zero section of the elliptic fibration and the vertical divisor of the (0)-curve become infinite in volume. Therefore in the limit $z_{dc} \rightarrow 0$ two B-periods diverge, and only $\text{rk} + 1$ affine B-periods remain finite. We choose a basis

$$(F_J), \quad J = 0, 1, \dots, \text{rk},\tag{2.16}$$

¹²They are the limit $z_{dc} \rightarrow 0$ of proper linear combinations of \hat{t}_i of the compact Calabi-Yau. Similarly F_J defined below are the limit $z_{dc} \rightarrow 0$ of linear combinations of \hat{F}_i .

which correspond to the divisor classes \mathfrak{D}_I with $I = 0, 1, \dots, \text{rk}$. The special geometry relation of the non-compact Calabi-Yau X dictates

$$F_J = \sum_{I=0, \dots, \text{rk}, B} C_{JI} \frac{\partial F_{(0,0)}}{\partial t_I}, \quad J = 0, 1, \dots, \text{rk}, \quad (2.17)$$

where C_{JI} are the components of the divisor-curve intersection matrix (2.2). The identity (2.17) allows the computation of the semi-classical components of $F_{(0,0)}(t)$ up to a term proportional to τ^3 ; in other words, the intersection number $\kappa_{\tau\tau\tau}$ can not be fixed by (2.17). Since τ is a mass parameter, the term τ^3 can always be factored out of the blowup equations and it is not of importance to us. Nevertheless, in appendix B we will introduce a normalisation scheme which fixes such a term in a reasonable way, and we adopt this normalisation scheme in the example section 4. In any case, for all the minimal 6d SCFTs with a pure gauge theory in the bulk which can be reduced to a 5d pure SYM, we find that up to τ^3

$$F_{(0,0)}^{\text{cls}}(t) = - \sum_{\alpha \in \Delta_+} \left(\frac{\langle \alpha, m \rangle^3}{6} + \frac{t_q}{2h_G^V} \langle \alpha, m \rangle^2 \right) - \frac{((n-2)\tau + t_q)^3 - t_q^3}{6n(n-2)} + \dots \quad (2.18)$$

where m, t_q are defined in (2.7) and (2.6) respectively. We recognise the sum over positive roots is from the Nekrasov prepotential of the 5d pure SYM. We can also massage (2.18) into a more suggestive form

$$F_{(0,0)}^{\text{cls}} = \frac{1}{2n} t_{\text{ell}}^2 \tau + \frac{1}{2} t_{\text{ell}}(m, m) - \frac{n-2}{4} \tau(m, m) + \dots \quad (2.19)$$

up to τ^3 and cubic terms in m_i . Here $(,)$ is the invariant bilinear form on $\mathfrak{h}_{\mathbb{C}}$. See appendix A for our convention.

As for $F_{(1,0)}^{\text{cls}}(t)$ and the intersection numbers with $c_2(X)$, we use the same formula (2.11) with $\hat{\kappa}_{ijk}$ replaced by the triple intersection numbers of the non-compact Calabi-Yau and $l^{(i)}$ replaced by the toric charges of $\Sigma_0, \dots, \Sigma_r, \Sigma_B$. In the example of $G = F_4$ discussed above, one has the toric charges $l_{F_4}^{(0)}, l_{F_4}^{(1)}, l_{F_4}^{(2)}, l_{F_4}^{(3)}, l_{F_4}^{(4)}, l_{F_4}^{(b)}$ as in table 2.

As in the case of the prepotential, one cannot determine the pure mass term proportional to τ which is irrelevant, although it can be fixed by the same normalisation scheme if one wishes. We have checked that $F_{(1,0)}^{\text{cls}}(t)$ computed in this way reduces correctly to the semiclassical Nekrasov partition function when the 6d SCFT is reduced to the 5d pure SYM.

2.3 Determination of r fields

In general, the r fields associated to a non-compact Calabi-Yau can be determined by the method in [15]. Here we give a brief description of it. As proved in [15], even without any assumption or constraint put on Λ (for instance it can depend on all Kähler moduli), the Λ as *defined* by (1.2) must be quasi-modular of weight zero under the Siegel modular transformations of $\tau_{ij} = \frac{\partial^2}{\partial t_i \partial t_j} F_{(0,0)}$, where t_i are the true Kähler parameters.¹³ Let us

¹³Note that this τ_{ij} parameter is different from the τ parameter of the elliptic fiber.

expand this Λ in terms of all the exponentiated Kähler moduli e^{t_i} . The leading terms, which come from Z^{cls} , read

$$\begin{aligned} \log \Lambda &\sim \log \left(Z^{\text{cls}}(\epsilon_1, \epsilon_2 - \epsilon_1) Z^{\text{cls}}(\epsilon_1 - \epsilon_2, \epsilon_2) / Z^{\text{cls}}(\epsilon_1, \epsilon_2) \right) \\ &= \left(-\frac{1}{6} \sum_{i,j,k=1}^{b_2^c} \kappa_{ijk} R_i R_j R_k + \sum_{i=1}^{b_2^c} (b_i^{\text{GV}} - b_i^{\text{NS}}) R_i \right) (\epsilon_1 + \epsilon_2) + \sum_{k=1}^{b_2^c} \left(-\frac{1}{2} \sum_{i,j=1}^{b_2^c} \kappa_{ijk} R_i R_j \right) t_k \\ &=: f_0(\underline{n})(\epsilon_1 + \epsilon_2) + \sum_{k=1}^{b_2^c} f_k(\underline{n}) t_k, \end{aligned} \tag{2.20}$$

which are linear in t_i . It implies that Λ can be expanded as a well-defined power series in e^{t_i} . Now let us assume that for appropriate choice of the r -field, this power series with all the instanton contributions taken into account truncates at finite orders for all true t_i ,¹⁴ which actually implies that the Λ does not depend on any of the true Kähler parameters as stated in the introduction, since otherwise it can not be of modular weight zero. Needless to say, this assumption puts very strong constraint on the choice of the r -field, and these are the r -fields we are interested in.

In our current cases of elliptic non-compact Calabi-Yau threefolds, there is only one mass parameter which is τ . The strong constraint then means that for the r fields we are interested in only the lowest order of Kähler parameters contributes. Then we can simply define Λ

$$\Lambda(\tau, \epsilon_1, \epsilon_2) = \sum_{\underline{n} \in \mathcal{I}} (-1)^{|\underline{n}|} e^{f_0(\underline{n})} e^{f_k(\underline{n}) t_k}, \tag{2.21}$$

where \mathcal{I} is the set of integral vectors \underline{n} that minimize all the $f_k(\underline{n})$ for true Kähler parameters *simultaneously* after subtracting mass parameters. If the minimal values for one r are not zero simultaneously, then it must be a vanishing r or an incorrect r .

In the case of elliptic non-compact Calabi-Yau threefolds associated to minimal 6d SCFTs, the τ parameter is always some combinations of Kähler parameters in the fiber direction and it will be a little bit subtle to subtract the mass parameter τ . We can first consider the τ irrelevant Kähler parameter t_{ell} , and the minimum of the associated $f_{\text{ell}}(\underline{n})$, and then check the solved r fields with the condition (2.21). As also shown in [25], the existence of the minimum of $f_{\text{ell}}(\underline{n})$, already suffices to fix all the r fields. In particular, similar to t , we decompose r into components $r_0, r_1, \dots, r_{\text{rk}}, r_B$. Then the weak consistency condition implies the *admissibility* condition, which we will prove shortly

$$r_\tau = \sum_{I=0}^{\text{rk}} a_I r_I = 0. \tag{2.22}$$

It means the component of r in the direction of the elliptic fiber must vanish. Recall that we only consider r modulo $2C \cdot \underline{n}$ for $\forall \underline{n} \in \mathbb{Z}^{b_4^c} = \mathbb{Z}^{\text{rk}+1}$. The intersection matrix C defines the injection $\mathbb{Z}^{\text{rk}+1} \hookrightarrow \mathbb{Z}^{\text{rk}+2}$. The r fields that satisfy (2.22) can only take value in the

¹⁴This assumption is the most natural consideration for the generalized blowup equations compared with the initial Nakajima-Yoshioka blowup equations.

n	3	4	5	6	8	12
$\#(r)$	9	16	5	18	16	12
$ P : Q^\vee $	3	4	1	3	2	1

Table 3. Numbers and sizes of groups of r -fields giving rise to the same embedding ϕ_λ .

dimension zero quotient lattice $\Gamma/\mathbb{Z}^{\text{rk}+1}$ with $\Gamma = \mathbb{Z}^{\text{rk}+2}|_{r_r=0}$, and they are thus finite in number. Finally we impose the BPS checkerboard pattern condition

$$2j_l + 2j_r + 1 \equiv r \cdot \beta \pmod{2} \quad (2.23)$$

to remove half of them. We comment that the checkerboard condition can be written down without computing any BPS invariants. As argued in [25], any rational curve β in the Calabi-Yau with normal bundle $\mathcal{O}(-n) \oplus \mathcal{O}(n-2)$ must have

$$2j_l + 2j_r + 1 \equiv n \pmod{2}. \quad (2.24)$$

This implies the r -field always satisfies

$$r \equiv (0, \dots, 0, \mathbf{n}) \pmod{2}, \quad (2.25)$$

for minimal 6d (1,0) SCFTs. The argument above not only establishes the finiteness of admissible r , but also provides a guideline on how to determine them. We find all of them for the $G = \text{SU}(3), \text{SO}(8)$ models in [25] and for the remaining models with $G = F_4, E_{6,7,8}$ in the example section 4 of this paper. In all these examples we checked that they satisfy the stronger consistency condition. Here we summarise their numbers in table 3.

Let us prove the admissibility condition (2.22). An important ingredient of the blowup equations (1.2) are the shifts of Kähler moduli m_i ($i = 1, \dots, \text{rk}$) by

$$R_i = \sum_{J=0}^{\text{rk}} C_{i,J} n_J + \frac{1}{2} r_i, \quad (2.26)$$

where n_J take value in \mathbb{Z} . We collect them into a single vector just like m

$$R = \sum_{i=1}^{\text{rk}} \left(- \sum_{J=0}^{\text{rk}} \hat{A}_{i,J} n_J + \frac{1}{2} r_i \right) \omega_i = -\alpha^\vee + \lambda + n_0 \theta, \quad (2.27)$$

where we have used the generic form (2.2) of the intersection matrix C and the following Lie algebraic notation

$$\lambda = \frac{1}{2} \sum_{i=1}^{\text{rk}} r_i \omega_i, \quad \alpha^\vee = \sum_{i=1}^{\text{rk}} n_i \alpha_i^\vee. \quad (2.28)$$

The function $f_{\text{ell}}(\underline{n})$, which we also denote by $P_G(\underline{n})$ due to its particular importance, then has the form

$$f_{\text{ell}}(\underline{n}) = \frac{1}{\mathbf{n}} R_{\text{ell}} R_\tau + \frac{1}{2} (R, R) =: P_G(\underline{n}), \quad (2.29)$$

where R_{ell}, R_τ are the shifts of t_{eff}, τ given respectively by

$$R_{\text{ell}} = -\mathbf{n}n_0 + \frac{1}{2}r_{\text{ell}}, \quad R_\tau = \frac{1}{2}r_\tau = \frac{1}{2} \sum_{I=0}^{\text{rk}} a_I r_I. \quad (2.30)$$

A little algebraic manipulation leads to

$$P_G(\underline{n}) = \frac{1}{2} \sum_{I,J=0}^{\text{rk}} n_I n_J (\alpha_I^\vee, \alpha_J^\vee) - \frac{1}{2} \sum_{I,J=0}^{\text{rk}} r_I n_J (\widehat{\omega}_I, \alpha_J^\vee) + \dots \quad (2.31)$$

where \dots denote the terms that are independent from \underline{n} . If we demand that this function have a minimum, all the derivatives $\partial_{n_I} P_G(\underline{n})$ must have a common zero. Multiplying each of them with comark a_I and adding them up, we immediately arrive at the admissibility condition (2.22).¹⁵ Let us make some remarks here. As we will see in section 4, all the admissible r -fields are such that the components r_i ($i = 1, \dots, \text{rk}$) are even integers. It is clear then that λ associated to r as well as the shift R have nice interpretation as weight vectors of \mathfrak{g} . And $P_G(\underline{n})$ is nothing else but half the norm square of R due to (2.22)

$$P_G(\underline{n}) = \frac{1}{2}(R, R). \quad (2.32)$$

Furthermore λ defines an embedding of Q^\vee into P

$$\begin{aligned} \phi_\lambda : Q^\vee &\hookrightarrow P \\ \alpha^\vee &\mapsto \phi_\lambda(\alpha^\vee) = -\alpha^\vee + \lambda \end{aligned} \quad (2.33)$$

and R takes value in the image $\phi_\lambda(Q^\vee)$ if $n_0 = 0$. The number of inequivalent embeddings is the index $|P : Q^\vee|$, which also happens to be the order of the automorphism group of the associated Dynkin diagram. We also list these numbers in table 3. The reader may notice the curious relation

$$\mathbf{n} \cdot |P : Q^\vee| = \#(r), \quad (2.34)$$

whose meaning will be clear in section 3.

2.4 De-affinisation

Once the semiclassical piece Z^{cls} and the r -fields are known, we can start solving refined BPS invariants by expanding the blowup equations (1.2) in terms of Kähler moduli and extracting constraint equations of BPS invariants at each order. Alternatively since the $Z^{\text{1-loop}}$ piece is rather easy to compute for the minimal 6d SCFTs, which reads [25]¹⁶

$$Z^{\text{1-loop}}(m, \epsilon_1, \epsilon_2) = \text{PE} \left[- \frac{q_R + q_R^{-1}}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})} \sum_{\alpha \in \Delta_+} \left(e^{\langle \alpha, m \rangle} + Q_\tau e^{-\langle \alpha, m \rangle} \right) \frac{1}{1 - Q_\tau} \right], \quad (2.36)$$

¹⁵This condition was found in [25] by an intuitive geometric argument for minimal theories with a gauge group G of the ADE type. Here we prove it for all gauge groups including the non-ADE types.

¹⁶Here PE is the plethystic exponent operator defined as

$$\text{PE}[f(x)] = \exp \left[\sum_{n=1}^{\infty} \frac{1}{n} f(x^n) \right]. \quad (2.35)$$

Note we only consider the contribution from vector multiplets to the one-loop partition function here. The tensor multiplets actually also contribute to one-loop partition function, but their contribution does not depend on the gauge parameter m , i.e. it is pure τ terms which will decouple from the blowup equations. Thus we do not consider them here. See more discussion in section 2.4 in [25].

we can also plug in this piece of information, expand the partition functions in the blowup equation (1.2) only in terms of Q_{ell} , and obtain recursion relations of elliptic genera

$$\begin{aligned} \sum_{P_G(\underline{n})+d_1+d_2=d} D'^{\text{cls}} D^{1\text{-loop}} \mathbb{E}_{d_1}(\tau, m + \epsilon_1 R_m(\underline{n}), \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_{d_2}(\tau, m + \epsilon_2 R_m(\underline{n}), \epsilon_1 - \epsilon_2, \epsilon_2) \\ = \Lambda \mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2), \end{aligned} \tag{2.37}$$

which allows the solution of the elliptic genera in compact formulas. Here

$$D'^{\text{cls}} = (-1)^{|\underline{n}|+(d_1+d_2-d)r_{\text{ell}}} \exp\left(f_0(\underline{n})(\epsilon_1 + \epsilon_2) + \sum_{I=0}^{\text{rk}} f_I(\underline{n})t_I + (d_1\epsilon_1 + d_2\epsilon_2)R_{\text{ell}}(\underline{n})\right) \tag{2.38}$$

collects contributions from the semiclassical partition function (as well as the shift of t_{ell}), and

$$D^{1\text{-loop}} = Z^{1\text{-loop}}(\tau, m + \epsilon_1 R_m(\underline{n}), \epsilon_1, \epsilon_2 - \epsilon_1) Z^{1\text{-loop}}(\tau, m + \epsilon_2 R_m(\underline{n}), \epsilon_1 - \epsilon_2, \epsilon_2) / Z^{1\text{-loop}}(\tau, m, \epsilon_1, \epsilon_2) \tag{2.39}$$

is the contribution from the one-loop partition function. The elliptic blowup equations can be put in an elegant form by partially resumming the left hand side of (2.37). With (2.31) and the admissibility condition (2.22), one can show that the polynomial $P_G(\underline{n})$ that characterises the summation index \underline{n} is invariant under the translation

$$n_I \rightarrow n_I + a_I^\vee k, \quad k \in \mathbb{Z}. \tag{2.40}$$

Besides, in the components $Z^{1\text{-loop}}$ and \mathbb{E}_d the dependence on \underline{n} only appears in the shifts

$$R_I = - \sum_{J=0}^{\text{rk}} \hat{A}_{I,J} n_J + \frac{1}{2} r_I, \quad I = 0, 1, \dots, \text{rk}, \tag{2.41}$$

which are also invariant under (2.40). As a result, we can decompose the summation index

$$\underline{n} = \hat{\underline{n}} + a^\vee k \tag{2.42}$$

with the zeroth component of $\hat{\underline{n}}$ fixed to zero, i.e. $\hat{\underline{n}} = (0, n_1, n_2, \dots, n_{\text{rk}})$, a step we call “de-affinisation”, and perform the infinite sum on the left hand side of (2.37) in two steps. In the first step, we factor out $D^{1\text{-loop}}$, \mathbb{E}_{d_1, d_2} and only sum D'^{cls} over $k \in \mathbb{Z}$. Due to the quadratic nature of the polynomials $f_0(\underline{n})$, $f_I(\underline{n})$, and the relation of t_I with τ (2.4), this first summation in fact produces a theta function with characteristics [25]. In the second step, we sum over d_1, d_2 and $\hat{\underline{n}}$ that satisfy

$$P_G(\hat{\underline{n}}) + d_1 + d_2 = d. \tag{2.43}$$

Instead of $\hat{\underline{n}}$ we can treat $R(\hat{\underline{n}})$ as the summation index, which as we argued before is now interpreted as a weight vector in $\phi_\lambda(Q^\vee)$ determined by r . In addition, the one-loop contribution $D^{1\text{-loop}}$ also turns out to be a quotient of theta functions [25], while the elliptic genera themselves are meromorphic Jacobi forms. Therefore in the end, the elliptic blowup equations can be presented as beautiful equations of Jacobi forms with a sum over the shifted coroot lattice $\phi_\lambda(Q^\vee)$. These equations are the highlights of the next section. We will present these equations in the beginning of the next section, and then discuss their properties and how to solve elliptic genera from them.

3 Elliptic blowup equations

In this section, we first present the elliptic blowup equations for 6d minimal $\mathcal{N} = (1, 0)$ SCFTs with $G = \text{SU}(3), \text{SO}(8), F_4, E_6, E_7, E_8$, and discuss their two interesting properties, the modularity and the universality. The first property in particular serves as a strong support for the validity of the elliptic blowup equations to arbitrary degrees. Then we distinguish two cases with $\Lambda(\epsilon_1, \epsilon_2, m, r)$ non-vanishing or identically vanishing, and discuss these two cases in detail. In particular, the blowup equations in the first case with non-vanishing Λ allow us to write down an exact and universal recursion formula for elliptic genera, thus offering a complete solution to the elliptic genera.

Let us first fix some conventions. In the following whenever there is no risk of confusion we will use the dot to denote both the invariant bilinear form on \mathfrak{h} or \mathfrak{h}^* and the natural inner product between \mathfrak{h} and \mathfrak{h}^*

$$\alpha \cdot \beta = (\alpha, \beta), \text{ or } \alpha \cdot \beta = \langle \alpha, \beta \rangle. \quad (3.1)$$

We define the norm square

$$\|\alpha\|^2 = \alpha \cdot \alpha. \quad (3.2)$$

For a coroot $\beta^\vee \in Q^\vee$, we also define

$$|\beta^\vee| = \sum_{i=1}^{\text{rk}} \beta_i^\vee, \quad \text{with } \beta^\vee = \sum_{i=1}^{\text{rk}} \beta_i^\vee \alpha_i^\vee. \quad (3.3)$$

Besides, for a vector m representing the Kähler parameters associated to a Lie algebra, we denote $m_\alpha = m \cdot \alpha$ for short.

Following the de-affinisation procedure described in section 2.4, we derive the *elliptic blowup equations* as

$$\begin{aligned} & \frac{1}{2} \|\omega\|^2 + d_1 + d_2 = d \\ & \sum_{\omega \in \phi_\lambda(Q^\vee), d_{1,2} \in \mathbb{N}} (-1)^{|\phi_\lambda^{-1}(\omega)|} \cdot \theta_i^{[a]} \left(\mathfrak{n}\tau, (\mathfrak{n}-2)(\epsilon_1 + \epsilon_2) - \mathfrak{n} \left(\left(\frac{1}{2} \|\omega\|^2 + d_1 \right) \epsilon_1 + \left(\frac{1}{2} \|\omega\|^2 + d_2 \right) \epsilon_2 - m_\omega \right) \right) \\ & \quad \times A_\omega(m) \cdot \mathbb{E}_{d_1}(\tau, m - \epsilon_1 \omega, \epsilon_1, \epsilon_2 - \epsilon_1) \cdot \mathbb{E}_{d_2}(\tau, m - \epsilon_2 \omega, \epsilon_1 - \epsilon_2, \epsilon_2) \\ & = \begin{cases} \theta_i^{[a]}(\mathfrak{n}\tau, (\mathfrak{n}-2)(\epsilon_1 + \epsilon_2)) \cdot \mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2), & \text{fixed } d \in \mathbb{N}, \\ 0, & \text{fixed } d \notin \mathbb{N}. \end{cases} \end{aligned} \quad (3.4)$$

Here the subscript of theta functions i is 4 if \mathfrak{n} is odd and 3 if \mathfrak{n} is even, and the characteristic $a = k/\mathfrak{n} - 1/2$, $k = 0, 1, \dots, \mathfrak{n} - 1$. The factor $A_\omega(m)$ is given by

$$A_\omega(m) = \prod_{\alpha \in \Delta_+} \check{\theta}(m_\alpha, -\alpha \cdot \omega), \quad (3.5)$$

where we denote for $R \geq 0$,

$$\check{\theta}(z, R) = \prod_{\substack{m, n \geq 0 \\ m+n \leq R-1}} \frac{\eta}{\theta_1(z + m\epsilon_1 + n\epsilon_2)} \prod_{\substack{m, n \geq 0 \\ m+n \leq R-2}} \frac{\eta}{\theta_1(z + (m+1)\epsilon_1 + (n+1)\epsilon_2)}, \quad (3.6)$$

and for $R \leq 0$, $\check{\theta}(z, R) = \check{\theta}(z, -R)|_{\epsilon_{1,2} \rightarrow -\epsilon_{1,2}}$. Note the $\alpha \cdot \omega$ in (3.5) is guaranteed to be an integer, as requested for the definition of $\check{\theta}(z, R)$. See more about the origin of $\check{\theta}(z, R)$ function in appendix D. Let us show some examples here:

$$\begin{aligned} \check{\theta}(z, 0) &= 1, \\ \check{\theta}(z, \pm 1) &= \frac{\eta}{\theta_1(z)}, \\ \check{\theta}(z, \pm 2) &= \frac{\eta^4}{\theta_1(z)\theta_1(z \pm \epsilon_1)\theta_1(z \pm \epsilon_2)\theta_1(z \pm \epsilon_1 \pm \epsilon_2)}. \end{aligned} \tag{3.7}$$

The dependence on the r -field in (3.4) is related to the choice of a . If we choose basis of Kähler moduli $(t_{\text{ell}}, \tau, m_i)$, $i = 1, \dots, \text{rk}$ as in (2.4), the corresponding components of the r -field are $(r_{\text{ell}}, r_\tau, r_i)$, $i = 1, \dots, \text{rk}$. The first component r_{ell} controls the characteristic a through $a = r_{\text{ell}}/2\mathbf{n}$, thus the latter can take any of the following values

$$a = -\frac{\mathbf{n} - 2k}{2\mathbf{n}}, \quad k = 0, 1, \dots, \mathbf{n} - 1. \tag{3.8}$$

The component r_τ vanishes due to the admissibility condition (2.22). The remaining components r_i , $i = 1, \dots, \text{rk}$ always correspond to a weight vector $\lambda \in P$ through (2.28), which in turn induces the embedding $\phi_\lambda : Q^\vee \hookrightarrow P$, and the summation index vector $\omega \in \phi_\lambda(Q^\vee)$ plays the role of the shift vector $-R$. The number of different embeddings is $|P : Q^\vee|$. The total number of different blowup equations is then $\mathbf{n}|P : Q^\vee|$, which explains the numerology found in (2.34).

3.1 Modularity of elliptic blowup equations

In this section, we provide evidence for the elliptic blowup equations (3.4) by showing that the components of the elliptic blowup equations transforms correctly as weak Jacobi forms. This is established by showing that the weight and the index, in general a quadratic polynomial, of the corresponding components in (3.4) match the predictions for the index and weight made from the 2d and the 6d anomaly polynomial or from the transformation properties of the refined topological string partition function under the S and T monodromies of the Calabi-Yau space X , see [22] and more generally [58]. In general the blowup equations give interesting identities for Jacobi forms, one example is proven in section 3.3.1, see also (3.48). In the fortuitous cases where the expressions of \mathbb{E}_d are already known, for instance the $G = \text{SU}(3), \text{SO}(8)$ models [8, 34], we can plug in their expressions, and verify these identities by small Q_τ expansion.

It is easy to see that each term in the summation of (3.4) has weight $1/2$ as both A_ω and \mathbb{E}_d are of weight zero and θ_i has weight $1/2$. The identification of the modular indices requires a bit of computation, which is independent from the characteristic a . The basic idea is to repeatedly use the fact that $\theta(N\tau, Nz)$ is of index $N/2$. Let us denote

$$d_0 = \frac{1}{2} \|\omega\|^2. \tag{3.9}$$

It is easy to see the theta function $\theta_i^{[a]}$ on the left hand side has modular index

$$\text{Ind}_{d_0}^G = \frac{\mathbf{n}}{2} \left(\left(d_0 + d_1 - \frac{\mathbf{n} - 2}{\mathbf{n}} \right) \epsilon_1 + \left(d_0 + d_2 - \frac{\mathbf{n} - 2}{\mathbf{n}} \right) \epsilon_2 - m_\omega \right)^2. \tag{3.10}$$

Using (D.16), the modular index polynomial of $A_\omega(m)$ can be computed as

$$\begin{aligned} \text{Ind}_A^G &= -\frac{\mathbf{n}}{2}m_\omega^2 - \frac{\mathbf{n}}{2}d_0m \cdot m + (\mathbf{n}d_0 + 2 - \mathbf{n})(\epsilon_1 + \epsilon_2)m_\omega \\ &\quad - \frac{d_0}{2}(\mathbf{n}d_0 + 2 - \mathbf{n})(\epsilon_1^2 + \epsilon_1\epsilon_2 + \epsilon_2^2). \end{aligned} \quad (3.11)$$

The elliptic genus $\mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2)$ is known to have the modular index [24, 37]

$$\text{Ind}_d^G = -\frac{1}{2}d(\mathbf{n} - 2)(\epsilon_1 + \epsilon_2)^2 + \frac{1}{2}d(\mathbf{n}d - \mathbf{n} + 2)\epsilon_1\epsilon_2 - \frac{1}{2}\mathbf{n}dm \cdot m. \quad (3.12)$$

Thus the modular index polynomials of $\mathbb{E}_{d_1}(\tau, m - \epsilon_1\omega, \epsilon_1, \epsilon_2 - \epsilon_1)$ and $\mathbb{E}_{d_2}(\tau, m - \epsilon_2\omega, \epsilon_1 - \epsilon_2, \epsilon_2)$ can be computed respectively as

$$\text{Ind}_{d_1}^G = \frac{1}{2} \left(-(\mathbf{n} - 2)d_1\epsilon_2^2 + (\mathbf{n}d_1 - \mathbf{n} + 2)d_1\epsilon_1(\epsilon_2 - \epsilon_1) - \mathbf{n}d_1(m - \epsilon_1\omega)^2 \right), \quad (3.13)$$

$$\text{Ind}_{d_2}^G = \frac{1}{2} \left(-(\mathbf{n} - 2)d_2\epsilon_1^2 + (\mathbf{n}d_2 - (\mathbf{n} - 2))d_2(\epsilon_1 - \epsilon_2)\epsilon_2 - \mathbf{n}d_2(m - \epsilon_2\omega)^2 \right). \quad (3.14)$$

Using $d = \|\omega\|^2/2 + d_1 + d_2$, we find that the four components on the left hand side of (3.4) has total modular index polynomial as

$$\text{Ind}_{d_0}^G + \text{Ind}_A^G + \text{Ind}_{d_1}^G + \text{Ind}_{d_2}^G = -\frac{(\mathbf{n} - 2)(\mathbf{n} - 2 + d\mathbf{n})}{2\mathbf{n}}(\epsilon_1 + \epsilon_2)^2 + \frac{d(d\mathbf{n} - \mathbf{n} + 2)}{2}\epsilon_1\epsilon_2 - \frac{d\mathbf{n}}{2}m \cdot m, \quad (3.15)$$

which is independent from ω, d_1, d_2 individually but only depends on their combination d ! This highly nontrivial fact guarantees the modularity of elliptic blowup equation, which means in summation of the left hand side of (3.4) all terms share the same modular index, thus transform as whole Jacobi form together! In the case that $d \notin \mathbb{N}$ where the right hand side of (3.4) vanishes, this is the end of the story. If $d \in \mathbb{N}$ so that the right hand side of (3.4) is non-vanishing, we still need to show the right hand side also shares the same index polynomial. The index polynomial of Λ is simply,

$$\text{Ind}_\Lambda^G = \frac{(\mathbf{n} - 2)^2}{2\mathbf{n}}(\epsilon_1 + \epsilon_2)^2, \quad (3.16)$$

which together with (3.12) indeed sum up to (3.15).

3.2 Universality of elliptic blowup equations

We demonstrate here an interesting property of the elliptic blowup equations. The \mathbf{n} blowup equations (3.4) with a fixed embedding ϕ_λ can be ordered by the characteristics a of the theta functions, where for two consecutive equations a differ by $1/\mathbf{n}$. We claim that if two consecutive unity blowup equations are valid, the other equations must hold automatically. We call this the universality of the elliptic blowup equations.¹⁷

¹⁷The following argument assumes the form of refined BPS expansion. Thus the universality here does not contradict with our statement that choosing arbitrary three unity r fields, i.e. three different characteristics a , one is able to use the blowup equations to solve out the elliptic genus.

Theta functions with characteristics have the following properties

$$\theta_3^{\left[\frac{m}{2n}\right]}(\mathbf{n}\tau, z) = e^{\pi i \tau \frac{m^2}{4n} + \pi i z \frac{m}{n}} \theta_3(\mathbf{n}\tau, z + \frac{1}{2}m\tau), \quad m \in \mathbb{Z}, \quad (3.17)$$

$$\theta_4^{\left[\frac{m}{2n}\right]}(\mathbf{n}\tau, z) = e^{\pi i \frac{m}{2n} + \pi i \tau \frac{m^2}{4n} + \pi i z \frac{m}{n}} \theta_4(\mathbf{n}\tau, z + \frac{1}{2}m\tau), \quad m \in \mathbb{Z}. \quad (3.18)$$

Therefore, shifting z by τ is equivalent to shifting the characteristic of these theta function by $1/n$. Then starting from one unity blowup equation, let us shift ϵ_1 by τ and check how various Jacobi forms in (3.4) change.

- $\theta_i^{[a]}(\mathbf{n}\tau, \dots)$: the elliptic parameter changes by $-2\tau + \text{integer} \cdot \mathbf{n}\tau$. The shift $\text{integer} \cdot \mathbf{n}\tau$ can be removed at the expense of an additional exponential factor due to quasi-periodicity of the theta function, while the shift -2τ , as we have argued, is equivalent to shifting a by $-2/n$.
- $A_\omega(m)$: it is a product of factors like $\theta_1(\tau, m_\alpha + m\epsilon_1 + n\epsilon_2)$, $m, n \in \mathbb{Z}$, therefore is invariant under this shift up to an exponential factor.
- \mathbb{E}_k : we first argue that under the shift $\epsilon_1 \rightarrow \epsilon_1 + 1$, $\mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2)$ is invariant up to an exponential factor. The refined BPS are defined from the topological string free energy as¹⁸

$$F^{\text{inst}}(Q, \epsilon_1, \epsilon_2) = \sum_{j_l, j_r \geq 0} \sum_{w \geq 1, \beta} (-1)^{2(j_l + j_r)} N_{j_l, j_r}^\beta \frac{\chi_{j_l}(q_l^w) \chi_{j_r}(q_r^w)}{w(q_1^{w/2} - q_1^{-w/2})(q_2^{w/2} - q_2^{-w/2})} Q^{w\beta}, \quad (3.19)$$

where

$$\chi_j(q) = \frac{q^{2j+1} - q^{-2j-1}}{q - q^{-1}}. \quad (3.20)$$

Exponentiating (3.19) the instanton partition function reads

$$Z^{\text{inst}}(Q, \epsilon_1, \epsilon_2) = \prod_{\beta} \prod_{j_l, j_r=0}^{\infty} \prod_{m_l, r=-j_l, r}^{\infty} \prod_{m_1, m_2=1}^{\infty} \left(1 - q_l^{m_l} q_r^{m_r} q_1^{m_1 - \frac{1}{2}} q_2^{m_2 - \frac{1}{2}} Q^\beta \right)^{(-1)^{2(j_l + j_r)} N_{j_l, j_r}^\beta}. \quad (3.21)$$

Using (3.19) and the checkerboard pattern identity

$$2j_l + 2j_r + 1 \equiv r \cdot \beta \pmod{2} \quad (3.22)$$

for non-vanishing BPS invariants one can show that the refined BPS partition function is invariant under the combined transformation

$$(\epsilon_1, \epsilon_2, t) \rightarrow (\epsilon_1 + 1, \epsilon_2, t + r). \quad (3.23)$$

Since the r -vector for the minimal 6d SCFTs in terms of the Kähler moduli t_I, t_B have components

$$r \equiv (0, \dots, 0, \mathbf{n}) \pmod{(2\mathbb{Z})^{\text{rk}+2}}, \quad (3.24)$$

¹⁸Here $q_j \equiv \exp(2\pi i \epsilon_j)$ $j = 1, 2$ and $q_l \equiv \sqrt{q_1/q_2}$, $q_r \equiv \sqrt{q_1 q_2}$. We will also use $v \equiv q_r$, $x \equiv q_l$ in section 4 and 5 to make contact with the literature such as [37, 38].

the elliptic genus $\mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2)$ is invariant under the shift $\epsilon_1 \rightarrow \epsilon_1 + 1$, at most up to a sign if \mathfrak{n} is odd. In fact

$$\mathbb{E}_k(\tau, m, \epsilon_1 + 1, \epsilon_2) = (-1)^{n\mathfrak{k}} \mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2). \quad (3.25)$$

Together with the modular property of \mathbb{E}_k , this implies $\mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2)$ is quasi-periodic for $\epsilon_1 \rightarrow \epsilon_1 + \tau$. Similarly, one can show that $\mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2)$ is quasi-periodic for $\epsilon_2 \rightarrow \epsilon_2 + \tau$ as well. As examples one could inspect the expressions of \mathbb{E}_k for the $\mathfrak{n} = 3, 4$ models [8, 34], which are composed of $\theta_1(\tau, \text{integer} \cdot \epsilon_1 + \text{integer} \cdot \epsilon_2 + \dots)$ and therefore are indeed quasi-periodic for $\epsilon_{1,2} \rightarrow \epsilon_{1,2} + \tau$. Now if we forget for the moment the shift on the mass parameters, the three instances of elliptic genera $\mathbb{E}_k(\tau, m - \epsilon_1\omega, \epsilon_1, \epsilon_2 - \epsilon_1)$, $\mathbb{E}_k(\tau, m - \epsilon_2\omega, \epsilon_1 - \epsilon_2, \epsilon_2)$, $\mathbb{E}_k(\tau, m, \epsilon_1, \epsilon_2)$ in (3.4) should already be invariant under $\epsilon_1 \rightarrow \epsilon_1 + \tau$ up to an exponential factor. The shift on the mass parameters $m - \epsilon_1\omega$ in the first instance of \mathbb{E}_k means that upon $\epsilon_1 \rightarrow \epsilon_1 + \tau$ its elliptic parameter is in addition shifted by τ times a weight vector, which can also be removed at the expense of an additional exponential factor [24].¹⁹

In summary, the shift $\epsilon_1 \rightarrow \epsilon_1 + \tau$ is equivalent to shifting the characteristic a of the theta functions $\theta_i^{[a]}(n\tau, \dots)$ by $-2/\mathfrak{n}$ and in addition multiplying each term in (3.4) by an exponential factor. These exponential factors are determined by the index polynomial of each term, which as a consequence of section 3.1, should be identical. Thus all the exponential factors can be factored out and removed, and we are left again with a unity blowup equation where the characteristic is shifted by $-2/\mathfrak{n}$. This immediately indicates that starting from two consecutive unity blowup equations, we can obtain all the other unity blowup equations, hence the universality property.

3.3 Unity blowup equations

The elliptic blowup equations depend on the choice of the weight vector λ and they take different forms depending on if $\lambda \in Q^\vee$ or not. We first consider the former case where $\phi_\lambda(Q^\vee)$ coincides with Q^\vee . We can denote ω as vector β^\vee in the coroot lattice. Then $\|\beta^\vee\|^2/2$ and thus d are always nonnegative integers, and as a result the right hand side of (3.4) does not vanish

$$\begin{aligned} & \sum_{\beta^\vee \in Q^\vee, d_{1,2} \in \mathbb{N}}^{\frac{1}{2}\|\beta^\vee\|^2 + d_1 + d_2 = d} (-1)^{|\beta^\vee|} \theta_i^{[a]} \left(n\tau, (n-2)(\epsilon_1 + \epsilon_2) - \mathfrak{n} \left(\left(\frac{1}{2} \|\beta^\vee\|^2 + d_1 \right) \epsilon_1 + \left(\frac{1}{2} \|\beta^\vee\|^2 + d_2 \right) \epsilon_2 - m_{\beta^\vee} \right) \right) \\ & \quad \times A_{\beta^\vee}(m) \mathbb{E}_{d_1}(\tau, m - \epsilon_1 \beta^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_{d_2}(\tau, m - \epsilon_2 \beta^\vee, \epsilon_1 - \epsilon_2, \epsilon_2) \\ & = \theta_i^{[a]}(n\tau, (n-2)(\epsilon_1 + \epsilon_2)) \cdot \mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2), \quad d \in \mathbb{N}. \end{aligned} \quad (3.26)$$

We say these elliptic blowup equations are of the *unity* type following the nomenclature in [15]. Since the number of embedding $Q^\vee \hookrightarrow \phi_\lambda(Q^\vee) = Q^\vee$ is unique, the number of unity

¹⁹It is established in section 4 of [24] that the elliptic genera of 6d SCFTs with a pure gauge bulk theory in fact consist of special Weyl invariant Jacobi forms, which, among other things, are quasi-periodic if the elliptic parameter is shifted by τ times a weight vector.

blowup equations is the same as the range of characteristics a , which is \mathfrak{n} . We also point out that, using the property that the leading Q_τ order of \mathbb{E}_d is $-dh_G^\vee/6$,²⁰ these equations in the leading order of Q_τ boil down to the identity

$$\sum_{\alpha \in \Delta^+(G)} (\beta \cdot \alpha)^2 = h_G^\vee \|\beta\|^2, \quad (3.27)$$

which is guaranteed by the Lie algebraic identity (A.5).

The unity blowup equations are particular interesting as they allow us to write down recursion formulas for the elliptic genera.

3.3.1 Recursion formulas for elliptic genera

The unity blowup equations (3.26) can be put in the following more suggestive form

$$\begin{aligned} & \theta_i^{[a]}(\mathfrak{n}\tau, (\mathfrak{n} - 2 - d\mathfrak{n})\epsilon_1 + (\mathfrak{n} - 2)\epsilon_2)\mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2 - \epsilon_1) \\ & + \theta_i^{[a]}(\mathfrak{n}\tau, (\mathfrak{n} - 2)\epsilon_1 + (\mathfrak{n} - 2 - d\mathfrak{n})\epsilon_2)\mathbb{E}_d(\tau, m, \epsilon_1 - \epsilon_2, \epsilon_2) \\ & - \theta_i^{[a]}(\mathfrak{n}\tau, (\mathfrak{n} - 2)(\epsilon_1 + \epsilon_2))\mathbb{E}_d(\tau, m, \epsilon_1, \epsilon_2) = I_d(\mathbb{E}_{<d}). \end{aligned} \quad (3.28)$$

where $I_d(\mathbb{E}_{<d})$ only contains the elliptic genera of degrees lower than d . Since three copies of \mathbb{E}_d on the left hand side do not depend on the characteristic a , if we have three such equations with different a , which is indeed the case for all the minimal 6d $\mathcal{N} = (1, 0)$ SCFTs with pure gauge bulk theory, we can solve $\mathbb{E}_d(\tau, \epsilon_1, \epsilon_2)$ in terms of elliptic genera with lower number of strings; in other words, we obtain recursion formulas for elliptic genera.

Let us use the short hand notation²¹

$$\theta_{i,\{d_0,d_1,d_2\}}^{[a]} = \theta_i^{[a]}(\mathfrak{n}\tau, \mathfrak{n}m_{\alpha^\vee} + (\mathfrak{n} - 2)(\epsilon_1 + \epsilon_2) - \mathfrak{n}((d_0 + d_1)\epsilon_1 + (d_0 + d_2)\epsilon_2)), \quad (3.29)$$

where $m_{\alpha^\vee} = m \cdot \alpha^\vee$, and furthermore define

$$D_d = \text{Det} \begin{pmatrix} \theta_{i,\{0,d,0\}}^{[a_1]} & \theta_{i,\{0,0,d\}}^{[a_1]} & \theta_{i,\{0,0,0\}}^{[a_1]} \\ \theta_{i,\{0,d,0\}}^{[a_2]} & \theta_{i,\{0,0,d\}}^{[a_2]} & \theta_{i,\{0,0,0\}}^{[a_2]} \\ \theta_{i,\{0,d,0\}}^{[a_3]} & \theta_{i,\{0,0,d\}}^{[a_3]} & \theta_{i,\{0,0,0\}}^{[a_3]} \end{pmatrix}, \quad (3.30)$$

as well as

$$D_{\{d_0,d_1,d_2\}}^{\alpha^\vee} = \text{Det} \begin{pmatrix} \theta_{i,\{0,d,0\}}^{[a_1]} & \theta_{i,\{0,0,d\}}^{[a_1]} & \theta_{i,\{d_0,d_1,d_2\}}^{[a_1]} \\ \theta_{i,\{0,d,0\}}^{[a_2]} & \theta_{i,\{0,0,d\}}^{[a_2]} & \theta_{i,\{d_0,d_1,d_2\}}^{[a_2]} \\ \theta_{i,\{0,d,0\}}^{[a_3]} & \theta_{i,\{0,0,d\}}^{[a_3]} & \theta_{i,\{d_0,d_1,d_2\}}^{[a_3]} \end{pmatrix}. \quad (3.31)$$

Note that $D_d = D_{\{0,0,0\}}^{\alpha^\vee}$ does not depend on α^\vee since $\alpha^\vee = 0$ when $d_0 = \|\alpha^\vee\|^2/2 = 0$. Then the recursion formulas of \mathbb{E}_d solved from (3.28) read

$$\mathbb{E}_d = \sum_{d_0=\frac{1}{2}\|\alpha^\vee\|^2, d_1, d_2 < d}^{d_0+d_1+d_2=d} (-1)^{|\alpha^\vee|} \frac{D_{\{d_0,d_1,d_2\}}^{\alpha^\vee}}{D_d} A_{\alpha^\vee}(m) \mathbb{E}_{d_1}(m - \epsilon_1 \alpha^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_{d_2}(m - \epsilon_2 \alpha^\vee, \epsilon_1 - \epsilon_2, \epsilon_2). \quad (3.32)$$

Here the τ dependence is implied.

²⁰See more discussion in section 4.1.

²¹Note $d_0 = \|\alpha^\vee\|^2/2$ is always implied. When $d_0 = 0$, $\theta_{i,\{d_0,d_1,d_2\}}^{[a]}$ does not depend on α^\vee since $\alpha^\vee = 0$.

Let us look at some examples. The one-string elliptic genus is given by

$$\mathbb{E}_1 = \sum_{\substack{\alpha \in \Delta^\vee \\ \|\alpha^\vee\|^2=2}} \frac{D_{\{1,0,0\}}^{\alpha^\vee}}{D_1} \frac{\eta^4}{\theta_1(m_\alpha)\theta_1(m_\alpha - \epsilon_1)\theta_1(m_\alpha - \epsilon_2)\theta_1(m_\alpha - \epsilon_1 - \epsilon_2)} \prod_{\substack{\beta \in \Delta \\ \alpha^\vee \cdot \beta=1}} \frac{\eta}{\theta_1(m_\beta)} \quad (3.33)$$

where $m_\beta = m \cdot \beta$. In particular, for ADE type algebras, the \mathbb{E}_1 formula can be further simplified due to the identification of roots and coroots. Indeed, for $A_2, D_4, E_{6,7,8}$, we have the following universal formula

$$\mathbb{E}_1 = \sum_{\alpha \in \Delta} \frac{D_\alpha}{D} \frac{\eta^4}{\theta_1(m_\alpha)\theta_1(m_\alpha - \epsilon_1)\theta_1(m_\alpha - \epsilon_2)\theta_1(m_\alpha - \epsilon_1 - \epsilon_2)} \prod_{\substack{\beta \in \Delta \\ \alpha \cdot \beta=1}} \frac{\eta}{\theta_1(m_\beta)}. \quad (3.34)$$

Here D_α and D are the short notations for $D_{\{1,0,0\}}^\alpha$ and D_1 .

In the $Q_\tau \rightarrow 0$ limit, \mathbb{E}_1 formula (3.33) reduces to the universal one-instanton partition function of 5d $\mathcal{N} = 1$ pure SYM theory [30, 59]

$$Z_1 = \frac{1}{(1-e^{-\epsilon_1})(1-e^{-\epsilon_2})} \sum_{\gamma \in \Delta_l} \frac{e^{(h_G^\vee - 1)m_\gamma/2}}{(1-e^{-\epsilon_1 - \epsilon_2 + m_\gamma})(e^{m_\gamma/2} - e^{-m_\gamma/2}) \prod_{\alpha \cdot \gamma=1} (e^{m_\alpha/2} - e^{-m_\alpha/2})}, \quad (3.35)$$

where Δ_l denotes the set of long roots which is the same with coroots with $\|\alpha^\vee\|^2 = 2$.

Furthermore, the two-string elliptic genus is given by

$$\begin{aligned} \mathbb{E}_2 = & \frac{D_{\{0,1,1\}}^{\alpha^\vee=0}}{D_2} \mathbb{E}_1(m, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_1(m, \epsilon_1 - \epsilon_2, \epsilon_2) + \sum_{\|\alpha^\vee\|^2=4} (-1)^{|\alpha^\vee|} A_{\alpha^\vee}(m) \frac{D_{\{2,0,0\}}^{\alpha^\vee}}{D_2} \\ & + \sum_{\|\alpha^\vee\|^2=2} (-1)^{|\alpha^\vee|} A_{\alpha^\vee}(m) \left(\frac{D_{\{1,1,0\}}^{\alpha^\vee}}{D_2} \mathbb{E}_1(m - \epsilon_1 \alpha^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) + \frac{D_{\{1,0,1\}}^{\alpha^\vee}}{D_2} \mathbb{E}_1(m - \epsilon_2 \alpha^\vee, \epsilon_1 - \epsilon_2, \epsilon_2) \right). \end{aligned} \quad (3.36)$$

Note in the bracket of the second line of (3.36), the two terms are symmetric in $\epsilon_1 \leftrightarrow \epsilon_2$. In the later section, we use this formula to compute the two-string elliptic genus of all 6d (1,0) minimal SCFTs with $G = A_2, D_4, F_4, E_{6,7,8}$. From (3.32), we can also easily write down the universal formula for three-string elliptic genus as

$$\begin{aligned} \mathbb{E}_3 = & \left[\frac{D_{\{0,1,2\}}^{\alpha^\vee=0}}{D_3} \mathbb{E}_1(m, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_2(m, \epsilon_1 - \epsilon_2, \epsilon_2) + (\epsilon_1 \leftrightarrow \epsilon_2) \right] + \sum_{\|\alpha^\vee\|^2=6} (-1)^{|\alpha^\vee|} A_{\alpha^\vee}(m) \frac{D_{\{3,0,0\}}^{\alpha^\vee}}{D_3} \\ & + \sum_{\|\alpha^\vee\|^2=4} (-1)^{|\alpha^\vee|} A_{\alpha^\vee}(m) \left[\frac{D_{\{2,1,0\}}^{\alpha^\vee}}{D_3} \mathbb{E}_1(m - \epsilon_1 \alpha^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) + (\epsilon_1 \leftrightarrow \epsilon_2) \right] \\ & + \sum_{\|\alpha^\vee\|^2=2} (-1)^{|\alpha^\vee|} A_{\alpha^\vee}(m) \left[\left(\frac{D_{\{1,2,0\}}^{\alpha^\vee}}{D_3} \mathbb{E}_2(m - \epsilon_1 \alpha^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) + (\epsilon_1 \leftrightarrow \epsilon_2) \right) \right. \\ & \left. + \frac{D_{\{1,1,1\}}^{\alpha^\vee}}{D_3} \mathbb{E}_1(m - \epsilon_1 \alpha^\vee, \epsilon_1, \epsilon_2 - \epsilon_1) \mathbb{E}_1(m - \epsilon_2 \alpha^\vee, \epsilon_1 - \epsilon_2, \epsilon_2) \right]. \end{aligned} \quad (3.37)$$

From the topological string point of view, the d -string elliptic genus \mathbb{E}_d encodes the BPS invariants N_{j_l, j_r}^β with base degree d as well as multi-wrapping contributions from lower base degree curves. Once all the elliptic genera up to certain base degree d are computed, all the BPS invariants up to base degree d and arbitrary degrees along other directions can be extracted. The recursion formulas (3.32) thus allow us to reproduce the genus zero Gopakumar-Vafa invariants for the Calabi-Yau threefolds associated to the minimal 6d SCFTs with $\mathfrak{n} = 5, 6, 8, 12$ [24], and to compute the refined BPS invariants for the first time in the literature.

Let us make a remark here concerning the validity of the recursion formula (3.32). Obviously, the recursion formula is only well-defined when $D_d \neq 0$. We have checked that this is indeed true for all the minimal models except for the model of $G = \text{SU}(3)$ with $d = 1$, where both $D_{\{1,0,0\}}^\alpha$ and D_1 vanish. This is a special situation since for the model with $G = \text{SU}(3)$ there are only three choices of the characteristics a_i and there may be certain symmetry enhancement for \mathbb{E}_1 such that only two of the three unity blowup equations are linearly independent. Note that this does *not* contradict with the fact that the universal one-instanton partition function Z_1 of 5d pure SYM theories [30, 59] works perfectly for the $\text{SU}(3)$ theory which is recovered from one-string elliptic genus in the 5d limit with $Q_\tau \rightarrow 0$. What happens in this limit is that the 6d unity r -field $r = (0, 0, 0, 3)$ splits to two 5d unity r -fields r_1 and r_2 , and the leading Q_τ order term of D_1 is the difference of two contributions associated to r_1 and r_2 respectively, both of which remain finite and identical. Nevertheless, it should be emphasize that though recursion formula does not work for $\mathbb{E}_1^{\text{SU}(3)}$, by assuming the refined BPS expansion, one can still use two unity blowup equations and one vanishing blowup equation to solve out all the refined BPS invariants, which is what we have done in [25]. If further assuming the knowledge on Z_0 , one can actually use one single unity blowup equations to solve out all refined BPS invariants. To compute $\mathbb{E}_2^{\text{SU}(3)}$ by recursion formula (3.36), practically one can use the exact formula $\mathbb{E}_1^{\text{SU}(3)}$ in [34]. For all $d > 1$, the recursion formula (3.32) works well for $\text{SU}(3)$.

The identity $D_1^{\text{SU}(3)} = 0$ here despite its simple form does not seem so trivial. In fact, we find it is a special case of the following series of identities. With $q = e^{\pi i \tau}$ and $y = e^{2\pi i z}$ we define the following double indexed θ functions

$$\begin{aligned} \theta_3^{[r/2n]}(n\tau, z) &= q^{\frac{r^2}{4n}} y^{\frac{r}{2n}} \sum_{l \in \mathbb{Z}} q^{l(nl+r)} y^l = \sum_{k=r(2n)} q^{\frac{k^2}{4n}} y^{\frac{k}{2n}}, \\ \theta_4^{[r/2n]}(n\tau, z) &= e^{\frac{\pi i r}{4n}} q^{\frac{r^2}{4n}} y^{\frac{r}{2n}} \sum_{l \in \mathbb{Z}} (-1)^l q^{l(nl+r)} y^l = \sum_{k=r(2n)} q^{\frac{k^2}{4n}} (-y)^{\frac{k}{2n}}, \end{aligned} \tag{3.38}$$

where the notations $k = r(m)$ in the second sums, mean that k runs over values of the form $r + ml$, for any $l \in \mathbb{Z}$. We claim that if

$$\sum_{j=0}^{n-1} z_j = 0 \quad \leftrightarrow \quad \prod_{j=0}^{n-1} y_j = 1, \tag{3.39}$$

the determinant of the $n \times n$ matrices defined from these θ functions

$$\begin{aligned} \det \left[\theta_3^{\left[\frac{i}{n}\right]}(n\tau, z_j)_{i,j=0,\dots,n-1} \right] &= 0, & \text{if } 2 \mid n \\ \det \left[\theta_4^{\left[\frac{2i+1}{2n}\right]}(n\tau, z_j)_{i,j=0,\dots,n-1} \right] &= 0, & \text{if } 2 \nmid n \end{aligned} \tag{3.40}$$

vanish. Note the $n = 1$ case is just the well-known fact $\theta_1(\tau, 0) = 0$, while $n = 3$ case implies the $D_1^{\text{SU}(3)} = 0$ identity in our previous context. The proof proceeds in both cases in (3.40) by showing that each term of the form $q^m \prod_{j=1}^n y_j^{k_j}$ that occur in the expansion of the determinant, constrained by (3.39), appears once with positive and once with negative sign.²² The first case is notationally simpler so we prove it explicitly. Using $2 \mid n$ and an irrelevant rescaling²³ $z_i \rightarrow \tilde{z}_i = nz_i$ we rewrite the first determinate in the statement in (3.40) as

$$\begin{aligned} \det \left[\left(\sum_{k=i} \binom{k^2}{n} q^{\frac{k^2}{n}} y_j^k \right)_{i,j=0,\dots,n-1} \right] &= \sum_{\Pi} \text{sgn}(\Pi) \sum_{\substack{k_0 = \Pi(0) \dots \\ \vdots \\ k_{n-1} = \Pi(n-1)}} q^{\frac{k_0^2 + \dots + k_{n-1}^2}{n}} y_0^{k_0} \dots y_{n-1}^{k_{n-1}} \\ &= \sum_{\substack{k_0, \dots, k_{n-1} \in \mathbb{Z} \\ k_i \neq k_l \pmod n}} \text{sgn}(P_k) q^{\frac{k_0^2 + \dots + k_{n-1}^2}{n}} y_0^{k_0} \dots y_{n-1}^{k_{n-1}}. \end{aligned} \tag{3.41}$$

Here P_k is defined by the maps $i \mapsto k_i \pmod n$, for $i=0, \dots, n-1$. Let $\bar{k} = (k_0 + \dots + k_{n-1})/n$ be the average of k_i and use it to define $k_i^* = k_i - 2\bar{k}$. Since $2 \mid n$, \bar{k} runs over $\bar{k} = \frac{1}{2} (1)$ and hence $k_i^* \in \mathbb{Z}$. Direct calculation shows that under the $*$ operation $k_i \mapsto k_i^*$ the terms $m_k := q^{(k_0^2 + \dots + k_{n-1}^2)/n} \prod_{j=0}^{n-1} y_j^{k_j}$ in (3.40) stay invariant $m_k = m_{k^*}$; the second factor due to (3.39). Let now P_k^* be defined by $i \mapsto k_i^* \pmod n$. It follows immediately that it likewise defines a permutation of the indices $\{0, \dots, n-1\}$, however with the opposite parity. Hence m_k appears twice with opposite sign and the sum (3.41) is zero. The proof of the second case in (3.40) proceeds analogously, with appropriate relabelling of the indices.

3.3.2 Uniqueness of recursion formulas

One important consequence of the universality property is that the recursion formula (3.32) does not depend on the choice of three different r -fields in its construction, as it should. Let us suppose that we already know $\mathbb{E}_{<d}$ and we wish to compute \mathbb{E}_d using the recursion formula (3.32) obtained from three consecutive unity blowup equations forming the linear system

$$\begin{pmatrix} \theta_{i,\{0,d,0\}}^{[a_1]} & \theta_{i,\{0,0,d\}}^{[a_1]} & \theta_{i,\{0,0,0\}}^{[a_1]} \\ \theta_{i,\{0,d,0\}}^{[a_2]} & \theta_{i,\{0,0,d\}}^{[a_2]} & \theta_{i,\{0,0,0\}}^{[a_2]} \\ \theta_{i,\{0,d,0\}}^{[a_3]} & \theta_{i,\{0,0,d\}}^{[a_3]} & \theta_{i,\{0,0,0\}}^{[a_3]} \end{pmatrix} \cdot \begin{pmatrix} \mathbb{E}_d(m, \epsilon_1, \epsilon_2 - \epsilon_1) \\ \mathbb{E}_d(m, \epsilon_1 - \epsilon_2, \epsilon_2) \\ -\mathbb{E}_d(m, \epsilon_1, \epsilon_2) \end{pmatrix} = - \begin{pmatrix} I_d^{[a_1]}(\mathbb{E}_{<d}) \\ I_d^{[a_2]}(\mathbb{E}_{<d}) \\ I_d^{[a_3]}(\mathbb{E}_{<d}) \end{pmatrix}, \tag{3.42}$$

²²We thank Don Zagier for pointing this mechanism out to us.

²³Followed by a renaming of the \tilde{z}_i to z_i again.

where $a_2 - a_1 = a_3 - a_2 = 1/n$, and the matrix of theta function on the l.h.s. is of full rank. If the $\mathbb{E}_d(m, \epsilon_1, \epsilon_2)$ solved from the linear system is correct, so should be $\mathbb{E}_d(m, \epsilon_1, \epsilon_2 - \epsilon_1)$, $\mathbb{E}_d(m, \epsilon_1 - \epsilon_2, \epsilon_2)$, and the three unity blowup equations in (3.42) should all be correct as well. Otherwise, the linear system could always be corrected by

$$\begin{pmatrix} \theta_{i,\{0,d,0\}}^{[a_1]} & \theta_{i,\{0,0,d\}}^{[a_1]} & \theta_{i,\{0,0,0\}}^{[a_1]} \\ \theta_{i,\{0,d,0\}}^{[a_2]} & \theta_{i,\{0,0,d\}}^{[a_2]} & \theta_{i,\{0,0,0\}}^{[a_2]} \\ \theta_{i,\{0,d,0\}}^{[a_3]} & \theta_{i,\{0,0,d\}}^{[a_3]} & \theta_{i,\{0,0,0\}}^{[a_3]} \end{pmatrix} \cdot \begin{pmatrix} \mathbb{E}_d(m, \epsilon_1, \epsilon_2 - \epsilon_1) \\ \mathbb{E}_d(m, \epsilon_1 - \epsilon_2, \epsilon_2) \\ -\mathbb{E}_d(m, \epsilon_1, \epsilon_2) \end{pmatrix} = - \begin{pmatrix} I_d^{[a_1]}(\mathbb{E}_{<d}) \\ I_d^{[a_2]}(\mathbb{E}_{<d}) \\ I_d^{[a_3]}(\mathbb{E}_{<d}) \end{pmatrix} + \begin{pmatrix} R_1 \\ R_2 \\ R_3 \end{pmatrix}. \quad (3.43)$$

By inverting the matrix of theta functions in (3.42) and (3.43) and subtracting the two equations from each other, we get

$$\begin{pmatrix} \theta_{i,\{0,d,0\}}^{[a_1]} & \theta_{i,\{0,0,d\}}^{[a_1]} & \theta_{i,\{0,0,0\}}^{[a_1]} \\ \theta_{i,\{0,d,0\}}^{[a_2]} & \theta_{i,\{0,0,d\}}^{[a_2]} & \theta_{i,\{0,0,0\}}^{[a_2]} \\ \theta_{i,\{0,d,0\}}^{[a_3]} & \theta_{i,\{0,0,d\}}^{[a_3]} & \theta_{i,\{0,0,0\}}^{[a_3]} \end{pmatrix}^{-1} \cdot \begin{pmatrix} R_1 \\ R_2 \\ R_3 \end{pmatrix} = 0, \quad (3.44)$$

which means the corrections $R_{1,2,3}$ must all vanish, as a consequence of $D_d \neq 0$. Once the validity of the three unity blowup equations in (3.42) is established, using the universality we can argue for the validity of all unity blowup equations. The recursion formula constructed out of any three unity blowup equations then should always gives the correct \mathbb{E}_d which coincides with the solution of (3.42).

3.4 Vanishing blowup equations

We consider here the case where $\lambda \in P \setminus Q^\vee$. This is only possible if $|P : Q^\vee| > 1$, i.e. for the minimal 6d $\mathcal{N} = (1, 0)$ SCFTs with $G = \text{SU}(3), \text{SO}(8), E_6, E_7$. In this case, $\phi_\lambda(Q^\vee) \neq Q^\vee$; $||\omega||^2/2$ for any $\omega \in \phi_\lambda(Q^\vee)$ and thus d is not an integer. In fact we find

$$\frac{||\omega||^2}{2} = \frac{\mathbf{n} - 2}{\mathbf{n}} + k, \quad k \in \mathbb{Z}_{\geq 0}, \quad (3.45)$$

where the minimum norm square $(\mathbf{n} - 2)/\mathbf{n}$ is reached if and only if ω is in a lowest dimensional irreducible representation²⁴ \square_G of G . As a consequence, the right hand side of the elliptic blowup equations (3.4) vanishes

$$\begin{aligned} & \frac{1}{2} ||\omega||^2 + d_1 + d_2 = d \\ & \sum_{\omega \in \phi_\lambda(Q^\vee), d_{1,2} \in \mathbb{N}} (-1)^{|\phi_\lambda^{-1}(\omega)|} \cdot \theta_i^{[a]}(\mathbf{n}\tau, (\mathbf{n}-2)(\epsilon_1 + \epsilon_2) - \mathbf{n}((\frac{1}{2}||\omega||^2 + d_1)\epsilon_1 + (\frac{1}{2}||\omega||^2 + d_2)\epsilon_2 - m_\omega)) \\ & \times A_\omega(m) \cdot \mathbb{E}_{d_1}(\tau, m - \epsilon_1\omega, \epsilon_1, \epsilon_2 - \epsilon_1) \cdot \mathbb{E}_{d_2}(\tau, m - \epsilon_2\omega, \epsilon_1 - \epsilon_2, \epsilon_2) = 0, \quad \lambda \notin Q^\vee \text{ fixed } d. \end{aligned} \quad (3.46)$$

Following the nomenclature of [15], we call these equations of the *vanishing* type.

The number of inequivalent embeddings ϕ_λ of this kind is $|P : Q^\vee| - 1$, which happens to be the order of the automorphism group of the Dynkin diagram Γ_G for

²⁴This can be the fundamental representation, the anti-fundamental representation, and in the case of $\text{SO}(8)$ the two spinor representations.

$G = \text{SU}(3), \text{SO}(8), E_6, E_7$. As representatives we can choose λ to be a fundamental weight²⁵ that generates \square_G as the highest weight. The total number of inequivalent vanishing blowup equations for each of these models is $\mathfrak{n}(|P : Q^\vee| - 1)$. Furthermore, using the property that the leading Q_τ order of \mathbb{E}_d is $-dh_G^\vee/6$, the equations (3.46) at the leading order requires that

$$\sum_{\alpha \in \Delta^+} ((\omega \cdot \alpha)^2 - (\omega_0 \cdot \alpha)^2) = 2kh_G^\vee, \quad \omega \in \phi_\lambda(Q^\vee), \omega_0 \in \square_G, \quad (3.47)$$

where the integer k is associated to the weight vector ω by (3.45), and it is again guaranteed by the Lie algebraic identity (A.5).

With the elliptic genus solved from the recursion formulas (3.32) plugged in, the vanishing blowup equations give rise to infinitely many nontrivial identities of Jacobi forms. In the lowest order, $d = (\mathfrak{n} - 2)/\mathfrak{n}$ and $\phi_\lambda(Q^\vee)$ is chosen to be one of the lowest dimensional representations

$$\sum_{\omega \in \square_G} (-1)^{|\phi_\lambda^{-1}(\omega)|} \theta_i^{[a]}(\mathfrak{n}\tau, \mathfrak{n}m_\omega) \prod_{\substack{\beta \in \Delta_+ \\ \omega \cdot \beta = \pm 1}} \frac{1}{\theta_1(m_\beta)} = 0. \quad (3.48)$$

This elegant formula specializing to $G = \text{SU}(3)$ and $\text{SO}(8)$ has been explicitly shown and checked in [25]. Here we further checked it for E_6 and E_7 for various characteristic a to higher order of Q_τ . For example, for E_6 , the relevant representation is **27**, with the weights encoded in the character

$$\chi_{\mathbf{27}}^{E_6} = \sum_{i=1}^{27} \prod_{j=1}^6 e^{m_j w_{ij}}. \quad (3.49)$$

Then for arbitrary $a \in \mathbb{Z}/6$, the following identity holds:²⁶

$$\sum_{i=1}^{27} (-1)^{|\phi_\lambda^{-1}(\omega_i)|} \theta_3^{[a]}(6\tau, 6m_{\omega_i}) \prod_{\substack{\beta \in \Delta_+(E_6) \\ \omega_i \cdot \beta = \pm 1}} \frac{1}{\theta_1(m_\beta)} = 0. \quad (3.51)$$

Note there actually exist two **27** representations due to the symmetry of Dynkin diagram of E_6 , i.e. $|\text{Aut}(\Gamma_{E_6})| = 2$, both of them make (3.51) holds. This also explains why there are two copies of vanishing r field for E_6 geometry, as we will see in next section in table 9. For higher base degree of the vanishing blowup equations, one can also write down some more complicated identities like (3.48). We checked them for all $G = \text{SU}(3), \text{SO}(8), E_{6,7}$ in the setting of refined BPS expansion to very high orders.

²⁵Not all the fundamental weights generate a lowest dimensional irreducible representation. For instance, the fundamental weight of $\text{SO}(8)$ corresponding to the central node in the Dynkin diagram $\Gamma_{\text{SO}(8)}$ generates the adjoint representation. The nodes associated to the \square_G -generating fundamental weights are permuted precisely by the automorphism group $\text{Aut}(\Gamma_G)$.

²⁶Let us write the notations in components in case of any misunderstanding.

$$|\phi_\lambda^{-1}(\omega_i)| = \sum_{j,k=1}^6 (C_{E_6}^{-1})_{jk} \omega_{ij}, \quad m_{\omega_i} = \sum_{j,k=1}^6 m_k (C_{E_6}^{-1})_{jk} \omega_{ij}, \quad \omega_i \cdot \beta = \sum_{j=1}^6 \omega_{ij} \beta_j, \quad m_\beta = \sum_{j=1}^6 m_j \beta_j, \quad (3.50)$$

where C_{E_6} is the Cartan matrix.

G	SU(3)	SO(8)	E_6	E_7
$\dim(\square)$	3	8	27	56
$\dim(\Delta_+)$	3	12	36	63
$\#\{\beta \in \Delta_+ : \omega \cdot \beta = 0, \forall \omega \in \square\}$	1	6	20	36
$\#\{\beta \in \Delta_+ : \omega \cdot \beta = \pm 1, \forall \omega \in \square\}$	2	6	16	27
$\#\{\beta \in \Delta_+ : \omega \cdot \beta = \pm 2, \forall \omega \in \square\}$	0	0	0	0

Table 4. Distribution of positive roots with respect to product with weights in \square_G .

We list various Lie theoretical data including the distribution of positive roots with respect to product with any weight in \square_G in table 4, from which one can check (3.47) indeed holds.

4 Elliptic genera for 6d (1, 0) minimal SCFTs

In this section we illustrate explicitly the solution of one-string and two-string elliptic genera of minimal 6d (1,0) SCFTs with $G = F_4, E_6, E_7, E_8$, using the elliptic blowup equations. The elliptic genera of the minimal theories with $G = \text{SU}(3), \text{SO}(8)$ have been computed in [8, 34], and we also reproduce some relevant results here. From these concrete results we summarise some universal features of the elliptic genera, including the expansion coefficients, the symmetric product approximation, and some additional symmetries. They are presented immediately in the first subsection, which one can then check in the following example subsections.

In this and the next sections, we work with the reduced elliptic genera which has the center-of-mass degree of freedom removed:²⁷

$$\mathbb{E}_{h_G^{(k)}} = \mathbb{E}_{\tilde{h}_G^{(k)}} / \mathbb{E}_{\text{c.m.}}, \quad (4.1)$$

where

$$\mathbb{E}_{\text{c.m.}} = -\frac{\theta_1(\epsilon_1)\theta_1(\epsilon_2)}{\eta^2}. \quad (4.2)$$

In the reduced version, elliptic genera normally obtain simplification. For example, the reduced one-string elliptic genus is independent from ϵ_- , i.e. $\text{SU}(2)_l$, as expected.

4.1 Universal behaviors of elliptic genera

4.1.1 Universal expansion

For all possible gauge group G , recall $v \equiv \exp(\pi i(\epsilon_1 + \epsilon_2))$ and $x \equiv \exp(\pi i(\epsilon_1 - \epsilon_2))$, we propose the following general ansatz for the reduced k -string elliptic genera

$$\mathbb{E}_{h_G^{(k)}}(v, x, Q_\tau, Q_{m_i}) = v^{kh_G^\vee - 1} Q_\tau^{-(kh_G^\vee - 1)/6} \sum_{n=0}^{\infty} Q_\tau^n g_{k,G}^{(n)}(v, x, Q_{m_i}). \quad (4.3)$$

²⁷We follow the notation of [37] where $\mathbb{E}_{\tilde{h}_G^{(k)}}$ is used to stress it is the RR elliptic genus of underlying 2d (0,4) CFT associated to the k -strings in 6d (1,0) minimal SCFT with gauge group G . It is the same with what we previously denoted as \mathbb{E}_k to emphasize that it is coefficient of base degree k in the topological string partition function.

Here all $g_{k,G}^{(n)}(v, x, Q_{m_i})$ are rational functions. In particular, $g_{1,G}^{(n)}$ is independent from x . One obvious symmetry for all $g_{k,G}^{(n)}$ is

$$g_{k,G}^{(n)}(v, x, Q_{m_i}) = g_{k,G}^{(n)}(v, x^{-1}, Q_{m_i}), \quad (4.4)$$

which comes from the symmetry between ϵ_1 and ϵ_2 in the Omega background, and can be understood as the Weyl symmetry of $SU(2)_x$. From now on we use $SU(2)_x$ to denote $SU(2)_l$ symmetry to stress the associated fugacity is x . We can further compute the v -expansion of each $g_{k,G}^{(n)}$ function where the coefficients are finite sum of products between the characters of $SU(2)_x$ and characters of G which respect Weyl symmetries of both groups. For example, $g_{k,G}^{(0)} = 1 + \dots$ gives the Hilbert series of the reduced k G -instanton moduli space. In fact, we find plenty of universal coefficients for the first a few order v -expansion of $g_{k,G}^{(n)}$.

It is known that the Hilbert series of the reduced one-instanton moduli space for any simple gauge group G has the expansion [41]

$$g_{1,G}^{(0)}(v, Q_{m_i}) = \sum_{k=0}^{\infty} \chi_{n\theta}^G v^{2n}, \quad (4.5)$$

where $\chi_{k\theta}$ is the character of the representation whose highest weight is k -multiple of the longest root θ ; in particular χ_θ is the character of the adjoint representation of G . In particular this is true for $G = SU(3), SO(8), F_4, E_{6,7,8}$ when $g_{1,G}^{(0)}$ serves as the leading contribution to one-string elliptic genus. As for sub- and subsub-leading contributions, we find that except for $G = SU(3)$ ²⁸

$$g_{1,G}^{(1)}(v, Q_{m_i}) = 1 + \chi_\theta + \left(1 + \chi_\theta + \chi_{2\theta} + \chi_{\text{Alt}^2\theta}\right)v^2 + \left(2\chi_{2\theta} + \chi_{\text{Alt}^2\theta} + \chi_{3\theta} + B_2(G)\right)v^4 + \mathcal{O}(v^6), \quad (4.6)$$

while except for $G = SU(3), SO(8)$,

$$g_{1,G}^{(2)}(v, Q_{m_i}) = 2 + 2\chi_\theta + \chi_{\text{Sym}^2\theta} + \mathcal{O}(v^2). \quad (4.7)$$

Here $B_2(G)$ are characters of some representations for which we do not find any universal expressions, and we list them in table 5.²⁹ The exceptions of $SU(3)$ and $SO(8)$ can be explained by the higher structures of \mathbb{E}_1 revealed by its intriguing relation with the Schur indices of certain rank one 4d SCFTs discovered in [37], which we will review and extend in section 5.

Furthermore, we find the Hilbert series of reduced two-instanton moduli space for any simple gauge group G has the expansion

$$\begin{aligned} g_{0,G}^{(2)}(v, x, Q_{m_i}) &= 1 + (\chi_\theta + \chi_3)v^2 + \chi_\theta\chi_2v^3 + (\chi_5 + \chi_\theta\chi_3 + \chi_{\text{Sym}^2\theta})v^4 + (\chi_\theta\chi_4 + (\chi_{2\theta} + \chi_{\text{Alt}^2\theta})\chi_2)v^5 \\ &+ (\chi_7 + \chi_5\chi_\theta + \chi_3(\chi_{\text{Sym}^2\theta} + \chi_{2\theta}) + \chi_{\text{Sym}^3\theta} - C_6(G))v^6 \\ &+ \left(\chi_\theta\chi_6 + (\chi_{2\theta} + \chi_{\text{Alt}^2\theta})\chi_4 + (\chi_{2\theta} + \chi_{3\theta} + \chi_{\text{Alt}^2\theta} + B_2(G) + C_7(G))\chi_2\right)v^7 \\ &+ \left(\chi_9 + \chi_7\chi_\theta + \chi_5(\chi_{\text{Sym}^2\theta} + \chi_{2\theta}) + \chi_3(\chi_{3\theta} + \chi_{2\theta} + B_2(G) + \chi_{\text{Sym}^3\theta} - C_6(G)) + \chi_{\text{Sym}^4\theta} - C_8(G)\right)v^8 \\ &+ \left(\chi_\theta\chi_8 + (\chi_{2\theta} + \chi_{\text{Alt}^2\theta})\chi_6 + (\chi_{2\theta} + 2\chi_{3\theta} + \chi_{\text{Alt}^2\theta} + B_2(G) + C_7(G))\chi_4 + \dots\right)v^9 + \dots \end{aligned} \quad (4.8)$$

²⁸From now on, to shorten formulas, we do not explicitly write G in each character.

²⁹The bold numbers mean the character of representations with dimension of such number. Note different representations can have the same dimension sometimes, for instance, the representations $\mathbf{35}_v$, $\mathbf{35}_s$ and $\mathbf{35}_c$ of $SO(8)$. To lighten the notation, we do not distinguish them in the table. Nevertheless, they can be recovered by taking into account the symmetry of Dynkin diagrams.

G	χ_θ	$\chi_{2\theta}$	$\chi_{3\theta}$	B_2	C_6	C_7	C_8
A_2	8	27	64	$2 \cdot \mathbf{35}$	1	27	8
D_4	28	300	1925	4096	$2 \cdot \mathbf{28}$	$3 \cdot \mathbf{567}$	$2 \cdot (\mathbf{300} + \mathbf{350} + \mathbf{1}) + 3 \cdot \mathbf{35}$
F_4	52	1053	12376	29172	273	10829	$\mathbf{8424} + \mathbf{4096} + \mathbf{324} + \mathbf{26}$
E_6	78	2430	43758	105600	650	34749	$\mathbf{34749} + 2 \cdot \mathbf{5824} + \mathbf{650} + \mathbf{78}$
E_7	133	7371	238602	573440	1463	152152	$\mathbf{150822} + \mathbf{40755} + \mathbf{1539}$
E_8	248	27000	1763125	4096000	0	779247	147250

Table 5. Certain representations appearing in the expansion of $g_{k,G}^{(n)}$ functions.

Here χ_n is the character of n -dimensional representation of $SU(2)_x$. The expansion coefficients up to v^6 were already observed in [30], and we further push the observation up to v^8 . We have checked this expression to be consistent with all the results on Hilbert series of reduced two G instanton moduli space in [42]. In particular it is true for $G = SU(3), SO(8), F_4, E_{6,7,8}$ when $g_{2,G}^{(0)}$ is the leading contribution to the two-string elliptic genera. Note that in this expression, $C_6(G), C_7(G)$ are characters of certain representations of G for which universal expressions are not found. They are collected for individual G in table 5. As for the subleading and subsubleading contribution to the two-string elliptic genera, we find there exists the following universal v -expansion: except for $G = SU(3)$,

$$\begin{aligned}
 g_{1,G}^{(2)}(v, x, Q_{m_i}) &= 1 + \chi_\theta + \chi_3 + (\chi_\theta + 1)\chi_2 v + (\chi_5 + (2\chi_\theta + 3)\chi_3 + (\chi_\theta + 1)^2)v^2 \\
 &\quad + \left((2\chi_\theta + 1)\chi_4 + (\chi_{2\theta} + (\chi_\theta + 1)^2 + \chi_{\text{Sym}^2\theta})\chi_2 \right) v^3 \\
 &\quad + (\chi_7 + (2\chi_\theta + 3)\chi_5 + \dots)v^4 + \mathcal{O}(v^5),
 \end{aligned} \tag{4.9}$$

while except for $G = SU(3)$ and $SO(8)$,

$$\begin{aligned}
 g_{2,G}^{(2)}(v, x, Q_{m_i}) &= (\chi_5 + (\chi_\theta + 2)\chi_3 + \chi_{\text{Sym}^2\theta} + 2\chi_\theta + 4) + \left((\chi_\theta + 1)\chi_4 + ((\chi_\theta + 1)^2 + 2(\chi_\theta + 1))\chi_2 \right) v \\
 &\quad + \left(\chi_7 + (2\chi_\theta + 4)\chi_5 + ((\chi_\theta + 1)^2 + 2\chi_{\text{Sym}^2\theta} + 6\chi_\theta + 9)\chi_3 + \dots \right) v^2 + \mathcal{O}(v^3).
 \end{aligned} \tag{4.10}$$

For the reduced three string elliptic genus $\mathbb{E}_{h_G^{(3)}}$, although we have not checked for all six G due to the complexity of computation, still we propose the following universal expansion:

$$\begin{aligned}
 g_{0,G}^{(3)}(v, x, Q_{m_i}) &= 1 + (\chi_3 + \chi_\theta)v^2 + (\chi_4 + \chi_\theta\chi_2)v^3 + \left(\chi_5 + \chi_\theta\chi_3 + \chi_{\text{Sym}^2\theta} + 1 \right) v^4 \\
 &\quad + \left(\chi_6 + (2\chi_\theta + 1)\chi_4 + 2\chi_{\text{Sym}^2\theta} \right) v^5 \\
 &\quad + \left(2\chi_7 + 3\chi_\theta\chi_5 + (\chi_{2\theta} + 3\chi_{\text{Sym}^2\theta} + 1)\chi_3 + \chi_{\text{Sym}^3\theta} + \chi_{\text{Sym}^2\theta} \right) v^6 + \mathcal{O}(v^7).
 \end{aligned} \tag{4.11}$$

We have checked this against the three-instanton Hilbert series for $SU(2), G_2, SO(7), Sp(4), Sp(6)$ in [43, 60], and against the three-string elliptic genus for $SU(3)$ [34]. Note the first

two terms also agree with the rank three E_6 Hall-Littlewood index ((A.14) in [45]). For the subleading Q_τ order, again except $SU(3)$, we propose

$$g_{1,G}^{(3)}(v, x, Q_{m_i}) = (\chi_3 + \chi_\theta + 1) + (\chi_4 + (\chi_\theta + 1)\chi_2)v + (\chi_5 + (3\chi_\theta + 4)\chi_3 + 2\chi_{\text{Sym}^2\theta} + \chi_\theta + 2)v^2 + \mathcal{O}(v^3). \quad (4.12)$$

As in rank one and two cases, for $SU(3)$, the higher contributions begin to merge in at Q_τ subleading order.

All above v -expansion coefficients can be easily obtained by setting $Q_m = 1$ in $g_{n,G}^{(k)}(v, x, Q_m)$. Thus the rational functions $g_{n,G}^{(k)}(v, x, 1)$ are very useful as they encode most information. For large k or n , such rational functions with generic x turn out to be too lengthy. One can take the unrefined limit $x = 1$ in $g_{n,G}^{(k)}(v, x, 1)$ to still store meaning information on arbitrary order coefficients of v -expansion. Indeed, when the fugacities of flavor as well as $SU(2)_x$ are turned off, we find

$$g_{1,G}^{(n)}(v) = \frac{1}{(1-v^2)^{2(h_G^\vee-1)}} \times P_{1,G}^{(n)}(v), \quad (4.13)$$

$$g_{2,G}^{(n)}(v) = \frac{1}{(1-v^2)^{2(h_G^\vee-1)}(1+v)^{2b_G}(1+v+v^2)^{2h_G^\vee-1}} \times P_{2,G}^{(n)}(v). \quad (4.14)$$

The exponents b_G are given by

G	$SU(3)$	$SO(8)$	F_4	E_6	E_7	E_8
b	3	6	11	16	26	46

We notice that $b = 5h_G^\vee/3 - 4$ except for $SU(3)$. The numerators $P_{1,G}^{(n)}(v)$ and $P_{2,G}^{(n)}(v)$ are palindromic Laurent polynomials. They have negative powers of v when n is large. Nevertheless $P_{1,G}^{(0)}(v), P_{2,G}^{(0)}(v)$ are both polynomials and their maximum degrees are $h_G^\vee - 1$ and $2(2h_G^\vee - 1) + 2b_G$ respectively. The explicit expressions of $P_{k,G}^{(n)}(v)$ for the minimal SCFTs with $G = SU(3), SO(8), F_4, E_6, E_7, E_8$ are presented in the following example subsections and also appendix F.

4.1.2 Symmetric product approximation

It was noticed both in [42] and [45] that the reduced two G -instanton Hilbert series can be realized as certain symmetric product of two one G -instantons as approximation:

$$\frac{1}{1-vx^{\pm 1}} g_{0,G}^{(2)}(v, x, a) = \frac{1}{2} \left(\left(g_{0,G}^{(1)}(v, a) \frac{1}{1-vx^{\pm 1}} \right)^2 + g_{0,G}^{(1)}(v^2, a^2) \frac{1}{1-v^2x^{\pm 2}} \right) + \mathcal{O}(v^4). \quad (4.15)$$

Here we adopt their notation $a = Q_{m_G}$ to lighten the notation. It also was noticed in [43] that the reduced three G -instanton Hilbert series can be realized as certain symmetric product of three one G -instantons as approximation:

$$\begin{aligned} \frac{1}{1-vx^{\pm 1}} g_{0,G}^{(3)}(v, x, a) &= \frac{1}{6} \left(\left(\frac{1}{1-vx^{\pm 1}} g_{0,G}^{(1)}(v, a) \right)^3 + \frac{3}{(1-vx^{\pm 1})(1-v^2x^{\pm 2})} g_{0,G}^{(1)}(v, a) g_{0,G}^{(1)}(v^2, a^2) \right. \\ &\quad \left. + \frac{2}{1-v^3x^{\pm 3}} g_{0,G}^{(1)}(v^3, a^3) \right) + \mathcal{O}(v^4). \end{aligned} \quad (4.16)$$

The above formulas have clear physical meaning. For example in (4.16), the first term represents the configuration where three instantons are far from each other, the second term represents the configuration where two instantons sit on the same site and the third one is far from them, while the third term represents the configuration where all three instantons sit on the same site. Note the triple symmetric product would give the coefficient of v^4 of $g_{0,G}^{(3)}$ as $\chi_5 + \chi_\theta \chi_3 + \chi_{2\theta} + \chi_{\text{Sym}^2 \theta} + 1$, one can see the difference with (4.11) begins to appear.

In fact, it is reasonable that arbitrary k G -instanton Hilbert series can be realized as symmetric product of k G -instantons as approximation:

$$\frac{1}{1 - vx^{\pm 1}} g_{0,G}^{(k)}(v, x, a) = \text{Sym}_{\mathcal{M}_{G,1}}^k(v, x, a) + \mathcal{O}(v^4), \quad (4.17)$$

where $\text{Sym}_{\mathcal{M}_{G,1}}^k(v, x, a)$ can be obtained from generating function

$$\sum_{k=1}^{\infty} \text{Sym}_{\mathcal{M}_{G,1}}^k(v, x, a) Q^k = \text{PE} \left[\frac{Q}{1 - vx^{\pm 1}} g_{0,G}^{(1)}(v, a) \right] \equiv \text{PE} \left[\tilde{g}_{0,G}^{(1)}(v, x, a) Q \right]. \quad (4.18)$$

For example,

$$\begin{aligned} \text{Sym}_{\mathcal{M}_{G,1}}^4(v, x, a) &= \frac{1}{24} \left(\left(\tilde{g}_{0,G}^{(1)}(v, x, a) \right)^4 + 6 \left(\tilde{g}_{0,G}^{(1)}(v, x, a) \right)^2 \tilde{g}_{0,G}^{(1)}(v^2, x^2, a^2) + 3 \left(\tilde{g}_{0,G}^{(1)}(v^2, x^2, a^2) \right)^2 \right. \\ &\quad \left. + 8 \tilde{g}_{0,G}^{(1)}(v, x, a) \tilde{g}_{0,G}^{(1)}(v^3, x^3, a^3) + 6 \tilde{g}_{0,G}^{(1)}(v^4, x^4, a^4) \right) \\ &= 1 + (\chi_3 + \chi_\theta) v^2 + (\chi_4 + \chi_\theta \chi_2) v^3 + \mathcal{O}(v^4). \end{aligned} \quad (4.19)$$

It is not hard to find that for all $k \geq 3$, the leading coefficients in v expansion of symmetric product are the same:

$$\begin{aligned} \text{Sym}_{\mathcal{M}_{G,1}}^k(v, x, a) &= \frac{1}{k!} \left(\left(\tilde{g}_{0,G}^{(1)}(v, x, a) \right)^k + C_2^k \left(\tilde{g}_{0,G}^{(1)}(v, x, a) \right)^{k-2} \tilde{g}_{0,G}^{(1)}(a^2, v^2, x^2) + \dots \right) \\ &= 1 + (\chi_3 + \chi_\theta) v^2 + (\chi_4 + \chi_\theta \chi_2) v^3 + \mathcal{O}(v^4). \end{aligned} \quad (4.20)$$

Here the first term represents all k instantons are far from each other, while the second term represents two instantons sit at the same site and the rest $k - 2$ instanton are far from them and each other... From v^4 , the interaction among instantons will contribute.

We can also include $g_{0,G}^{(k)}$ into the elliptic genus to write down the above symmetric product approximation. For example in the reduced three-string elliptic genus, since

$$\mathbb{E}_{\tilde{h}_G^{(3)}}(v, x, a, Q_\tau) = \frac{v^{3h_G^\vee}}{1 - vx^{\pm 1}} g_{0,G}^{(3)}(v, x, a) Q_\tau^{-h_G^\vee/2} + \mathcal{O}(Q_\tau^{-h_G^\vee/2+1}), \quad (4.21)$$

combining (4.16), we obtain

$$\begin{aligned} \mathbb{E}_{\tilde{h}_G^{(3)}}(v, x, a, Q_\tau) &= \frac{1}{6} \left(\mathbb{E}_{\tilde{h}_G^{(1)}}(v, x, a, Q_\tau)^3 + 3 \mathbb{E}_{\tilde{h}_G^{(1)}}(v, x, a, Q_\tau) \mathbb{E}_{\tilde{h}_G^{(1)}}(v^2, x^2, a^2, Q_\tau^2) \right. \\ &\quad \left. + 2 \mathbb{E}_{\tilde{h}_G^{(1)}}(v^3, x^3, a^3, Q_\tau^3) \right) + \dots, \end{aligned} \quad (4.22)$$

which holds for the leading Q_τ order and the first four v -expansion coefficients. For arbitrary k -strings elliptic genus, it is better to use Hecke transformation. Neglecting the

interaction among strings, the resulting k -strings elliptic genus $\mathbb{E}_{\text{Sym}_G^{(k)}}(v, x, a, Q_\tau)$ can be generated from

$$\sum_{k=0} \mathbb{E}_{\text{Sym}_G^{(k)}}(v, x, a, Q_\tau) Q^k = \exp \left[\sum_{n \geq 0} Q^n \frac{1}{n} \sum_{\substack{cd=n \\ c, d > 0}} \sum_{b \pmod{d}} \mathbb{E}_{h_G^{(1)}} \left(\frac{c\tau + b}{d}, c\epsilon_i, cm_G \right) \right]. \quad (4.23)$$

Note this relies on the Jacobi form nature of $\mathbb{E}_{h_G^{(1)}}(\tau, \epsilon_i, m_G)$. Also take $d = 1$ in (4.23), one will go back to instanton formula (4.18) where there is no modularity. Finally, we obtain

$$\mathbb{E}_{h_G^{(k)}}(v, x, a, Q_\tau) = \mathbb{E}_{\text{Sym}_G^{(k)}}(v, x, a, Q_\tau) + \mathcal{O}(Q_\tau^{-kh_G^\vee/6+1}) + \mathcal{O}(v^{kh_G^\vee+4}). \quad (4.24)$$

As we have checked this symmetric product approximation does not give exact subleading Q_τ orders $g_{1,G}^{(k)}$ even for its leading v -expansion coefficient. This means all subleading Q_τ orders involves interaction among strings.

4.1.3 Symmetries

Besides the obvious symmetry

$$\mathbb{E}_{h_G^{(k)}}(v, x, Q_\tau, Q_m) = \mathbb{E}_{h_G^{(k)}}(v, x^{-1}, Q_\tau, Q_m), \quad (4.25)$$

which comes from the symmetry between ϵ_1 and ϵ_2 in Omega background, it was found in [37] that the reduced one-string elliptic genus $\mathbb{E}_{h_G^{(1)}}(v, Q_\tau)$ satisfies an additional symmetry

$$\mathbb{E}_{h_G^{(1)}}(Q_\tau^{1/2}/v, Q_\tau) = (-1)^{n+1} v^{2(1-h_G^\vee/3)} Q_\tau^{(h_G^\vee/3-1)/2} \mathbb{E}_{h_G^{(1)}}(v, Q_\tau). \quad (4.26)$$

Here the dependence on m_G is implicit. This symmetry was later interpreted as a spectral flow symmetry in [38]. The left hand side of (4.26) actually computes the NS-R elliptic genus, which should be equal to the R-R elliptic genus on the right hand side due to the lack of chiral fermions in the minimal SCFT in consideration. See section 6.4 of [37] for a detailed discussion.

We extend the symmetry (4.26) to arbitrary k -string elliptic genus $\mathbb{E}_{h_G^{(k)}}(v, x, Q_\tau)$:

$$\mathbb{E}_{h_G^{(k)}} \left(\frac{Q_\tau^{1/2}}{v}, \frac{Q_\tau^{1/2}}{x}, Q_\tau \right) = (-1)^{nk+1} v^{\frac{k(k-5)}{6} h_G^\vee + k^2 + 1} x^{-\frac{k(k-1)}{6} h_G^\vee - k^2 + 1} Q_\tau^{(kh_G^\vee - 3)/6} \mathbb{E}_{h_G^{(k)}}(v, x, Q_\tau), \quad (4.27)$$

which can be derived by combining (3.25) and the modular anomaly of elliptic genera (3.12). For the situation where 2d quiver description is known, i.e. $G = \text{SU}(3)$ and $\text{SO}(8)$, the above symmetry can also be obtained by looking into the transformation of integrand of localization with the quasi-periodicity of Jacobi theta function (D.5), (D.6). Note symmetry (4.27) is a nonperturbative symmetry, which can not be seen from the Q_τ expansion of the elliptic genus, except for the one-string case that is (4.26).³⁰ This means (4.27) should be seen as

³⁰Practically, we find that for the two-string elliptic genus, when Q_τ order is enough high, for one order of Q_τ goes up, the leading v order goes down for 3. Thus, if one naively does the transformation for the left hand side of (4.27) in Q_τ expansion, one would get infinite negative order of Q_τ . Similar situation also happens for three-string elliptic genus. But for one-string elliptic genus, luckily for one order of Q_τ goes up, the leading v order goes down for 2, which only result in finite negative order of Q_τ .

	r	old r	fundamental weights
unity	(0, 0, 0, 1)	(0, 0, 0, 1)	$\omega_1 + \omega_2$
	(0, 0, 0, 3)	(0, 0, 0, 3)	
	(0, 0, 0, 5)	(0, 0, 0, -1)	
vanishing	(-2, 2, 0, 1)	(-2, 2, 0, 1)	ω_1
	(-2, 2, 0, 3)	(0, -2, 2, 1)	
	(-2, 2, 0, 5)	(2, 0, -2, 1)	
vanishing	(-2, 0, 2, 1)	(-2, 0, 2, 1)	ω_2
	(-2, 0, 2, 3)	(0, 2, -2, 1)	
	(-2, 0, 2, 5)	(2, -2, 0, 1)	

Table 6. The r -fields of the $\mathfrak{n} = 3$ model and the fundamental weights of \mathfrak{a}_2 which induce the embedding $\phi : Q^\vee \hookrightarrow P$. The old r -fields are from our previous paper [25]. They are equivalent with the r in the second column by $2C \cdot \underline{n}$ shift.

the symmetry of the chiral algebra associated to the underlying (0, 4) 2d CFT, as suggested in [37].

4.2 Revisiting $G = \text{SU}(3)$ and $\text{SO}(8)$

With the new understanding on the structure of r fields of 6d (1, 0) minimal SCFTs for all G , we now can reproduce all r fields for $\text{SU}(3)$ and $\text{SO}(8)$ found in [25] using just the fundamental weights of the Lie algebras. We summarize the correspondence between the r fields given in [25] and fundamental weights in tables 6 and 7.

The elliptic genera of 6d (1, 0) SCFT with $G = \text{SU}(3)$ were computed using Jeffrey-Kirwan residue in [34], and were checked to satisfy the elliptic blowup equations [25]. Following the general proposal (4.3), the reduced two-string elliptic genus for $\text{SU}(3)$ model can be written as

$$\mathbb{E}_{h_{A_2}^{(2)}}(v, x, Q_\tau, m_i) = v^5 Q_\tau^{-5/6} \sum_{n=0}^{\infty} Q_\tau^n g_{2,A_2}^{(n)}(v, x, Q_{m_i}), \tag{4.28}$$

where $g_{2,A_2}^{(n)}(v, x, Q_{m_i})$ are rational functions. We computed $g_{2,A_2}^{(n)}(v, x, Q_{m_i}=1)$ up to $n=10$. Let us turn off the fugacities of both $\text{SU}(3)$ and $\text{SU}(2)_x$, we obtain

$$g_{2,A_2}^{(n)}(v, x=1, Q_{m_i}=1) = \frac{1}{(1-v)^{10}(1+v)^6(1+v+v^2)^5} \times P_{2,A_2}^{(n)}(v), \tag{4.29}$$

where all $P_{2,A_2}^{(n)}(v)$ are palindromic Laurent polynomials, in which only $P_{2,A_2}^{(0)}(v)$ is a true polynomial:

$$P_{2,A_2}^{(0)}(v) = 1 + v + 6v^2 + 17v^3 + 31v^4 + 52v^5 + 92v^6 + 110v^7 + 112v^8 + 110v^9 + \dots + v^{16}. \tag{4.30}$$

Here the ellipsis is completed by making the expression palindromic. This agrees with the Hilbert series of reduced two $\text{SU}(3)$ -instanton moduli space in [42]. For the subleading

	r	old r	fundamental weights
unity	(0, 0, 0, 0, 0, 0)	(0, 0, 0, 0, 0, 0)	ω_c
	(0, 0, 0, 0, 0, 2)	(0, 0, 0, 0, 0, 2)	
	(0, 0, 0, 0, 0, 4)	(0, 0, 0, 0, 0, 4)	
	(0, 0, 0, 0, 0, 6)	(0, 0, 0, 0, 0, -2)	
vanishing	(-2, 2, 0, 0, 0, 0)	(-2, 2, 0, 0, 0, 0)	ω_1
	(-2, 2, 0, 0, 0, 2)	(-2, -2, 0, 0, 2, 2)	
	(-2, 2, 0, 0, 0, 4)	(0, 0, -2, 2, 0, 0)	
	(-2, 2, 0, 0, 0, 6)	(0, 0, -2, -2, 2, 2)	
vanishing	(-2, 0, 2, 0, 0, 0)	(-2, 0, 2, 0, 0, 0)	ω_2
	(-2, 0, 2, 0, 0, 2)	(-2, 0, 2, 0, 2, 2)	
	(-2, 0, 2, 0, 0, 4)	(0, -2, 0, 2, 0, 0)	
	(-2, 0, 2, 0, 0, 6)	(0, -2, 0, -2, 2, 2)	
vanishing	(-2, 0, 0, 2, 0, 0)	(-2, 0, 0, 2, 0, 0)	ω_3
	(-2, 0, 0, 2, 0, 2)	(-2, 0, 0, -2, 2, 2)	
	(-2, 0, 0, 2, 0, 4)	(0, -2, 2, 0, 0, 0)	
	(-2, 0, 0, 2, 0, 6)	(0, -2, -2, 0, 2, 2)	

Table 7. The r -fields of the $n = 4$ model and the fundamental weights of \mathfrak{d}_4 which induce the embedding $\phi : Q^\vee \hookrightarrow P$. The old r -fields are from our previous paper [25]. They are equivalent with the r in the second column by $2C \cdot \underline{n}$ shift.

order, $P_{2,A_2}^{(1)}(v)$ starts with negative power of v , which is *different* from all the other minimal SCFTs.³¹ Indeed,

$$P_{2,A_2}^{(1)}(v) = v^{-4}(1 + 3v + 8v^2 + 11v^3 + 18v^4 + 13v^5 + 55v^6 + 238v^7 + 601v^8 + 1121v^9 + 1777v^{10} + 2262v^{11} + 2424v^{12} + 2262v^{13} + \dots + v^{24}). \quad (4.31)$$

More results on $P_{2,A_2}^{(n)}(v)$ with $n > 1$ can be found in appendix F. Let us also show some results with generic fugacities, for example,

$$g_{0,A_2}^{(2)}(v, x, m_i) = 1 + (\mathbf{8} + \chi_3)v^2 + \mathbf{8}\chi_2v^3 + (\chi_5 + \mathbf{8}\chi_3 + \text{Sym}^2\mathbf{8})v^4 + (\mathbf{8}\chi_4 + (\mathbf{27} + \text{Alt}^2\mathbf{8})\chi_2)v^5 + (\chi_7 + \mathbf{8}\chi_5 + (\text{Sym}^2\mathbf{8} + \mathbf{27})\chi_3 + \text{Sym}^3\mathbf{8} - \mathbf{1})v^6 + \mathcal{O}(v^7), \quad (4.32)$$

$$v^4 g_{1,A_2}^{(2)}(v, x, m_i) = 1 + \chi_2v + (\chi_3 + \mathbf{8})v^2 + (\chi_4 + \mathbf{8}\chi_2)v^3 + (\chi_5 + (\mathbf{8} + \mathbf{1})\chi_3 + \text{Sym}^2\mathbf{8})v^4 + (\chi_6 + \mathbf{8}\chi_4 + \text{Sym}^2\mathbf{8}\chi_2)v^5 + \mathcal{O}(v^6). \quad (4.33)$$

Note (4.32) agrees with our universal expansion formula (4.8).

³¹This phenomenon as also occurring in one-string elliptic genus, will be discussed in detail in section 5.

Similarly, the reduced three-string elliptic genus for SU(3) model can be written as

$$\mathbb{E}_{h_{A_2}^{(2)}}(v, x, Q_\tau, m_i) = v^8 Q_\tau^{-4/3} \sum_{n=0}^{\infty} Q_\tau^n g_{3,A_2}^{(n)}(v, x, Q_{m_i}), \quad (4.34)$$

where all $g_{3,A_2}^{(n)}(v, x, Q_{m_i})$ are rational functions. We computed $g_{3,A_2}^{(n)}(v, x, Q_{m_i} = 1)$ up to $n = 6$. Turning off the fugacities of both SU(3) and SU(2)_x, we obtain

$$g_{3,A_2}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{16}(1+v)^{10}(1+v^2)^5(1+v+v^2)^6} \times P_{3,A_2}^{(n)}(v), \quad (4.35)$$

where all $P_{3,A_2}^{(n)}(v)$ are palindromic Laurent polynomials, in which only $P_{3,A_2}^{(0)}(v)$ is a true polynomial:

$$P_{3,A_2}^{(0)}(v) = 1 + 6v^2 + 14v^3 + 40v^4 + 82v^5 + 213v^6 + 388v^7 + 772v^8 + 1260v^9 + 2079v^{10} + 2986v^{11} + 4226v^{12} + 5226v^{13} + 6384v^{14} + 6940v^{15} + 7334v^{16} + 6940v^{17} + \dots + v^{32}, \quad (4.36)$$

$$P_{3,A_2}^{(1)}(v) = v^{-4}(1+v^2)(1+2v+8v^2+24v^3+62v^4+114v^5+242v^6+456v^7+964v^8+1926v^9 + 4225v^{10} + 8448v^{11} + 16317v^{12} + 28038v^{13} + 44954v^{14} + 64960v^{15} + 87437v^{16} + 106636v^{17} + 121046v^{18} + 125368v^{19} + 121046v^{20} + \dots + v^{38}). \quad (4.37)$$

Note $g_{3,A_2}^{(0)}$ agrees with our universal expansion formula (4.11). More higher $P_{3,A_2}^{(n)}(v)$ can be found in appendix F.

The elliptic genera for the 6d (1,0) SCFT with $G = \text{SO}(8)$ were computed using Jeffrey-Kirwan residue in [24], and they were checked to satisfy elliptic blowup equations in [25]. Let us write the reduced two-string elliptic genus as

$$\mathbb{E}_{h_{D_4}^{(2)}}(v, x, Q_\tau, m_i) = v^{11} Q_\tau^{-11/6} \sum_{n=0}^{\infty} Q_\tau^n g_{2,D_4}^{(n)}(v, x, m_i). \quad (4.38)$$

We computed $g_{2,D_4}^{(2)}(v, x, m_i = 0)$ up to $n = 6$. In particular, $g_{2,D_4}^{(2)}$ for $n = 0, 1, 2$ agree with our universal expansion formulas (4.8), (4.9). Turning off the SU(2)_x fugacity, we have

$$g_{2,D_4}^{(n)}(v, x = 1, m_i = 0) = \frac{1}{(1-v)^{22}(1+v)^{12}(1+v+v^2)^{11}} \times P_{2,D_4}^{(n)}(v). \quad (4.39)$$

Here $P_{2,D_4}^{(n)}(v)$ are palindromic Laurent polynomials. In particular, only for $n = 0, 1$, they are true polynomials:

$$P_{2,D_4}^{(0)}(v) = 1 + v + 20v^2 + 65v^3 + 254v^4 + 841v^5 + 2435v^6 + 6116v^7 + 14290v^8 + 29700v^9 + 55947v^{10} + 96519v^{11} + 152749v^{12} + 220408v^{13} + 293226v^{14} + 359742v^{15} + 406014v^{16} + 421960v^{17} + 406014v^{18} + \dots + v^{34}. \quad (4.40)$$

$$P_{2,D_4}^{(1)}(v) = (1+v^2)(32+90v+697v^2+2913v^3+10582v^4+34415v^5+97961v^6+242492v^7 + 540749v^8 + 1085137v^9 + 1958185v^{10} + 3205774v^{11} + 4789888v^{12} + 6522178v^{13} + 8110633v^{14} + 9248825v^{15} + 9668450v^{16} + 9248825v^{17} + \dots + v^{32}). \quad (4.41)$$

Note the above $g_{0,D_4}^{(2)}$ agrees with the SO(8) two-instanton Hilbert series in [42]. More results on $P_{2,D_4}^{(n)}(v)$ with $n > 1$ can be found in appendix F.

4.3 $G = F_4$

The divisors and curves of the non-compact $n = 5$ geometry are explained in [53]. There are five compact divisors, all of which are Hirzebruch surfaces \mathbb{F}_{n_i} of various degrees n_i . We denote them by \mathfrak{D}_I ($I = 0, 1, \dots, 5$). They intersect with each other like the affine dynkin diagram of \mathfrak{f}_4

$$\begin{array}{cccccc}
 \textcircled{\mathbb{F}_3} & \textcircled{\mathbb{F}_1} & \textcircled{\mathbb{F}_1} & \textcircled{\mathbb{F}_6} & \textcircled{\mathbb{F}_8} \\
 (0:1:1) & (1:2:2) & (2:3:3) & (3:4:2) & (4:2:1)
 \end{array}$$

where each node corresponds to a Hirzebruch surface, and two nodes are connected if the corresponding Hirzebruch surfaces intersect at a \mathbb{P}^1 normal to their respective \mathbb{P}^1 fibers. In the diagram above we also give the ordering of the nodes I and the associated marks/comarks a_I/a_I^\vee with the notation $(I : a_I : a_I^\vee)$ following [61]. The \mathbb{F}_3 denoted by a dashed circle corresponds to the affine node and it intersects with the base at the \mathbb{P}^1 with normal bundle $\mathcal{O}(-5) \oplus \mathcal{O}(3) \rightarrow \mathbb{P}^1$. The arrow with double line means the \mathbb{F}_1 and \mathbb{F}_6 intersect at a \mathbb{P}^1 which is the double cover of the $(+1)$ curve in \mathbb{F}_1 . See the illustration in figure 1. There are six linearly independent curves, which we choose for the moment to be the \mathbb{P}^1 fibers Σ_I of the divisors \mathfrak{D}_I and the (-5) curve in the base denoted by Σ_B . Denoting their complexified Kähler moduli by t_I and t_B , the linear combination

$$\sum_{I=0}^5 a_I t_I = \tau, \tag{4.42}$$

with a_I the marks of \mathfrak{f}_4 , is the volume of the elliptic fiber. Since we will be interested in the extraction of BPS invariants from the partition function, we would like to also identify among the compact curves the Mori cone generators. They include the \mathbb{P}^1 fibers Σ_I ($I = 0, 1, \dots, 4$), as well as the \mathbb{P}^1 base of the \mathbb{F}_1 surface that intersects with \mathbb{F}_6 (see the Dynkin diagram above and the figure 7 in [53], which we reproduce in figure 1). We denote the latter curve by Σ_b , and it is related to Σ_B by

$$[\Sigma_B] = [\Sigma_b] + 3[\Sigma_0] + [\Sigma_1]. \tag{4.43}$$

This implies the relation between their Kähler moduli

$$t_B = t_b + 3t_0 + t_1. \tag{4.44}$$

The C -matrix of intersection between Σ_I, Σ_b and D_I ³²

$$C = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 \\ 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & -2 & 2 & 0 \\ 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 1 & -2 \\ 0 & -1 & -1 & 0 & 0 \end{pmatrix}. \tag{4.45}$$

³²Note that here we use Σ_b instead of Σ_B , which is why the matrix C does not follow exactly the pattern (2.2).

	r	fundamental weights
	$(0, 0, 0, 0, 0, 1)$	
	$(0, 0, 0, 0, 0, 3)$	
unity	$(0, 0, 0, 0, 0, 5)$	ω_i ($i = 1, \dots, 4$)
	$(0, 0, 0, 0, 0, 7)$	
	$(0, 0, 0, 0, 0, 9)$	

Table 8. The r -fields of the $\mathfrak{n} = 5$ model and the fundamental weights of \mathfrak{f}_4 which induce the same embedding $\phi : Q^\vee \hookrightarrow P$. All the r -fields and all the fundamental weights induce the same embedding as $Q^\vee = P$ for \mathfrak{f}_4 .

The semiclassical components of the partition function can be computed using the prescription in section 2.2 with the normalisation scheme in appendix B. We obtain

$$\begin{aligned}
 -F_{(0,0)}^{\text{cls}} = & \left(\frac{t_0}{10} + \frac{t_1}{5} + \frac{3t_2}{10} + \frac{2t_3}{5} + \frac{t_4}{5} \right) t_b^2 + \left(\frac{3t_0^2}{10} + \frac{t_0 t_1}{5} + \frac{t_1^2}{5} + \frac{3t_2^2}{10} + \frac{6t_3^2}{5} + \frac{4t_4^2}{5} + \frac{4t_2 t_3}{5} \right. \\
 & \left. + \frac{2t_2 t_4}{5} + \frac{6t_3 t_4}{5} \right) t_b + \frac{3t_0^3}{10} + \frac{3t_0^2 t_1}{10} + \frac{t_0 t_1^2}{10} + \frac{t_1^3}{15} + \frac{t_2^3}{10} + \frac{6t_3^3}{5} + \frac{16t_4^3}{15} + \frac{6t_2 t_3^2}{5} \\
 & + \frac{4t_2 t_4^2}{5} + \frac{12t_3 t_4^2}{5} + \frac{2t_2^2 t_3}{5} + \frac{t_2^2 t_4}{5} + \frac{9t_3^2 t_4}{5} + \frac{6t_2 t_3 t_4}{5},
 \end{aligned} \tag{4.46}$$

which is consistent with the universal formula (2.18). Furthermore, using the relations (2.4), (4.44) and (4.42), $F_{(0,0)}^{\text{cls}}$ can be more succinctly written as

$$-F^{(0,0)} = \frac{1}{10} t_{\text{ell}}^2 \tau + \frac{1}{2} t_{\text{ell}}(m, m) - \frac{3}{4} \tau(m, m) + \dots \tag{4.47}$$

up to τ^3 and terms cubic in m_i , which agrees with the universal formula (2.19). Therefore the analysis in section 2.3 goes through. Here for F_4 ,

$$m = \sum_{i=1}^4 m_i \omega_i^\vee, \tag{4.48}$$

and

$$(m, m) = m_1^2 + 3m_1 m_2 + 3m_2^2 + 4m_1 m_3 + 8m_2 m_3 + 6m_3^2 + 2m_1 m_4 + 4m_2 m_4 + 6m_3 m_4 + 2m_4^2. \tag{4.49}$$

We also find

$$F_{(1,0)}^{\text{cls}} = \frac{18}{5} t_0 + \frac{16}{5} t_1 + 6t_2 + 12t_3 + 8t_4 + \frac{36}{5} t_b. \tag{4.50}$$

Imposing the admissibility condition (2.22) and the BPS checkerboard pattern condition (2.25), which specialises to

$$r \equiv (0, 0, 0, 0, 0, 1) \pmod{2}, \tag{4.51}$$

there are only five inequivalent r -fields, and we list their representatives in table 8. According to the discussion in section 3, we should classify them according to the embedding

$\phi_\lambda : Q^\vee \hookrightarrow P$ induced by the reduced r -vector λ defined in (2.28). In the case of the $\mathfrak{n} = 5$ model, all the r -fields have the same reduced $\lambda = (0, 0, 0, 0)$, which induces the unique embedding $\phi_{(0,0,0,0)} : Q^\vee \hookrightarrow P = Q^\vee$. As a consequence, this model has no blowup equation of the vanishing type. We notice that all the fundamental weights ω_i also induce the same embedding (which is not the case in all the other models.)

We use (3.33) and (3.36) to compute the one-string and two-string elliptic genera. The one-string elliptic genus does not depend on $SU(2)_x$. Its expansion in Q_τ reads

$$\mathbb{E}_{h_{F_4}^{(1)}}(v, Q_\tau, m_i) = v^8 Q_\tau^{-4/3} \sum_{n=0}^{\infty} g_{1,F_4}^{(n)}(v, Q_{m_i}) Q_\tau^n, \quad (4.52)$$

where $g_{1,F_4}^{(n)}(v, Q_{m_i})$ are rational functions. Turning off all flavor fugacities,

$$g_{1,F_4}^{(n)}(v, Q_{m_i} = 1) = \frac{1}{(1-v^2)^{16}} \times P_{1,F_4}^{(n)}(v), \quad (4.53)$$

where

$$\begin{aligned} P_{1,F_4}^{(0)}(v) &= 1 + 36v^2 + 341v^4 + 1208v^6 + 1820v^8 + 1208v^{10} + 341v^{12} + 36v^{14} + v^{16}, \\ P_{1,F_4}^{(1)}(v) &= (1+v^2)^2 (53 + 1478v^2 + 9419v^4 + 18036v^6 + 9419v^8 + 1478v^{10} + 53v^{12}), \\ P_{1,F_4}^{(2)}(v) &= 1484 + 36252v^2 + 241608v^4 + 663716v^6 + 909400v^8 + \dots + 1484v^{16}. \end{aligned} \quad (4.54)$$

The ellipsis in $P_{1,F_4}^{(2)}(v)$ is completed by making the expression palindromic. Here the leading order expression $g_{1,F_4}^{(0)}$ agrees with the Hilbert series of the reduced moduli space of one F_4 -instanton in [41], which is not surprising since the one-string formula (3.33) reduces to the one-instanton partition function (3.35) in the $Q_\tau \rightarrow 0$ limit. Furthermore, higher order expressions agree with [37].

The Q_τ expansion of the two-string elliptic genus reads

$$\mathbb{E}_{h_{F_4}^{(2)}}(v, x, Q_\tau, m_i) = v^{17} Q_\tau^{-17/6} \sum_{n=0}^{\infty} g_{2,F_4}^{(n)}(v, x, Q_{m_i}) Q_\tau^n, \quad (4.55)$$

where $g_{2,F_4}^{(n)}(v, x, Q_{m_i})$ are rational functions. Turning off flavor fugacities and $SU(2)_x$, we find

$$g_{2,F_4}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{34}(1+v)^{22}(1+v+v^2)^{17}} \times P_{2,F_4}^{(n)}(v), \quad (4.56)$$

where

$$\begin{aligned} P_{2,F_4}^{(0)}(v) &= 1 + 5v + 48v^2 + 287v^3 + 1560v^4 + 7503v^5 + 32316v^6 + 125355v^7 + 444325v^8 \\ &+ 1443572v^9 + 4322993v^{10} + 11989241v^{11} + 30913094v^{12} + 74321701v^{13} + 167106519v^{14} \\ &+ 352245510v^{15} + 697557618v^{16} + 1300152932v^{17} + 2284606168v^{18} + 3790004228v^{19} \\ &+ 5943020899v^{20} + 8818128233v^{21} + 12392104012v^{22} + 16505926853v^{23} \\ &+ 20851379873v^{24} + 24994963144v^{25} + 28442119825v^{26} + 30731161887v^{27} \\ &+ 31533797982v^{28} + 30731161887v^{29} + \dots + v^{56}. \end{aligned} \quad (4.57)$$

$$\begin{aligned}
 P_{2,F_4}^{(1)}(v) = & (1+v^2)(56+386v+3217v^2+20295v^3+110327v^4+529286v^5+2266151v^6 \\
 & +8718327v^7+30479449v^8+97433532v^9+286304088v^{10}+777049966v^{11} \\
 & +1956035588v^{12}+4581942186v^{13}+10017235514v^{14}+20492637094v^{15} \\
 & +39315499928v^{16}+70871529676v^{17}+120240591034v^{18}+192278945658v^{19} \\
 & +290168035137v^{20}+413676858801v^{21}+557641624668v^{22}+711294838217v^{23} \\
 & +859008747683v^{24}+982638991174v^{25}+1065069893896v^{26}+1094033908456v^{27} \\
 & +1065069893896v^{28}+\dots+v^{56}). \tag{4.58}
 \end{aligned}$$

and the ellipses are again completed by making the expressions palindromic. Here the leading order expression $g_{2,F_4}^{(0)}$ agrees with the Hilbert series of the reduced moduli space of two F_4 -instanton in [42]. Some polynomials $P_{2,F_4}^{(n)}(v)$ of higher order n can be found in the appendix F.


We also use the expressions of $\mathbb{E}_1, \mathbb{E}_2$ to extract the BPS invariants N_{j_l, j_r}^β . For this purpose, we need to use instead the Kähler moduli t_b, t_I ($I = 0, 1, \dots, 4$) associated to the Mori cone generators. The results are tabulated in appendix G. They display the proper checkerboard pattern, and reproduce the known genus 0 Gopakumar-Vafa invariants [24]. At base degree one, we also notice a pattern that the only non-vanishing BPS invariants for the curve classes $\beta = (0, k, 0, 0, 0, 1), (0, 0, k, 0, 0, 1)$ are

$$N_{0,k}^{(0,k,0,0,0,1)} = N_{0,k}^{(0,0,k,0,0,1)} = 1, \quad k = 0, 1, \dots, \tag{4.59}$$

which in fact can be proved from the one-string formula (3.33) with the help of (D.7), (D.8) and (D.9).

4.4 $G = E_6$

The divisors and curves of the non-compact $\mathfrak{n} = 6$ geometry is explained in [53]. There are seven compact divisors, which are Hirzebruch surfaces \mathbb{F}_{n_i} of various degrees n_i . We denote them by D_I ($I = 0, 1, \dots, 6$). They intersect with each other like the affine dynkin diagram of \mathfrak{e}_6



where each node corresponds to a Hirzebruch surface and two nodes are connected if the corresponding Hirzebruch surfaces intersect (see figure 5 in [53]). In the diagram above we also give the ordering of the nodes I and the associated marks a_I with the notation $(I : a_I)$ following [61]. The \mathbb{F}_4 denoted by a dashed circle corresponds to the affine node and it

intersects with the base at the \mathbb{P}^1 with normal bundle $\mathcal{O}(-6) \oplus \mathcal{O}(4) \rightarrow \mathbb{P}^1$. There are eight linearly independent curves, which we choose for the moment to be the \mathbb{P}^1 fibers Σ_I of the divisors D_I and the (-6) curve in the base denoted by Σ_B . Denoting their complexified Kähler moduli by t_I and t_B , the linear combination

$$\sum_{I=0}^6 a_I t_I = \tau \tag{4.60}$$

with a_I the marks of \mathfrak{e}_6 , is the volume of the elliptic fiber. We also identify the Mori cone generators. They include the \mathbb{P}^1 fibers Σ_I ($I = 0, 1, \dots, 6$), as well as the \mathbb{P}^1 base of the \mathbb{F}_0 surface in the center. We denote the last curve by Σ_b , and it is related to Σ_B by

$$[\Sigma_B] = [\Sigma_b] + 4[\Sigma_0] + 2[\Sigma_1]. \tag{4.61}$$

This implies the following relation of their Kähler moduli

$$t_B = t_b + 4t_0 + 2t_1. \tag{4.62}$$

The C -matrix of intersection between Σ_I, Σ_b and D_I is

$$C = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 \end{pmatrix}. \tag{4.63}$$

The semiclassical components of the partition function can be computed using the prescription in section 2.2 with the normalisation scheme in appendix B. We obtain

$$\begin{aligned} -F_{(0,0)}^{\text{cls}} &= \left(\frac{t_0}{12} + \frac{t_1}{6} + \frac{t_2}{12} + \frac{t_3}{6} + \frac{t_4}{4} + \frac{t_5}{6} + \frac{t_6}{12} \right) t_b^2 \\ &\quad \left(\frac{t_0^2}{3} + \frac{t_1^2}{3} + \frac{t_2^2}{3} + \frac{t_3^2}{3} + \frac{t_5^2}{3} + \frac{t_6^2}{3} + \frac{t_0 t_1}{3} + \frac{t_2 t_3}{3} + \frac{t_5 t_6}{3} \right) t_b \\ &\quad + \frac{4t_0^3}{9} + \frac{2t_1^3}{9} + \frac{4t_2^3}{9} + \frac{2t_3^3}{9} + \frac{2t_5^3}{9} + \frac{4t_6^3}{9} + \frac{2t_0^2 t_1}{3} + \frac{t_0 t_1^2}{3} + \frac{t_2 t_3^2}{3} + \frac{2t_2^2 t_3}{3} + \frac{2t_5 t_6^2}{3} + \frac{t_5^2 t_6}{3}, \end{aligned} \tag{4.64}$$

which is consistent with the universal formula (2.19). Using the relations (2.4), (4.62), (4.60), we can express $F^{(0,0)}(t)$ in terms of the Kähler moduli $t_{\text{ell}}, \tau, m_i$ ($i = 1, \dots, 6$) and find

$$-F^{(0,0)} = \frac{1}{12} t_{\text{ell}}^2 \tau + \frac{1}{2} t_{\text{ell}}(m, m) - \tau(m, m) + \dots \tag{4.65}$$

up to τ^3 and terms cubic in m_i , where

$$m = \sum_{i=1}^6 m_i \omega_i^\vee. \tag{4.66}$$

	r		fundamental weights
unity	$(0, 0, 0, 0, 0, 0, 0, 0)$	$(0, 0, 0, 0, 0, 0, 0, 2)$	ω_1, ω_4
	$(0, 0, 0, 0, 0, 0, 0, 4)$	$(0, 0, 0, 0, 0, 0, 0, 6)$	
	$(0, 0, 0, 0, 0, 0, 0, 8)$	$(0, 0, 0, 0, 0, 0, 0, 10)$	
vanishing	$(-2, 0, 2, 0, 0, 0, 0, 0)$	$(-2, 0, 2, 0, 0, 0, 0, 2)$	ω_2, ω_5
	$(-2, 0, 2, 0, 0, 0, 0, 4)$	$(-2, 0, 2, 0, 0, 0, 0, 6)$	
	$(-2, 0, 2, 0, 0, 0, 0, 8)$	$(-2, 0, 2, 0, 0, 0, 0, 10)$	
vanishing	$(-2, 0, 0, 0, 0, 0, 2, 0)$	$(-2, 0, 0, 0, 0, 0, 2, 2)$	ω_3, ω_6
	$(-2, 0, 0, 0, 0, 0, 2, 4)$	$(-2, 0, 0, 0, 0, 0, 2, 6)$	
	$(-2, 0, 0, 0, 0, 0, 2, 8)$	$(-2, 0, 0, 0, 0, 0, 2, 10)$	

Table 9. The r -fields of the $\mathfrak{n} = 6$ model and the fundamental weights of \mathfrak{e}_6 which induce the same embedding $\phi : Q^\vee \hookrightarrow P$. They can be divided into three groups; inside each group r -fields or fundamental weights induce the same embedding.

It is in agreement with the universal expression (2.19), and thus the analysis in section 2.3 goes through. We also find

$$F_{(1,0)}^{\text{cls}} = \frac{9}{2}t_0 + 5t_1 + \frac{9}{2}t_2 + 5t_3 + \frac{3}{2}t_4 + 5t_5 + \frac{9}{2}t_6 + 8t_b. \tag{4.67}$$

Imposing the admissibility condition (2.22) and the BPS checkerboard pattern condition (2.25), which specialises to

$$r \equiv (0, 0, 0, 0, 0, 0, 0, 0) \pmod{2}, \tag{4.68}$$

there are in total 18 inequivalent r -fields, and we list their representatives in table 9. We classify them according to the embeddings $\phi_\lambda : Q^\vee \hookrightarrow P$ induced by the reduced r -field λ defined in (2.28), (2.33). We also list in the table the fundamental weights which induce the same embedding.

We use (3.33) and (3.36) to compute the one-string and two-string elliptic genera. The results are again presented in terms of the reduced elliptic genera defined in (4.1).

The reduced one-string elliptic genus does not depend on $SU(2)_x$. The expansion in Q_τ reads

$$\mathbb{E}_{h_{E_6}^{(1)}}(v, Q_\tau, m_i) = v^{11} Q_\tau^{-11/6} \sum_{n=0}^{\infty} g_{1,E_6}^{(n)}(v, Q_{m_i}) Q_\tau^n, \tag{4.69}$$

where $g_{1,E_6}^{(n)}(v, Q_{m_i})$ are rational functions. Turning off all flavor fugacities

$$g_{1,E_6}^{(n)}(v, Q_{m_i} = 1) = \frac{1}{(1 - v^2)^{22}} \times P_{1,E_6}^{(n)}(v), \tag{4.70}$$

where the first few orders are

$$\begin{aligned}
 P_{1,E_6}^{(0)}(v) &= 1 + 56v^2 + 945v^4 + 6776v^6 + 23815v^8 + 43989v^{10} + \dots + v^{22}, \\
 P_{1,E_6}^{(1)}(v) &= 79 + 3774v^2 + 54206v^4 + 337457v^6 + 1067286v^8 + 1862806v^{10} + \dots + 79v^{22}, \\
 P_{1,E_6}^{(2)}(v) &= 3239 + 130034v^2 + 1603334v^4 + 8798601v^6 + 25393522v^8 + 42223058v^{10} + \dots + 3239v^{22}.
 \end{aligned} \tag{4.71}$$

The ellipses are completed by making the expression palindromic. Here $g_{1,E_6}^{(0)}$ agrees with the Hilbert series of reduced one E_6 -instanton moduli space [41], while higher order contributions agree with [37].

The Q_τ expansion of the two-string elliptic genus reads

$$\mathbb{E}_{h_{E_6}^{(2)}}(v, x, Q_\tau, m_i) = v^{23} Q_\tau^{-23/6} \sum_{n=0}^{\infty} g_{2,E_6}^{(n)}(v, x, Q_{m_i}) Q_\tau^n. \tag{4.72}$$

Turning off flavor fugacities and $SU(2)_x$, we obtain

$$g_{2,E_6}^{(n)}(v, x, Q_{m_i} = 1) = \frac{1}{(1-v)^{46}(1+v)^{32}(1+v+v^2)^{23}} \times P_{2,E_6}^{(n)}(v), \tag{4.73}$$

where the first few orders are

$$\begin{aligned}
 P_{2,E_6}^{(0)}(v) &= 1 + 9v + 94v^2 + 739v^3 + 5121v^4 + 31432v^5 + 173895v^6 + 874485v^7 + 4036298v^8 \\
 &+ 17200367v^9 + 68039474v^{10} + 250943933v^{11} + 866242068v^{12} + 2807705547v^{13} \\
 &+ 8569454706v^{14} + 24690503239v^{15} + 67304396959v^{16} + 173919980352v^{17} \\
 &+ 426790882149v^{18} + 996158535441v^{19} + 2214670938701v^{20} + 4695878015170v^{21} \\
 &+ 9507297417908v^{22} + 18398716114730v^{23} + 34066083855696v^{24} + 60399840583490v^{25} \\
 &+ 102628223553496v^{26} + 167232472484542v^{27} + 261500117384417v^{28} + 392614934492341v^{29} \\
 &+ 566271723784347v^{30} + 784947220008032v^{31} + 1046126546231772v^{32} + 1340924322289616v^{33} \\
 &+ 1653587141756229v^{34} + 1962268356880815v^{35} + 2241216639463322v^{36} + 2464163123099051v^{37} \\
 &+ 2608327634962043v^{38} + 2658213934310966v^{39} + \dots + v^{78}.
 \end{aligned} \tag{4.74}$$

Note that $g_{2,E_6}^{(0)}$ agrees with the Hilbert series of reduced two E_6 -instanton moduli space [42].


We also use the expressions of $\mathbb{E}_1, \mathbb{E}_2$ to extract the BPS invariants N_{j_l, j_r}^β . For this purpose, we need to use the Kähler moduli t_I, t_b ($I = 0, 1, \dots, 6$) associated to the Mori cone generators. The results are tabulated in appendix G. They display the proper checkerboard pattern, and reproduce the known genus 0 Gopakumar-Vafa invariants [24]. At base degree one, we notice the interesting pattern that the only non-vanishing BPS invariants for the curve classes $\beta = (0, k, 0, 0, 0, 0, 1)$, $(0, 0, 0, k, 0, 0, 1)$, $(0, 0, 0, 0, 0, k, 0, 1)$, $(0, 0, 0, 0, k, 0, 0, 1)$ are

$$N_{0, k-1/2}^{(0, k, 0, 0, 0, 0, 1)} = N_{0, k-1/2}^{(0, 0, 0, k, 0, 0, 1)} = N_{0, k-1/2}^{(0, 0, 0, 0, 0, k, 0, 1)} = N_{0, k+1/2}^{(0, 0, 0, 0, k, 0, 0, 1)} = 1, \quad k = 1, 2, \dots, \tag{4.75}$$

which in fact can be proved from the one-string formula (3.33) with the help of (D.7), (D.8) and (D.9).

4.5 $G = E_7$

The divisors and curves of the non-compact $\mathfrak{n} = 8$ geometry is explained in [53]. There are eight compact divisors, which are Hirzebruch surfaces \mathbb{F}_{n_i} of various degrees n_i . We denote them by D_I ($I = 0, 1, \dots, 7$). They intersect with each other like the affine dynkin diagram of \mathfrak{e}_7



In the diagram above we also give the ordering of the nodes I and the associated marks a_I with the notation $(I : a_I)$. The \mathbb{F}_6 denoted by a dashed circle corresponds to the affine node and it intersects with the base at the \mathbb{P}^1 with normal bundle $\mathcal{O}(-8) \oplus \mathcal{O}(6) \rightarrow \mathbb{P}^1$. There are nine linearly independent curves, which we choose for the moment to be the \mathbb{P}^1 fibers Σ_I of the divisors D_I and the (-8) curve in the base denoted by Σ_B . Denoting their complexified Kähler moduli by t_I and t_B , the linear combination

$$\sum_{I=0}^7 a_I t_I = \tau \tag{4.76}$$

with a_I the marks of \mathfrak{e}_7 , is the volume of the elliptic fiber. We identify the Mori cone generators. They include the \mathbb{P}^1 fibers Σ_I ($I = 0, 1, \dots, 7$), as well as the \mathbb{P}^1 base of the \mathbb{F}_0 surface in the middle. We denote the last curve by Σ_b , which is related to Σ_B by

$$[\Sigma_B] = [\Sigma_b] + 6[\Sigma_0] + 4[\Sigma_1] + 2[\Sigma_2]. \tag{4.77}$$

Their Kähler moduli are consequently related by

$$t_B = t_b + 6t_0 + 4t_1 + 2t_2. \tag{4.78}$$

The C -matrix of intersection between Σ_I, Σ_b and D_I is

$$C = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{4.79}$$

The semiclassical components of the partition function can be computed using the prescription in section 2.2 with the normalisation scheme in appendix B. We obtain

$$\begin{aligned}
 -F_{(0,0)}^{\text{cls}} = & \left(\frac{t_0}{16} + \frac{t_1}{8} + \frac{3t_2}{16} + \frac{t_3}{4} + \frac{3t_4}{16} + \frac{t_5}{8} + \frac{t_6}{16} + \frac{t_7}{8} \right) t_b^2 + \left(\frac{3t_0^2}{8} + \frac{t_1^2}{2} + \frac{3t_2^2}{8} + \frac{3t_4^2}{8} + \frac{t_5^2}{2} + \frac{3t_6^2}{8} \right. \\
 & + \left. \frac{t_7^2}{4} + \frac{t_0 t_1}{2} + \frac{t_0 t_2}{4} + \frac{t_1 t_2}{2} + \frac{t_4 t_5}{2} + \frac{t_4 t_6}{4} + \frac{t_5 t_6}{2} \right) t_b + \frac{3t_0^3}{4} + \frac{2t_1^3}{3} + \frac{t_2^3}{4} + \frac{t_4^3}{4} + \frac{2t_5^3}{3} \\
 & + \frac{3t_6^3}{4} + \frac{t_7^3}{6} + \frac{3t_0^2 t_1}{2} + \frac{3t_0^2 t_2}{4} + t_0 t_1^2 + \frac{t_0 t_2^2}{4} + t_0 t_1 t_2 + \frac{t_1 t_2^2}{2} + t_4 t_5^2 + \frac{3t_4 t_6^2}{4} + \frac{3t_5 t_6^2}{2} \\
 & + t_1^2 t_2 + \frac{t_4^2 t_5}{2} + \frac{t_4^2 t_6}{4} + t_5^2 t_6 + t_4 t_5 t_6,
 \end{aligned} \tag{4.80}$$

which is consistent with the universal formula (2.19). Using the relations (2.4), (4.78) and (4.76), we can express $F^{(0,0)}(t)$ in terms of the Kähler moduli $t_{\text{ell}}, \tau, m_i$ ($i = 1, \dots, 7$) and find

$$-F^{(0,0)} = \frac{1}{16} t_{\text{ell}}^2 \tau + \frac{1}{2} t_{\text{ell}}(m, m) - \frac{3}{2} \tau(m, m) + \dots \tag{4.81}$$

up to τ^3 and terms cubic in m_i , where

$$m = \sum_{i=1}^7 m_i \omega_i^\vee. \tag{4.82}$$

It is in agreement with the universal expression (2.19), and therefore the analysis in section 2.3 goes through, which then leads to the elliptic blowup equations (3.4). We also find

$$F_{(1,0)}^{\text{cls}} = \frac{51}{8} t_0 + \frac{35}{4} t_1 + \frac{57}{8} t_2 + \frac{3}{2} t_3 + \frac{57}{8} t_4 + \frac{35}{4} t_5 + \frac{51}{8} t_6 + \frac{11}{4} t_b. \tag{4.83}$$

Imposing the admissibility condition (2.22) and the BPS checkerboard pattern condition (2.25), which specialises to

$$r \equiv (0, 0, 0, 0, 0, 0, 0, 0, 0) \pmod{2}, \tag{4.84}$$

there are in total 16 inequivalent r -fields, and we list their representatives in table 10. We classify them according to the embeddings $\phi_\lambda : Q^\vee \hookrightarrow P$ induced by the reduced r -field λ defined in (2.28), (2.33). We also list in the table the fundamental weights which induce the same embedding.

We use (3.33) and (3.36) to compute the one-string and two-string elliptic genera and convert them to reduced versions. The one string elliptic genus when expanded in Q_τ reads

$$\mathbb{E}_{h_{E_7}}^{(1)}(v, Q_\tau, m_i) = v^{17} Q_\tau^{-17/6} \sum_{n=0}^{\infty} g_{1,E_7}^{(n)}(v, Q_{m_i}) Q_\tau^n, \tag{4.85}$$

where $g_{1,E_7}^{(n)}(v, Q_{m_i})$ are rational functions. When all flavor fugacities are turned off

$$g_{1,E_7}^{(n)}(v, Q_{m_i} = 1) = \frac{1}{(1-v^2)^{34}} \times P_{n,E_7}^{(1)}(v), \tag{4.86}$$

	r	fundamental weights
unity	$(0, 0, 0, 0, 0, 0, 0, 0, 0)$	$(0, 0, 0, 0, 0, 0, 0, 0, 2)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 4)$	$(0, 0, 0, 0, 0, 0, 0, 0, 6)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 8)$	$(0, 0, 0, 0, 0, 0, 0, 0, 10)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 12)$	$(0, 0, 0, 0, 0, 0, 0, 0, 14)$
vanishing	$(-2, 0, 0, 0, 0, 0, 2, 0, 0)$	$(-2, 0, 0, 0, 0, 0, 2, 0, 2)$
	$(-2, 0, 0, 0, 0, 0, 2, 0, 4)$	$(-2, 0, 0, 0, 0, 0, 2, 0, 6)$
	$(-2, 0, 0, 0, 0, 0, 2, 0, 8)$	$(-2, 0, 0, 0, 0, 0, 2, 0, 10)$
	$(-2, 0, 0, 0, 0, 0, 2, 0, 12)$	$(-2, 0, 0, 0, 0, 0, 2, 0, 14)$

Table 10. The r -fields of the $\mathfrak{n} = 8$ model and the fundamental weights of \mathfrak{e}_7 which induce the same embedding $\phi : Q^\vee \hookrightarrow P$. They can be divided into two groups; inside each group r -fields or fundamental weights induce the same embedding.

where the leading order contributions are

$$\begin{aligned}
 P_{1,E_7}^{(0)}(v) &= (1+v^2)(1+98v^2+3312v^4+53305v^6+468612v^8+2421286v^{10}+7664780v^{12} \\
 &\quad +15203076v^{14}+19086400v^{16}+15203076v^{18}+\dots+v^{32}), \\
 P_{1,E_7}^{(1)}(v) &= (1+v^2)(134+11593v^2+345521v^4+4931707v^6+38850151v^8+182614170v^{10} \\
 &\quad +536726278v^{12}+1014596958v^{14}+1252490096v^{16}+1014596958v^{18}+\dots+v^{32}), \\
 P_{1,E_7}^{(2)}(v) &= (1+v^2)(9179+693316v^2+18210733v^4+231525774v^6+1645739978v^8 \\
 &\quad +7093827388v^{10}+19507715662v^{12}+35350906224v^{14}+43009574252v^{16} \\
 &\quad +35350906224v^{18}\dots+v^{32}). \tag{4.87}
 \end{aligned}$$

where the ellipses are completed by palindrome. Here $g_{1,E_7}^{(0)}$ agrees with the Hilbert series of reduced one E_7 -instanton moduli space in [41], while higher order contributions agree with [37].

The Q_τ expansion of the two-string elliptic genus reads

$$\mathbb{E}_{h_{E_7}^{(2)}}(v, x, Q_\tau, m_i) = v^{35} Q_\tau^{-35/6} \sum_{n=0}^{\infty} g_{2,E_7}^{(n)}(v, x, Q_{m_i}) Q_\tau^n. \tag{4.88}$$

Turning off $SU(2)_x$ and flavours, we have

$$g_{2,E_7}^{(n)}(v, x, Q_{m_i} = 1) = \frac{1}{(1-v)^{70}(1+v)^{52}(1+v+v^2)^{35}} \times P_{n,E_7}^{(2)}(v). \tag{4.89}$$

We have computed $P_{n,E_7}^{(2)}(v)$ for $n = 0, 1$ which we put in the appendix F. Indeed, our $g_{2,E_7}^{(0)}$ agrees with the Hilbert series of reduced two E_7 -instanton moduli space in [42].

We also use the expressions of $\mathbb{E}_1, \mathbb{E}_2$ to extract the BPS invariants N_{j_l, j_r}^β . For this purpose, we need to use the Kähler moduli t_I, t_b ($I = 0, 1, \dots, 7$) associated to the Mori cone generators. The results are tabulated in appendix G. They display the proper checkerboard


pattern, and reproduce the known genus zero Gopakumar-Vafa invariants [24]. At base degree one, we notice the interesting pattern that the only non-vanishing BPS invariants for the curve classes $\beta = (0, 0, k, 0, 0, 0, 0, 0, 1)$, $(0, 0, 0, 0, k, 0, 0, 0, 1)$, $(0, 0, 0, 0, 0, 0, k, 1)$, $(0, 0, 0, k, 0, 0, 0, 0, 1)$ are

$$N_{0,k-1/2}^{(0,0,k,0,0,0,0,0,1)} = N_{0,k-1/2}^{(0,0,0,k,0,0,0,1)} = N_{0,k-1/2}^{(0,0,0,0,0,0,k,1)} = N_{0,k+1/2}^{(0,0,0,k,0,0,0,1)} = 1, \quad k = 1, 2, \dots, \quad (4.90)$$

which in fact can be proved from the one-string formula (3.33) with the help of (D.7), (D.8) and (D.9).

4.6 $G = E_8$

The divisors and curves of the non-compact $\mathfrak{n} = 12$ geometry is explained in [53]. There are nine compact divisors, which are Hirzebruch surfaces of various degrees. We denote them by D_I ($I = 0, 1, \dots, 8$). They intersect with each other like the affine dynkin diagram of \mathfrak{e}_8



In the diagram above we also give the ordering of the nodes I and the associated marks a_I with the notation $(I : a_I)$. The \mathbb{F}_{10} denoted by a dashed circle corresponds to the affine node and it intersects with the base at the \mathbb{P}^1 with normal bundle $\mathcal{O}(-12) \oplus \mathcal{O}(10) \rightarrow \mathbb{P}^1$. There are ten linearly independent curves, which we choose for the moment to be the \mathbb{P}^1 fibers Σ_I of the divisors D_I and the (-12) curve in the base denoted by Σ_B . Let t_I and t_B be their complexified Kähler moduli. The linear combination

$$\sum_{I=0}^8 a_I t_I = \tau \quad (4.91)$$

with a_I marks of \mathfrak{e}_8 , is the volume of the elliptic fiber. We identify the Mori cone generators. They include \mathbb{P}^1 fibers Σ_I ($I = 0, 1, \dots, 8$) as well as the \mathbb{P}^1 base of the \mathbb{F}_0 surface in the center. We denote the latter by Σ_b , which is related to Σ_B by

$$[\Sigma_B] = [\Sigma_b] + 10[\Sigma_0] + 8[\Sigma_1] + 6[\Sigma_2] + 4[\Sigma_3] + 2[\Sigma_4], \quad (4.92)$$

which implies

$$t_B = t_b + 10t_0 + 8t_1 + 6t_2 + 4t_3 + 2t_4. \quad (4.93)$$

The C -matrix of intersection between Σ_I, Σ_b and D_I is

$$C = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 \end{pmatrix}. \quad (4.94)$$

The semiclassical components of the partition function can be computed using the prescription in section 2.2 with the normalisation scheme in appendix B. We obtain

$$\begin{aligned} -F_{(0,0)}^{\text{cls}} = & \left(\frac{t_0}{24} + \frac{t_1}{12} + \frac{t_2}{8} + \frac{t_3}{6} + \frac{5t_4}{24} + \frac{t_5}{4} + \frac{t_6}{6} + \frac{t_7}{12} + \frac{t_8}{8} \right) t_b^2 \\ & + \left(\frac{5t_0^2}{12} + \frac{2t_1^2}{3} + \frac{3t_2^2}{4} + \frac{2t_3^2}{3} + \frac{5t_4^2}{12} + \frac{t_6^2}{3} + \frac{t_7^2}{3} + \frac{t_8^2}{4} + \frac{2t_0t_1}{3} + \frac{t_0t_2}{2} + \frac{t_0t_3}{3} + \frac{t_0t_4}{6} \right. \\ & + t_1t_2 + \frac{2t_1t_3}{3} + t_2t_3 + \frac{t_1t_4}{3} + \frac{t_2t_4}{2} + \frac{2t_3t_4}{3} + \left. \frac{t_6t_7}{3} \right) t_b \\ & + \frac{25t_0^3}{18} + \frac{16t_1^3}{9} + \frac{3t_2^3}{2} + \frac{8t_3^3}{9} + \frac{5t_4^3}{18} + \frac{2t_6^3}{9} + \frac{4t_7^3}{9} + \frac{t_8^3}{6} + \frac{10t_0^2t_1}{3} + \frac{5t_0^2t_2}{2} + \frac{5t_0^2t_3}{3} + \frac{5t_0^2t_4}{6} \\ & + \frac{8t_0t_1^2}{3} + \frac{3t_0t_2^2}{2} + \frac{2t_0t_3^2}{3} + \frac{t_0t_4^2}{6} + 4t_0t_1t_2 + \frac{8t_0t_1t_3}{3} + 2t_0t_2t_3 + \frac{4}{3}t_0t_1t_4 + t_0t_2t_4 + \frac{2t_0t_3t_4}{3} \\ & + 3t_1t_2^2 + \frac{4t_1t_3^2}{3} + 2t_2t_3^2 + \frac{t_1t_4^2}{3} + \frac{t_2t_4^2}{2} + \frac{2t_3t_4^2}{3} + \frac{2t_6t_7^2}{3} + 4t_1^2t_2 + \frac{8t_1^2t_3}{3} + 3t_2^2t_3 + 4t_1t_2t_3 \\ & + \frac{4t_1^2t_4}{3} + \frac{3t_2^2t_4}{2} + \frac{4t_3^2t_4}{3} + 2t_1t_2t_4 + \frac{4t_1t_3t_4}{3} + 2t_2t_3t_4 + \frac{t_6^2t_7}{3}, \end{aligned} \quad (4.95)$$

which is consistent with (2.19). Using the relations (2.4), (4.93), (4.91), we can express $F^{(0,0)}(\underline{t})$ in terms of the Kähler moduli $t_{\text{ell}}, \tau, m_i$ ($i = 1, \dots, 8$) and find

$$-F^{(0,0)} = \frac{1}{24}t_{\text{ell}}^2\tau + \frac{1}{2}t_{\text{ell}}(m, m) - \frac{5}{2}\tau(m, m) + \dots \quad (4.96)$$

up to τ^3 and terms cubic in m_i , where

$$m = \sum_{i=1}^8 m_i \omega_i^\vee. \quad (4.97)$$

It is in agreement with the universal expression (2.19), and therefore the analysis in section 2.3 goes through leading to the elliptic blowup equations (3.4). We also find

$$F_{(1,0)}^{\text{cls}} = \frac{41}{4}t_0 + \frac{33}{2}t_1 + \frac{75}{4}t_2 + 17t_3 + \frac{45}{4}t_4 + \frac{3}{2}t_5 + 5t_6 + \frac{9}{2}t_7 + \frac{11}{4}t_8 + 10t_b. \quad (4.98)$$

	r	fundamental weights
unity	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 0)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 2)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 4)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 6)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 8)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 10)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 12)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 14)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 16)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 18)$
	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 20)$	$(0, 0, 0, 0, 0, 0, 0, 0, 0, 22)$

Table 11. The r -fields of the $\mathfrak{n} = 12$ model and the fundamental weights of \mathfrak{e}_8 which induce the same embedding $\phi : Q^\vee \hookrightarrow P$. All the r -fields and all the fundamental weights induce the same embedding as $Q^\vee = P$ for \mathfrak{e}_8 .

Imposing the admissibility condition (2.22) and the BPS checkerboard pattern condition (2.25), which specialises to

$$r \equiv (0, 0, 0, 0, 0, 0, 0, 0, 0, 0) \pmod{2}, \tag{4.99}$$

there are in total 12 inequivalent r -fields, and we list their representatives in table 11. All of them have the same reduced λ which induces the same embedding $\phi_\lambda : Q^\vee \hookrightarrow P = Q^\vee$. In this special case, there is no vanishing blowup equations.

We use (3.33) and (3.36) to compute the one-string and two-string elliptic genera and convert them to reduced versions. The one-string reduced elliptic genus in Q_τ expansion reads

$$\mathbb{E}_{h_{E_8}^{(1)}}(v, Q_\tau, m_i) = v^{29} Q_\tau^{-29/6} \sum_{n=0}^{\infty} g_{1,E_8}^{(n)}(v, Q_{m_i}) Q_\tau^n, \tag{4.100}$$

where $g_{1,E_8}^{(n)}$ are rational functions. Turning off flavor fugacities

$$g_{1,E_8}^{(n)}(v, Q_{m_i} = 1) = \frac{1}{(1-v^2)^{58}} \times P_{n,E_8}^{(1)}(v), \tag{4.101}$$

where the leading orders are

$$\begin{aligned} P_{0,E_8}^{(1)}(v) = & (1+v^2)(1+189v^2+14080v^4+562133v^6+13722599v^8+220731150v^{10} \\ & +2454952400v^{12}+19517762786v^{14}+113608689871v^{16}+492718282457v^{18} \\ & +1612836871168v^{20}+4022154098447v^{22}+7692605013883v^{24}+11332578013712v^{26} \\ & +12891341012848v^{28}+11332578013712v^{30}+\dots+v^{56}), \end{aligned} \tag{4.102}$$

$$\begin{aligned} P_{1,E_8}^{(1)}(v) = & 249+43435v^2+2998484v^4+111587988v^6+2558096217v^8+38985250263v^{10} \\ & +415090167480v^{12}+3197400818096v^{14}+18281159666407v^{16}+79099752469353v^{18} \\ & +262872507223458v^{20}+678620928038790v^{22}+1372471431431505v^{24} \\ & +2187800775100695v^{26}+2759575276449180v^{28}+2759575276449180v^{30}+\dots+v^{58}, \end{aligned} \tag{4.103}$$

$$\begin{aligned}
P_{2,E_8}^{(1)}(v) = & 31374 + 4996185v^2 + 316301853v^4 + 10844316461v^6 + 230109165319v^8 \\
& + 3262175735364v^{10} + 32482207865920v^{12} + 235331998114532v^{14} \\
& + 1273365718136904v^{16} + 5249113972780491v^{18} + 16738824444898167v^{20} \\
& + 41781447040327605v^{22} + 82360817736515085v^{24} + 129037047832755990v^{26} \\
& + 161349436368883950v^{28} + 161349436368883950v^{30} + \dots + v^{58}.
\end{aligned} \tag{4.104}$$

where the ellipses are completed by palindome. Note $g_{1,E_8}^{(0)}$ indeed agrees with the Hilbert series of reduced one E_8 -instanton moduli space in [41]. Higher order contributions agree with [37].

The two-string reduced elliptic genus in Q_τ expansion reads

$$\mathbb{E}_{h_{E_8}^{(2)}}(v, x, Q_\tau, m_i) = v^{59} Q_\tau^{-59/6} \sum_{n=0}^{\infty} g_{2,E_8}^{(n)}(v, x, Q_{m_i}) Q_\tau^n, \tag{4.105}$$

where $g_{2,E_8}^{(n)}(v, x, Q_{m_i})$ are rational functions. Turning off flavor fugacities and $SU(2)_x$

$$g_{2,E_8}^{(n)}(v, x, Q_{m_i} = 1) = \frac{1}{(1-v)^{118}(1+v)^{92}(1+v+v^2)^{59}} \times P_{2,E_8}^{(n)}(v). \tag{4.106}$$

We have computed $g_{2,E_8}^{(0)}$ which indeed agrees with the Hilbert series of two E_8 -instanton reduced moduli space in [42].

We also use the expressions of $\mathbb{E}_1, \mathbb{E}_2$ to extract the BPS invariants N_{j_l, j_r}^β . For this purpose, we need to use the Kähler moduli t_I, t_b ($I = 0, 1, \dots, 8$) associated to the Mori cone generators. The results are tabulated in appendix G. They display the proper checkerboard pattern, and reproduce the known genus zero Gopakumar-Vafa invariants [24]. At base degree one, we notice a pattern that the only non-vanishing BPS invariants for the curve classes $\beta = (0, 0, 0, 0, k, 0, 0, 0, 0, 1)$, $(0, 0, 0, 0, 0, 0, k, 0, 0, 1)$, $(0, 0, 0, 0, 0, 0, 0, 0, k, 1)$, $(0, 0, 0, 0, 0, k, 0, 0, 0, 1)$ are

$$N_{0,k-1/2}^{(0,0,0,0,k,0,0,0,1)} = N_{0,k-1/2}^{(0,0,0,0,0,k,0,0,1)} = N_{0,k-1/2}^{(0,0,0,0,0,0,0,k,1)} = N_{0,k+1/2}^{(0,0,0,0,k,0,0,0,1)} = 1, \quad k = 1, 2, \dots, \tag{4.107}$$

which in fact can be proved from the one-string formula (3.33) with the help of (D.7), (D.8) and (D.9).

5 On the relation with 4d SCFTs of type $H_G^{(k)}$

The purpose of this section is to connect the k -string elliptic genera $\mathbb{E}_{h_G^{(k)}}$ for the minimal $\mathcal{N} = (1, 0)$ 6d SCFTs with $G = A_2, D_4, F_4, E_{6,7,8}$ discussed above to the superconformal indices of the $\mathcal{N} = 2$ 4d SCFTs of rank k denoted by $H_G^{(k)}$. The simplest series of $\mathcal{N} = 2$ SCFTs namely $H_G^{(1)}$ can be obtained by geometric engineering on non-compact del Pezzo geometries and contains the Minahan-Nemeschansky theories. The main result is an extension of a surprising conjecture by Del Zotto-Lockhart from the rank one case [37] to the higher rank cases. To be precise [37] recognised that the one-string elliptic

genus $\mathbb{E}_{h_G^{(1)}}(Q_\tau, v)$ can be decomposed in terms of a seemingly more fundamental function $L_G(Q_\tau, v)$, which for special choices of Q_τ and v specialises to the Hall-Littlewood index or the Schur index of the $H_G^{(1)}$ theories. With the two string elliptic genera computed in our previous sections, we are able to study this conjectural relation at rank two and in principle at arbitrary rank, and find indeed that similar striking relations exist.

We first review some basic properties of 4d rank k type $H_G^{(k)}$ — and $\tilde{H}_G^{(k)}$ theories, including their class \mathcal{S} theory construction, and then review the superconformal indices of 4d SCFTs in various physically motivated limits as well as the methods to compute them. Next we state the conjectural relation at rank one from [37], and explain in some detail the new relations at rank two for all G . We also extend the analysis to some rank three cases. For all choices of rank and G we analysed, the surprising relation between elliptic genera and superconformal indices exists. We define an intermediate function at rank k called $L_G^{(k)}$.³³ This function is on the one hand the ingredient of k -string elliptic genus, on the other hand gives the Hall-Littlewood index and Schur index of $H_G^{(k)}$ theories at special choices of parameters. This general structure allows us to calculate the latter indices efficiently from the $\mathbb{E}_{h_G^{(k)}}$ that are determined from the elliptic blowup equations.

5.1 Rank k H_G theories

The 4d $\mathcal{N} = 2$ SCFTs $H_G^{(k)}$ are well known to exist for $G = \emptyset, A_1, A_2, D_4, E_{6,7,8}$ and $k = 1, 2, 3, \dots$ [62–66].³⁴ In type IIB superstring theory, they are realized as the world-volume theory for k multiple D3-branes probing a stack of exotic seven-branes. Such seven-branes in F-theory are defined as codimension one singularities with Kodaira type $II, III, IV, I_0^*, IV^*, III^*$, and II^* , which give the gauge symmetries G for the low energy 8d SYM theories. The number k is usually called the rank of H_G theories. For example, the rank one H_{\emptyset, A_1, A_2} theories appear as certain limit of $SU(2)$ gauge theory with $N_f = 1, 2, 3$ respectively [62]. The rank one H_{D_4} theory is well known to be the $SU(2)$ gauge theory with $N_f = 4$, while the higher rank cases with $k > 1$ are equivalent to $USp(2k)$ gauge theories with four fundamental hypermultiplets and one antisymmetric hypermultiplet, which are all Lagrangian theories. The rank one $H_{E_{6,7,8}}$ are also known as the Minahan-Nemeschansky theories [65, 66], where the simplest example rank one E_6 theory is in S-duality with $SU(3)$, $N_f = 6$ theory [67].

All $H_G^{(k)}$ theory can be coupled with a free hypermultiplet associated to the center of mass motion of the instantons. We follow [37] and denote these theories as $\tilde{H}_G^{(k)}$. As was observed in [45], for higher rank cases, \tilde{H}_G are sometimes more natural than H_G theories. One major difference between rank one and higher rank H_G theories is the flavour symmetry. Besides the flavour G given by the strings stretched between D3-branes and exotic seven-brane, for $k > 1$ there is one more $SU(2)$ symmetry coming from the transverse space in the seven-brane. By coupling a free hypermultiplet, all $\tilde{H}_G^{(k)}$ theories share flavour symmetry $G \times SU(2)$.

³³The L_G function in [37] becomes $L_G^{(1)}$ here.

³⁴The $G = \emptyset, A_1, A_2$ type theories are also traditionally denoted as $H_{0,1,2}$ theories. Here we follow the notations in [37].

The $H_G^{(k)}$ theories of interest in this paper are $G = A_2, D_4, E_{6,7,8}$ as they are directly related to 6d minimal $(1, 0)$ SCFTs with corresponding gauge group G . To be precise, the RR elliptic genus is identified as the β -twisted $T^2 \times S^2$ partition function of the 4d SCFTs:

$$\mathbb{E}_{h_G^{(k)}} = Z_{(T^2 \times S^2)_\beta}(H_G^{(k)}). \tag{5.1}$$

Adding the “tildes”, one can also obtain the equality with the free hypermultiplet coupled. Here the β -twist was introduced by Kapustin in [68] to preserve half of the supersymmetries on the backgrounds such as $T^2 \times S^2$. See a good description of such twist in for example section 3.2 of [37]. The identification (5.1) makes it sometimes possible to compute the elliptic genus from 4d setting, in which cases the S-duality with a Lagrangian theory is invoked and one can use certain analogy of Spiridonov-Warnaar inverse formula [69] to compute the $T^2 \times S^2$ partition function. This was indeed achieved for one string elliptic genus with $G = D_4, E_{6,7}$ [37, 46–48]. For example, the elliptic genus of one E_7 instanton string was obtained in [48] via $SU(4)$ gauge theory $N_f = 8$ and appropriate Higgsing as

$$\begin{aligned} Z_{(T^2 \times S^2)_\beta}(H_{E_7}^{(1)}) &= 1 + \chi_{133}^{E_7} v^2 + \chi_{7371}^{E_7} v^4 + \chi_{238602}^{E_7} v^6 + \chi_{5248750}^{E_7} v^8 + \dots \\ &+ Q_\tau \left(1 + \chi_{133}^{E_7} + (1 + 2\chi_{133}^{E_7} + \chi_{7331}^{E_7} + \chi_{8645}^{E_7}) v^2 \right. \\ &\quad \left. + (\chi_{133}^{E_7} + 2\chi_{7371}^{E_7} + \chi_{8645}^{E_7} + \chi_{238602}^{E_7} + \chi_{573440}^{E_7}) v^4 + \dots \right) \\ &+ Q_\tau^2 \left(3 + 2\chi_{133}^{E_7} + \chi_{1539}^{E_7} + \chi_{7371}^{E_7} + \dots \right) + \mathcal{O}(Q_\tau^3), \end{aligned} \tag{5.2}$$

which completely agrees with our universal expansion formula (4.5), (4.6) and (4.7).³⁵ We also checked for D_4 and E_6 , where the agreement holds to all known orders.

Another important feature of $H_G^{(k)}$ theories is that they all admit 6d construction. It is well known all rank k $H_{D_4, E_{6,7,8}}$ theories can be realized by compactifying a 6d A_{N-1} $(2,0)$ SCFT on some punctured sphere with regular singularities [70], i.e. they are class \mathcal{S} theories. The regular singularities are classified by embeddings of $SU(2)$ in $SU(N)$, thus can be denoted as Young diagrams. Such punctures with associated Young diagram represent how the $SU(N)$ decomposes and what is the residual flavour symmetry. For example, the rank one $H_{SO(8)}$ theory is obtained by compactifying 6d A_1 $(2,0)$ SCFT on a sphere with four full punctures $\{1^2\}$, i.e. the residual flavour symmetry is $SU(2)$. Thus the resulting 4d theory has gauge symmetry $SU(2)$ and four fundamentals, as was mentioned already above. We summarize the gauge algebras and punctures for the 6d construction of all $H_G^{(k)}$ theories with $G = D_4, E_{6,7,8}$ in table 12. The 6d construction for rank k H_{A_2} theories however involves irregular punctures. For example, they can be realized by compactifying 6d A_{2k-1} theory on a sphere with one regular puncture with Young diagram $\{k^2\}$ and one irregular puncture of form

$$\Phi = \frac{1}{z^3} \text{diag}(1, \dots, 1_{k_{\text{th}}}, -1, \dots, -1_{k_{\text{th}}}) + \dots \tag{5.3}$$

³⁵In the coefficients of Q_τ^2 , one also need to use the Joseph relation $\text{Sym}^2 \mathbf{133} = 1 + \mathbf{133} + \mathbf{7371}$ to obtain the identification.

G	6d (2, 0) A_{N-1}	punctures Λ_i
D_4	A_{2k-1}	four $\{k^2\}$
E_6	A_{3k-1}	three $\{k^3\}$
E_7	A_{4k-1}	$\{(2k)^2\}$ and two $\{k^4\}$
E_8	A_{6k-1}	$\{(3k)^2\}, \{(2k)^3\}$ and $\{k^6\}$

Table 12. 6d construction for rank k H_G theory.

where the coefficients of z^{-2} and z^{-1} have the same type of matrix [71]. In particular, the rank one H_{A_2} theory coincides with (A_1, D_4) Argyres-Douglas theory. See also the 6d construction involving irregular punctures in [72].

Class \mathcal{S} 4d SCFTs are also known to be connected to 2d vertex operator algebra, i.e. chiral algebra [73, 74]. This correspondence relies directly on the class \mathcal{S} construction and can be understood from certain generalized TQFT structure on the punctured Riemann surface. This relation sometimes gives a new approach to compute the indices of 4d SCFT by realizing them as the vacuum character of associated chiral algebra. For example, the chiral algebras associated to rank one H_{D_4} and H_{E_6} theories are identified as $\mathfrak{so}(8)$ affine Lie algebra at level $k_{2d} = -2$ and $\mathfrak{e}(6)$ affine Lie algebra at level $k_{2d} = -3$ in [73]. See some recent works trying to explain VOA/SCFT correspondence [75–79]. Besides, the rank one H_{D_4, E_6, E_7} theories are also connected with the curved $\beta\gamma$ systems on cones over the complex Grassmannian $\text{Gr}(2, 4)$, the complex orthogonal Grassmannian $\text{OG}^+(5, 10)$, and the complex Cayley plane $\mathbb{O}\mathbb{P}^2$ respectively in [80].

5.2 Hall-Littlewood and Schur indices

The superconformal index of 4d $\mathcal{N} = 2$ SCFT is defined as [81, 82]

$$\mathcal{I}(p, q, t) = \text{Tr} (-1)^F \left(\frac{t}{pq} \right)^r p^{j_{12}} q^{j_{34}} t^R \prod_i a_i^{f_i}, \tag{5.4}$$

where $j_{12} = j_2 + j_1$ and $j_{34} = j_2 - j_1$ denote the rotation generators in \mathbb{C}^2 with $j_{1,2}$ representing each $\text{SU}(2)$ Lorentz symmetry, and r and R denote the $\text{U}(1)_r$ and $\text{SU}(2)_R$ generators respectively. Besides, a_i are the fugacities for the flavour generators f_i which sometimes are set to be zero for simplicity. For generic 4d SCFT, the full superconformal indices with (p, q, t) are difficult to compute. For example, among all $H_G^{(k)}$ theories, the full superconformal indices to our knowledge are only computable so far for $H_{\text{SO}(8)}$ with arbitrary rank owing to their Lagrangian nature and H_{E_6, E_7} for rank one owing to the existence of certain $\mathcal{N} = 1$ Lagrangian flow [47, 48].

Certain limits of superconformal index are particularly interesting due to symmetry enhancement. The name of limit comes from the observation that the resulting indices involve corresponding symmetric polynomial known in mathematics literature. Following [44], we list three of them here:

- (Macdonald) $p \rightarrow 0$. Superconformal index when taking the Macdonald limit is computable for all class \mathcal{S} theory with regular punctures. For a genus \mathfrak{g} theory with

s punctures compactified from 6d A_{N-1} (2,0) SCFT, the Macdonald index is given in [44] as

$$\mathcal{I}_{\mathfrak{g},s}^M(\mathbf{a}, q, t) = \prod_{j=2}^N (t^j; q)^{2\mathfrak{g}-2+s} \frac{(t; q)^{(k-1)(1-\mathfrak{g})+s}}{(q; q)^{(k-1)(1-\mathfrak{g})}} \sum_{\lambda} \frac{\prod_{i=1}^s \hat{\mathcal{K}}_{\Lambda_i}(\mathbf{a}_i) P^{\lambda}(\mathbf{a}_i(\Lambda_i)|q, t)}{\left[P^{\lambda}(t^{\frac{k-1}{2}}, t^{\frac{k-3}{2}}, \dots, t^{\frac{1-k}{2}} | q, t) \right]^{2\mathfrak{g}-2+s}}. \quad (5.5)$$

Here $P^{\lambda}(\mathbf{a}_i(\Lambda_i)|q, t)$ are Macdonald polynomials and the summation is over all possible Young diagrams $\lambda = \{\lambda_1, \lambda_2, \dots, \lambda_{N-1}, 0\}$. The Pochhammer symbol $(a; b)$ is defined by

$$(a; b) = \prod_{i=0}^{\infty} (1 - ab^i). \quad (5.6)$$

The $\hat{\mathcal{K}}_{\Lambda_i}$ factors are defined by

$$\hat{\mathcal{K}}_{\Lambda}(\mathbf{a}) = \prod_{i=1}^{row(\Lambda)} \prod_{j,k=1}^{l_i} \text{PE} \left[\frac{a_j^i \bar{a}_k^i}{1 - q} \right]_{\mathbf{a}_i, q}, \quad (5.7)$$

with the coefficients a_k^i associated to the Young diagram as

$$a_j^i = c_j v^{\lambda_j + 1 - i} \quad \text{and} \quad \bar{a}_k^i = c_k^{-1} v^{\lambda_k + 1 - i}, \quad (5.8)$$

with $v^2 = t$. Here these c_j parameterize the residual flavour symmetry and are subject to constraint $\prod_{i=1}^{row(\Lambda)} \prod_j^{l_i} c_j = 1$ to preserve the traceless condition of $SU(N)$. The association of the flavour fugacities for a puncture $a(\Lambda)$ in Macdonald polynomial is defined similarly as $c_j v^{-\lambda_j - 1 + 2i}$. Some good figures to visualize these definitions can be found in [44, 45].

- (Hall-Littlewood) $p, q \rightarrow 0$. By taking limit in (5.5), it is easy to obtain the Hall-Littlewood index for all class \mathcal{S} theories. As only genus zero theories are of concern in this paper, we only write down the formulas with $\mathfrak{g} = 0$. For example, the Hall-Littlewood index of 4d SCFT compactified from 6d A_{N-1} theory is

$$\mathcal{I}^{\text{HL}} = \mathcal{N}_{N,s} \sum_{\lambda} \frac{\prod_{i=1}^s \hat{\mathcal{K}}_{\Lambda_i}(\mathbf{a}_i) \psi^{\lambda}(\mathbf{a}_i(\Lambda_i)|v)}{\left[\psi^{\lambda}(v^{N-1}, v^{N-3}, \dots, v^{1-N}|v) \right]^{s-2}}, \quad (5.9)$$

where

$$\mathcal{N}_{N,s} = (1 - v^2)^{N-1+s} \prod_{j=2}^N (1 - v^{2j})^{s-2}, \quad (5.10)$$

and ψ^{λ} is the Hall-Littlewood polynomials defined as

$$\psi^{\lambda}(x_1, \dots, x_N|v) = \mathcal{N}_{\lambda}(v) \sum_{\sigma \in S_N} x_{\sigma(1)}^{\lambda_1} \dots x_{\sigma(N)}^{\lambda_N} \prod_{i < j} \frac{x_{\sigma(i)} - v^2 x_{\sigma(j)}}{x_{\sigma(i)} - x_{\sigma(j)}}, \quad (5.11)$$

with

$$\mathcal{N}_{\lambda}(v) = \prod_{i=0}^{\infty} \prod_{j=1}^{m(i)} \left(\frac{1 - v^{2j}}{1 - v^2} \right)^{-1/2}, \quad (5.12)$$

where $m(i)$ is the number of rows in the Young diagram $\lambda = (\lambda_1, \dots, \lambda_N)$ of length i . Here we have made the substitution $t = v^2$ for convenience.

It is argued in [44] that for linear quiver theories the HL index is equivalent to the Hilbert series of the Higgs branch. In particular, this is true for all H_G theories. It is well-known the Higgs branch of $H_G^{(k)}$ theories are the reduced moduli space of k G -instantons, which can be understood from the probing picture that the k D3-branes dissolving into the seven-branes resemble k instantons in the transverse space. Thus the HL index of $H_G^{(k)}$ theory are supposed to be equal to the Hilbert series of reduced moduli space of k G -instantons. On the other hand, the Hilbert series can also be obtained from the 5d Nekrasov partition function with pure gauge group G , which are just the 5d limit of elliptic genus of 6d minimal $(1, 0)$ SCFT with type G . Therefore, we arrive at the relation:

$$\mathcal{I}_{H_G^{(k)}}^{\text{HL}} = \text{Hilb}_G^k = g_{k,G}^{(0)}, \tag{5.13}$$

where $g_{k,G}^{(0)}$ as we defined previously in (4.3) is the coefficient of leading Q_τ order of k -string elliptic genus $\mathbb{E}_{h_G^{(k)}}$. One can also add “tildes” to get the equality with a free hypermultiplet coupled, in which situation one encounters the full Hilbert series other than the reduced. We have checked relation (5.13) for $k = 1, 2$ for all possible G and $k = 3$ for $\text{SU}(3)$.³⁶

- (Schur) $q = t$ with p arbitrary. In fact, it can be shown in such specialization the index is independent of p . Thus, taking $p \rightarrow 0$, Schur index is actually a limit of Macdonald index. Using (5.5), the Schur index for a class \mathcal{S} theory is given by

$$\mathcal{I}^{\text{Schur}} = \hat{\mathcal{N}}_{N,s} \frac{\prod_{i=1}^s \hat{\mathcal{K}}_{\Lambda_i}(\mathbf{a}_i) \chi^\lambda(\mathbf{a}_i(\Lambda_i))}{[\chi^\lambda(v^{N-1}, v^{N-3}, \dots, v^{1-N})]^{s-2}}, \tag{5.14}$$

where³⁷

$$\hat{\mathcal{N}}_{N,s} = (v^2; v^2)^s \prod_{j=2}^N (v^{2j}; v^2)^{s-2}, \tag{5.15}$$

and χ^λ is the Schur polynomials defined as

$$\chi_\lambda(\mathbf{a}) = \frac{\det(a_i^{\lambda_j+k-j})}{\det(a_i^{k-j})}. \tag{5.16}$$

At last, one replaces back $v^2 \rightarrow q$.

The Schur indices in some sense are more interesting than the Hall-Littlewood indices. For instance, for class \mathcal{S} theories, Schur indices equal the q -deformed topological 2d

³⁶For $\text{SU}(3)$ and F_4 , we are not aware how to compute the HL indices directly. Still, the Hilbert series are well-defined and computed in [41, 42], which are in perfect agreement with our computation for elliptic genus from blowup equations.

³⁷As in this paper we only deal with the cases with three or four punctures, we also shorten $\mathcal{N}_{N,3}$ as \mathcal{N}_N and $\mathcal{N}_{N,4}$ as \mathcal{N}'_N in the latter subsections, and same for those with hat.

Yang-Mills partition function on the punctured Riemann surface [83], and also equal the vacuum character of the associated chiral algebra [73, 74]. Furthermore, Schur indices can be computed in IR via wall crossing for theories even beyond class \mathcal{S} , such as certain Argyres-Douglas theories [84] including rank one H_{A_2} theory.

The full superconformal indices of rank one $H_{D_4, E_6, 7}$ theories have been computed in [46, 48, 85]. The Schur index of rank one H_{E_8} was given in [37] and the Schur index of rank one H_{A_2} was given in [84]. To compute the Hall-Littlewood indices and Schur indices of higher rank $H_{D_4, E_6, 7, 8}$ theories one will encounter certain subtle issues. Directly using the general formulas (5.9) and (5.14) fails to give correct results, because at a given order of v infinite number of Young diagrams λ contribute. To cure such divergence, it was suggested in [45] that one reduces the flavor symmetry “one box at a time”, that is to change one specific puncture by moving one box down in the associated Young diagram. The physical meaning of such operation is interpreted as coupling a free hypermultiplet to $H_G^{(k)}$ theory, which in our notation is just $\tilde{H}_G^{(k)}$ theory. In the terminology of [45], $H_G^{(k)}$ are “bad” theories, while $\tilde{H}_G^{(k)}$ are “good” theories. One can directly use (5.9) and (5.14) to compute the indices of $\tilde{H}_G^{(k)}$, then divide by the index of a free hypermultiplet which is well defined, finally one will obtain the finite indices of $H_G^{(k)}$. Following this procedure, the Hall-Littlewood indices of rank two $H_{D_4, E_6, 7, 8}$ theories were computed in [45]. Similarly, we computed the Schur indices of rank two and three $H_{D_4, E_6, 7, 8}$ theories which will be shown in details in later sections. For higher rank H_{A_2} we are not aware how to compute its Schur indices due to the irregular punctures of 6d construction. Although there exist no H_G^k theory for $G = F_4$, we suspect certain analogy can be constructed such that Hall-Littlewood indices still make sense as the Hilbert series of moduli space of $k F_4$ instantons, and the Schur indices can be associated with affine \mathfrak{f}_4 algebra. One support for such speculation is that the Hilbert series for arbitrary $k F_4$ instantons has been constructed from certain folding from E_6 [43]. Thus we sometimes informally denote the analogy as $H_{F_4}^{(k)}$ theories.

5.3 Rank one: Del Zotto-Lockhart’s conjecture

In [37], Del Zotto-Lockhart found an intriguing structure of one string elliptic genera of 6d minimal $(1, 0)$ SCFTs and a surprising relation between the elliptic genera and the supersymmetric indices of rank one H_G theories. Let us rephrase their conjecture here:

Conjecture (Del Zotto-Lockhart). There exists a function $L_G^{(1)}(v, m_G, Q_\tau) = \sum_{i,j=0}^{\infty} b_{i,j}^G Q_\tau^i v^j$ such that

1. $b_{i,j}^G$ can be written as the sum of characters of irreducible representations of G with integral coefficients.
2. $L_G^{(1)}(v, m_G, 0)$ is the Hilbert series of the reduced moduli space of one G -instanton, i.e. the Hall-Littlewood index of the $H_G^{(1)}$ theory.
3. $L_G^{(1)}(q^{1/2}, m_G, q^2)$ is the Schur index of the $H_G^{(1)}$ theory.

i, j	0	2	4	6	8	10	12	14
0	1	28	300	1925	8918	32928	102816	282150
1	0	0	29	707	6999	42889	193102	699762
2	-1	0	0	2 · 1	463 + 1	9947	92391	544786
3	0	-28	-29	-2 · 1	1 - 1	2 · 29	5280 + 29 + 2 · 28	101850
4	0	0	-300	-707	-463 - 1	-2 · 29	29 - 29	2 · 463 + 2 · 1
5	0	0	0	-1925	-6999	-9947	-5280 - 29 - 2 · 28	-2 · 463 - 2 · 1
6	0	0	0	0	-8918	-42889	-92391	-101850
7	0	0	0	0	0	-32928	-193102	-544786
8	0	0	0	0	0	0	-102816	-699762
9	0	0	0	0	0	0	0	-282150

Table 13. Expansion coefficients $c_{i,j}^{\text{SO}(8)}$ for one SO(8) instanton string.

4. The reduced one-string elliptic genus $\mathbb{E}_{h_G^{(1)}}(v)$ can be generated from $L_G^{(1)}(v)$ by the following formula in which the symmetry (4.26) is manifest:³⁸

$$\begin{aligned}
 \mathbb{E}_{h_G^{(1)}}(v) = & v^{2h-1} Q_\tau^{1/6} \sum_{n \geq 0} Q_\tau^{2n} \left[u^{4h} L_G(Q_\tau^n v) - (-1)^{2h} u^{-4h} L_G(Q_\tau^{n+1/2}/v) \right. \\
 & + (1 + (-1)^{2h}) Q_\tau^{h+1/2} \left(u^2 L_G(Q_\tau^{n+1/2} v) - u^{-2} L_G(Q_\tau^{n+1}/v) \right) \\
 & \left. + Q_\tau^2 \left((-1)^{2h} u^{4(1-h)} L_G(Q_\tau^{n+1} v) - u^{-4(1-h)} L_G(Q_\tau^{n+3/2}/v) \right) \right] \quad (5.17)
 \end{aligned}$$

where $h = h_G^\vee/6$, $u = v/Q_\tau^{1/4}$.

The conjectural formula (5.17) is quite intricate. Roughly speaking, it means the coefficient matrix of reduced one-string elliptic genus contains several “blocks”, overlapping or non-overlapping, and each block contains infinite copies of the $L_G^{(1)}$ function. The number of blocks turns out to be 2 for SU(3), 4 for F_4 and 6 for the other G . In the following we show the coefficient matrix of one-string elliptic genus of SO(8) in a way consistent with our later higher rank discussion. The coefficient matrix of elliptic genus and the $L_G^{(1)}$ functions for other G can be found in [37]. Let us denote

$$\mathbb{E}_{h_{\text{SO}(8)}^{(1)}}(v, Q_\tau, m_i = 0) = v^5 Q_\tau^{-5/6} \sum_{i,j=0}^{\infty} c_{i,j}^{\text{SO}(8)} v^j (Q_\tau v^{-4})^i. \quad (5.18)$$

Then we have table 13 for the coefficients $c_{i,j}^{\text{SO}(8)}$ where each “block” is colored differently: the coefficients coming from the first term in the square bracket in (5.17) is colored red, the second black, the third blue, the fourth orange, the fifth cyan and the last magenta. As we can see from the table, the reduced one-string elliptic genus indeed depends on v^2 . One can also see the symmetry (4.26) on the two sides of the ray with slope $-1/2$. Here the

³⁸Here the dependence on Q_τ and Q_m are implied.

$L_{\text{SO}(8)}^{(1)}(v, Q_\tau)$ function can be defined by all the red numbers in table 13 with the red +1 and +29 moving out, as they come from $n = 1$ term in the summation. Thus we have

$$\begin{aligned}
 L_{\text{SO}(8)}^{(1)}(v, Q_\tau) &= (1+28v^2+300v^4+1925v^6+8918v^8+32928v^{10}+102816v^{12}+282150v^{14}+\mathcal{O}(v^{16})) \\
 &+ (29+707v^2+6999v^4+42889v^6+193102v^8+699762v^{10}+\mathcal{O}(v^{12}))Q_\tau \\
 &+ (463+9947v^2+92391v^4+544786v^6+\mathcal{O}(v^8))Q_\tau^2 \\
 &+ (5280+101850v^2+\mathcal{O}(v^4))Q_\tau^3+\mathcal{O}(Q_\tau^4). \tag{5.19}
 \end{aligned}$$

Clearly, the first row in table 13 gives the well-known Hilbert series for the reduced moduli space for one SO(8) instanton, i.e. the Hall-Littlewood index for rank one $H_{\text{SO}(8)}$ theory:

$$L_{\text{SO}(8)}^{(1)}(v, 0) = \sum_{n=0}^{\infty} \chi_{n\theta}^{\text{SO}(8)} v^{2n} = 1+28v^2+300v^4+1925v^6+8918v^8+32928v^{10}+102816v^{12}+\mathcal{O}(v^{14}) \tag{5.20}$$

Adding the red numbers from $L_{\text{SO}(8)}^{(1)}(v, Q_\tau)$ in each column of table 13 together, one expects to obtain the Schur index of rank one $H_{\text{SO}(8)}$ theory. Indeed, by making $v \rightarrow q^{1/2}$ to make contact with the literature, we obtain

$$L_{\text{SO}(8)}(q^{1/2}, q^2) = 1+28q+329q^2+2632q^3+16380q^4+85764q^5+393589q^6+1628548q^7+\mathcal{O}(q^8). \tag{5.21}$$

Such series was actually already obtained by a lot of methods. For example, from the viewpoint of VOA/SCFT correspondence, it equals the vacuum character of affine Lie algebra $\mathfrak{so}(8)_{k=-2}$ [73]. From the nature that rank one $H_{\text{SO}(8)}$ theory is actually just SU(2) gauge theory with $N_f = 4$, the Schur index can be computed both from UV Lagrangian and IR wall-crossing formula [86]. See the Schur series from vacuum character up to q^{14} in the end of the appendix of [86].

Such comparison between the reduced elliptic genus and Schur index for all other rank one H_G theory except $G = F_4$ has been done in [37]. In particular, all $L_G^{(1)}(v, Q_\tau, m_G = 0)$ functions are identified, and the conjectural formula (5.17) holds to substantial orders. Similarly, one can also couple a free hypermultiplet to establish the relation between original one-string elliptic genus $\mathbb{E}_{\tilde{h}_G^{(1)}}(v)$ and the Hall-Littlewood and Schur indices of $\tilde{H}_G^{(k)}$ theory. Indeed, the Schur index of a 4d hypermultiplet is known to be [86]

$$\mathcal{I}_{h.m.}^{\text{Schur}} = \text{PE} \left[\frac{q^{1/2}}{1-q} (x + x^{-1}) \right], \tag{5.22}$$

which can also be obtained by taking limit $\mathbb{E}_{\text{c.m.}}(v, x, Q_\tau) \rightarrow \mathbb{E}_{\text{cm}}(q^{1/2}, x, q^2)$. The Hall-Littlewood index of a 4d hypermultiplet i.e. the Hilbert series of \mathbb{C}^2 is well-known to be

$$\mathcal{I}_{h.m.}^{\text{HL}} = \frac{1}{(1-vx^{\pm 1})}, \tag{5.23}$$

which can also be obviously obtained by taking limit $\mathbb{E}_{\text{c.m.}}(v, x, Q_\tau \rightarrow 0)$, with a factor $vQ_\tau^{-1/6}$ absorbed into the overall factor of (4.3). This makes the whole story consistent.

In the viewpoint of pure 4d, this intriguing conjecture indicates there exists certain precise relation between the β -twisted partition function on $T^2 \times S^2$ and the partition function on $S^3 \times S^1$. We suspect the connection may be established by realizing one S^1 of T^2 as the Hopf fibration over S^2 to get $S^3 \times S^1$. To find the consequence of such realization one has to go into the details of localization which is beyond the scope of current paper.³⁹

5.4 Rank two

We would like to generalize Del Zotto-Lockhart’s conjecture to the rank two cases, where there exist more flavour symmetry that is $SU(2)_x$ in H_G theories. To be precise, we want to find some functions $L_G^{(2)}(v, x, m_G, Q_\tau) = \sum_{i,j=0}^\infty b_{i,j}^G Q_\tau^i v^j$ such that

1. $b_{i,j}^G$ can be written as the sum of products between the characters of irreducible representations of $SU(2)_x$ and the characters of irreducible representations of G with integral coefficients.
2. $L_G^{(2)}(v, x, m_G, 0)$ is the Hilbert series of the reduced moduli space of two G -instanton, i.e. the Hall-Littlewood index of the $H_G^{(2)}$ theory.
3. $L_G^{(2)}(q^{1/2}, x, m_G, q^2)$ is the Schur index of the $H_G^{(2)}$ theory.
4. The reduced two-string elliptic genus $\mathbb{E}_{h_G^{(2)}}(v, x, m_G, Q_\tau)$ can be generated from $L_G^{(2)}(v, x, m_G, Q_\tau)$ and $L_G^{(1)}(v, x, m_G, Q_\tau)$ functions.

It turns out the rank two cases are much more complicated than the rank one cases, one reason for which is that we can not rely on the additional symmetry (4.27). Although we have not achieved an exact formula to generate the two string elliptic genus, we successfully manage to identify the $L_G^{(2)}$ functions to substantial orders, which we will elaborate on later for each example. In fact, the leading and subleading Q_τ order of $L_G^{(2)}(v, x, m_G, Q_\tau)$ are just given by $g_{2,G}^{(0)}(v, x, m_G)$ in (4.8) and $g_{2,G}^{(1)}(v, x, m_G)$ in (4.9), while the subsubleading order is given by

$$(\chi_5 + (\chi_\theta + 2)\chi_3 + \chi_{S_{\text{Sym}^2\theta}} + 2\chi_\theta + 3) + ((\chi_\theta + 1)\chi_4 + ((\chi_\theta + 1)^2 + (2\chi_\theta + 1))\chi_2)v + \dots, \quad (5.24)$$

which differs from $g_{2,G}^{(2)}(v, x, m_G)$ in (4.10) by $1 + \chi_2 v + \dots$. Such difference is recognized as what we call “blue” series in contrast to the red $L_G^{(2)}$ functions. Indeed, the reason we also include $L_G^{(1)}$ in the last condition is that we observe a “blue” series appearing multiple times in the coefficient matrix of $\mathbb{E}_{h_G^{(2)}}$:

$$\begin{aligned} M_G^{(2),\text{blue}}(v, x) &= \sum_{n=0}^\infty v^n \sum_{i+2j=n+1} \chi_i \chi_j \theta = \frac{1}{(1-vx)(1-v/x)} g_{1,G}^{(0)}(v) \\ &= 1 + \chi_2 v + (\chi_3 + \chi_\theta)v^2 + (\chi_4 + \chi_\theta \chi_2)v^3 + (\chi_5 + \chi_\theta \chi_3 + \chi_{2\theta})v^4 + (\chi_6 + \chi_\theta \chi_4 + \chi_{2\theta} \chi_2)v^5 + \dots \end{aligned} \quad (5.25)$$

³⁹Guglielmo Lockhart came up independently with a similar idea (private communication).

i, j	-2	-1	0	1	2	3	4	5	6	7	8
0	0	0	1	0	11	16	65	142	335	700	1542
1	0	0	1	2	11	20	12+56	18+92	143+192	356+292	1091+517
2	-1	0	1	-2	2	0	51	150	473	1032	90+2225

i, j	9	10	11	12
0	2788	5350	9288	16184
1	2676+742	6387+1183	13476+1624	28204+2408
2	232+4024	8589	15552	30469

Table 14. Unrefined coefficients $c_{i,j}^{A_2}$ for the elliptic genus of two SU(3) instanton strings.

For example, the blue series always appears at Q_τ order $h_G^\vee/3$ with leading v order $-2h_G^\vee/3$ (comparing to the leading Q_τ order). The reason for such phenomenon is yet not clear to us.

On the other hand, from the technique of class \mathcal{S} theory, we can compute the Schur index of $H_G^{(2)}$ theories for $G = D_4, E_{6,7,8}$. All of them are in agreement with our expectation from elliptic genera up to quite high orders. For example, from the $L_G^{(2)}$ functions, we are able to write down the following general formula for the Schur indices up to $q^{7/2}$:

$$\begin{aligned}
 \mathcal{I}_{H_G^{(2)}}^{\text{Schur}} &= \mathcal{I}_{\tilde{H}_G^{(2)}}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} = 1 + (\chi_3 + \chi_\theta)q + \chi_\theta \chi_2 q^{3/2} + (\chi_5 + (\chi_\theta + 1)\chi_3 + \chi_{\text{Sym}^2 \theta} + \chi_\theta + 1)q^2 \\
 &\quad + (\chi_\theta \chi_4 + (\chi_{2\theta} + \chi_{\text{Sym}^2 \theta} + 1)\chi_2)q^{5/2} + (\chi_7 + (\chi_\theta + 1)\chi_5 + (\chi_{2\theta} + \chi_{\text{Sym}^2 \theta} + 2\chi_\theta + 3)\chi_3 \\
 &\quad + \chi_{\text{Sym}^3 \theta} + (\chi_\theta + 1)^2 - C_6(G))q^3 + (\chi_\theta \chi_6 + (\chi_{2\theta} + \chi_{\text{Alt}^2 \theta} + 2\chi_\theta + 1)\chi_4 \\
 &\quad + (\chi_{3\theta} + 2\chi_{2\theta} + (\chi_\theta + 1)^2 + \chi_{\text{Sym}^2 \theta} + \chi_{\text{Alt}^2 \theta} + B_2(G) + C_7(G))\chi_2)q^{7/2} + \dots \quad (5.26)
 \end{aligned}$$

In the following, we show the striking comparison between elliptic genus and indices at rank two for all symmetry group G .

SU(3). For SU(3), let us denote the two-string elliptic genus as

$$\mathbb{E}_{h_{A_2}^{(2)}}(v, x, Q_\tau, Q_m) = v^5 Q_\tau^{-5/6} \sum_{i,j=0}^{\infty} c_{i,j}^{A_2}(x, Q_m) v^i (Q_\tau v^{-4})^j. \quad (5.27)$$

Then we have the unrefined coefficients $c_{i,j}^{A_2}(x = 1, Q_m = 1)$ listed in table 14. Keeping in mind that all such numbers can be refined to incorporate SU(2)_x, we show the unrefined coefficients just to make them look clearer. The red numbers give the definition of $L_G^{(2)}$ functions. In particular, they are in agreement with the universal expansion (4.8), (4.9) and (5.24). Note the red numbers in the first row agrees with the Hilbert series for reduced moduli space of two A_2 instantons in [42]. The two red numbers in the $i = 2$ rows are predicted from (5.24). Besides, the blue numbers agree with our proposal (5.25). Adding the red numbers in each column together, we expect to obtain a series that is equal to the Schur index of rank two H_{A_2} 4d SCFT.

The construction of $H_{A_2}^{(2)}$ theory from 6d involves irregular punctures. We are not aware how to directly compute its indices. We write our prediction from elliptic genus

here: the Hall-Littlewood index of rank two H_{A_2} theory is

$$\begin{aligned} \mathcal{I}_{H_{A_2}^{(2)}}^{\text{HL}} &= 1 + (\chi_3 + 8)q + 8\chi_2 q^{3/2} + (\chi_5 + 8\chi_3 + 36)q^2 + (8\chi_4 + 55\chi_2)q^{5/2} \\ &\quad + (\chi_7 + 8\chi_5 + 63\chi_3 + 119)q^3 + (8\chi_6 + 55\chi_4 + 216\chi_2)q^{7/2} \\ &\quad + (\chi_9 + 8\chi_7 + 63\chi_5 + 280\chi_3 + 322)q^4 + (8\chi_8 + 55\chi_6 + 280\chi_4 + 637\chi_2)q^{9/2} + \mathcal{O}(q^5), \end{aligned} \tag{5.28}$$

which agrees with the Hilbert series of reduced moduli space of two $SU(3)$ instantons [42], and the Schur index of rank two H_{A_2} theory is

$$\begin{aligned} \mathcal{I}_{H_{A_2}^{(2)}}^{\text{Schur}} &= 1 + (\chi_3 + 8)q + 8\chi_2 q^{3/2} + (\chi_5 + 9\chi_3 + 45)q^2 + (8\chi_4 + 64\chi_2)q^{5/2} \\ &\quad + (\chi_7 + 9\chi_5 + 82\chi_3 + 200)q^3 + (8\chi_6 + 72\chi_4 + 360\chi_2)q^{7/2} \\ &\quad + (\chi_9 + 9\chi_7 + 83\chi_5 + 479\chi_3 + 799)q^4 + (8\chi_8 + 72\chi_6 + 496\chi_4 + 1608\chi_2)q^{9/2} + \mathcal{O}(q^5). \end{aligned} \tag{5.29}$$

Taking $x = 1$ in (5.29), we have the unrefined Schur index as

$$1 + 11q + 16q^{3/2} + 77q^2 + 160q^{5/2} + 498q^3 + 1056q^{7/2} + 2723q^4 + 5696q^{9/2} + \mathcal{O}(q^5). \tag{5.30}$$

This is in complete agreement with Beem-Rastalli's to appear computation from chiral algebra!⁴⁰

SO(8). The $H_{D_4}^{(2)}$ theory can be constructed by compactifying $A_3(2,0)$ 6d SCFT on a sphere with four square punctures $\{2^2\}$, i.e. 2222 theory, which is expected to be a $usp(4)$ gauge theory with four fundamental hypermultiplets and one anti-fundamental. On the other hand, the $\tilde{H}_{D_4}^{(2)}$ theory can be constructed as a 222L theory, i.e. we replace one $\{2^2\}$ puncture to $\{2, 1^2\}$. In [45], the Hall-Littlewood indices of both 222L theory and $usp(4) + 4f + 1a$ theory were computed, which are in relation

$$\mathcal{I}_{222L}(v, x, m_i) = \frac{1}{1 - vx^{\pm 1}} \mathcal{I}_{usp(4)+4f+1a}(v, x, m_i). \tag{5.31}$$

We expect and indeed checked to high orders

$$\mathcal{I}_{usp(4)+4f+1a}(v, x, m_i) = g_{0, D_4}^{(2)}(\tau, a, m_i). \tag{5.32}$$

For example, one can directly see the series coefficients in (A.12) of [45] agree with the Q_τ^0 entries in table 15.

The Schur index of 222L theory can be obtained in a similar manner. Following the general formula in [44], we obtain

$$\begin{aligned} \mathcal{I}_{222L}^{\text{Schur}}(c, d, e; a, b) &= \hat{\mathcal{N}}'_4 \hat{\mathcal{K}}_1(c) \hat{\mathcal{K}}_1(d) \hat{\mathcal{K}}_1(e) \hat{\mathcal{K}}_2(a, b) \sum_{\lambda} \frac{\chi_{\lambda}(vb, v^{-1}b, b^{-1}a, b^{-1}a^{-1})}{\chi_{\lambda}^2(v^{-3}, v^{-1}, v, v^3)} \\ &\quad \times \chi_{\lambda}(vc, v^{-1}c, vc^{-1}, v^{-1}c^{-1}) \chi_{\lambda}(vd, v^{-1}d, vd^{-1}, v^{-1}d^{-1}) \chi_{\lambda}(ve, v^{-1}e, ve^{-1}, v^{-1}e^{-1}), \end{aligned} \tag{5.33}$$

⁴⁰We thank Beem and Rastelli for providing us their unpublished results on the unrefined Schur index of rank two $SU(3)$ theory.

where $(b_1 = b, b_2 = 1/b)$. The summation is over Young diagrams $\lambda = (\lambda_1, \lambda_2, \lambda_3, 0)$. The $\hat{\mathcal{N}}$ and $\hat{\mathcal{K}}$ factors are given by

$$\begin{aligned} \hat{\mathcal{N}}_4 &= (v^2; v^2)^4 \prod_{j=2}^4 (v^{2j}; v^2)^2, \\ \hat{\mathcal{K}}_1(b) &= \text{PE} \left[\frac{(v^2 + v^4)(b^2 + b^{-2} + 2)}{1 - v^2} \right], \\ \hat{\mathcal{K}}_2(a, b) &= \text{PE} \left[\frac{3v^2 + v^4 + v^3 b^{\pm 2} a^{\pm 1} + v^2 a^{\pm 2}}{1 - v^2} \right]. \end{aligned} \tag{5.34}$$

At last, one usually replaces $v \rightarrow q^{1/2}$ to make contact with literature. From the above formula, we computed the Schur index up to v^{20} as

$$\mathcal{I}_{222L}^{\text{Schur}} = 1 + \chi_2 v + (2\chi_3 + 28)v^2 + (2\chi_4 + 58\chi_2)v^3 + (3\chi_5 + 87\chi_3 + 465)v^4 + \dots \tag{5.35}$$

Decoupling the free hypermultiplet, we obtain the Schur index of $H_{D_4}^{(2)}$ theory

$$\mathcal{I}_{H_{D_4}^{(2)}}^{\text{Schur}} = \mathcal{I}_{usp(4)+4f+1a}^{\text{Schur}} = \mathcal{I}_{222L}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} \tag{5.36}$$

up to q^{10} . The first 12 terms with full $SU(2)_x$ fugacity are

$$\begin{aligned} \mathcal{I}_{H_{D_4}^{(2)}}^{\text{Schur}} &= 1 + (\chi_3 + 28)q + 28\chi_2 q^{3/2} + (\chi_5 + 29\chi_3 + 435)q^2 + (28\chi_4 + 707\chi_2)q^{5/2} \\ &\quad + (\chi_7 + 29\chi_5 + 765\chi_3 + 4845)q^3 + (28\chi_6 + 735\chi_4 + 9947\chi_2)q^{7/2} \\ &\quad + (\chi_9 + 29\chi_7 + 766\chi_5 + 12337\chi_3 + 43353)q^4 \\ &\quad + (28\chi_8 + 735\chi_6 + 12607\chi_4 + 101878\chi_2)q^{9/2} \\ &\quad + (\chi_{11} + 29\chi_9 + 766\chi_7 + 12667\chi_5 + 141518\chi_3 + 330360)q^5 \\ &\quad + (28\chi_{10} + 735\chi_8 + 12635\chi_6 + 155449\chi_4 + 845225\chi_2)q^{11/2} + \mathcal{O}(q^6). \end{aligned} \tag{5.37}$$

We can compare this with elliptic genus up to $q^{11/2}$. Let us denote the $SO(8)$ two-string elliptic genus as

$$\mathbb{E}_{h_{D_4}^{(2)}}(v, x, \tau, m_i = 0) = v^{11} Q_\tau^{-11/6} \sum_{i,j=0}^{\infty} c_{i,j}^{\text{SO}(8)}(x) v^j (Q_\tau v^{-4})^i. \tag{5.38}$$

Then we have table 15 for the coefficients $c_{i,j}^{\text{SO}(8)}(x = 1)$. Here the red numbers are from the $L_{D_4}^{(2)}$ series. Add the red numbers in each column together, we expect to obtain a series that is equal to the Schur index of rank two H_{D_4} 4d SCFT. Indeed, we have

$$\begin{aligned} L_{D_4}^{(2)}(q^{1/2}, x = 1, m_{D_4} = 0, q^2) &= 1 + 31q + 56q^{3/2} + 527q^2 + 1526q^{5/2} + 7292q^3 + 23002q^{7/2} \\ &\quad + 84406q^8 + 258818q^{9/2} + 823883q^5 + 2394216q^{11/2} + \dots \end{aligned} \tag{5.39}$$

On the other hand, by taking the unrefined limit $x = 1$ in (5.36), we obtain the unrefined Schur series

$$\begin{aligned} &1 + 31q + 56q^{3/2} + 527q^2 + 1526q^{5/2} + 7292q^3 + 23002q^{7/2} + 84406q^8 + 258818q^{9/2} + 823883q^5 \\ &\quad + 2394216q^{11/2} + 6943434q^6 + 19082748q^{13/2} + 51665849q^7 + 134888730q^{15/2} + 345764537q^8 \\ &\quad + 862482876q^{17/2} + 2112344321q^9 + 5061362222q^{19/2} + 11921262927q^{10} + \mathcal{O}(q^{21/2}). \end{aligned} \tag{5.40}$$

One can see the two series match perfectly up to $q^{11/2}$!

i, j	0	1	2	3	4	5	6	7	8
0	1	0	31	56	495	1468	6269	19680	64768
1	0	0	0	0	32	58	1023	3322	19078
2	-1	-2	-31	-60	-389	-718	$-2972 + 2 \cdot 1$	$-5226 + 2 \cdot 2$	$560 - 16398 + 2 \cdot 31 + 1$

i, j	9	10	11
0	187792	537021	1424526
1	69114	266799	886104
2	$1912 - 27570 + 2 \cdot 60 + 2$	$20063 - 71670 + 2 \cdot 389 + 31$	$83586 - 115770 + 2 \cdot 718 + 60$

Table 15. Series coefficients $c_{i,j}^{\text{SO}(8)}$ for the elliptic genus of two SO(8) instanton strings.

i, j	0	1	2	3	4	5	6	7	8	9
0	1	0	55	104	1539	4966	32091	119340	542109	1973088
1	0	0	0	0	56	106	3135	10900	97125	405480
2	0	0	0	0	0	0	0	0	$1652 + 1$	$6040 + 2$
3	1	2	55	108	1214	2320	15802	29284	$143542 - 1$	$257800 - 2$

i, j	10	11	12
0	7460100	25288640	84766812
1	2210027	9075756	38900537
2	$99611 + 55$	$466860 + 108$	$3399668 + 1214$
3	$999970 - 55$	$1742140 - 108$	5704242

Table 16. Series coefficients $c_{i,j}^{F_4}$ for the elliptic genus of two F_4 instanton strings.

F_4 . Let us denote the two-string elliptic genus with gauge symmetry F_4 as

$$\mathbb{E}_{h_{F_4}^{(2)}}(v, x = 1, \tau, m_i = 0) = v^{17} Q_\tau^{-17/6} \sum_{i,j=0}^{\infty} c_{i,j}^{F_4} v^j (Q_\tau v^{-4})^i. \quad (5.41)$$

Then we have table 16 for the unrefined coefficients $c_{i,j}^{F_4}$. The red numbers in the first row agree with the Hilbert series for reduced moduli space of two F_4 instantons in [42]. By summing over the red numbers in each column, we obtain certain analogy of Schur index of rank two H_G theory for F_4 up to $q^{11/2}$. The unrefined version is

$$1 + 55q + 104q^{3/2} + 1595q^2 + 5072q^{5/2} + 35226q^3 + 130240q^{7/2} + 640886q^4 + 2384608q^{9/2} + 9769738q^5 + 34831256q^{11/2} + \mathcal{O}(q^6). \quad (5.42)$$

This is in complete agreement with Beem-Rastalli's to appear computation from chiral algebra!⁴¹

⁴¹We thank Beem and Rastelli for providing us their unpublished results on the unrefined Schur index of rank two F_4 theory.

E₆. The $H_{E_6}^{(2)}$ theory can be constructed by compactifying A_5 (2, 0) 6d SCFT on a sphere with three $\{2^3\}$ punctures, which is a “bad” theory. One can change one of the punctures to $\{2^2, 1^2\}$ to add a decoupled hypermultiplet, i.e. the $\tilde{H}_{E_6}^{(2)}$ theory. The Hall-Littlewood index of this theory was computed in [45]. We expect and indeed checked

$$\mathcal{I}_{H_{E_6}^{(2)}}^{\text{HL}}(v, x, m_{E_6}) = g_{0, E_6}^{(2)}(v, x, m_{E_6}). \tag{5.43}$$

The Schur index can be obtained in a similar manner. Following the general formula in [44], we obtain

$$\begin{aligned} \mathcal{I}_{\tilde{H}_{E_6}^{(2)}}^{\text{Schur}} &= \hat{\mathcal{N}}_6 \hat{\mathcal{K}}_1(a_1, a_2) \hat{\mathcal{K}}_1(a_3, a_4) \hat{\mathcal{K}}_2(a_5, a_6, x) \sum_{\lambda} \frac{\chi_{\lambda}(va_5, v^{-1}a_5, va_6, v^{-1}a_6, \frac{x}{a_5a_6}, \frac{x^{-1}}{a_5a_6})}{\chi_{\lambda}(v^{-5}, v^{-3}, v^{-1}, v^1, v^3, v^5)} \\ &\times \chi_{\lambda}\left(v a_1, v^{-1} a_1, v a_2, v^{-1} a_2, v \frac{1}{a_1 a_2}, v^{-1} \frac{1}{a_1 a_2}\right) \chi_{\lambda}\left(v a_3, v^{-1} a_3, v a_4, v^{-1} a_4, v \frac{1}{a_3 a_4}, v^{-1} \frac{1}{a_3 a_4}\right). \end{aligned} \tag{5.44}$$

Here $\lambda = (\lambda_1, \dots, \lambda_5, 0)$ and $(b_3 \equiv \frac{1}{b_1 b_2})$

$$\begin{aligned} \hat{\mathcal{N}}_6 &= (v^2; v^2)^3 \prod_{j=2}^6 (v^{2j}; v^2), \\ \hat{\mathcal{K}}_1(b_1, b_2) &= \prod_{\ell=1}^2 \prod_{i,j=1}^3 \text{PE} \left[\frac{v^{2\ell} b_i / b_j}{1 - v^2} \right], \\ \hat{\mathcal{K}}_2(b_1, b_2, x) &= \prod_{\ell=1}^2 \prod_{i,j=1}^2 \text{PE} \left[\frac{v^{2\ell} b_i / b_j}{1 - v^2} \right] \times \text{PE} \left[\frac{2v^2 + v^2 x^{\pm 2} + v^3 (b_1^2 b_2 x^{\pm 1})^{\pm 1} + v^3 (b_1 b_2^2 x^{\pm 1})^{\pm 1}}{1 - v^2} \right]. \end{aligned} \tag{5.45}$$

At last, one needs to replace $v \rightarrow q^{1/2}$. We computed the Schur index up to q^7 :

$$\begin{aligned} \mathcal{I}_{H_{E_6}^{(2)}}^{\text{Schur}} &= 1 + (\chi_3 + 78)q + 78\chi_2 q^{3/2} + (\chi_5 + 79\chi_3 + 3160)q^2 + (78\chi_4 + 5512\chi_2)q^{5/2} \\ &+ (\chi_7 + 79\chi_5 + 5670\chi_3 + 87751)q^3 + (78\chi_6 + 5590\chi_4 + 201292\chi_2)q^{7/2} + (\chi_9 + 79\chi_7 \\ &+ 5671\chi_5 + 248290\chi_3 + 1871196)q^4 + (78\chi_8 + 5590\chi_6 + 250640\chi_4 + 5048654\chi_2)q^{9/2} \\ &+ (\chi_{11} + 79\chi_9 + 5671\chi_7 + 250400\chi_5 + 7248975\chi_3 + 32615793)q^5 \\ &+ (78\chi_{10} + 5590\chi_8 + 250718\chi_6 + 7900243\chi_4 + 97665932\chi_2)q^{11/2} \\ &+ (\chi_{13} + 79\chi_{11} + 5671\chi_9 + 250801\chi_7 + 7949911\chi_5 + 157280287\chi_3 + 483480405)q^6 \\ &+ (78\chi_{12} + 5590\chi_{10} + 250718\chi_8 + 7949591\chi_6 + 186447755\chi_4 + 1552411211\chi_2)q^{13/2} \\ &+ (\chi_{15} + 79\chi_{13} + 5671\chi_{11} + 250801\chi_9 + 7952421\chi_7 + 193661181\chi_5 + 2725694921\chi_3 \\ &+ 6263699772)q^7 + \dots \end{aligned} \tag{5.46}$$

Note the leading terms up to $q^{7/2}$ agree with our general proposal (5.26).

Let us denote the two-string elliptic genus as

$$\mathbb{E}_{h_{E_6}^{(2)}}(v, x = 1, \tau, m_i = 0) = v^{23} Q_{\tau}^{-23/6} \sum_{i,j=0}^{\infty} c_{i,j}^{E_6} v^j (Q_{\tau} v^{-4})^i. \tag{5.47}$$

i, j	0	1	2	3	4	5	6	7	8	9
0	1	0	81	156	3320	11178	98440	401280	2344619	9785226
1	0	0	0	0	82	158	6723	24132	296879	1335694
2	0	0	0	0	0	0	0	0	3485+1	13112+2
3	0	0	0	0	0	0	0	0	-1	-2
4	-1	-2	-81	-160	-2669	-5178	-51445	-97712	-681945	-1266178

i, j	10	11	12	13
0	45870686	182872426	746229150	2782158570
1	9484963	44112702	236141466	1042037420
2	301488+81	1497516+160	14405643+2669	75613998+5178
3	-81	-160	102090-2669+83	563580-5178+322
4	-6819518 + 2 · 1	-12372858 + 2 · 2	-54611704 + 2 · 81 - 83	-96850550 + 2 · 160 - 322

i, j	14	15
0	10261780870	35695088906
1	4709271558	19202312882
2	486421964+51445	2415319754+97712
3	9603627-51445+7039	58071366-97712+24620
4	-365050846 + 2 · 2669 - 7039	-633251142 + 2 · 5178 - 24620

Table 17. Series coefficients $c_{i,j}^{E_6}$ for the unrefined elliptic genus of two E_6 instanton strings.

Then we have table 17 for the coefficients $c_{i,j}^{E_6}$. Here the red numbers are from the $L_{E_6}^{(2)}$ series. Add the red numbers in each column together, we expect to obtain a series that is equal to the Schur index of rank two H_{E_6} 4d SCFT. Indeed, we have

$$\begin{aligned}
 & L_{E_6}^{(2)}(q^{1/2}, x=1, m_{E_6}=0, q^2) \\
 &= 1 + 81q + 156q^{3/2} + 3402q^2 + 11336q^{5/2} + 105163q^3 + 425412q^{7/2} \\
 &\quad + 2644983q^4 + 11134032q^{9/2} + 55655137q^5 + 228482644q^{11/2} + 996878349q^6 \\
 &\quad + 3900373568q^{13/2} + 15467078019q^7 + 57370792908q^{15/2} + \dots
 \end{aligned} \tag{5.48}$$

On the other hand, taking $x=1$ in (5.46), the unrefined Schur index is

$$\begin{aligned}
 & 1 + 81q + 156q^{3/2} + 3402q^2 + 11336q^{5/2} + 105163q^3 + 425412q^{7/2} + 2644983q^4 + 11134032q^{9/2} \\
 &\quad + 55655137q^5 + 228482644q^{11/2} + 996878349q^6 + 3900373568q^{13/2} + 15467078019q^7 + \dots
 \end{aligned} \tag{5.49}$$

We can see the two series match perfectly up to q^7 !

E_7 . The $H_{E_7}^{(2)}$ theory can be constructed by compactifying $A_7(2,0)$ 6d SCFT on a sphere with one $\{4^2\}$ puncture and two $\{2^4\}$ punctures, which is a “bad” theory. One can change one of the $\{2^4\}$ punctures to $\{2^3, 1^2\}$ to add a decoupled hypermultiplet, i.e. the $\tilde{H}_{E_7}^{(2)}$ theory. The Hall-Littlewood index of this theory was computed in [45]. We find it agrees with our computation for $g_{0,E_7}^{(2)}(\tau, x, m_{E_7})$. The Schur index can be obtained in a similar

i, j	0	1	2	3	4	5	6	7	8	9
0	1	0	136	266	9315	32830	449050	2026080	17179899	84195608
1	0	0	0	0	137	268	18768	69544	1349005	6575250
2	0	0	0	0	0	0	0	0	9590+1	36982+2

Table 18. Series coefficients $c_{i,j}^{E_7}$ for the unrefined elliptic genus of two E_7 instanton.

manner. Following the general formula in [44], we obtain

$$\begin{aligned}
 \mathcal{I}_{\tilde{H}_{E_7}^{(2)}}^{\text{Schur}} &= \hat{\mathcal{N}}_8 \hat{\mathcal{K}}_1(a_1, a_2, a_3) \hat{\mathcal{K}}_2(a_4, a_5, a_6, x) \hat{\mathcal{K}}_3(a_7) \\
 &\times \sum_{\lambda} \frac{\chi_{\lambda}(v^3 a_7, v^{-3} a_7, v a_7, v^{-1} a_7, v^3 a_7^{-1}, v^{-3} a_7^{-1}, v a_7^{-1}, v^{-1} a_7^{-1})}{\chi_{\lambda}(v^{-7}, v^{-5}, v^{-3}, v^{-1}, v, v^3, v^5, v^7)} \\
 &\times \chi_{\lambda} \left(v a_1, v^{-1} a_1, v a_2, v^{-1} a_2, v a_3, v^{-1} a_3, v \frac{1}{a_1 a_2 a_3}, v^{-1} \frac{1}{a_1 a_2 a_3} \right) \\
 &\times \chi_{\lambda} \left(v a_4, v^{-1} a_4, v a_5, v^{-1} a_5, v a_6, v^{-1} a_6, \frac{x}{a_4 a_5 a_6}, \frac{x^{-1}}{a_4 a_5 a_6} \right).
 \end{aligned} \tag{5.50}$$

Here

$$\begin{aligned}
 \hat{\mathcal{K}}_3(b) &= \text{PE} \left[\frac{2(v^2 + v^4 + v^6 + v^8)}{1 - v^2} \right] \prod_{\ell=1}^4 \text{PE} \left[\frac{v^{2\ell} b^{\pm 2}}{1 - v^2} \right], \\
 \hat{\mathcal{K}}_2(b_1, b_2, b_3, x) &= \text{PE} \left[\frac{2v^2 + v^2 x^{\pm 2}}{1 - v^2} \right] \left[\prod_{\ell=1}^2 \prod_{i,j=1}^3 \text{PE} \left[\frac{v^{2\ell} b_i / b_j}{1 - v^2} \right] \right], \prod_{i=1}^3 \left[\frac{v^3 (x^{-1} b_i / b_4)^{\pm 1} + v^3 (x b_i / b_4)^{\pm 1}}{1 - v^2} \right], \\
 \hat{\mathcal{K}}_1(b_1, b_2, b_3) &= \prod_{\ell=1}^2 \prod_{i,j=1}^4 \text{PE} \left[\frac{v^{2\ell} b_i / b_j}{1 - v^2} \right].
 \end{aligned} \tag{5.51}$$

At last, one needs to replace $v \rightarrow q^{1/2}$. We computed the Schur index up to q^2 order.

After decoupling the free hypermultiplet, the Schur index of $H_{E_7}^{(2)}$ theory is given by

$$\mathcal{I}_{\tilde{H}_{E_7}^{(2)}}^{\text{Schur}} = \mathcal{I}_{\tilde{H}_{E_7}^{(2)}}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} = 1 + (\chi_3 + \mathbf{133})q + \mathbf{133}\chi_2 q^{3/2} + (\chi_5 + (\mathbf{133} + 1)\chi_3 + \text{Sym}^2 \mathbf{133} + \mathbf{133} + 1)q^2 + \dots \tag{5.52}$$

Let us denote the two-string elliptic genus as

$$\mathbb{E}_{h_{E_7}^{(2)}}(v, x = 1, \tau, m_i = 0) = v^{35} Q_{\tau}^{-35/6} \sum_{i,j=0}^{\infty} c_{i,j}^{E_7} v^j (Q_{\tau} v^{-4})^i. \tag{5.53}$$

Then we have table 18 for the coefficients $c_{i,j}^{E_7}$. Here the red numbers are from the $L_{E_7}^{(2)}$ series. Add the red numbers in each column together, we expect to obtain a series that is equal to the Schur index of rank two H_{E_7} 4d SCFT. Thus, we predict the unrefined Schur index as

$$\begin{aligned}
 &1 + 136q + 266q^{3/2} + 9452q^2 + 33098q^{5/2} + 467818q^3 + 2095624q^{7/2} + 18538494q^4 \\
 &+ 90807840q^{9/2} + \mathcal{O}(q^5).
 \end{aligned} \tag{5.54}$$

Indeed, taking $x = 1$ in (5.52), the unrefined Schur index is given by

$$1 + 136q + 266q^{3/2} + 9452q^2 + \mathcal{O}(q^{5/2}). \quad (5.55)$$

We can see the two series match perfectly!

E_8 . The $H_{E_8}^{(2)}$ theory can be constructed by compactifying A_{11} (2,0) 6d SCFT on a sphere with three $\{6^2\}$, $\{4^3\}$ and $\{2^6\}$, which is a “bad” theory. One can change the $\{2^6\}$ puncture to $\{2^5, 1^2\}$ to add a decoupled free hypermultiplet, i.e. the $\tilde{H}_{E_8}^{(2)}$ theory. Following the general formula in [44], we obtain its Schur index as

$$\begin{aligned} \mathcal{I}_{\tilde{H}_{E_8}^{(2)}}^{\text{Schur}} &= \hat{\mathcal{N}}_{12} \hat{\mathcal{K}}_1(a_1, a_2, a_3, a_4, a_5, x) \hat{\mathcal{K}}_2(a_6, a_7) \hat{\mathcal{K}}_3(a_8) \sum_{\lambda} \frac{\chi_{\lambda}(va_1, v^{-1}a_1, \dots, va_5, v^{-1}a_5, \frac{x}{a_1 \dots a_5}, \frac{x^{-1}}{a_1 \dots a_5})}{\chi_{\lambda}(v^{-11}, v^{-9}, \dots, v^9, v^{11})} \\ &\quad \times \chi_{\lambda} \left(v^3 a_6, v^{-3} a_6, va_6, v^{-1} a_6, v^3 a_7, v^{-3} a_7, va_7, v^{-1} a_7, v^3 \frac{1}{a_6 a_7}, v^{-3} \frac{1}{a_6 a_7}, v \frac{1}{a_6 a_7}, v^{-1} \frac{1}{a_6 a_7} \right) \\ &\quad \times \chi_{\lambda} \left(v^{-5} a_8, v^{-3} a_8, \dots, v^3 a_8, v^5 a_8, v^{-5} a_8^{-1}, v^{-3} a_8^{-1}, \dots, v^3 a_8^{-1}, v^5 a_8^{-1} \right). \end{aligned} \quad (5.56)$$

Here $\lambda = (\lambda_1, \dots, \lambda_{11}, 0)$ and

$$\begin{aligned} \hat{\mathcal{K}}_3(b) &= \prod_{\ell=1}^6 \text{PE} \left[\frac{2v^{2\ell} + v^{2\ell} b^{\pm 2}}{1 - v^2} \right], \\ \hat{\mathcal{K}}_2(b_1, b_2) &= \prod_{\ell=1}^4 \prod_{i,j=1}^3 \text{PE} \left[\frac{v^{2\ell} b_i / b_j}{1 - v^2} \right], \\ \hat{\mathcal{K}}_1(c_1, c_2, c_3, c_4, c_5, x) &= \text{PE} \left[\frac{2v^2 + v^2 x^{\pm 2}}{1 - v^2} \right] \left[\prod_{\ell=1}^2 \prod_{i,j=1}^5 \text{PE} \left[\frac{v^{2\ell} c_i / c_j}{1 - v^2} \right] \right] \\ &\quad \times \prod_{i=1}^5 \text{PE} \left[\frac{v^3 (x^{-1} c_i / c_6)^{\pm 1} + v^3 (x c_i / c_6)^{\pm 1}}{1 - v^2} \right], \end{aligned} \quad (5.57)$$

where $b_3 \equiv \frac{1}{b_1 b_2}$ and $c_6 \equiv \frac{1}{c_1 c_2 c_3 c_4 c_5}$. At last, one needs to replace $v \rightarrow q^{1/2}$. As the leading terms up to $q^{3/2}$ are contributed from rank one theory, the Schur index is given by

$$\mathcal{I}_{\tilde{H}_{E_8}^{(2)}}^{\text{Schur}} = 1 + \chi_2 q^{1/2} + (2\chi_3 + \mathbf{248})q + (2\chi_4 + 2(\mathbf{248} + 1)\chi_2)q^{3/2} + \dots \quad (5.58)$$

After decoupling the free hypermultiplet, the Schur index of $H_{E_8}^{(2)}$ theory is

$$\mathcal{I}_{H_{E_8}^{(2)}}^{\text{Schur}} = \mathcal{I}_{\tilde{H}_{E_7}^{(2)}}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} = 1 + (\chi_3 + \mathbf{248})q + \mathbf{248}\chi_2 q^{3/2} + \dots \quad (5.59)$$

Let us denote the two-string elliptic genus as

$$\mathbb{E}_{h_{E_8}^{(2)}}(v, x = 1, \tau, m_i = 0) = v^{59} Q_{\tau}^{-59/6} \sum_{i,j=0}^{\infty} c_{i,j}^{E_8} v^j (Q_{\tau} v^{-4})^i. \quad (5.60)$$

Then we have table 19 for the coefficients $c_{i,j}^{E_8}$. Here the red numbers are from the $L_{E_8}^{(2)}$ series. Add the red numbers in each column together, we expect to obtain a series that is

i, j	0	1	2	3	4	5	6	7
0	1	0	251	496	31625	116248	2747875	13624000
1	0	0	0	0	252	498	63503	241742

Table 19. Series coefficients $c_{i,j}^{E_8}$ for the unrefined elliptic genus of two E_8 instanton strings.

equal to the Schur index of rank two H_{E_8} 4d SCFT. Thus, we predict from the general formula (5.26) for the unrefined Schur index as

$$1 + 251q + 496q^{3/2} + 31877q^2 + 116746q^{5/2} + 2811378q^3 + 13865742q^{7/2} + \mathcal{O}(q^4). \quad (5.61)$$

Indeed, taking $x = 1$ in (5.59), the unrefined Schur index is given by

$$1 + 251q + 496q^{3/2} + \mathcal{O}(q^2). \quad (5.62)$$

Indeed, the two series match perfectly!

5.5 Rank three and higher

We expect the Del Zotto-Lockhart’s conjecture can be generalized to rank three and higher. From the universal leading expansion for three-string elliptic genus (4.11) and (4.12), we are able to predict the Schur index of rank three H_G SCFT up to order q^3 :

$$\begin{aligned} \mathcal{I}_{H_G^{(3)}}^{\text{Schur}} &= 1 + (\chi_3 + \chi_\theta)q + (\chi_4 + \chi_\theta\chi_2)q^{3/2} + \left(\chi_5 + (\chi_\theta + 1)\chi_3 + \chi_{\text{Sym}^2\theta} + \chi_\theta + 2\right)q^2 \\ &\quad + \left(\chi_6 + (2\chi_\theta + 2)\chi_4 + 2\chi_{\text{Sym}^2\theta} + \chi_\theta + 1\right)q^{5/2} \\ &\quad + \left(2\chi_7 + (3\chi_\theta + 1)\chi_5 + (\chi_{2\theta} + 3\chi_{\text{Sym}^2\theta} + 3\chi_\theta + 5)\chi_3 + \chi_{\text{Sym}^3\theta} + 3\chi_{\text{Sym}^2\theta} + \chi_\theta + 2\right)q^3 \\ &\quad + \mathcal{O}(q^{7/2}). \end{aligned} \quad (5.63)$$

This is actually because (4.11) and (4.12) are also the definition of leading and subleading Q_τ order of $L_G^{(3)}$ functions. Besides, we observe in the coefficient matrix of reduced three string elliptic genus, other than the $L_G^{(3)}$ function that appears as expected, the blue series also appears as in the rank two. The difference is that here the blue series is generated from the leading Q_τ order of *two* string elliptic genus!

$$M_G^{(3),\text{blue}}(v, x) = \frac{1}{(1 - vx)(1 - v/x)} g_{0,G}^{(2)}(v, x). \quad (5.64)$$

Note $g_{0,G}^{(2)}(v, x)$ is also the leading Q_τ order of $L_G^{(2)}$. In the following, we show the relation between reduced elliptic genus of three strings and the Schur index of $H_G^{(3)}$ theories for each G .

SU(3). The formula for the elliptic genus of three SU(3) string has been written down via Jeffrey-Kirwan residues in [34], using which we computed $\mathbb{E}_{h_{A_2}^{(3)}}$ up to Q_τ^6 order. Denote

$$\mathbb{E}_{h_{A_2}^{(3)}}(v, x, \tau, m_i = 0) = v^8 Q_\tau^{-4/3} \sum_{i,j=0}^{\infty} c_{i,j}^{\text{SU}(3)}(x) v^j (Q_\tau v^{-4})^i. \quad (5.65)$$

i, j	0	1	2	3	4	5	6	7
0	1	0	11	20	90	218	698	1618
1	1	2	14	22	135+12	370+22	960+171	2250+502

Table 20. Coefficients $c_{i,j}^{\text{SU}(3)}$ for the unrefined elliptic genus of three SU(3) instanton strings.

Then the unrefined $L_G^{(3)}$ function is shown red in the coefficient matrix of $\mathbb{E}_{h_{A_2}^{(3)}}$ in table 20. Note the red numbers are in agreement with our universal expansion (4.11) and (4.12), while the blue numbers are in agreement with our proposal (5.64).

The construction for rank three H_{A_2} theory from 6d involves certain irregular punctures as the rank two case. We are not aware how to compute its indices directly. We write down our prediction for the Schur index of rank three H_{A_2} theory here:

$$\begin{aligned} \mathcal{I}_{H_{A_2}^{(3)}}^{\text{Schur}} = & 1 + (\chi_3 + 8)q + (\chi_4 + 8\chi_2)q^{3/2} + (\chi_5 + 17\chi_3 + 46)q^2 + (\chi_6 + 18\chi_4 + 81\chi_2)q^{5/2} \\ & + (2\chi_7 + 25\chi_5 + 164\chi_3 + 248)q^3 + (\chi_8 + 27\chi_6 + 209\chi_4 + 557\chi_2)q^{7/2} + \mathcal{O}(q^4). \end{aligned} \quad (5.66)$$

The unrefined limit is

$$\mathcal{I}_{H_{A_2}^{(3)}}^{\text{Schur}}(x=1) = 1 + 11q + 20q^{3/2} + 102q^2 + 240q^{5/2} + 869q^3 + 2120q^{7/2} + \mathcal{O}(q^4). \quad (5.67)$$

SO(8). We can use class \mathcal{S} theory technique to compute the HL and Schur index of rank three H_{D_4} 4d SCFT. The $H_{D_4}^{(3)}$ theory can be constructed by compactifying A_5 (2, 0) 6d SCFT on a sphere with four $\{3^2\}$ punctures, which is a “bad” theory. We need instead to consider $\tilde{H}_{D_4}^{(3)}$ theory obtained from three $\{3^2\}$ punctures and one $\{3, 2, 1\}$ puncture. We compute the Schur index as

$$\begin{aligned} & \mathcal{I}_{\tilde{H}_{D_4}^{(3)}}^{\text{Schur}}(c_1, c_2, c_3; x, b) \\ & = \hat{\mathcal{N}}'_6 \hat{\mathcal{K}}_1(c) \hat{\mathcal{K}}_1(d) \hat{\mathcal{K}}_1(e) \hat{\mathcal{K}}_2(x, b) \sum_{\lambda} \frac{\chi_{\lambda}(v^2 b, b, v^{-2} b, v b^{-1} x, v^{-1} b^{-1} x, b^{-1} x^{-2})}{\chi_{\lambda}^2(v^{-5}, v^{-3}, v^{-1}, v, v^3, v^5)} \\ & \quad \times \prod_{i=1,2,3} \chi_{\lambda}(v^2 c_i, c_i, v^{-2} c_i, v^2 c_i^{-1}, c_i^{-1}, v^{-2} c_i^{-1}), \end{aligned} \quad (5.68)$$

with $(b_1 = b, b_2 = 1/b)$

$$\begin{aligned} \hat{\mathcal{N}}'_6 & = (v^2; v^2)^4 \prod_{j=2}^6 (v^{2j}; v^2)^2, \\ \hat{\mathcal{K}}_1(b) & = \text{PE} \left[\frac{(v^2 + v^4 + v^6)(b^2 + b^{-2} + 2)}{1 - v^2} \right], \\ \hat{\mathcal{K}}_2(a, b) & = \text{PE} \left[\frac{3v^2 + 2v^4 + v^6 + (v^3 + v^5)(b^2 a^{-1})^{\pm 1} + v^3 a^{\pm 3} + v^4 (b^2 a^2)^{\pm 1}}{1 - v^2} \right]. \end{aligned} \quad (5.69)$$

i, j	0	1	2	3	4	5	6
0	1	0	31	60	580	1858	9457
1	0	0	0	0	32	62	1111

Table 21. Expected coefficients $c_{i,j}^{\text{SO}(8)}$ for the unrefined elliptic genus of three SO(8) instanton strings.

From the above formula, we compute the Schur index up to $q^{11/2}$. After decoupling the free hypermultiplet, we obtain

$$\begin{aligned}
 \mathcal{I}_{H_{D_4}^{(3)}}^{\text{Schur}} &= \mathcal{I}_{\tilde{H}_{D_4}^{(3)}}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} = 1 + (\chi_3 + 28)q + (\chi_4 + 28\chi_2)q^{3/2} + (\chi_5 + 57\chi_3 + 436)q^2 \\
 &\quad + (\chi_6 + 58\chi_4 + 841\chi_2)q^{5/2} + (2\chi_7 + 85\chi_5 + 1607\chi_3 + 5308)q^3 \\
 &\quad + (\chi_8 + 87\chi_6 + 2042\chi_4 + 14135\chi_2)q^{7/2} + (2\chi_9 + 115\chi_7 + 2806\chi_5 + 29042\chi_3 + 55871)q^4 \\
 &\quad + (2\chi_{10} + 115\chi_8 + 3242\chi_6 + 43166\chi_4 + 177896\chi_2)q^{9/2} \\
 &\quad + (2\chi_{11} + 144\chi_9 + 4008\chi_7 + 60673\chi_5 + 392233\chi_3 + 527217)q^5 \\
 &\quad + (2\chi_{12} + 145\chi_{10} + 4441\chi_8 + 75128\chi_6 + 649112\chi_4 + 1857119\chi_2)q^{11/2} + \mathcal{O}(q^6). \quad (5.70)
 \end{aligned}$$

The unrefined limit is

$$\begin{aligned}
 \mathcal{I}_{H_{D_4}^{(3)}}^{\text{Schur}}(x=1) &= 1 + 31q + 60q^{3/2} + 612q^2 + 1920q^{5/2} + 10568q^3 + 36968q^{7/2} + 157850q^4 \\
 &\quad + 548848q^{9/2} + 2036655q^5 + 6798456q^{11/2} + \mathcal{O}(q^6). \quad (5.71)
 \end{aligned}$$

Let us denote the reduced three-string elliptic genus as

$$\mathbb{E}_{h_{D_4}^{(3)}}(v, x, \tau, m_i = 0) = v^{17} Q_\tau^{-17/6} \sum_{i,j=0}^{\infty} c_{i,j}^{\text{SO}(8)} v^j (Q_\tau v^{-4})^i. \quad (5.72)$$

Then from (4.11) and (4.12), we expect to have table 21 for the unrefined coefficients $c_{i,j}^{\text{SO}(8)}$. Here the red numbers are from the $L_{D_4}^{(3)}$ series. Add the red numbers in each column together, we expect to obtain a series that is equal to the Schur index of rank three H_{D_4} 4d SCFT. Indeed, we have

$$L_{D_4}^{(3)}(q^{1/2}, x=1, Q_m=1, q^2) = 1 + 31q + 60q^{3/2} + 612q^2 + 1920q^{5/2} + 10568q^3 + \dots \quad (5.73)$$

One can see the two series match perfectly up to q^3 !

E₆. The formula to compute the Hall-Littlewood index of rank three H_{E_6} SCFT has been written down in [45]. Similarly, we compute the Schur index as

$$\begin{aligned}
 \mathcal{I}_{\tilde{H}_{E_6}^{(3)}}^{\text{Schur}} &= \hat{\mathcal{N}}_9 \hat{\mathcal{K}}_1(a_1, a_2) \hat{\mathcal{K}}_1(a_3, a_4) \hat{\mathcal{K}}_2(a_5, a_6, x) \times \\
 &\quad \sum_{\lambda} \frac{\chi_{\lambda} \left(v^2 a_5, v^{-2} a_5, a_5, v^2 a_6, v^{-2} a_6, a_6, v \frac{x}{a_5 a_6}, v^{-1} \frac{x}{a_5 a_6}, \frac{x^{-2}}{a_5 a_6} \right)}{\chi_{\lambda}(v^{-8}, v^{-6}, v^{-4}, v^{-2}, 1, v^2, v^4, v^6, v^8)} \times \\
 &\quad \times \chi_{\lambda} \left(v^2 a_1, v^{-2} a_1, a_1, v^2 a_2, v^{-2} a_2, a_2, v^2 \frac{1}{a_1 a_2}, v^{-2} \frac{1}{a_1 a_2}, \frac{1}{a_1 a_2} \right) \\
 &\quad \times \chi_{\lambda} \left(v^2 a_3, v^{-2} a_3, a_3, v^2 a_4, v^{-2} a_4, a_4, v^2 \frac{1}{a_3 a_4}, v^{-2} \frac{1}{a_3 a_4}, \frac{1}{a_3 a_4} \right).
 \end{aligned}$$

i, j	0	1	2	3	4	5	6
0	1	0	81	160	3555	12958	121447
1	0	0	0	0	82	162	6961

Table 22. Coefficients $c_{i,j}^{E_6}$ for the unrefined elliptic genus of three E_6 instanton strings.

with

$$\hat{\mathcal{K}}_1(b_1, b_2) = \text{PE} \left[\sum_{i,j=1}^3 \frac{(v^2 + v^4 + v^6)b_i/b_j}{1 - v^2} \right],$$

$$\hat{\mathcal{K}}_2(b_1, b_2, x) = \text{PE} \left[\frac{1}{1 - v^2} \left((v^2 + v^4 + v^6) \left(\sum_{i,j=1}^2 b_i/b_j \right) + 2v^2 + v^4 + v^3x^{\pm 3} \right. \right. \\ \left. \left. + (b_1 + b_2)((v^3 + v^5)(b_3x)^{\pm 1} + v^4(b_3x^{-2})^{\pm 1}) \right) \right],$$

where $b_1b_2b_3 = 1$. From the above formula, we computed the Schur index up to q^2 . After decoupling the free hypermultiplet, we obtain

$$\mathcal{I}_{H_{E_6}^{(3)}}^{\text{Schur}} = \mathcal{I}_{\tilde{H}_{E_6}^{(3)}}^{\text{Schur}} / \mathcal{I}_{h.m.}^{\text{Schur}} = 1 + (\chi_3 + 78)q + (\chi_4 + 78\chi_2)q^{3/2} + (\chi_5 + 157\chi_3 + 3161)q^2 \\ + (\chi_6 + 158\chi_4 + 6241\chi_2)q^{5/2} + (2\chi_7 + 235\chi_5 + 11912\chi_3 + 91483)q^3 \\ + (\chi_8 + 237\chi_6 + 15072\chi_4 + 260821\chi_2)q^{7/2} + \mathcal{O}(q^4). \quad (5.74)$$

The unrefined limit is

$$\mathcal{I}_{H_{E_6}^{(3)}}^{\text{Schur}}(x = 1) = 1 + 81q + 160q^{3/2} + 3637q^2 + 13120q^{5/2} + 128408q^3 + 583360q^{7/2} + \mathcal{O}(q^4). \quad (5.75)$$

On the other hand, the universal leading expansion (4.11) and (4.12) indicate the following table 22 for the coefficients of reduced three-string elliptic genus for E_6 . By adding the red numbers in each column together, one can indeed obtain the same unrefined Schur series as (5.75) up to q^3 .

F_4, E_7, E_8 . The Schur indices with generic $SU(2)_x$ fugacity for rank three H_G theories can be predicted from (5.63) up to q^3 order. Let us just mark the unrefined series here:

$$\mathcal{I}_{H_{F_4}^{(3)}}^{\text{Schur}} = 1 + 55q + 108q^{3/2} + 1752q^2 + 6048q^{5/2} + 45835q^3 + \mathcal{O}(q^{7/2}),$$

$$\mathcal{I}_{H_{E_7}^{(3)}}^{\text{Schur}} = 1 + 136q + 270q^{3/2} + 9852q^2 + 36990q^{5/2} + 533401q^3 + \mathcal{O}(q^{7/2}), \quad (5.76)$$

$$\mathcal{I}_{H_{E_8}^{(3)}}^{\text{Schur}} = 1 + 251q + 500q^{3/2} + 32622q^2 + 126000q^{5/2} + 3030748q^3 + \mathcal{O}(q^{7/2}).$$

Note for F_4 , we always mean the analogy for H_G theories.

In summary, we arrive at the final conjecture for arbitrary rank:

Conjecture. There exists an infinite series of functions $L_G^{(n)}(v, x, m_G, Q_\tau) = \sum_{i,j=0}^{\infty} b_{i,j}^{G,n} Q_\tau^i v^j$, $n = 1, 2, \dots$ such that

1. $b_{i,j}^{G,n}$ can be written as the sum of products between the characters of irreducible representations of $SU(2)_x$ and the characters of irreducible representations of G with integral coefficients.
2. $L_G^{(n)}(v, x, m_G, 0)$ is the Hilbert series of the reduced moduli space of n G -instantons, i.e. the Hall-Littlewood index of the $H_G^{(n)}$ theory.
3. $L_G^{(n)}(q^{1/2}, x, m_G, q^2)$ is the Schur index of the $H_G^{(n)}$ theory.
4. The n -string elliptic genus $\mathbb{E}_{h_G^{(n)}}(v, x, m_G, Q_\tau)$ can be generated from the first n L_G functions, i.e. $L_G^{(r)}(v, x, m_G, Q_\tau)$, $r = 1, 2, \dots, n$.

6 Conclusion and outlook

In this paper we study the elliptic blowup equations for minimal 6d (1,0) SCFTs with all six possible gauge groups $G = SU(3), SO(8), F_4, E_{6,7,8}$. The study is twofold, topological string partition function on elliptic non-compact Calabi-Yau and elliptic genera for 6d (1,0) SCFTs. From the viewpoint of Calabi-Yau, we use the geometric construction in [8] and the generalized blowup equations in [15] to solve the refined BPS invariants to high base degrees, which in turn serve as numerous nontrivial checks for the blowup equations, both unity and vanishing ones. From the viewpoint of 6d SCFTs, we use the de-affinisation procedure to derive some elegant functional equations for the elliptic genera, from which we obtain an exact and universal recursion formula for the elliptic genera of arbitrary number of strings and arbitrary gauge group. In particular, we explicitly compute the one and two-string elliptic genera for all G , which recover all previous partial results from refined topological string, modular bootstrap, Hilbert series, 2d quiver gauge theories and the β -twisted partition function of $\mathcal{N} = 2$ superconformal H_G theories. We also prove the modularity of the elliptic blowup equations which is a strong support that they hold for arbitrary number of strings.

The elliptic genera we solve from blowup equations could be useful in many aspects. For example, they would help to identify the 2d quiver description of the 6d minimal SCFT with exceptional gauge symmetry, see some attempts for $G = E_7$ in [87]. They also serve as the calibration to determine modular ansatz for higher-string elliptic genus and the web of topological vertex for the associated non-toric Calabi-Yau threefolds [40]. The elliptic genera of 2d (0,4) SCFTs we studied also play a role in the context of certain compact elliptic Calabi-Yau threefolds [88, 89]. For example, the $SO(8)$ and E_6 minimal SCFT serve as the constituents of the 2d quiver gauge theories associated to the $T^6/\mathbb{Z}_2 \times \mathbb{Z}_2$ and $T^6/\mathbb{Z}_3 \times \mathbb{Z}_3$ geometries respectively, and their elliptic genera are useful to compute the degeneracies of 5d spinning BPS black holes in the dual gravity picture, as suggested in [89]. We hope our exact formulas of the elliptic genera for exceptional minimal SCFTs would contribute to this subject.

It is also interesting to investigate the K-theoretic blowup equations for all possible 5d SYM theories. The K-theoretic blowup equations are quite different from the elliptic ones in that they exist for all simple Lie groups. Part of the unity K-theoretic blowup equations were already conjectured by Nakajima-Yoshioka [28] and explicitly checked by Keller-Song [30]. One may suggest to use dimensional reduction i.e. $Q_\tau \rightarrow 0$ to obtain the K-theoretic blowup equations for the six gauge groups G . However, intriguingly we find that the 5d reduction of elliptic blowup equations does *not* produce all non-equivalent K-theoretic blowup equations for these six gauge groups. We leave these issues and the complete set of K-theoretic blowup equations for all simple Lie group to future work.

The elliptic blowup equations for 6d (1,0) minimal SCFTs allow us to further study more complicated examples. Some immediate models are the non-Higgsable clusters with matters which we will investigate in a subsequent paper [51]. There are four of them: one belongs to minimal (1,0) SCFT with $\mathfrak{n} = 7$, and the other three have more than one dimensional tensor branch. See some primary results on the elliptic genera for such theories in [8, 38, 87]. One can also use blowup equations to study elliptic Calabi-Yau with multi-sections. See some discussion on such geometries in [90, 91]. Our final goal is to find an exact, explicit and universal formula for the elliptic genera of all 6d (1,0) SCFTs in the atomic classification [6], see also a good review on the classification in [92]. It is known there are two approaches to classifying 6d (1,0) SCFTs: top down and bottom up. We expect blowup equations make sense in both settings. In the top down approach, given the explicit description of a elliptic non-compact Calabi-Yau, i.e. base and elliptic fibration, one should be able to use the generalized blowup equations in [15] to solve the partition function of refined topological strings. On the other hand, in the bottom up approach, given the explicit content of 6d multiplets which satisfy the anomaly cancellations, one should also be able to directly write down the elliptic blowup equations for the elliptic genera of such 6d SCFT, as a generalization of the current paper. The two pictures are related by geometric engineering, as the two formalism of blowup equations are related by de-affinisation.

The K-theoretic Nekrasov partition function inspired the study on K-theoretic invariants for general 4-manifolds, specially complex surfaces [17]. Since the elliptic genus of 6d (1,0) SCFTs in 5d limit gives K-theoretic Nekrasov partition function, naturally one wonders if elliptic genus can be used to construct some elliptic version of 4-manifold invariants, such as Donaldson invariants. Besides, the Nekrasov partition function is known to relate to W-algebras. In 4d, the equality between the universal one-instanton Nekrasov partition function and the norm of Gaiotto-Whittaker vectors in W-algebra has been checked in [59]. See proof in [93]. In 5d, the relation between K-theoretic $SU(N)$ Nekrasov partition function and q -deformed W_N -algebra was also studied in [94]. It seems natural to extend such relation to 6d where the elliptic genus should be related to the elliptic W-algebras. The elliptic W-algebra associated to general Lie algebras is very difficult to study. We hope our exact formula on the elliptic genus could shed some new light. For example, it would be nice to see if one can use elliptic W-algebra to make comparison with our universal one-string elliptic genus formula (3.33) like those comparison done in [59]. One can even ask whether the structure of blowup equations itself can find some origin in pure algebras. See a possible direction [95].

One major remaining question is of course how to prove the elliptic blowup equations, or more general, the blowup equations for all local Calabi-Yau in [15, 23, 29]. As the refined BPS invariants for non-compact Calabi-Yau threefolds have been rigorously defined via refined stable pairs in [11], these functional equations for the partition functions are indeed well-formulated mathematical conjectures. See also the definition of refined invariants in [12]. The proof of Göttsche-Nakajima-Yoshioka K-theoretic blowup equations [18] relies deeply on the structure of gauge theories, which may not be exactly suitable for Calabi-Yau setting, as the latter does not necessarily engineer a gauge theory. As emphasized before, the formalism of generalized blowup equations is not sensitive to additional structures of non-compact Calabi-Yau threefolds, be they toric or elliptic. Let us also point out there is even no physical proof for the generalized blowup equations in [15, 23, 29]. In particular, it would be good to see if one can connect the blowup equations with refined holomorphic anomaly equations. Specializing to elliptic blowup equations studied in this paper, we suspect by using the Kac-Weyl character formulas and following the 4d derivation in [59], one may be able to derive the universal \mathbb{E}_1 formulas (3.33) and the identities from the leading degree of vanishing blowup equations (3.48). We leave these for future studies.

Another major question is how to explain the surprising relation between the elliptic genera of 6d (1,0) SCFT and the Schur indices of 4d $\mathcal{N} = 2$ H_G theories. Despite the striking relation for rank two cases and even some rank three cases shown in this paper, we do not find the exact formulas connecting the two and three-string elliptic genera and those L_G functions like (5.17) in rank one cases found in [37]. To obtain such fascinating formulas for arbitrary rank, it seems one has to answer some questions first. For example, what is the physical meaning for the $L_G^{(k)}$ functions?⁴² How to interpret and make use of those nonperturbative symmetries (4.27)? In 4d SCFT H_G , precisely how should the β -twisted partition function on $T^2 \times S^2$ be related to the superconformal indices on $S^3 \times S^1$? And how should the SCFT/VOA correspondence be put in this picture? One possible direction is to look into the localization on the 4d backgrounds following the recent works [75–79].

⁴²Naively one may tempt to identify L_G functions as Macdonald indices, since they both have two parameters, and both serve as an unification of Hall-Littlewood indices and Schur indices. However, this seems not true. For example, the Macdonald index of rank one H_{E_7} can be easily obtained by taking limit in the full superconformal index in [48] as

$$\begin{aligned} \mathcal{I}_{H_{E_7}^{(1)}}^M(q, t) &= 1 + 133t + (134tq + 7371t^2) + (134tq^2 + 16149t^2q + 238602t^3) \\ &\quad + (134tq^3 + 25193t^2q^2 + 819413t^3q + 5248750t^4) + \dots \end{aligned} \tag{6.1}$$

While the $L_{E_7}^{(1)}$ function is determined in [37] as

$$\begin{aligned} L_{E_7}^{(1)}(Q_\tau, v) &= 1 + 133v^2 + 7371v^4 + 238602v^6 + 5248750v^8 + \dots \\ &\quad + Q_\tau(134 + 16283v^2 + 835562v^4 + \dots) + Q_\tau^2(31373 + \dots) + \dots \end{aligned} \tag{6.2}$$

One can see they are indeed not the same, albeit $\mathcal{I}_{H_{E_7}^{(1)}}^M(0, v^2) = L_{E_7}^{(1)}(0, v)$ and $\mathcal{I}_{H_{E_7}^{(1)}}^M(q, q) = L_{E_7}^{(1)}(q^2, q^{1/2})$.

Acknowledgments

We thank Babak Haghighat for the early participation of this project. We also thank Christopher Beem, Giulio Bonelli, Michele Del Zotto, Lothar Göttsche, Min-xin Huang, Guglielmo Lockhart, Joonho Kim, Seok Kim, Hiraku Nakajima, Leonardo Rastelli, Alessandro Tanzini, Wenbin Yan, Don Zagier and Rui-Dong Zhu for useful discussion. JG is supported by the Fonds National Suisse, subsidiary 200020-175539 (project “Quantum mechanics, geometry and strings”).

A Lie algebraic convention

We collect some definitions in (affine) Lie algebras and fix our convention used throughout the paper. Given a simple Lie algebra \mathfrak{g} of rank $\text{rk}(G)$, there are four $\text{rk}(G)$ -dimensional lattices of importance, the root and coroot lattices Q, Q^\vee , the weight and coweight lattices P, P^\vee . They satisfy

$$Q^\vee \subset P \subset \mathfrak{h}_\mathbb{C}, \tag{A.1}$$

$$Q \subset P^\vee \subset \mathfrak{h}_\mathbb{C}^*. \tag{A.2}$$

Here $\mathfrak{h}_\mathbb{C}, \mathfrak{h}_\mathbb{C}^* \cong \mathbb{C}^{\text{rk}}$ are the complexified Cartan subalgebra and its dual. They are isomorphic to each other via the natural inner product

$$\langle \bullet, \bullet \rangle : \mathfrak{h}_\mathbb{C}^* \times \mathfrak{h}_\mathbb{C} \rightarrow \mathbb{C}. \tag{A.3}$$

The Cartan matrix is then defined by

$$A_{ij} = \langle \alpha_i, \alpha_j^\vee \rangle, \tag{A.4}$$

where α_i are simple roots. Consider the invariant bilinear form (\bullet, \bullet) on the coroot lattice Q^\vee normalized so that the norm square of the shortest coroot θ^\vee is two. It can be generalized to a bilinear form on $\mathfrak{h}_\mathbb{C}$ in which Q^\vee is embedded. By the isomorphism between $\mathfrak{h}_\mathbb{C}$ and $\mathfrak{h}_\mathbb{C}^*$, it induces also an invariant bilinear form with the same notation on the latter vector space. With our normalization, the bilinear form satisfies

$$(k, k) = \frac{1}{2h_g^\vee} \sum_{\alpha \in \Delta} \langle \alpha, k \rangle^2, \quad k \in \mathfrak{h}, \tag{A.5}$$

where h_g^\vee is the dual Coxeter number, and Δ the set of all roots.

We also define the fundamental weights $\omega_i \in P$ and fundamental coweights $\omega_i^\vee \in P^\vee$ ($i = 1, \dots, \text{rk}$) through

$$\langle \alpha_i, \omega_j \rangle = \langle \omega_i^\vee, \alpha_j^\vee \rangle = \delta_{ij}. \tag{A.6}$$

They are related to roots and coroots by

$$\alpha_i = \sum_{j=1}^{\text{rk}} A_{ij} \omega_j^\vee, \quad \alpha_i^\vee = \sum_{j=1}^{\text{rk}} \omega_j A_{ji}. \tag{A.7}$$

Most of these definitions can be generalized to the affine Lie algebra $\hat{\mathfrak{g}}$. We add an additional simple root α_0 satisfying

$$\alpha_0 = \alpha_0^\vee, \quad (\alpha_0, \alpha_0) = (\alpha_0^\vee, \alpha_0^\vee) = 2. \quad (\text{A.8})$$

The affine Cartan matrix is defined to be

$$\hat{A}_{IJ} = \langle \alpha_I, \alpha_J^\vee \rangle, \quad I, J = 0, 1, \dots, \text{rk}, \quad (\text{A.9})$$

where a_I, a_J^\vee are the marks and the comarks of the affine Lie algebra $\hat{\mathfrak{g}}$ respectively. The affine Cartan matrix satisfies

$$\sum_{I=0}^{\text{rk}} a_I \hat{A}_{IJ} = \sum_{J=0}^{\text{rk}} \hat{A}_{IJ} a_J^\vee = 0. \quad (\text{A.10})$$

Note that α_0 can be written in terms of the longest root θ and the imaginary root δ , which annihilates anything in \mathfrak{h} or \mathfrak{h}^* and has a vanishing norm square, by

$$\alpha_0 = \delta - \theta. \quad (\text{A.11})$$

Similarly we can also define the fundamental weights $\hat{\omega}_I$ and coweights $\hat{\omega}_I^\vee$ in the affine Lie algebra $\hat{\mathfrak{g}}$ by

$$\langle \alpha_I, \hat{\omega}_J \rangle = \langle \hat{\omega}_I^\vee, \alpha_J^\vee \rangle = \delta_{IJ}, \quad I, J = 0, \dots, \text{rk}. \quad (\text{A.12})$$

For $i = 1, \dots, \text{rk}$ the fundamental (co-)weights in $\hat{\mathfrak{g}}$ are related those in \mathfrak{g} by

$$\hat{\omega}_i = \omega_i + a_i \hat{\omega}_0, \quad (\text{A.13})$$

$$\hat{\omega}_i^\vee = \omega_i^\vee + a_i^\vee \hat{\omega}_0^\vee, \quad (\text{A.14})$$

while $\hat{\omega}_0 = \hat{\omega}_0^\vee$ is imaginary, and it satisfies

$$\langle \alpha_i, \hat{\omega}_0 \rangle = \langle \hat{\omega}_0, \alpha_i^\vee \rangle = 0, \quad (\delta, \hat{\omega}_0) = 1. \quad (\text{A.15})$$

Using these relations together with (A.7), we find the affine version of (A.7)

$$\alpha_I = \sum_{J=0}^{\text{rk}} \hat{A}_{IJ} \hat{\omega}_J^\vee + \delta \cdot \delta_{I,0}, \quad \alpha_I^\vee = \sum_{J=0}^{\text{rk}} \hat{\omega}_J \hat{A}_{JI} + \delta \cdot \delta_{I,0}. \quad (\text{A.16})$$

B Mirror symmetry for elliptic non-compact Calabi-Yau three-folds

The prescription in section 2.2 can determine all the triple intersection numbers in the non-compact Calabi-Yau X associated to a minimal 6d SCFT except for the number $\kappa_{\tau\tau\tau}$. Given the non-compactness of X we do not expect all the triple intersection numbers to be computable, and the number $\kappa_{\tau\tau\tau}$ is irrelevant for the blowup equations in any case. Nevertheless we propose here a reasonable normalisation scheme for $\kappa_{\tau\tau\tau}$, which involves a local version of mirror symmetry. We use this normalisation scheme in the example section 4.

For the compact Calabi-Yau \hat{X} where X is embedded, one can define for every toric charge $l^{(i)}$ a Picard-Fuchs operator $\hat{\mathcal{L}}_i$ which annihilates the homogeneous periods $\hat{\omega}_0, \hat{\Pi}_i^{(1)}, \hat{\Pi}_i^{(2)}$. In the decompactification limit $z_{de} \rightarrow 0$, the Picard-Fuchs (PF) operator associated to the (0)-curve in the base vanishes, while the other operators \mathcal{L}_I remain well-defined and non-trivial, and they annihilate all the finite homogeneous periods⁴³ $\omega_0, \omega_0 t_I, \omega_0 F_J$ of the resulting local Calabi-Yau. These operators, however, do not form a PF complete system, in the sense that they have extra independent solutions. To cure this problem, we define in addition the PF operator \mathcal{L}_τ from the toric charge of the elliptic fiber $\tau : (-6, 2, 3, 0, \dots, 1, \dots)$.⁴⁴ It annihilates all the finite homogeneous periods, but not the other superfluous solutions, thus making the PF system complete.

The number $\kappa_{\tau\tau\tau}$ is contained in the homogeneous B-period $\omega_0 F_\tau = \omega_0 \partial_\tau F_{(0,0)}$ which corresponds to the zero section. We find that it is completely fixed by the normalisation condition

$$\mathcal{L}_\tau(\omega_0 \partial_\tau F_{(0,0)}) = 0. \tag{B.1}$$

Note that the resulting homogeneous B-period is not a solution to the complete PF system. When it is acted upon by the other PF operators it does not vanish but produces ω_0 up to a scaling factor, which may have some open string interpretation.

There are several ways to understand this normalisation scheme. Once all the triple intersection numbers are known, the normalised Euler characteristic can be computed by⁴⁵

$$\chi = \int_X c_3(X) = \frac{1}{3} \sum_{ijk} \kappa_{ijk} l_n^{(i)} l_n^{(j)} l_n^{(k)}, \tag{B.2}$$

we list the results of all the minimal 6CFTs except for the cases of $\mathbf{n} = 3, 7$ in table 23. Note that the calculation of the normalised Euler characteristic for the case $\mathbf{n} > 3$ is different from that for the first three cases. For $\mathbf{n} > 2$ the elliptic singularity is constant over the $-\mathbf{n}$ curve in the base. Except for the $\mathbf{n} = 3$ case, the resolved geometry can be described as configuration of Hirzebruch surfaces inside the compact CY-3-fold with only even Betti numbers and $\chi(X) = 1 + b_2 + b_4$. The case of $\mathbf{n} = 3, G = \text{SU}(3)$ is special as the reduction from the compact geometry to the non-compact geometry also involves a flop operation. The geometry of $\mathbf{n} = 2$ has more supersymmetry and odd Betti numbers $b_0 = b_4 = 1, b_1 = b_2 = b_3 = 2$. In the case $\mathbf{n} = 1$ we normalise the Euler characteristic using the formula for the compact threefold [96] with E_8 elliptic fibre type $\chi(X_{\text{comp}}) = -2 \cdot 30 \times \int_B c_1^2(M)$. The effect of blowing up a $C^2 = -\mathbf{n} = 1$ curve decreases $\int_B c_1^2(M)$ by one. Hence the contribution of the non-compact geometry should be $\chi(M) = 60$.

The Euler characteristics thus computed for the theories with a pure gauge bulk agree with the naive definition in terms of the numbers b_n of compact n -cycles $\chi = \sum_n (-1)^n b_n$.

⁴³For these elliptic Calabi-Yau threefolds decompactified in the horizontal direction, the fundamental period $\hat{\omega}_0$ does not become a constant but remains a non-trivial holomorphic function ω_0 .

⁴⁴The charge 1 corresponds to the zero section of the elliptic fibration.

⁴⁵Here it is understood that we omit the toric charge of the (0)-curve in the summation. If one wishes the charge entry associated to the pullback of the base curve (denoted by S in table 2) can also be ignored as it is only nonzero for the (0)-curve. See for instance table 2 for the model with $G = F_4$.

n	1	2	4	5	6	8	12
G	–	–	$SO(8)$	F_4	E_6	E_7	E_8
b_2	1		6	6	8	9	10
b_4	10		5	5	7	8	9
$\chi(X)$	60	0	12	12	16	18	20

Table 23. The normalised Euler characteristics of the non-compact elliptic Calabi-Yau threefolds associated to minimal 6d SCFTs.

Furthermore, by integrating the B-period $\partial_\tau F_{(0,0)}$ we can compute the genus 0 GW invariants in the τ direction, which should be the same as the Euler characteristics. We checked this for the $\mathbf{n} = 1, 2, 5, 6$ models.⁴⁶

In the following, we illustrate this idea with two examples.

n = 1.

D	ν_i^*				$l^{(1)}$	$l^{(2)}$	$l^{(3)}$
D_0	0	0	0	0	–6	0	0
D_1	–1	0	0	0	2	0	0
D_2	0	–1	0	0	3	0	0
S'	2	3	0	–1	0	–1	1
K	2	3	0	0	1	–1	–2
F	2	3	–1	–1	0	1	0
S	2	3	0	1	0	0	1
F	2	3	1	0	0	1	0

(B.3)

The Picard-Fuchs operators of the compact geometry are

$$\begin{aligned}
 \hat{\mathcal{L}}_1 &= \theta_1 (\theta_1 - \theta_2 - 2\theta_3) - 12z_1 (6\theta_1 + 1) (6\theta_1 + 5), \\
 \hat{\mathcal{L}}_2 &= \theta_2^2 - z_2 (\theta_1 - \theta_2 - 2\theta_3) (\theta_3 - \theta_2), \\
 \hat{\mathcal{L}}_3 &= \theta_3 (\theta_3 - \theta_2) - z_3 (\theta_1 - \theta_2 - 2\theta_3 - 1) (\theta_1 - \theta_2 - 2\theta_3),
 \end{aligned}
 \tag{B.4}$$

where $\theta_i := z_i \frac{\partial}{\partial z_i}$. Denote \hat{F}_0 the compact genus zero free energy, we have the periods:

$$\begin{aligned}
 \hat{X}_0 &= \hat{\omega}_0, \quad \hat{X}_1 = \hat{\omega}_0 t_1, \quad \hat{X}_2 = \hat{\omega}_0 t_2, \\
 \hat{X}_3 &= \left(\frac{\partial}{\partial \rho_2 \partial \rho_3} + \frac{1}{2} \frac{\partial}{\partial \rho_3^2} \right) \hat{\omega}_0(\rho)|_{\rho=0}, \\
 \hat{X}_4 &= \left(-\frac{1}{2} \frac{\partial}{\partial \rho_1^2} - \frac{\partial}{\partial \rho_1 \partial \rho_2} \right) \hat{\omega}_0(\rho)|_{\rho=0}, \\
 \hat{X}_5 &= \left(-\frac{\partial}{\partial \rho_1^2} - \frac{\partial}{\partial \rho_1 \partial \rho_3} \right) \hat{\omega}_0(\rho)|_{\rho=0},
 \end{aligned}
 \tag{B.5}$$

⁴⁶For the remaining models the first non-vanishing invariants appear at very high degree and we fail to obtain them within a reasonable period of time.

and another period with triple logarithmic singularity. Here $\hat{\omega}_0(\rho)$ is the deformed fundamental period

$$\hat{\omega}_0(\rho) = \sum_{\substack{n_1, n_2, n_3 \in \mathbb{Z}_{\geq 0} \\ x_i = n_i + \rho_i}} \frac{\Gamma(x_1+1)}{\Gamma(x_1+1)\Gamma(x_1+1)\Gamma(x_1-x_2-2x_3+1)\Gamma(-x_2+x_3+1)\Gamma(x_3+1)\Gamma(x_2+1)^2} z_1^{x_1} z_2^{x_2} z_3^{x_3}, \tag{B.6}$$

and $\hat{\omega}_0 = \hat{\omega}_0(0)$. The non-compact geometry is related to the compact geometry by setting $z_{dc} := z_3 \rightarrow 0$. Then the Picard-Fuchs operator $\hat{\mathcal{L}}_3$ vanishes while the other two become:

$$\begin{aligned} \mathcal{L}_1 &= \theta_1(\theta_1 - \theta_2) - 12z_1(6\theta_1 + 1)(6\theta_1 + 5), \\ \mathcal{L}_2 &= \theta_2^2 + z_2(\theta_1 - \theta_2)\theta_2. \end{aligned} \tag{B.7}$$

The deformed fundamental period $\hat{\omega}_0(\rho)$ becomes $\omega_0(\rho) = \hat{\omega}_0(\rho)|_{z_3 \rightarrow 0}$. There are one period $X_0 = \omega_0 = \hat{\omega}_0|_{z_3 \rightarrow 0}$ without singularity, two periods with logarithmic singularities

$$X_1 = \hat{X}_1|_{z_3 \rightarrow 0}, \quad X_2 = \hat{X}_2|_{z_3 \rightarrow 0}, \tag{B.8}$$

one with double logarithmic singularities

$$X_3 = \left(-\frac{1}{2} \frac{\partial}{\partial \rho_1^2} - \frac{\partial}{\partial \rho_1 \partial \rho_2} \right) \omega_0(\rho)|_{\rho=0} = \omega_0 \frac{\partial}{\partial t_2} F_0 = \hat{X}_3|_{z_3 \rightarrow 0}, \tag{B.9}$$

and no solution with triple logarithmic singularities.

The reason we cannot fix the τ terms is because we do not know the $\tau = t_1$ derivative of the free energy, however, there is an interesting ‘‘period’’ $X_4 = \omega_0 \frac{\partial}{\partial \tau} F_{(0,0)}$ which satisfies (up to all the orders we have checked)

$$\mathcal{L}_1 X_4 = 0, \quad \mathcal{L}_2 X_4 = 2\omega_0. \tag{B.10}$$

Note that \mathcal{L}_1 is precisely the PF operator in the τ -direction. If X_4 is a special period, we can integrate both X_3, X_4 and fix the full triple intersection ring

$$\mathcal{R} = -J_1^3 - J_2 J_1^2 - J_2^2 J_1, \tag{B.11}$$

and then proceed to compute the Euler number $\chi = 60$ as well as $\int c_2 J_1 = -10$, $\int c_2 J_1 = -12$ using (B.2) and (2.11).

Note that X_4 indeed descends from a period of the compact geometry. In the non-compact geometry it is the properly normalised integral over a non-compact cycle in the mirror Calabi-Yau.

n = 5. The computation for the geometry with $\mathbf{n} = 1$ is kind of trivial. Let us now consider a more complicated model with $\mathbf{n} = 5$. Notice in the compact cases, we embed our elliptic Calabi-Yau 3-fold into a toric variety described by a reflexive polytope and its star triangulation. Then the Mori cone generators $l^{(i)}$, which are also known as toric charges, are related to the star triangulation directly. In the de-compactification limit, a point of the polytope is missing, leaving a non-reflexive polytope. The dual polytope now in principle have infinite size. As depicted in section 2.2, the limit happens to take the

variable z_{dc} to 0. This is equivalent to deleting one Mori generator, and keeping the others. From the polytope point of view, we delete a sub-polytope from it, and keep the same triangulation on the remaining part. We may assume that the standard method of mirror symmetry for a compact hyper-surface embedded in a compact toric variety still holds.

For now, we try to triangulate the non-reflexive polytope, it has 16 star triangulations. For one of them, the toric charges are $l^{(i)}$, $i = 1, 2, 3, 4, 6, 7$ in table 2, we say the associated curves form the toric basis of compact curves. One can in principle write down the Picard-Fuchs equations, and then try to find solutions. For this model, it is possible to change the variables of complex structure parameters z_i so that the solutions do not change, and the mirror maps have expansions with positive powers of z_i . The charges of the new basis $l_{F_4}^{(i)}$ can be found in table 2, which correspond exactly to nodes in the Dynkin diagram. Then the complete Picard-Fuchs operators are given in (B.12). There are five B-periods solved from these operators, and an extra one $X_0 \frac{\partial}{\partial \tau} F_{(0,0)}$ annihilated by all the operators except for \mathcal{L}_b , with $\mathcal{L}_b(\omega_0 \frac{\partial}{\partial \tau} F_{(0,0)}) \sim \omega_0$. The Euler number can be predicted from the τ direction genus zero invariant as 12.

$$\begin{aligned}
 \mathcal{L}_0 &= \theta_1 (\theta_1 - 2\theta_2 + \theta_3 - \theta_6) - z_1 (2\theta_1 - \theta_2) (2\theta_1 - \theta_2 + 1) , \\
 \mathcal{L}_1 &= - (2\theta_1 - \theta_2) (\theta_2 - 2\theta_3 + \theta_4 - \theta_6) - z_2 (\theta_1 - 2\theta_2 + \theta_3 - \theta_6 - 1) (\theta_1 - 2\theta_2 + \theta_3 - \theta_6) , \\
 \mathcal{L}_2 &= (2\theta_3 - 2\theta_4 + \theta_5 - 1) (2\theta_3 - 2\theta_4 + \theta_5) (\theta_1 - 2\theta_2 + \theta_3 - \theta_6) , \\
 &\quad - 2z_3 (2\theta_3 + 1) (\theta_2 - 2\theta_3 + \theta_4 - \theta_6 - 1) (\theta_2 - 2\theta_3 + \theta_4 - \theta_6) , \\
 \mathcal{L}_3 &= (\theta_4 - 2\theta_5) (\theta_2 - 2\theta_3 + \theta_4 - \theta_6) - z_4 (2\theta_3 - 2\theta_4 + \theta_5 - 1) (2\theta_3 - 2\theta_4 + \theta_5) , \\
 \mathcal{L}_4 &= \theta_5 (2\theta_3 - 2\theta_4 + \theta_5) - z_5 (\theta_4 - 2\theta_5 - 1) (\theta_4 - 2\theta_5) , \\
 \mathcal{L}_b &= \theta_6^2 - z_6 (\theta_1 - 2\theta_2 + \theta_3 - \theta_6) (\theta_2 - 2\theta_3 + \theta_4 - \theta_6) , \\
 \mathcal{L}_\tau &= \theta_1 (\theta_5 - 1) \theta_5 - 8z_1 z_2^2 z_3^3 z_4^4 z_5^2 (2\theta_3 + 1) (2\theta_3 + 3) (2\theta_3 + 5) .
 \end{aligned} \tag{B.12}$$

C Geometric data

We express here the Mori cone generators of the elliptic non-compact Calabi-Yau threefolds X associated to the minimal 6d SCFTs in terms of the Mori cone generators $l^{(i)}$ of the compact Calabi-Yau \hat{X} given in [8].

$\mathbf{G} = \mathbf{SO}(8)$. $l^{(5)}$ is the direction of decompactification.

$$\begin{aligned}
 \Sigma_b &= l^{(4)} , & \Sigma_0 &= l^{(6)} , & \Sigma_1 &= 2l^{(1)} + l^{(6)} + 2l^{(7)} , \\
 \Sigma_2 &= 2l^{(1)} + l^{(6)} , & \Sigma_3 &= l^{(3)} + l^{(6)} , & \Sigma_4 &= l^{(1)} + l^{(2)} + 2l^{(7)} .
 \end{aligned} \tag{C.1}$$

$\mathbf{G} = \mathbf{F}_4$. $l^{(5)}$ is the direction of decompactification.

$$\Sigma_b = l^{(4)} , \quad \Sigma_0 = l^{(3)} , \quad \Sigma_1 = l^{(2)} + l^{(6)} + 2l^{(7)} , \quad \Sigma_2 = l^{(1)} , \quad \Sigma_3 = l^{(6)} , \quad \Sigma_4 = l^{(7)} . \tag{C.2}$$

$\mathbf{G} = \mathbf{E}_6$. $l^{(3)}$ is the direction of decompactification.

$$\begin{aligned}
 \Sigma_b &= l^{(5)} , \quad \Sigma_0 = l^{(4)} , & \Sigma_1 &= l^{(1)} + l^{(6)} + 2l^{(7)} + 2l^{(9)} , & \Sigma_2 &= l^{(8)} + l^{(9)} , \\
 \Sigma_3 &= l^{(6)} + 2l^{(7)} + l^{(8)} , & \Sigma_4 &= l^{(2)} + l^{(7)} + l^{(8)} , & \Sigma_5 &= l^{(6)} , & \Sigma_6 &= l^{(7)} + l^{(9)} .
 \end{aligned} \tag{C.3}$$

$G = E_7$. $l^{(4)}$ is the direction of decompactification.

$$\begin{aligned} \Sigma_b = l^{(3)}, \quad \Sigma_0 = l^{(5)}, \quad \Sigma_1 = l^{(6)}, \quad \Sigma_2 = l^{(2)} + l^{(7)} + 2l^{(8)} + l^{(9)}, \quad \Sigma_3 = l^{(1)} + l^{(7)} + l^{(10)}, \\ \Sigma_4 = l^{(7)}, \quad \Sigma_5 = l^{(8)} + l^{(10)}, \quad \Sigma_6 = l^{(9)}, \quad \Sigma_7 = l^{(7)} + 2l^{(8)} + l^{(9)}. \end{aligned} \quad (C.4)$$

$G = E_8$. $l^{(3)}$ is the direction for decompactification.

$$\begin{aligned} \Sigma_b = l^{(8)}, \quad \Sigma_0 = l^{(4)}, \quad \Sigma_1 = l^{(5)}, \quad \Sigma_2 = l^{(6)}, \quad \Sigma_3 = l^{(7)}, \quad \Sigma_4 = l^{(1)} + l^{(2)} + l^{(10)} + 2l^{(11)}, \\ \Sigma_5 = l^{(9)}, \quad \Sigma_6 = l^{(10)}, \quad \Sigma_7 = l^{(11)}, \quad \Sigma_8 = l^{(1)} + l^{(10)} + 2l^{(11)}. \end{aligned} \quad (C.5)$$

D Useful identities

Jacobi theta functions with characteristics are defined as

$$\begin{aligned} \theta_1^{[a]}(\tau, z) &= -i \sum_{k \in \mathbb{Z}} (-1)^{k+a} Q_\tau^{(k+1/2+a)^2/2} Q_z^{k+1/2+a}, \\ \theta_2^{[a]}(\tau, z) &= \sum_{k \in \mathbb{Z}} Q_\tau^{(k+1/2+a)^2/2} Q_z^{k+1/2+a}, \\ \theta_3^{[a]}(\tau, z) &= \sum_{k \in \mathbb{Z}} Q_\tau^{(k+a)^2/2} Q_z^{k+a}, \\ \theta_4^{[a]}(\tau, z) &= \sum_{k \in \mathbb{Z}} (-1)^{k+a} Q_\tau^{(k+a)^2/2} Q_z^{k+a}, \end{aligned} \quad (D.1)$$

which satisfy the well-known addition formulas

$$\begin{aligned} \theta_3^{[a_1]}(\tau, z_1) \theta_3^{[a_2]}(\tau, z_2) &= \sum_{i=2,3} \theta_i^{[\frac{a_1+a_2}{2}]}(2\tau, z_1+z_2) \theta_i^{[\frac{a_1-a_2}{2}]}(2\tau, z_1-z_2), \\ \theta_4^{[a_1]}(\tau, z_1) \theta_4^{[a_2]}(\tau, z_2) &= \sum_{i=1,4} \theta_i^{[\frac{a_1+a_2}{2}]}(2\tau, z_1+z_2) \theta_i^{[\frac{a_1-a_2}{2}]}(2\tau, z_1-z_2). \end{aligned} \quad (D.2)$$

Jacobi theta function θ_1 can be defined as triple products

$$\theta_1(\tau, z) = i Q_\tau^{\frac{1}{12}} Q_z^{-\frac{1}{2}} \eta(\tau) \prod_{n=1}^{\infty} (1 - Q_z Q_\tau^{n-1}) \left(1 - \frac{Q_\tau^n}{Q_z} \right), \quad (D.3)$$

which satisfies the quasi-periodicity

$$\theta_1(\tau, z+1) = -\theta_1(\tau, z), \quad (D.4)$$

$$\theta_1(\tau, z+\tau) = -Q_\tau^{-1/2} Q_z^{-1} \theta_1(\tau, z), \quad (D.5)$$

$$\theta_1(\tau, z-\tau) = -Q_\tau^{-1/2} Q_z \theta_1(\tau, z). \quad (D.6)$$

For a cluster of refined BPS invariants $N_{(0,k)}^k = 1$ for all $k \geq 0$, the total contribution to BPS partition function is

$$\sum_{k=0}^{\infty} \frac{Q^k \chi_{2k+1}((q_1 q_2)^{1/2})}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})} = \frac{1+Q}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})(1 - Q q_1 q_2)(1 - Q q_1^{-1} q_2^{-1})}. \quad (D.7)$$

Similarly, for a cluster of refined BPS invariants $N_{(0,k+1/2)}^k = -1$ for all $k \geq 0$, the total contribution to BPS partition function is

$$\sum_{k=0}^{\infty} \frac{Q^k \chi_{2k+2}((q_1 q_2)^{1/2})}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})} = \frac{(q_1 q_2)^{1/2} + (q_1 q_2)^{-1/2}}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})(1 - Q q_1 q_2)(1 - Q q_1^{-1} q_2^{-1})}. \quad (\text{D.8})$$

We also often encounter the case where a cluster of refined BPS invariants $N_{(0,k-1/2)}^k = -1$ for all $k \geq 0$ are combined with a “zero” degree invariants $N_{(0,1/2)}^0 = -1$. In such case, the total contribution is

$$\begin{aligned} & \frac{1}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})} + \sum_{k=1}^{\infty} \frac{Q^k \chi_{2k}((q_1 q_2)^{1/2})}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})} \\ &= \frac{((q_1 q_2)^{1/2} + (q_1 q_2)^{-1/2})(1 + Q + Q^2 - (q_1 q_2 + q_1^{-1} q_2^{-1}) Q)}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})(1 - Q q_1 q_2)(1 - Q q_1^{-1} q_2^{-1})}. \end{aligned} \quad (\text{D.9})$$

In the computation of vector multiplets, we often encounter the following expressions:

$$\text{PE} \left[\frac{Q}{1 - Q_\tau} \right] = \prod_{n=0}^{\infty} \frac{1}{1 - Q Q_\tau^n}, \quad (\text{D.10})$$

and

$$\text{PE} \left[\left(Q_z + \frac{Q_\tau}{Q_z} \right) \left(\frac{1}{1 - Q_\tau} \right) \right] = \frac{i Q_\tau^{1/2} Q_z^{-1/2} \eta(\tau)}{\theta_1(\tau, z)}. \quad (\text{D.11})$$

In counting the total index quadratic form of the contribution from vector multiplets, we often encounter the following expression:

$$\text{PE} \left(- \left(Bl_{(0,1/2,R)}(q_1, q_2) Q_z + Bl_{(0,1/2,-R)}(q_1, q_2) \frac{Q_\tau}{Q_z} \right) \left(\frac{1}{1 - Q_\tau} \right) \right). \quad (\text{D.12})$$

Here

$$Bl_{(j_l, j_r, R)}(q_1, q_2) = f_{(j_l, j_r)}(q_1, q_2/q_1) q_1^R + f_{(j_l, j_r)}(q_1/q_2, q_2) q_2^R - f_{(j_l, j_r)}(q_1, q_2), \quad (\text{D.13})$$

and

$$f_{(j_l, j_r)}(q_1, q_2) = \frac{\chi_{j_l}(q_L) \chi_{j_r}(q_R)}{(q_1^{1/2} - q_1^{-1/2})(q_2^{1/2} - q_2^{-1/2})}. \quad (\text{D.14})$$

Supposing $R \geq 2$, the expression (D.12) can be written as

$$\begin{aligned} & \prod_{\substack{m, n \geq 0 \\ m+n \leq R-1}} \frac{i Q_\tau^{1/12} \eta(Q_z q_1^m q_2^n)^{-1/2}}{\theta_1(z + m \epsilon_1 + n \epsilon_2)} \prod_{\substack{m, n \geq 0 \\ m+n \leq R-2}} \frac{i Q_\tau^{1/12} \eta(Q_z q_1^{m+1} q_2^{n+1})^{-1/2}}{\theta_1(z + (m+1) \epsilon_1 + (n+1) \epsilon_2)} \\ &= (i Q_\tau^{1/12} Q_z^{-1/2})^{R^2} (q_1 q_2)^{-\frac{(R-1)R(R+1)}{6}} \prod_{\substack{m, n \geq 0 \\ m+n \leq R-1}} \frac{\eta}{\theta_1(z + m \epsilon_1 + n \epsilon_2)} \prod_{\substack{m, n \geq 0 \\ m+n \leq R-2}} \frac{\eta}{\theta_1(z + (m+1) \epsilon_1 + (n+1) \epsilon_2)}. \end{aligned} \quad (\text{D.15})$$

We normally denote the modular part of (D.12), i.e. those θ_1 and η functions in the above expression together as $\check{\theta}$. The index quadratic form of $\check{\theta}$ can be computed as

$$\text{Ind}_{\check{\theta}}(z, R) = -\frac{R^2 z^2}{2} - \frac{(R-1)R(R+1)}{3} z(\epsilon_1 + \epsilon_2) - \frac{(R-1)R^2(R+1)}{12} (\epsilon_1^2 + \epsilon_1 \epsilon_2 + \epsilon_2^2), \quad (\text{D.16})$$

which actually holds for all $R \in \mathbb{Z}$. See more details in the appendix A in [25].

E Relation with modular ansatz

In this appendix, we show how the modular ansatz, or its denominator to be specific, for the elliptic genus emerges from our exact formulas. Simply speaking, the denominator of the modular ansatz comes from summing over all α^\vee with a fixed norm square in the recursion formula.

It was proposed in [24, 37] the k -string elliptic genus satisfies the following ansatz

$$\mathbb{E}_{h_G^{(k)}}(\tau, \epsilon_1, \epsilon_2, m_\alpha) = \frac{\mathcal{N}_{G,k}(\tau, \epsilon_1, \epsilon_2, m_\alpha)}{\mathcal{D}_{G,k}(\tau, \epsilon_1, \epsilon_2, m_\alpha)}, \quad (\text{E.1})$$

where both the numerator and the denominator are Weyl invariant Jacobi forms. Furthermore the denominator has the following unique structure as a Weyl invariant Jacobi form which reproduces the poles of the Hilbert series of the moduli space of k G -gauge instantons [43] and the correct leading order of Q_τ :

$$\mathcal{D}_{G,k} = \eta(\tau)^{4kh_G^\vee} \prod_{i=1}^k \varphi_{-1, \frac{1}{2}}(i\epsilon_1) \varphi_{-1, \frac{1}{2}}(i\epsilon_2) \tilde{D}_{G,k} \quad (\text{E.2})$$

with the gauge group related factor

$$\tilde{D}_{G,k} = \prod_{\alpha \in \Delta} \prod_{i=1}^k \prod_{\ell=0}^{i-1} \varphi_{-1, \frac{1}{2}}((i+1)\epsilon_+ + (i-1-2\ell)\epsilon_- + m_\alpha), \quad (\text{E.3})$$

multiplying over the set of roots. Later [97] claims that \mathbb{E}_k has actually fewer poles and as a consequence the denominator is smaller (see also [59]). It can be written as⁴⁷

$$\mathcal{D}_{G,k}^{\text{red}} = \eta(\tau)^{4kh_G^\vee} \prod_{i=1}^k \varphi_{-1, \frac{1}{2}}(i\epsilon_1) \varphi_{-1, \frac{1}{2}}(i\epsilon_2) \tilde{D}_{G,k}^{\text{red}} \quad (\text{E.4})$$

with the gauge group related factor

$$\tilde{D}_{G,k}^{\text{red}} = \prod_{\alpha \in \Delta^l} \mathcal{D}_{k,\alpha}^{\text{SU}(2)}(\tau, m) \prod_{\alpha \in \Delta^s} \mathcal{D}_{\lfloor k/c_\alpha \rfloor, \alpha}^{\text{SU}(2)}(\tau, m), \quad (\text{E.5})$$

where Δ^l, Δ^s are the set of long roots and short roots respectively, the constants c_α are

$$\begin{aligned} c_\alpha &= 2 & \text{if } G = \text{Sp}(N), \text{SO}(2N+1), F_4, \\ c_\alpha &= 3 & \text{if } G = G_2, \end{aligned} \quad (\text{E.6})$$

⁴⁷Here “red” means “reduced”, i.e. the number of poles reduces. The “tilde” and “reduced” in this section should not be confused with them in the main text where they mean a free hypermultiplet is coupled or decoupled.

and that

$$\mathcal{D}_{k,\alpha}^{\text{SU}(2)}(\tau, m) = \prod_{\substack{a,b \leq k \\ a,b > 0}} \varphi_{-1, \frac{1}{2}}(a\epsilon_1 + b\epsilon_2 + m_\alpha). \quad (\text{E.7})$$

To see that $\mathcal{D}_{G,k}^{\text{red}}$ is actually smaller than $\mathcal{D}_{G,k}$, we spell out explicitly the components of $\tilde{\mathcal{D}}_{G,k}, \tilde{\mathcal{D}}_{G,k}^{\text{red}}$ for some small values of k . To be concrete, we take the model of $\mathfrak{n} = 5$ with $G = F_4$. On the one hand,

$$\tilde{\mathcal{D}}_1 = \prod_{\alpha \in \Delta} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha), \quad (\text{E.8})$$

$$\tilde{\mathcal{D}}_2 = \prod_{\alpha \in \Delta} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(2\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 2\epsilon_2 + m_\alpha), \quad (\text{E.9})$$

$$\begin{aligned} \tilde{\mathcal{D}}_3 = \prod_{\alpha \in \Delta} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(2\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 2\epsilon_2 + m_\alpha) \\ \varphi_{-1, \frac{1}{2}}(3\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(2\epsilon_1 + 2\epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 3\epsilon_2 + m_\alpha). \end{aligned} \quad (\text{E.10})$$

On the other hand,

$$\tilde{\mathcal{D}}_1^{\text{red}} = \prod_{\alpha \in \Delta^l} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha), \quad (\text{E.11})$$

$$\begin{aligned} \tilde{\mathcal{D}}_2^{\text{red}} = \prod_{\alpha \in \Delta^s} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \times \\ \prod_{\alpha \in \Delta^l} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(2\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 2\epsilon_2 + m_\alpha), \end{aligned} \quad (\text{E.12})$$

$$\begin{aligned} \tilde{\mathcal{D}}_3^{\text{red}} = \prod_{\alpha \in \Delta^s} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \times \\ \prod_{\alpha \in \Delta^l} \varphi_{-1, \frac{1}{2}}(\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(2\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 2\epsilon_2 + m_\alpha) \\ \varphi_{-1, \frac{1}{2}}(3\epsilon_1 + \epsilon_2 + m_\alpha) \varphi_{-1, \frac{1}{2}}(\epsilon_1 + 3\epsilon_2 + m_\alpha). \end{aligned} \quad (\text{E.13})$$

We will demonstrate that our recursion formulas (3.32) are consistent with (E.4) and (E.5) rather than (E.2), (E.3).

Let us first take a look at the case of $k = 1$ where the recursion formulas (3.32) simply read

$$\mathbb{E}_1 = \sum_{\|\alpha^\vee\|^2=2} (-1)^{|\alpha^\vee|} \frac{D_{\{1,0,0\}}^{\alpha^\vee}}{D_1} A_{\alpha^\vee}. \quad (\text{E.14})$$

We consider the poles contributed by each component. Suppose we choose three unity r fields with $a_{1,2,3}$, which differ from each other by $a_i - a_j = s_{ij}/\mathfrak{n}$. According to the requirement for $a_{1,2,3}$, we know that all s_{ij} are intergers and $0 < |s_{ij}| < n$. Using (D.2), it is not difficult to show that both $D_{1,0,0}^{\alpha^\vee}$ and D_1 contain the zero $\epsilon_1 - \epsilon_2 = 0$ of order $\min(|s_{ij}|, n - |s_{ij}|)$. For example, the minor

$$\Delta_{1,2} = \det \begin{pmatrix} \theta_3^{[a_1]}(n\tau, -2\epsilon_1 + (n-2)\epsilon_2) & \theta_3^{[a_1]}(n\tau, (n-2)\epsilon_1 - 2\epsilon_2) \\ \theta_3^{[a_2]}(n\tau, -2\epsilon_1 + (n-2)\epsilon_2) & \theta_3^{[a_2]}(n\tau, (n-2)\epsilon_1 - 2\epsilon_2) \end{pmatrix}, \quad (\text{E.15})$$

can be rewritten as

$$\Delta_{1,2} = \sum_{i=2,3} \theta_i^{\lfloor \frac{a_1 - a_2}{2} \rfloor} (2n\tau, n(\epsilon_2 - \epsilon_1)) - \theta_i^{\lfloor \frac{a_1 - a_2}{2} \rfloor} (2n\tau, n(\epsilon_1 - \epsilon_2)), \quad (\text{E.16})$$

which clearly contain zeros $\epsilon_1 - \epsilon_2 = 0$ of order $\min(|s_{ij}|, n - |s_{ij}|)$. Now by the universality argument in section 3.2, we can choose arbitrary three $a_{1,2,3}$ in recursion formulas. Let us choose three successive ones with $a_3 - a_2 = a_2 - a_1 = 1/n$. Both $D_{1,0,0}^{\alpha^\vee}$ and D_1 have the simple zero $\epsilon_1 - \epsilon_2 = 0$. In fact, more is true, the determinant

$$D(z_1, z_2, z_3) = \det \begin{pmatrix} \theta_i^{[a_1]}(n\tau, nz_1) & \theta_i^{[a_1]}(n\tau, nz_2) & \theta_i^{[a_1]}(n\tau, nz_3) \\ \theta_i^{[a_2]}(n\tau, nz_1) & \theta_i^{[a_2]}(n\tau, nz_2) & \theta_i^{[a_2]}(n\tau, nz_3) \\ \theta_i^{[a_3]}(n\tau, nz_1) & \theta_i^{[a_3]}(n\tau, nz_2) & \theta_i^{[a_3]}(n\tau, nz_3) \end{pmatrix} \quad (\text{E.17})$$

with characteristics $a_{1,2,3}$ chosen as above and $i = 3, 4$ always has simple zeros at $z_1 - z_2 = z_2 - z_3 = z_3 - z_1 = 0$. Therefore $D_{1,0,0}^{\alpha^\vee}/D_1$ has zeros/poles

$$\frac{(\epsilon_1 - \epsilon_2)(m_\alpha - \epsilon_1)(m_\alpha - \epsilon_2)}{(\epsilon_1 - \epsilon_2)\epsilon_1\epsilon_2}, \quad (\text{E.18})$$

which can be boosted to the modular object

$$\frac{\theta_1(m_\alpha - \epsilon_1)\theta_1(m_\alpha - \epsilon_2)}{\theta_1(\epsilon_1)\theta_1(\epsilon_2)}. \quad (\text{E.19})$$

The other component in (E.14) for \mathbb{E}_1 is

$$A_{\alpha^\vee}(my) = \frac{\eta^4}{\theta_1(m_\alpha)\theta_1(m_\alpha - \epsilon_1)\theta_1(m_\alpha - \epsilon_2)\theta_1(m_\alpha - \epsilon_1 - \epsilon_2)} \prod_{\substack{\beta \in \Delta \\ \beta \cdot \alpha^\vee = 1}} \frac{\eta}{\theta_1(m_\beta)}, \quad (\text{E.20})$$

where $\alpha = \alpha^\vee \cdot 2/|\alpha^\vee|^2$ in the root associated to the coroot α^\vee . The components $\theta_1(m_\alpha - \epsilon_1)\theta_1(m_\alpha - \epsilon_2)$ cancel with corresponding components in the numerator of $D_{1,0,0}^{\alpha^\vee}/D_1$. Thus naively A_{α^\vee} should contribute poles⁴⁸ $m_\beta = 0$ for $\beta \in \Delta$ and $m_\alpha - \epsilon_1 - \epsilon_2 = 0$ for $\alpha \in \Delta^l$ to \mathbb{E}_1 . The former poles, however, get canceled with some factors in the numerator after performing the summation over short coroots in (E.14), which can either be seen in explicit calculations,⁴⁹ or be argued formally from the Hilbert series of gauge instanton moduli space [43] as well as from W-algebra via AGT correspondence [59]. The remaining genuine poles are $m_\alpha - \epsilon_1 - \epsilon_2 = 0$ for $\alpha \in \Delta^l$, which are consistent with the slimmer expression (E.11) rather than (E.8).

The discussion above indicates that the recursion formulas (3.32) for \mathbb{E}_k naively contain both genuine poles and spurious poles, while the cancellation of the latter is not obvious. Nevertheless, all true poles should already be visible in the recursion formulas, which allows us to distinguish (E.4) from (E.2). In the following, we will argue in favor of (E.4) by pointing out that extra poles indicated in (E.2) are not present in the recursion formulas (3.32),

⁴⁸There are also poles at positions shifted by 1 or τ of course.

⁴⁹This cancellation can be checked in all the minimal models, not only in the $n = 5$ model.

and assume along the way that the spurious poles not consistent with either (E.2) or (E.4) cancel automatically. With this comment in mind, let us now look at the denominators of $\mathbb{E}_2, \mathbb{E}_3$ from the recursion formulas, and consider only *relevant* poles, the poles which appear also in either (E.2) or (E.4).

The right hand side of (3.32) for \mathbb{E}_2 has three sectors with

$$(\|\alpha^\vee\|^2, d_1, d_2) = (4, 0, 0), (2, 1, 0)/(2, 0, 1), (0, 1, 1) \quad (\text{E.21})$$

respectively. With an explicit calculation in the $\mathfrak{n} = 5$ models, we find the following relevant poles in each of the three sectors⁵⁰

$(\ \alpha^\vee\ ^2, d_1, d_2)$	relevant poles
(4,0,0)	$m_\alpha - \epsilon_1 - \epsilon_2, \alpha \in \Delta$
(2,1,0)/(2,0,1)	$m_\alpha - \epsilon_1 - \epsilon_2, m_\alpha - 2\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 2\epsilon_2, \alpha \in \Delta^l$
(0,1,1)	–

They are clearly consistent with (E.12). Likewise, for \mathbb{E}_3 there are five sectors on the right hand side of (3.32), and we find the relevant poles as follows

$(\ \alpha^\vee\ ^2, d_1, d_2)$	relevant poles
(6,0,0)	$m_\alpha - \epsilon_1 - \epsilon_2, \alpha \in \Delta;$ $m_\alpha - 2\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 2\epsilon_2, \alpha \in \Delta^l$
(4,1,0)/(4,0,1)	$m_\alpha - \epsilon_1 - \epsilon_2, m_\alpha - 2\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 2\epsilon_2, \alpha \in \Delta^l$
(2,2,0)/(2,0,2)	$m_\alpha - \epsilon_1 - \epsilon_2, \alpha \in \Delta;$ $m_\alpha - 2\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 2\epsilon_2,$ $m_\alpha - 3\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 3\epsilon_2, \alpha \in \Delta^l$
(2,1,1)	$m_\alpha - \epsilon_1 - \epsilon_2, m_\alpha - 2\epsilon_1 - \epsilon_2, m_\alpha - \epsilon_1 - 2\epsilon_2, \alpha \in \Delta^l$
(0,2,1)	$m_\alpha - \epsilon_1 - \epsilon_2, \alpha \in \Delta^l$

which are consistent with (E.13).

F Elliptic genera

We record some high Q_τ order results for the reduced elliptic genus here. Recall the k -string elliptic genus when expanded with respect to Q_τ can be written as

$$\mathbb{E}_{h_G^{(k)}}(v, x, Q_\tau, Q_{m_i}) = v^{kh_G^\vee - 1} Q_\tau^{-(kh_G^\vee - 1)/6} \sum_{n=0}^{\infty} Q_\tau^n g_{k,G}^{(n)}(v, x, Q_{m_i}).$$

We are interested in the v -expansion of $g_{k,G}^{(n)}(v, x, Q_{m_i})$. Usually, the leading v power becomes more and more negative when Q_τ order n goes up. When n is enough high, we

⁵⁰We also suppress the poles $\epsilon_1 = \epsilon_2 = 0$, which are guaranteed by the $D_{\{d_0, d_1, d_2\}}^{\alpha^\vee}/D_d$ structure similar to the previous discussion.

observe some patterns for the leading v -expansion behavior. For two-strings elliptic genus, we observe for $i \geq 2(\mathbf{n} - 2)$,

$$g_{i,G}^{(2)}(v, x, Q_{m_i}) = -v^{-3(\mathbf{n}+i)-5}(\chi_{(i-2(\mathbf{n}-2))\theta}\chi_{i-(\mathbf{n}-2)}) - \mathcal{O}(v^{-3(\mathbf{n}+i)-4}) \dots \quad (\text{F.1})$$

As high as the Q_τ order we have reached, this is true. It is nice to see how to explain this phenomenon.

A₂. For the reduced two-string elliptic genus for $G = \text{SU}(3)$ model, recall

$$g_{2,A_2}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{10}(1+v)^6(1+v+v^2)^5} \times P_{2,A_2}^{(n)}(v).$$

We have

$$\begin{aligned} P_{2,A_2}^{(2)}(v) &= v^{-10}(-1-v+6v^2+9v^3-10v^4-33v^5+41v^6+256v^7+428v^8+220v^9-347v^{10}-823v^{11}-131v^{12} \\ &\quad +2652v^{13}+7721v^{14}+14419v^{15}+21826v^{16}+27125v^{17}+28966v^{18}+27125v^{19}+\dots+v^{36}), \\ P_{2,A_2}^{(3)}(v) &= v^{-13}(-16-28v+50v^2+198v^3+138v^4-399v^5-963v^6-419v^7+1716v^8+4316v^9+5014v^{10} \\ &\quad +2174v^{11}-4110v^{12}-10701v^{13}-12583v^{14}-2128v^{15}+27073v^{16}+75426v^{17}+136089v^{18}+198723v^{19} \\ &\quad +244336v^{20}+260628v^{21}+244336v^{22}+\dots+v^{42}), \\ P_{2,A_2}^{(4)}(v) &= v^{-16}(-81-223v-5v^2+880v^3+1695v^4+577v^5-3110v^6-6735v^7-5803v^8+2582v^9+17019v^{10} \\ &\quad +31735v^{11}+38096v^{12}+25783v^{13}-11462v^{14}-68637v^{15}-118109v^{16}-116300v^{17}-11102v^{18} \\ &\quad +231810v^{19}+605425v^{20}+1054375v^{21}+1497688v^{22}+1816145v^{23}+1930514v^{24}+\dots+v^{48}), \\ P_{2,A_2}^{(5)}(v) &= v^{-19}(-256-874v-810v^2+1432v^3+5510v^4+7553v^5+2483v^6-10671v^7-26644v^8-34874v^9 \\ &\quad -23600v^{10}+16620v^{11}+90316v^{12}+185964v^{13}+260250v^{14}+241846v^{15}+62988v^{16}-279728v^{17} \\ &\quad -692866v^{18}-963287v^{19}-815649v^{20}+30457v^{21}+1729760v^{22}+4189517v^{23}+7044615v^{24}+9777494v^{25} \\ &\quad +11727151v^{26}+12428414v^{27}+11727151v^{28}+\dots+v^{54}), \\ &\dots \\ P_{2,A_2}^{(10)}(v) &= v^{-34}(-6561-31129v-72335v^2-120018v^3-168868v^4-206886v^5-222250v^6-170936v^7+98441v^8 \\ &\quad +773013v^9+1959712v^{10}+3612563v^{11}+5410138v^{12}+6633691v^{13}+5994478v^{14}+1510639v^{15} \\ &\quad -9064153v^{16}-27314880v^{17}-52091127v^{18}-77137569v^{19}-89646012v^{20}-69861741v^{21}+5587892v^{22} \\ &\quad +154737705v^{23}+377669387v^{24}+641219584v^{25}+868425362v^{26}+940214412v^{27}+718317837v^{28} \\ &\quad +93293293v^{29}-951170969v^{30}-2275980208v^{31}-3545719630v^{32}-4250592736v^{33}-3785597548v^{34} \\ &\quad -1612094548v^{35}+2569710910v^{36}+8659446500v^{37}+16075828305v^{38}+23808502207v^{39} \\ &\quad +30625221095v^{40}+35302136340v^{41}+36965592032v^{42}+35302136340v^{43}+\dots+v^{84}). \end{aligned}$$

Note that the leading power of v in $P_{2,A_2}^{(n)}(v)$ becomes more and more negative as n increases. In fact, we notice for $n \geq 2$,

$$P_{2,A_2}^{(n)}(v) = v^{-3n-4}(- (n-1)^4 - \dots + \text{palindrome up to } v^{6(n+4)}). \quad (\text{F.2})$$

With generic x , we also observe the following general expansion: for $n \geq 2$,

$$v^{3n+4}g_{2,A_2}^{(n)}(v, x, Q_{m_i}) = -\chi_{(n-2)\theta}^{A_2}\chi_{n-1} - (\chi_{(n-3)\theta}^{A_2}\chi_n + \dots)v - (\chi_{(n-4)\theta}^{A_2}\chi_{n+1} + \dots)v^2 + \dots \quad (\text{F.3})$$

For the reduced elliptic genus of three-strings, recall

$$g_{3,A_2}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{16}(1+v)^{10}(1+v^2)^5(1+v+v^2)^6} \times P_{3,A_2}^{(n)}(v). \quad (\text{F.4})$$

We obtain

$$P_{3,A_2}^{(2)}(v) = v^{-10}(-1-2v-5v^2-4v^3+11v^4+30v^5+74v^6+84v^7+267v^8+880v^9+2718v^{10}+6130v^{11} \\ +11512v^{12}+17594v^{13}+24774v^{14}+35306v^{15}+64404v^{16}+133960v^{17}+290609v^{18}+569846v^{19} \\ +1020364v^{20}+1628376v^{21}+2376984v^{22}+3145582v^{23}+3848955v^{24}+4318298v^{25}+4501676v^{26} \\ +4318298v^{27}+\dots+v^{52}),$$

$$P_{3,A_2}^{(3)}(v) = v^{-17}(2-10v^2-9v^3+12v^4+44v^5-48v^6-367v^7-622v^8-183v^9+1740v^{10}+4336v^{11}+4488v^{12} \\ -1116v^{13}-12504v^{14}-17424v^{15}+4576v^{16}+77007v^{17}+199630v^{18}+337366v^{19}+417362v^{20} \\ +382080v^{21}+255946v^{22}+265962v^{23}+839446v^{24}+2662413v^{25}+6368124v^{26}+12524957v^{27} \\ +21034734v^{28}+31405824v^{29}+42159674v^{30}+51803451v^{31}+58351834v^{32}+60785174v^{33} \\ +58351834v^{34}+\dots+v^{66}),$$

$$P_{3,A_2}^{(4)}(v) = v^{-21}(32+40v-84v^2-339v^3-426v^4+487v^5+2242v^6+2876v^7-1500v^8-12817v^9-23440v^{10} \\ -16831v^{11}+25630v^{12}+97493v^{13}+150494v^{14}+103012v^{15}-112582v^{16}-448583v^{17}-700938v^{18} \\ -495362v^{19}+512476v^{20}+2411835v^{21}+4741886v^{22}+6461123v^{23}+6254586v^{24}+3310035v^{25} \\ -1672684v^{26}-5447913v^{27}-2195270v^{28}+16177917v^{29}+57171170v^{30}+126228382v^{31}+222042686v^{32} \\ +337620393v^{33}+457167512v^{34}+563129124v^{35}+635332178v^{36}+661625872v^{37}+635332178v^{38}+\dots+v^{74}),$$

$$P_{3,A_2}^{(5)}(v) = v^{-25}(162+412v+276v^2-1295v^3-4518v^4-5771v^5+1292v^6+21793v^7+44712v^8+38030v^9 \\ -38688v^{10}-192767v^{11}-341004v^{12}-309898v^{13}+101910v^{14}+927411v^{15}+1862626v^{16}+2189163v^{17} \\ +1001340v^{18}-2210183v^{19}-6887108v^{20}-10727974v^{21}-10030208v^{22}-703145v^{23}+19133352v^{24} \\ +46719280v^{25}+72876216v^{26}+83251991v^{27}+63444248v^{28}+7529926v^{29}-72005488v^{30}-136950773v^{31} \\ -124639134v^{32}+43752233v^{33}+438669558v^{34}+1103016208v^{35}+2019390168v^{36}+3111882801v^{37} \\ +4234930428v^{38}+5220665928v^{39}+5892000612v^{40}+6133530828v^{41}+5892000612v^{42}+\dots+v^{82}),$$

$$P_{3,A_2}^{(6)}(v) = v^{-29}(512+1690v+3044v^2+766v^3-10596v^4-31191v^5-46930v^6-19288v^7+89188v^8+274511v^9 \\ +419176v^{10}+288140v^{11}-374722v^{12}-1614464v^{13}-2963132v^{14}-3347473v^{15}-1280918v^{16} \\ +4247141v^{17}+12688484v^{18}+20686636v^{19}+22001562v^{20}+9184210v^{21}-22497646v^{22}-69062433v^{23} \\ -113553734v^{24}-124631077v^{25}-65395112v^{26}+91413273v^{27}+339731228v^{28}+622477442v^{29} \\ +828007584v^{30}+813952151v^{31}+463211226v^{32}-241280313v^{33}-1140297436v^{34}-1857927117v^{35} \\ -1829759686v^{36}-398271211v^{37}+2994640544v^{38}+8636915664v^{39}+16332876522v^{40}+25397471432v^{41} \\ +34642293136v^{42}+42690290417v^{43}+48157593258v^{44}+50110832268v^{45}+48157593258v^{46}+\dots+v^{90}).$$

For $n \geq 3$, we notice the following universal leading behavior

$$g_{n,A_2}^{(3)}(v, x) = v^{-4n-5}(\chi_{2(n-2)}\chi_{(n-3)}\theta + \mathcal{O}(v)). \quad (\text{F.5})$$

D₄. For the reduced two-string elliptic genus of SO(8) 6d SCFT, recall

$$g_{2,D_4}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{22}(1+v)^{12}(1+v+v^2)^{11}} \times P_{2,D_4}^{(n)}(v).$$

We have

$$P_{2,D_4}^{(2)}(v) = -v^{-8}(1+3v+22v^2+47v^3+108v^4+29v^5-184v^6-861v^7-1762v^8-3305v^9-13965v^{10}-65210v^{11}-260932v^{12}-884324v^{13}-2589008v^{14}-6645978v^{15}-15280924v^{16}-31673046v^{17}-59506626v^{18}-101871752v^{19}-159566189v^{20}-229273231v^{21}-303095099v^{22}-369460130v^{23}-415787266v^{24}-432409780v^{25}-415787266v^{26}+\dots+v^{50}),$$

$$P_{2,D_4}^{(3)}(v) = v^{-11}(1+v^2)(2+3v-23v^2-281v^3-1338v^4-3786v^5-5794v^6-384v^7+22410v^8+57106v^9+65200v^{10}-8878v^{11}-148110v^{12}-82859v^{13}+911821v^{14}+4473554v^{15}+14488449v^{16}+39847612v^{17}+97908991v^{18}+217119419v^{19}+434067674v^{20}+782158341v^{21}+1274332193v^{22}+1885916954v^{23}+2545465687v^{24}+3144719815v^{25}+3566407238v^{26}+3718703248v^{27}+3566407238v^{28}+\dots+v^{54}),$$

$$P_{2,D_4}^{(4)}(v) = -v^{-19}(2+2v-22v^2-42v^3+88v^4+329v^5-11v^6-1411v^7-1930v^8+1877v^9+12149v^{10}+38843v^{11}+113904v^{12}+240220v^{13}+226932v^{14}-364906v^{15}-1770622v^{16}-3067485v^{17}-1974439v^{18}+3262797v^{19}+10272490v^{20}+10304550v^{21}-10767498v^{22}-74227039v^{23}-228511951v^{24}-601571798v^{25}-1479273049v^{26}-3376185023v^{27}-7028420238v^{28}-13251946148v^{29}-22656311766v^{30}-35281600040v^{31}-50298462814v^{32}-65951082224v^{33}-79826800694v^{34}-89424037262v^{35}-92856436862v^{36}-89424037262v^{37}+\dots+v^{72}),$$

$$P_{2,D_4}^{(5)}(v) = -v^{-22}(1+v^2)(84+146v-831v^2-2495v^3+2027v^4+16874v^5+12844v^6-55230v^7-118023v^8+45251v^9+412658v^{10}+306961v^{11}-747606v^{12}-1085075v^{13}+2514527v^{14}+8955862v^{15}+7677132v^{16}-13638464v^{17}-48153754v^{18}-58843606v^{19}-3166625v^{20}+114419603v^{21}+211444789v^{22}+164548302v^{23}-98377407v^{24}-542911427v^{25}-1148224939v^{26}-2306012154v^{27}-5445277331v^{28}-13585439053v^{29}-31372682848v^{30}-64179900027v^{31}-116221363135v^{32}-188200853472v^{33}-275432340276v^{34}-367411722852v^{35}-449506711376v^{36}-506517773112v^{37}-526950799964v^{38}-506517773112v^{39}+\dots+v^{76}),$$

$$P_{2,D_4}^{(6)}(v) = -v^{-25}(1200+3374v-6724v^2-40296v^3-31436v^4+159328v^5+410682v^6+15592v^7-1467224v^8-2279805v^9+1088019v^{10}+8007211v^{11}+8314028v^{12}-9090084v^{13}-32918254v^{14}-19726713v^{15}+67719233v^{16}+186559606v^{17}+183703487v^{18}-107579194v^{19}-661966959v^{20}-1107115384v^{21}-832956121v^{22}+557841530v^{23}+2672286699v^{24}+4109758618v^{25}+3046056345v^{26}-1483241349v^{27}-8623040702v^{28}-16826056321v^{29}-28634625983v^{30}-59027415291v^{31}-144463390006v^{32}-347331281609v^{33}-749797256633v^{34}-1433909622002v^{35}-2450540661913v^{36}-3786292890317v^{37}-5341633935638v^{38}-6932828077068v^{39}-8322655072044v^{40}-9274316155458v^{41}-9612992092064v^{42}-9274316155458v^{43}+\dots+v^{84}).$$

For $n \geq 4$, we observe the following general leading order behavior

$$v^{3n+7}g_{2,D_4}^{(n)}(v, x, Q_{m_i}) = -\chi_{(n-4)\theta}^{D_4}\chi_{n-2}(x) + \mathcal{O}(v). \tag{F.6}$$

F₄. For the reduced two-string elliptic genus of F_4 6d SCFT, recall

$$g_{2,F_4}^{(n)}(v, x = 1, Q_{m_i} = 1) = \frac{1}{(1-v)^{34}(1+v)^{22}(1+v+v^2)^{17}} \times P_{2,F_4}^{(n)}(v). \tag{F.7}$$

We have

$$\begin{aligned}
 P_{2,F_4}^{(2)}(v) = & 1653 + 14307v + 118305v^2 + 770928v^3 + 4293754v^4 + 20938534v^5 + 90874761v^6 + 354378137v^7 \\
 & + 1254181600v^8 + 4057153817v^9 + 12068000241v^{10} + 33173318084v^{11} + 84638789902v^{12} + 201171570880v^{13} \\
 & + 446852450528v^{14} + 930177353330v^{15} + 1818950197662v^{16} + 3348446417048v^{17} + 5813458948881v^{18} \\
 & + 9534395062259v^{19} + 14791974953829v^{20} + 21734938030680v^{21} + 30278882934513v^{22} + 40026745086453v^{23} \\
 & + 50246339190488v^{24} + 59931545438647v^{25} + 67951300040637v^{26} + 73260270949890v^{27} \\
 & + 75118982210308v^{28} + 73260270949890v^{29} + \dots + v^{56},
 \end{aligned}$$

$$\begin{aligned}
 P_{2,F_4}^{(3)}(v) = & v^{-12}(1 + 7v + 58v^2 + 277v^3 + 1071v^4 + 2976v^5 + 5964v^6 + 5832v^7 - 9266v^8 - 57418v^9 - 135552v^{10} \\
 & - 169110v^{11} + 55432v^{12} + 957484v^{13} + 4373263v^{14} + 21374265v^{15} + 111764209v^{16} + 546538072v^{17} \\
 & + 2392703794v^{18} + 9374157248v^{19} + 33193123730v^{20} + 107176864396v^{21} + 317784470985v^{22} \\
 & + 870042414425v^{23} + 2209544249477v^{24} + 5224968453408v^{25} + 11543089455336v^{26} + 23893104753132v^{27} \\
 & + 46455227531026v^{28} + 85029824340612v^{29} + 146800695692191v^{30} + 239466759024769v^{31} \\
 & + 369631201685833v^{32} + 540578567158590v^{33} + 749891181508407v^{34} + 987634558444181v^{35} \\
 & + 1235929372818009v^{36} + 1470501793161916v^{37} + 1664270696390466v^{38} + 1792334859311106v^{39} \\
 & + 1837134386523548v^{40} + 1792334859311106v^{41} + \dots + v^{80}),
 \end{aligned}$$

$$\begin{aligned}
 P_{2,F_4}^{(4)}(v) = & -v^{-15}(2 + 11v - 9v^2 - 729v^3 - 6740v^4 - 37612v^5 - 146242v^6 - 411322v^7 - 786424v^8 - 683123v^9 \\
 & + 1519991v^{10} + 7863641v^{11} + 17299644v^{12} + 19396808v^{13} - 7574716v^{14} - 81039308v^{15} - 177701238v^{16} \\
 & - 228893241v^{17} - 328644969v^{18} - 1704624732v^{19} - 10239254139v^{20} - 48437902734v^{21} - 193165328741v^{22} \\
 & - 683657580163v^{23} - 2198459381202v^{24} - 6491341536008v^{25} - 17700027945928v^{26} - 44757889196479v^{27} \\
 & - 105345430184255v^{28} - 231557373968430v^{29} - 476759190688969v^{30} - 921916213048446v^{31} \\
 & - 1678228390978831v^{32} - 2881832101851776v^{33} - 4676542813241321v^{34} - 7182956156002759v^{35} \\
 & - 10456822688468996v^{36} - 14445439874041454v^{37} - 18955600691329352v^{38} - 23647746652092629v^{39} \\
 & - 28066464498494385v^{40} - 31707630617443838v^{41} - 34110133094328607v^{42} - 34949875241183086v^{43} \\
 & - 34110133094328607v^{44} + \dots + v^{86}),
 \end{aligned}$$

$$\begin{aligned}
 P_{2,F_4}^{(5)}(v) = & v^{-18}(1 + v - 187v^2 - 1942v^3 - 8588v^4 + 388v^5 + 256583v^6 + 1875127v^7 + 8097680v^8 + 23995511v^9 \\
 & + 47743204v^{10} + 45403031v^{11} - 77639509v^{12} - 441768092v^{13} - 990656365v^{14} - 1141697391v^{15} \\
 & + 314912356v^{16} + 4322862537v^{17} + 9413229443v^{18} + 9932344114v^{19} - 1369097445v^{20} - 22028614537v^{21} \\
 & - 15689824140v^{22} + 143994570705v^{23} + 853983234794v^{24} + 3391440788051v^{25} + 11730377038184v^{26} \\
 & + 37232002426385v^{27} + 109311773773871v^{28} + 297084435895112v^{29} + 748700764659463v^{30} \\
 & + 1754971776388633v^{31} + 3839341467548316v^{32} + 7864675977003847v^{33} + 15128265661342010v^{34} \\
 & + 27394204368470677v^{35} + 46797420093685421v^{36} + 75560202097716654v^{37} + 115501696318370322v^{38} \\
 & + 167393226212698984v^{39} + 230299469534789175v^{40} + 301112810847585121v^{41} + 374497086121044331v^{42} \\
 & + 443382800033202476v^{43} + 500006416441373208v^{44} + 537304885313273980v^{45} + 550330680323572096v^{46} \\
 & + 537304885313273980v^{47} + \dots + v^{92}).
 \end{aligned}$$

For $n \geq 6$, we observe the following general leading order behavior

$$v^{3n+10} g_{2,F_4}^{(n)}(v, x, Q_{m_i}) = -\chi_{(n-6)\theta}^{F_4} \chi_{n-3}(x) + \mathcal{O}(v). \quad (\text{F.8})$$

E₆. For the reduced two-string elliptic genus of E_6 6d SCFT, recall

$$g_{2,E_6}^{(n)}(v, x, Q_{m_i} = 1) = \frac{1}{(1-v)^{46}(1+v)^{32}(1+v+v^2)^{23}} \times P_{2,E_6}^{(n)}(v).$$

We have

$$\begin{aligned}
 P_{2,E_6}^{(1)}(v) = & (1+v^2)(82+896v+9129v^2+73825v^3+515477v^4+3176394v^5+17567385v^6+88082527v^7 \\
 & +404122599v^8+1707996910v^9+6687039606v^{10}+24365673656v^{11}+82957003626v^{12} \\
 & +264812209428v^{13}+794925309293v^{14}+2249848989493v^{15}+6017588149603v^{16}+15241390482586v^{17} \\
 & +36623148751459v^{18}+83623554563863v^{19}+181712020504595v^{20}+376267731853770v^{21} \\
 & +743340720549339v^{22}+1402570753853399v^{23}+2530053857442778v^{24}+4367001323365453v^{25} \\
 & +7218179887542376v^{26}+11433257908228549v^{27}+17365401325615558v^{28}+25305594210396759v^{29} \\
 & +35398201343930359v^{30}+47551931562200552v^{31}+61367940071565626v^{32} \\
 & +76109936363599780v^{33}+90737018750916024v^{34}+104007721490984500v^{35} \\
 & +114645634265369518v^{36}+121537998101131452v^{37}+123925354694394472v^{38}+\dots+v^{76}).
 \end{aligned}$$

$$\begin{aligned}
 P_{2,E_6}^{(2)}(v) = & 3486+44488v+464913v^2+3873323v^3+27606333v^4+172731506v^5+966630727v^6+4894721995v^7 \\
 & +22642609833v^8+96385144324v^9+379809898432v^{10}+1392335683050v^{11}+4768406721146v^{12} \\
 & +15311706805952v^{13}+46244599549903v^{14}+131729726893973v^{15}+354773291170080v^{16} \\
 & +905322588772629v^{17}+2193217132886674v^{18}+5052883393549785v^{19}+11088020153427871v^{20} \\
 & +23207949295452654v^{21}+46391596457503471v^{22}+88666687652950697v^{23}+162200225646883046v^{24} \\
 & +28426293556020849v^{25}+477678248920949928v^{26}+770246305708025175v^{27} \\
 & +1192619227946678339v^{28}+1774271549538256254v^{29}+2537602212267590587v^{30} \\
 & +3490792045588793343v^{31}+4620696016056616197v^{32}+5887558970641741644v^{33} \\
 & +7223464858978994614v^{34}+8535984817703595260v^{35}+9717463741648588196v^{36} \\
 & +10658977913348459838v^{37}+11266596547576742504v^{38}+11476648364287371362v^{39} \\
 & +11266596547576742504v^{40}+\dots+v^{78}.
 \end{aligned}$$

$$\begin{aligned}
 P_{2,E_6}^{(3)}(v) = & v^{-4}(-1-11v-112v^2-769v^3+97959v^4+1465146v^5+15949836v^6+136827854v^7+992270036v^8 \\
 & +6276463714v^9+35347238184v^{10}+179602878296v^{11}+831787432544v^{12}+3538777989264v^{13} \\
 & +13918341585911v^{14}+50873087148945v^{15}+173571288630315v^{16}+554884515982156v^{17} \\
 & +1667591219224745v^{18}+4724849885190467v^{19}+12653182331604747v^{20}+32099832942977106v^{21} \\
 & +77298235563217848v^{22}+177003542946746952v^{23}+386049290828201197v^{24} \\
 & +803129445851193549v^{25}+1595815215747637914v^{26}+3032155653955325045v^{27} \\
 & +5515192004170661557v^{28}+9612446692952872376v^{29}+16067738341715188425v^{30} \\
 & +25779182779041415519v^{31}+39727277739048609244v^{32}+58842809938209124305v^{33} \\
 & +83817103245449704330v^{34}+114875661886259702901v^{35}+151556399981844604507v^{36} \\
 & +192548720900049944088v^{37}+235652392733234663776v^{38}+277900332002367574858v^{39} \\
 & +315856987228959670753v^{40}+346060446910493617699v^{41}+365533661213463473501v^{42} \\
 & +372262200765577638648v^{43}+\dots+v^{86}).
 \end{aligned}$$

$$\begin{aligned}
 P_{2,E_6}^{(4)}(v) = & v^{-16}(-1-11v-112v^2-769v^3-4214v^4-18313v^5-64197v^6-177594v^7-364431v^8-421607v^9 \\
 & +420751v^{10}+3722444v^{11}+10737460v^{12}+18191652v^{13}+11206753v^{14}-38487665v^{15} \\
 & -147892027v^{16}-235567050v^{17}+204232919v^{18}+4014329887v^{19}+28747116555v^{20} \\
 & +177469181418v^{21}+994624050267v^{22}+5061754204737v^{23}+23475175955326v^{24} \\
 & +99852192764195v^{25}+392058843196059v^{26}+1428966813600884v^{27}+4857570355921361v^{28} \\
 & +15462381593811917v^{29}+46247090390273044v^{30}+130358984643118795v^{31} \\
 & +347206472133377093v^{32}+875864441137943176v^{33}+2096960972906450647v^{34}
 \end{aligned}$$

$$\begin{aligned}
&+4773710145181408321v^{35} + 10350471755632365352v^{36} + 21407072068973867721v^{37} \\
&+ 42290120178103689822v^{38} + 79898824171145741885v^{39} + 144526191934600175231v^{40} \\
&+ 250551336753503603832v^{41} + 416666541458353995971v^{42} + 665245395666408218767v^{43} \\
&+ 1020471334680763094257v^{44} + 1505001425501016329934v^{45} + 2135260459421424616916v^{46} \\
&+ 2915910779649516764294v^{47} + 3834519574899765264582v^{48} + 4857787518331415856898v^{49} \\
&+ 5930740363027659407709v^{50} + 6979905103120498497197v^{51} + 7920702637212573264754v^{52} \\
&+ 8668249061573197611391v^{53} + 9149752299658818747593v^{54} + 9316044470623822160548v^{55} + \dots + v^{110}).
\end{aligned}$$

E7. For the reduced two-string elliptic genus of E_7 6d SCFT, recall

$$g_{2,E_7}^{(n)}(v, x, Q_{m_i} = 1) = \frac{1}{(1-v)^{70}(1+v)^{52}(1+v+v^2)^{35}} \times P_{n,E_7}^{(2)}(v). \quad (\text{F.9})$$

We have

$$\begin{aligned}
P_{0,E_7}^{(2)}(v) = &1 + 17v + 237v^2 + 2628v^3 + 25193v^4 + 213819v^5 + 1638666v^6 + 11476871v^7 + 74152233v^8 \\
&+ 445070980v^9 + 2495671432v^{10} + 13133928036v^{11} + 65121712327v^{12} + 305215505275v^{13} \\
&+ 1356033968529v^{14} + 5725284334978v^{15} + 23021851542594v^{16} + 88338636956104v^{17} \\
&+ 324035139906700v^{18} + 1138031848052668v^{19} + 3832341391241046v^{20} + 12390621413785440v^{21} \\
&+ 38509222288582663v^{22} + 115175603408208175v^{23} + 331836472263902521v^{24} + 921861932483495244v^{25} \\
&+ 2471530433876763846v^{26} + 6399961693050532054v^{27} + 16018745367471142680v^{28} \\
&+ 38781560068496818142v^{29} + 90876821066275028695v^{30} + 206242719899419463791v^{31} \\
&+ 453576963793872584712v^{32} + 967171231109021529977v^{33} + 2000571291562232590513v^{34} \\
&+ 4016126507767354504238v^{35} + 7828073649219480743672v^{36} + 14820947289312246349740v^{37} \\
&+ 27267076918737091016348v^{38} + 48764087264312469202730v^{39} + 84802326792798968389732v^{40} \\
&+ 143449590902653729399624v^{41} + 236104043071240448693797v^{42} + 378216261606533139497461v^{43} \\
&+ 589822792928957883073617v^{44} + 895677339869346647226824v^{45} + 1324728639658651633703727v^{46} \\
&+ 1908697079658876873038411v^{47} + 2679565476854052143878502v^{48} + 3665936157860425562998541v^{49} \\
&+ 4888414479465062757831170v^{50} + 6354435158683924634396271v^{51} + 8053206553397859455383003v^{52} \\
&+ 9951646269406905770095206v^{53} + 11992251412402642586454948v^{54} \\
&+ 14093734406768042617860546v^{55} + 16154939755233169917249815v^{56} \\
&+ 18062065264884658609927825v^{57} + 19698620890606501833935055v^{58} \\
&+ 20956986683280640928389866v^{59} + 21750009714684524653667914v^{60} \\
&+ 22020920210850484561094012v^{61} + 21750009714684524653667914v^{62} + \dots + v^{122}.
\end{aligned}$$

$$\begin{aligned}
P_{1,E_7}^{(2)}(v) = &(1+v^2)(137 + 2597v + 37024v^2 + 419921v^3 + 4077137v^4 + 34901534v^5 + 268811177v^6 + 1887255497v^7 \\
&+ 12196657853v^8 + 73094300214v^9 + 408614442098v^{10} + 2140990474296v^{11} + 10556715862964v^{12} \\
&+ 49151597538306v^{13} + 216730904533865v^{14} + 907396069059573v^{15} + 3615374924636545v^{16} \\
&+ 13736293007916068v^{17} + 49857926256318138v^{18} + 173164585658174276v^{19} + 576354715341079126v^{20} \\
&+ 1840835604225541174v^{21} + 5649018624246617909v^{22} + 16674709176092326437v^{23} \\
&+ 47394303096706259811v^{24} + 129836595656234291790v^{25} + 343131359707453293583v^{26} \\
&+ 875542039936399623515v^{27} + 2158650362542725175948v^{28} + 5146221002718346735055v^{29} \\
&+ 11870970192394860758359v^{30} + 26512271515436823962474v^{31} + 57361948999125457686102v^{32} \\
&+ 120296712068566252009120v^{33} + 244657061843538883914723v^{34} + 482772800850075541856889v^{35} \\
&+ 924703269912581018952608v^{36} + 1719956874446161848789295v^{37} + 3107832107172475492688890v^{38} \\
&+ 5457334614794588632799143v^{39} + 9316106764452824593797657v^{40} + 15465256039202958794051688v^{41}
\end{aligned}$$

$$\begin{aligned}
&+24973370386295921380753761v^{42} + 39238673033178891558314265v^{43} + 60003949644181883287996554v^{44} \\
&+ 89325800153382434388949763v^{45} + 129479490500199449940430040v^{46} + 182784785431420765743008945v^{47} \\
&+ 251347720581234682951528991v^{48} + 336728327510605097378060508v^{49} + 439563294320255738140535927v^{50} \\
&+ 559192119429259737303598283v^{51} + 693350791574808124298361559v^{52} + 838003231433873171234645238v^{53} \\
&+ 987373409206497489976018270v^{54} + 1134218396741161783456978908v^{55} \\
&+ 1270346407634715071774123344v^{56} + 1387339824356009883032251758v^{57} \\
&+ 1477400106011059794784293700v^{58} + 1534199301083129786878770830v^{59} \\
&+ 1553610262702054425407310320v^{60} + \dots + v^{120} .
\end{aligned}$$

Note $g_{0,E_7}^{(2)}$ agree with the two-instanton E_7 Hilbert series in [42].

G Refined BPS invariants

The refined BPS invariants are solved from the generalised blowup equations with the following initial input: the triple intersection numbers κ_{ijk} of divisors, the intersection numbers b_i^{GV} of divisors with $c_2(X)$ (these are two ingredients of Z^{cls}), the curve-divisor intersection matrix C , one unity r field with nonzero r_b , as well as the one-loop partition function $Z^{1\text{-loop}}$ and the bounds $j_{l,r}^{\text{max}}$.

In the case of $\mathfrak{n} = 5, 12$ models, there is no vanishing r -fields, the unity r do not have enough constraints on $Z^{1\text{-loop}}$. On the other hand, as seen in (2.36), $Z^{1\text{-loop}}$ is easily computed, we simply input $Z^{1\text{-loop}}$ for all the models.

The input of $j_{l,r}^{\text{max}}$ is strictly speaking also not necessary, as the bounds can be generated from the blowup equations with the other input data, but the inclusion of the bounds in the program makes the computation much faster. In any case, for $d_b \geq 1$, we observe an experimental formula for $j_{l,r}^{\text{max}}$. For F_4 , we observed for $d + d_b \leq 14$

$$\begin{aligned}
j_r^{\text{max}}(d, d_b) &= d(d_b + 1) - d_b(d_b - 1)/2, \\
j_l^{\text{max}}(d, d_b) &= (d - 1)(d_b - 1) - d_b(d_b - 1)/2 + \delta_{d_b,1} \lfloor d/(1 + d_\tau) \rfloor,
\end{aligned} \tag{G.1}$$

where d is the total degree of the fibers, d_b is the degree of the base, and d_τ is the first total degree of fibers when τ appears. For $E_{6,7,8}$, we observed for $d + d_b \leq 12 \leq d_\tau$

$$\begin{aligned}
j_r^{\text{max}}(d, d_b) &= d_b + d(d_b + 1), \\
j_l^{\text{max}}(d, d_b) &= (d - 1)(d_b - 1).
\end{aligned} \tag{G.2}$$

For F_4 model, we compute all the refined BPS invariants up to total degree 14, with 4777 non-vanishing. For $E_{6,7,8}$, we compute all the BPS invariants up to total degree 12, with 10383, 10491, 10068 non-vanishing respectively. We list part of the refined BPS invariants in the affine Lie algebra bases in tables 24–31, for complete lists, one can find them at [31]. It is worthwhile to point out that unlike the elliptic genus [24], the Weyl symmetry of gauge group G is not manifest in the refined BPS invariants. This is simply because the Weyl symmetry will change the sign of some Kähler parameters, while the refined BPS expansion is always in positive degrees. Note this should not be confused with the situation where the refined BPS invariants of E-strings do have manifest E_8 symmetry, in which case the E_8 is a global symmetry other than the gauge symmetry G we considered in this paper.

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0, 0, 0, 0, 0, 1)	(0, 0)	(0, 0, 1, 0, 0, 1)	(0, 1)
(0, 0, 1, 1, 0, 1)	(0, 0) \oplus (0, 1)	(0, 0, 1, 1, 1, 1)	(0, 0) \oplus (0, 1)
(0, 0, 1, 2, 0, 1)	(0, 1)	(0, 0, 1, 2, 1, 1)	(0, 0) \oplus (0, 1)
(0, 0, 1, 2, 2, 1)	(0, 1)	(0, 0, 2, 0, 0, 1)	(0, 2)
(0, 0, 2, 1, 0, 1)	(0, 1) \oplus (0, 2)	(0, 0, 2, 1, 1, 1)	(0, 1) \oplus (0, 2)
(0, 0, 2, 2, 0, 1)	(0, 0) \oplus (0, 1) \oplus (0, 2)	(0, 0, 2, 2, 1, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(0, 0, 2, 3, 0, 1)	(0, 1) \oplus (0, 2)	(0, 0, 3, 0, 0, 1)	(0, 3)
(0, 0, 3, 1, 0, 1)	(0, 2) \oplus (0, 3)	(0, 0, 3, 1, 1, 1)	(0, 2) \oplus (0, 3)
(0, 0, 3, 2, 0, 1)	(0, 1) \oplus (0, 2) \oplus (0, 3)	(0, 0, 4, 0, 0, 1)	(0, 4)
(0, 0, 4, 1, 0, 1)	(0, 3) \oplus (0, 4)	(0, 0, 5, 0, 0, 1)	(0, 5)
(0, 1, 0, 0, 0, 1)	(0, 1)	(0, 1, 1, 0, 0, 1)	(0, 0) \oplus (0, 1)
(0, 1, 1, 1, 0, 1)	(0, 0) \oplus (0, 1)	(0, 1, 1, 1, 1, 1)	(0, 0) \oplus (0, 1)
(0, 1, 1, 2, 0, 1)	(0, 0) \oplus (0, 1)	(0, 1, 1, 2, 1, 1)	(0, 0) \oplus (0, 1)
(0, 1, 2, 0, 0, 1)	(0, 1) \oplus (0, 2)	(0, 1, 2, 1, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(0, 1, 2, 1, 1, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)	(0, 1, 2, 2, 0, 1)	2(0, 0) \oplus 3(0, 1) \oplus (0, 2)
(0, 1, 3, 0, 0, 1)	(0, 2) \oplus (0, 3)	(0, 1, 3, 1, 0, 1)	(0, 1) \oplus 2(0, 2) \oplus (0, 3)
(0, 1, 4, 0, 0, 1)	(0, 3) \oplus (0, 4)	(0, 2, 0, 0, 0, 1)	(0, 2)
(0, 2, 1, 0, 0, 1)	(0, 1) \oplus (0, 2)	(0, 2, 1, 1, 0, 1)	(0, 1) \oplus (0, 2)
(0, 2, 1, 1, 1, 1)	(0, 1) \oplus (0, 2)	(0, 2, 1, 2, 0, 1)	(0, 1) \oplus (0, 2)
(0, 2, 2, 0, 0, 1)	(0, 0) \oplus (0, 1) \oplus (0, 2)	(0, 2, 2, 1, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(0, 2, 3, 0, 0, 1)	(0, 1) \oplus (0, 2) \oplus (0, 3)	(0, 3, 0, 0, 0, 1)	(0, 3)
(0, 3, 1, 0, 0, 1)	(0, 2) \oplus (0, 3)	(0, 3, 1, 1, 0, 1)	(0, 2) \oplus (0, 3)
(0, 3, 2, 0, 0, 1)	(0, 1) \oplus (0, 2) \oplus (0, 3)	(0, 4, 0, 0, 0, 1)	(0, 4)
(0, 4, 1, 0, 0, 1)	(0, 3) \oplus (0, 4)	(0, 5, 0, 0, 0, 1)	(0, 5)
(1, 1, 0, 0, 0, 1)	(0, 0) \oplus (0, 1)	(1, 1, 1, 0, 0, 1)	(0, 0) \oplus (0, 1)
(1, 1, 1, 1, 0, 1)	(0, 0) \oplus (0, 1)	(1, 1, 1, 1, 1, 1)	(0, 0) \oplus (0, 1)
(1, 1, 1, 2, 0, 1)	(0, 0) \oplus (0, 1)	(1, 1, 2, 0, 0, 1)	(0, 1) \oplus (0, 2)
(1, 1, 2, 1, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)	(1, 1, 3, 0, 0, 1)	(0, 2) \oplus (0, 3)
(1, 2, 0, 0, 0, 1)	(0, 1) \oplus (0, 2)	(1, 2, 1, 0, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(1, 2, 1, 1, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)	(1, 2, 2, 0, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(1, 3, 0, 0, 0, 1)	(0, 2) \oplus (0, 3)	(1, 3, 1, 0, 0, 1)	(0, 1) \oplus 2(0, 2) \oplus (0, 3)
(1, 4, 0, 0, 0, 1)	(0, 3) \oplus (0, 4)	(2, 1, 0, 0, 0, 1)	(0, 1)
(2, 2, 0, 0, 0, 1)	(0, 0) \oplus (0, 1) \oplus (0, 2)	(2, 2, 1, 0, 0, 1)	(0, 0) \oplus 2(0, 1) \oplus (0, 2)
(2, 3, 0, 0, 0, 1)	(0, 1) \oplus (0, 2) \oplus (0, 3)	(3, 1, 0, 0, 0, 1)	(0, 2)
(3, 2, 0, 0, 0, 1)	(0, 1) \oplus (0, 2)	(4, 1, 0, 0, 0, 1)	(0, 3)
(0, 0, 2, 0, 0, 2)	(0, 5/2)	(0, 0, 2, 1, 0, 2)	(0, 3/2) \oplus (0, 5/2)
(0, 0, 2, 1, 1, 2)	(0, 3/2) \oplus (0, 5/2)	(0, 0, 2, 2, 0, 2)	(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)
(0, 0, 3, 0, 0, 2)	(0, 5/2) \oplus (0, 7/2) \oplus (1/2, 4)	(0, 0, 3, 1, 0, 2)	(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)
(0, 0, 4, 0, 0, 2)	(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus (1, 11/2)	(0, 1, 2, 0, 0, 2)	(0, 3/2) \oplus (0, 5/2)
(0, 1, 2, 1, 0, 2)	(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)	(0, 1, 3, 0, 0, 2)	(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)
(0, 2, 0, 0, 0, 2)	(0, 5/2)	(0, 2, 1, 0, 0, 2)	(0, 3/2) \oplus (0, 5/2)
(0, 2, 1, 1, 0, 2)	(0, 3/2) \oplus (0, 5/2)	(0, 2, 2, 0, 0, 2)	2(0, 1/2) \oplus 2(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)
(0, 3, 0, 0, 0, 2)	(0, 5/2) \oplus (0, 7/2) \oplus (1/2, 4)	(0, 3, 1, 0, 0, 2)	(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)
(0, 4, 0, 0, 0, 2)	(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus (1, 11/2)	(1, 1, 2, 0, 0, 2)	(0, 3/2) \oplus (0, 5/2)

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(1, 2, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(1, 2, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(1, 3, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(2, 2, 0, 0, 0, 2)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 0, 3, 0, 0, 3)	$(0, 3) \oplus (1/2, 9/2)$	(0, 3, 0, 0, 0, 3)	$(0, 3) \oplus (1/2, 9/2)$

Table 24. Refined BPS invariants of 6d F_4 minimal SCFT.

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0, 0, 0, 0, 0, 0, 1)	(0, 1/2)	(0, 1, 0, 0, 0, 0, 0, 1)	(0, 1/2)
(0, 2, 0, 0, 0, 0, 0, 1)	(0, 3/2)	(0, 3, 0, 0, 0, 0, 0, 1)	(0, 5/2)
(0, 4, 0, 0, 0, 0, 0, 1)	(0, 7/2)	(0, 5, 0, 0, 0, 0, 0, 1)	(0, 9/2)
(0, 0, 0, 0, 0, 1, 0, 1)	(0, 1/2)	(0, 0, 0, 0, 0, 1, 1, 1)	(0, 1/2)
(0, 0, 0, 0, 0, 2, 0, 1)	(0, 3/2)	(0, 0, 0, 0, 0, 2, 1, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 0, 0, 0, 0, 2, 2, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 0, 0, 0, 0, 2, 3, 1)	(0, 3/2)
(0, 0, 0, 0, 0, 3, 0, 1)	(0, 5/2)	(0, 0, 0, 0, 0, 3, 1, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 0, 3, 2, 1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 0, 0, 4, 0, 1)	(0, 7/2)
(0, 0, 0, 0, 0, 4, 1, 1)	$(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 0, 5, 0, 1)	(0, 9/2)
(0, 0, 0, 0, 1, 0, 0, 1)	(0, 3/2)	(0, 1, 0, 0, 1, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 2, 0, 0, 1, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 3, 0, 0, 1, 0, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 4, 0, 0, 1, 0, 0, 1)	$(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 1, 1, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 1, 0, 0, 1, 1, 0, 1)	$2(0, 1/2) \oplus (0, 3/2)$	(0, 2, 0, 0, 1, 1, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 3, 0, 0, 1, 1, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 0, 1, 1, 1, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 1, 0, 0, 1, 1, 1, 1)	$2(0, 1/2) \oplus (0, 3/2)$	(0, 2, 0, 0, 1, 1, 1, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 0, 0, 0, 1, 2, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 1, 0, 0, 1, 2, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 0, 0, 0, 1, 2, 1, 1)	$2(0, 1/2) \oplus (0, 3/2)$	(0, 1, 0, 0, 1, 2, 1, 1)	$2(0, 1/2) \oplus (0, 3/2)$
(0, 0, 0, 0, 1, 2, 2, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 0, 0, 0, 1, 3, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 1, 0, 0, 1, 3, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 0, 1, 3, 1, 1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 1, 4, 0, 1)	$(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 2, 0, 0, 1)	(0, 5/2)
(0, 1, 0, 0, 2, 0, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$	(0, 2, 0, 0, 2, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 3, 0, 0, 2, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 0, 2, 1, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 1, 0, 0, 2, 1, 0, 1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 2, 0, 0, 2, 1, 0, 1)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 2, 1, 1, 1)	$(0, 3/2) \oplus (0, 5/2)$	(0, 1, 0, 0, 2, 1, 1, 1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 2, 2, 0, 1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$	(0, 1, 0, 0, 2, 2, 0, 1)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 2, 2, 1, 1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 0, 2, 3, 0, 1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 3, 0, 0, 1)	(0, 7/2)	(0, 1, 0, 0, 3, 0, 0, 1)	$(0, 5/2) \oplus (0, 7/2)$
(0, 2, 0, 0, 3, 0, 0, 1)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 3, 1, 0, 1)	$(0, 5/2) \oplus (0, 7/2)$
(0, 1, 0, 0, 3, 1, 0, 1)	$(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 3, 1, 1, 1)	$(0, 5/2) \oplus (0, 7/2)$
(0, 0, 0, 0, 3, 2, 0, 1)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 4, 0, 0, 1)	(0, 9/2)
(0, 1, 0, 0, 4, 0, 0, 1)	$(0, 7/2) \oplus (0, 9/2)$	(0, 0, 0, 0, 4, 1, 0, 1)	$(0, 7/2) \oplus (0, 9/2)$
(0, 0, 0, 0, 5, 0, 0, 1)	(0, 11/2)	(0, 0, 0, 1, 0, 0, 0, 1)	(0, 1/2)
(0, 0, 0, 1, 1, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 1, 0, 1, 1, 0, 0, 1)	$2(0, 1/2) \oplus (0, 3/2)$
(0, 2, 0, 1, 1, 0, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 3, 0, 1, 1, 0, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 1, 1, 0, 1)	$2(0, 1/2) \oplus (0, 3/2)$	(0, 1, 0, 1, 1, 1, 0, 1)	$3(0, 1/2) \oplus (0, 3/2)$
(0, 2, 0, 1, 1, 1, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 0, 0, 1, 1, 1, 1, 1)	$2(0, 1/2) \oplus (0, 3/2)$
(0, 1, 0, 1, 1, 1, 1, 1)	$3(0, 1/2) \oplus (0, 3/2)$	(0, 0, 0, 1, 1, 2, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$
(0, 1, 0, 1, 1, 2, 0, 1)	$(0, 1/2) \oplus (0, 3/2)$	(0, 0, 0, 1, 1, 2, 1, 1)	$2(0, 1/2) \oplus (0, 3/2)$
(0, 0, 0, 1, 1, 3, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 2, 0, 0, 1)	$(0, 3/2) \oplus (0, 5/2)$
(0, 1, 0, 1, 2, 0, 0, 1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 2, 0, 1, 2, 0, 0, 1)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,1,2,1,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,1,0,1,2,1,0,1)	$4(0,1/2) \oplus 4(0,3/2) \oplus (0,5/2)$
(0,0,0,1,2,1,1,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,2,2,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,1,3,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,1,0,1,3,0,0,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,1,3,1,0,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,1,4,0,0,1)	$(0,7/2) \oplus (0,9/2)$
(0,0,0,2,0,0,0,1)	$(0,3/2)$	(0,0,0,2,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,1,0,2,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,2,1,1,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,1,0,2,1,1,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,2,1,1,1,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,2,2,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,1,0,2,2,0,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,2,2,1,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,2,3,0,0,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$
(0,0,0,3,0,0,0,1)	$(0,5/2)$	(0,0,0,3,1,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,1,0,3,1,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,3,1,1,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,3,2,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,4,0,0,0,1)	$(0,7/2)$
(0,0,0,4,1,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,5,0,0,0,1)	$(0,9/2)$
(0,0,1,1,0,0,0,1)	$(0,1/2)$	(0,0,1,1,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,2,1,1,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,1,1,1,1,1,0,1)	$3(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,1,1,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,1,1,2,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,1,1,2,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,1,1,1,2,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,1,1,2,1,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,1,1,3,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,1,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,1,2,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,1,1,2,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,2,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,1,2,2,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,1,3,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,1,3,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,1,4,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,2,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,2,2,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,2,3,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,3,2,0,0,0,1)	$(0,3/2)$
(1,1,0,0,0,0,0,1)	$(0,1/2)$	(1,2,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(1,3,0,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(1,4,0,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(1,1,0,0,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(1,2,0,0,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(1,3,0,0,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(1,1,0,0,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$
(1,2,0,0,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$	(1,1,0,0,1,1,1,1)	$2(0,1/2) \oplus (0,3/2)$
(1,1,0,0,1,2,0,1)	$(0,1/2) \oplus (0,3/2)$	(1,1,0,0,2,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(1,2,0,0,2,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(1,1,0,0,2,1,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(1,1,0,0,3,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(1,1,0,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(1,2,0,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(1,1,0,1,1,1,0,1)	$3(0,1/2) \oplus (0,3/2)$
(1,1,0,1,2,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(1,1,0,2,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(1,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(2,2,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(2,3,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(2,2,0,0,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(3,2,0,0,0,0,0,1)	$(0,3/2)$	(0,3,0,0,0,0,0,2)	$(0,5/2)$
(0,4,0,0,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$	(0,0,0,0,3,0,2)	$(0,5/2)$
(0,0,0,0,0,3,1,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,4,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$
(0,0,0,0,1,0,0,2)	$(0,5/2)$	(0,1,0,0,1,0,0,2)	$(0,3/2) \oplus (0,5/2)$
(0,2,0,0,1,0,0,2)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,3,0,0,1,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,0,1,1,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,1,0,0,1,1,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,2,0,0,1,1,0,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,1,1,2)	$(0,3/2) \oplus (0,5/2)$
(0,1,0,0,1,1,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,0,2)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,1,0,0,1,2,0,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,3,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,2,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0, 1, 0, 0, 2, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 2, 0, 0, 2, 0, 0, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 0, 2, 1, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 1, 0, 0, 2, 1, 0, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 0, 2, 1, 1, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 0, 2, 2, 0, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 0, 3, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus (1, 11/2)$	(0, 1, 0, 0, 3, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$
(0, 0, 0, 0, 3, 1, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0, 0, 0, 0, 4, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus 2(0, 11/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus 2(1/2, 6) \oplus (1, 11/2) \oplus (1, 13/2) \oplus (3/2, 7)$
(0, 0, 0, 1, 1, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 1, 0, 1, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 2, 0, 1, 1, 0, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 1, 1, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 1, 0, 1, 1, 1, 0, 2)	$3(0, 1/2) \oplus 3(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 1, 1, 1, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 1, 2, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 2, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 1, 0, 1, 2, 0, 0, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 1, 2, 1, 0, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 1, 3, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0, 0, 0, 2, 1, 0, 0, 2)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 1, 0, 2, 1, 0, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 2, 1, 1, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 2, 2, 0, 0, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 3, 0, 0, 0, 2)	$(0, 5/2)$
(0, 0, 0, 3, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 4, 0, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus (1/2, 4)$
(0, 0, 1, 1, 1, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 1, 1, 1, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 1, 1, 1, 1, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 1, 2, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 1, 2, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 3, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$
(1, 3, 0, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(1, 1, 0, 0, 1, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$
(1, 2, 0, 0, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(1, 1, 0, 0, 1, 1, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(1, 1, 0, 0, 2, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(1, 1, 0, 1, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 0, 1, 0, 0, 3)	$(0, 7/2)$	(0, 1, 0, 0, 1, 0, 0, 3)	$(0, 5/2) \oplus (0, 7/2)$
(0, 2, 0, 0, 1, 0, 0, 3)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 1, 1, 0, 3)	$(0, 5/2) \oplus (0, 7/2)$
(0, 1, 0, 0, 1, 1, 0, 3)	$(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 1, 1, 1, 3)	$(0, 5/2) \oplus (0, 7/2)$
(0, 0, 0, 0, 1, 2, 0, 3)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 0, 2, 0, 0, 3)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus (1, 11/2)$
(0, 1, 0, 0, 2, 0, 0, 3)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0, 0, 0, 0, 2, 1, 0, 3)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,3,0,0,3)	$(0, 3/2) \oplus (0, 5/2) \oplus 3(0, 7/2) \oplus 3(0, 9/2) \oplus 4(0, 11/2) \oplus (1/2, 3) \oplus 2(1/2, 4) \oplus 3(1/2, 5) \oplus 3(1/2, 6) \oplus (1/2, 7) \oplus (1, 9/2) \oplus 2(1, 11/2) \oplus 3(1, 13/2) \oplus (3/2, 6) \oplus (3/2, 7) \oplus (2, 15/2)$	(0,0,0,1,1,0,0,3)	$(0, 5/2) \oplus (0, 7/2)$
(0,1,0,1,1,0,0,3)	$(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0,0,0,1,1,1,0,3)	$(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$
(0,0,0,1,2,0,0,3)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0,0,0,2,1,0,0,3)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$
(0,0,1,1,1,0,0,3)	$(0, 5/2) \oplus (0, 7/2)$	(1,1,0,0,1,0,0,3)	$(0, 5/2) \oplus (0, 7/2)$
(0,0,0,0,1,0,0,4)	$(0, 9/2)$	(0,1,0,0,1,0,0,4)	$(0, 7/2) \oplus (0, 9/2)$
(0,0,0,0,1,1,0,4)	$(0, 7/2) \oplus (0, 9/2)$	(0,0,0,0,2,0,0,4)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus 2(0, 11/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus 2(1/2, 6) \oplus (1, 11/2) \oplus (1, 13/2) \oplus (3/2, 7)$
(0,0,0,1,1,0,0,4)	$(0, 7/2) \oplus (0, 9/2)$	(0,0,0,0,1,0,0,5)	$(0, 11/2)$

Table 25. Refined BPS invariants of 6d E_6 minimal SCFT.

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,0,0,0,0,1)	$(0, 1/2)$	(0,0,0,0,0,0,0,1,1)	$(0, 1/2)$
(0,0,0,0,0,0,0,2,1)	$(0, 3/2)$	(0,0,0,0,0,0,0,3,1)	$(0, 5/2)$
(0,0,0,0,0,0,0,4,1)	$(0, 7/2)$	(0,0,0,0,0,0,0,5,1)	$(0, 9/2)$
(0,0,0,0,1,0,0,0,1)	$(0, 1/2)$	(0,0,0,0,1,1,0,0,1)	$(0, 1/2)$
(0,0,0,0,1,1,1,0,1)	$(0, 1/2)$	(0,0,0,0,2,0,0,0,1)	$(0, 3/2)$
(0,0,0,0,2,1,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$	(0,0,0,0,2,1,1,0,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,0,2,2,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$	(0,0,0,0,2,2,1,0,1)	$2(0, 1/2) \oplus (0, 3/2)$
(0,0,0,0,2,3,0,0,1)	$(0, 3/2)$	(0,0,0,0,3,0,0,0,1)	$(0, 5/2)$
(0,0,0,0,3,1,0,0,1)	$(0, 3/2) \oplus (0, 5/2)$	(0,0,0,0,3,1,1,0,1)	$(0, 3/2) \oplus (0, 5/2)$
(0,0,0,0,3,2,0,0,1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$	(0,0,0,0,4,0,0,0,1)	$(0, 7/2)$
(0,0,0,0,4,1,0,0,1)	$(0, 5/2) \oplus (0, 7/2)$	(0,0,0,0,5,0,0,0,1)	$(0, 9/2)$
(0,0,0,1,0,0,0,0,1)	$(0, 3/2)$	(0,0,0,1,0,0,0,1,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,0,0,0,2,1)	$(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,0,0,0,3,1)	$(0, 3/2) \oplus (0, 5/2)$
(0,0,0,1,0,0,0,4,1)	$(0, 5/2) \oplus (0, 7/2)$	(0,0,0,1,1,0,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,1,0,0,1,1)	$2(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,1,0,0,2,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,1,0,0,3,1)	$(0, 3/2) \oplus (0, 5/2)$	(0,0,0,1,1,1,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,1,1,0,1,1)	$2(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,1,1,0,2,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,1,1,1,0,1)	$(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,1,1,1,1,1)	$2(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,2,0,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,2,0,0,1,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,2,1,0,0,1)	$2(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,2,1,0,1,1)	$2(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,2,1,1,0,1)	$2(0, 1/2) \oplus (0, 3/2)$	(0,0,0,1,2,2,0,0,1)	$(0, 1/2) \oplus (0, 3/2)$
(0,0,0,1,3,0,0,0,1)	$(0, 3/2) \oplus (0, 5/2)$	(0,0,0,1,3,0,0,1,1)	$(0, 3/2) \oplus (0, 5/2)$
(0,0,0,1,3,1,0,0,1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0,0,0,1,4,0,0,0,1)	$(0, 5/2) \oplus (0, 7/2)$
(0,0,0,2,0,0,0,0,1)	$(0, 5/2)$	(0,0,0,2,0,0,0,1,1)	$(0, 3/2) \oplus (0, 5/2)$
(0,0,0,2,0,0,0,2,1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$	(0,0,0,2,0,0,0,3,1)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0,0,0,2,1,0,0,0,1)	$(0, 3/2) \oplus (0, 5/2)$	(0,0,0,2,1,0,0,1,1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0,0,0,2,1,0,0,2,1)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0,0,0,2,1,1,0,0,1)	$(0, 3/2) \oplus (0, 5/2)$
(0,0,0,2,1,1,0,1,1)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0,0,0,2,1,1,1,0,1)	$(0, 3/2) \oplus (0, 5/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,2,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,2,2,0,0,1,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,2,2,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,2,3,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,0,3,0,0,0,0,1)	$(0,7/2)$	(0,0,0,3,0,0,0,1,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,3,0,0,0,2,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,0,3,1,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,3,1,0,0,1,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,3,1,1,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,3,2,0,0,0,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,0,4,0,0,0,0,1)	$(0,9/2)$
(0,0,0,4,0,0,0,1,1)	$(0,7/2) \oplus (0,9/2)$	(0,0,0,4,1,0,0,0,1)	$(0,7/2) \oplus (0,9/2)$
(0,0,0,5,0,0,0,0,1)	$(0,11/2)$	(0,0,1,0,0,0,0,0,1)	$(0,1/2)$
(0,0,1,1,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,1,1,0,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,1,1,0,0,0,2,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,1,1,0,0,0,3,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,1,1,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,1,1,0,0,1,1)	$3(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,0,0,2,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,1,0,1,1)	$3(0,1/2) \oplus (0,3/2)$	(0,0,1,1,1,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,1,1,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,1,1,2,0,0,1,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,1,1,2,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,1,3,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,1,2,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,1,2,0,0,0,1,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,1,2,0,0,0,2,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,1,2,1,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,1,2,1,0,0,1,1)	$4(0,1/2) \oplus 4(0,3/2) \oplus (0,5/2)$	(0,0,1,2,1,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,1,2,2,0,0,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,1,3,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,1,3,0,0,0,1,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,1,3,1,0,0,0,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,1,4,0,0,0,0,1)	$(0,7/2) \oplus (0,9/2)$	(0,0,2,0,0,0,0,0,1)	$(0,3/2)$
(0,0,2,1,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,2,1,0,0,0,1,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,2,1,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,2,1,1,0,0,1,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,2,1,1,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,2,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,2,2,0,0,0,1,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,2,2,1,0,0,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,2,3,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,3,0,0,0,0,0,1)	$(0,5/2)$
(0,0,3,1,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,3,1,0,0,0,1,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,3,1,1,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,3,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,4,0,0,0,0,0,1)	$(0,7/2)$	(0,0,4,1,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,5,0,0,0,0,0,1)	$(0,9/2)$	(0,1,1,0,0,0,0,0,1)	$(0,1/2)$
(0,1,1,1,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,1,1,1,0,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$
(0,1,1,1,0,0,0,2,1)	$(0,1/2) \oplus (0,3/2)$	(0,1,1,1,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,1,1,1,1,0,0,1,1)	$3(0,1/2) \oplus (0,3/2)$	(0,1,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,1,1,1,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,1,1,2,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,1,1,2,0,0,0,1,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,1,1,2,1,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,1,1,3,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,1,2,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,1,2,1,0,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,1,2,1,0,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$
(0,1,2,1,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,1,2,2,0,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,1,3,0,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,1,3,1,0,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,1,4,0,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,2,2,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,2,2,1,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,2,3,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,3,2,0,0,0,0,0,1)	$(0,3/2)$	(1,1,1,0,0,0,0,0,1)	$(0,1/2)$
(1,1,1,1,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(1,1,1,1,0,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$
(1,1,1,1,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(1,1,1,2,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(1,1,2,0,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(1,1,2,1,0,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(1,1,3,0,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(1,2,2,0,0,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,0,0,0,0,3,2)	$(0,5/2)$	(0,0,0,0,0,0,0,4,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$
(0,0,0,0,3,0,0,0,2)	$(0,5/2)$	(0,0,0,0,3,1,0,0,2)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,4,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$	(0,0,0,1,0,0,0,0,2)	$(0,5/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0, 0, 0, 1, 0, 0, 0, 1, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 0, 0, 0, 2, 2)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 0, 0, 0, 3, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 1, 1, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 1, 0, 0, 1, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 1, 0, 0, 2, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 1, 1, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 1, 1, 0, 1, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 1, 1, 1, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 2, 0, 0, 0, 2)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 2, 0, 0, 1, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 2, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 0, 1, 3, 0, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 2, 0, 0, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus (1/2, 4)$
(0, 0, 0, 2, 0, 0, 0, 1, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 2, 0, 0, 0, 2, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 2, 1, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 2, 1, 0, 0, 1, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 2, 1, 1, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 0, 2, 2, 0, 0, 0, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 0, 3, 0, 0, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus (1, 11/2)$	(0, 0, 0, 3, 0, 0, 0, 1, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$
(0, 0, 0, 3, 1, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0, 0, 0, 4, 0, 0, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus 2(0, 9/2) \oplus 2(0, 11/2) \oplus (1/2, 4) \oplus (1/2, 5) \oplus 2(1/2, 6) \oplus (1, 11/2) \oplus (1, 13/2) \oplus (3/2, 7)$
(0, 0, 1, 1, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 1, 0, 0, 0, 1, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 1, 1, 0, 0, 0, 2, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 1, 1, 0, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 1, 1, 1, 0, 0, 1, 2)	$3(0, 1/2) \oplus 3(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 1, 1, 1, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 1, 1, 2, 0, 0, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 1, 2, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 0, 1, 2, 0, 0, 0, 1, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$	(0, 0, 1, 2, 1, 0, 0, 0, 2)	$(0, 1/2) \oplus 5(0, 3/2) \oplus 7(0, 5/2) \oplus 3(0, 7/2) \oplus (1/2, 2) \oplus 2(1/2, 3) \oplus (1/2, 4)$
(0, 0, 1, 3, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 5(0, 7/2) \oplus 3(0, 9/2) \oplus (1/2, 3) \oplus 3(1/2, 4) \oplus 2(1/2, 5) \oplus (1, 9/2) \oplus (1, 11/2)$	(0, 0, 2, 1, 0, 0, 0, 0, 2)	$(0, 1/2) \oplus (0, 3/2) \oplus (0, 5/2)$
(0, 0, 2, 1, 0, 0, 0, 1, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 0, 2, 1, 1, 0, 0, 0, 2)	$2(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 0, 2, 2, 0, 0, 0, 0, 2)	$(0, 1/2) \oplus 3(0, 3/2) \oplus 4(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 2) \oplus (1/2, 3) \oplus (1/2, 4)$	(0, 0, 3, 0, 0, 0, 0, 0, 2)	$(0, 5/2)$
(0, 0, 3, 1, 0, 0, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$	(0, 0, 4, 0, 0, 0, 0, 0, 2)	$(0, 5/2) \oplus (0, 7/2) \oplus (1/2, 4)$
(0, 1, 1, 1, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 1, 1, 1, 0, 0, 0, 1, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$
(0, 1, 1, 1, 1, 0, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 1, 1, 2, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus 3(0, 5/2) \oplus 2(0, 7/2) \oplus (1/2, 3) \oplus (1/2, 4)$
(0, 1, 2, 1, 0, 0, 0, 0, 2)	$(0, 1/2) \oplus 2(0, 3/2) \oplus (0, 5/2)$	(0, 1, 3, 0, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$
(1, 1, 1, 1, 0, 0, 0, 0, 2)	$(0, 3/2) \oplus (0, 5/2)$	(0, 0, 0, 1, 0, 0, 0, 0, 3)	$(0, 7/2)$
(0, 0, 0, 1, 0, 0, 0, 1, 3)	$(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 1, 0, 0, 0, 2, 3)	$(0, 3/2) \oplus (0, 5/2) \oplus (0, 7/2)$
(0, 0, 0, 1, 1, 0, 0, 0, 3)	$(0, 5/2) \oplus (0, 7/2)$	(0, 0, 0, 1, 1, 0, 0, 1, 3)	$(0, 3/2) \oplus 2(0, 5/2) \oplus (0, 7/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,1,1,1,0,0,3)	(0,5/2)⊕(0,7/2)	(0,0,0,1,2,0,0,0,3)	(0,3/2)⊕(0,5/2)⊕(0,7/2)
(0,0,0,2,0,0,0,0,3)	(0,5/2)⊕(0,7/2)⊕2(0,9/2)⊕ (1/2,4)⊕(1/2,5)⊕(1,11/2)	(0,0,0,2,0,0,0,1,3)	(0,3/2)⊕3(0,5/2)⊕5(0,7/2)⊕ 3(0,9/2)⊕(1/2,3)⊕3(1/2,4)⊕ 2(1/2,5)⊕(1,9/2)⊕(1,11/2)
(0,0,0,2,1,0,0,0,3)	(0,3/2)⊕3(0,5/2)⊕5(0,7/2)⊕ 3(0,9/2)⊕(1/2,3)⊕3(1/2,4)⊕ 2(1/2,5)⊕(1,9/2)⊕(1,11/2)	(0,0,0,3,0,0,0,0,3)	(0,3/2)⊕(0,5/2)⊕3(0,7/2)⊕ 3(0,9/2)⊕4(0,11/2)⊕ (1/2,3)⊕2(1/2,4)⊕3(1/2,5)⊕ 3(1/2,6)⊕(1/2,7)⊕(1,9/2)⊕ 2(1,11/2)⊕3(1,13/2)⊕ (3/2,6)⊕(3/2,7)⊕(2,15/2)
(0,0,1,1,0,0,0,0,3)	(0,5/2)⊕(0,7/2)	(0,0,1,1,0,0,0,1,3)	(0,3/2)⊕2(0,5/2)⊕(0,7/2)
(0,0,1,1,1,0,0,0,3)	(0,3/2)⊕2(0,5/2)⊕(0,7/2)	(0,0,1,2,0,0,0,0,3)	(0,3/2)⊕3(0,5/2)⊕5(0,7/2)⊕ 3(0,9/2)⊕(1/2,3)⊕3(1/2,4)⊕ 2(1/2,5)⊕(1,9/2)⊕(1,11/2)
(0,0,2,1,0,0,0,0,3)	(0,3/2)⊕(0,5/2)⊕(0,7/2)	(0,1,1,1,0,0,0,0,3)	(0,5/2)⊕(0,7/2)
(0,0,0,1,0,0,0,0,4)	(0,9/2)	(0,0,0,1,0,0,0,1,4)	(0,7/2)⊕(0,9/2)
(0,0,0,1,1,0,0,0,4)	(0,7/2)⊕(0,9/2)	(0,0,0,2,0,0,0,0,4)	(0,5/2)⊕(0,7/2)⊕2(0,9/2)⊕ 2(0,11/2)⊕(1/2,4)⊕ (1/2,5)⊕2(1/2,6)⊕(1,11/2)⊕ (1,13/2)⊕(3/2,7)
(0,0,1,1,0,0,0,0,4)	(0,7/2)⊕(0,9/2)	(0,0,0,1,0,0,0,0,5)	(0,11/2)

Table 26. Refined BPS invariants of 6d E_7 minimal SCFT.

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,0,0,0,0,1)	(0,1/2)	(0,0,0,0,0,0,0,0,1,1)	(0,1/2)
(0,0,0,0,0,0,0,0,2,1)	(0,3/2)	(0,0,0,0,0,0,0,0,3,1)	(0,5/2)
(0,0,0,0,0,0,0,0,4,1)	(0,7/2)	(0,0,0,0,0,0,0,0,5,1)	(0,9/2)
(0,0,0,0,0,0,1,0,0,1)	(0,1/2)	(0,0,0,0,0,0,1,1,0,1)	(0,1/2)
(0,0,0,0,0,0,2,0,0,1)	(0,3/2)	(0,0,0,0,0,0,2,1,0,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,0,2,2,0,1)	(0,1/2)⊕(0,3/2)	(0,0,0,0,0,0,2,3,0,1)	(0,3/2)
(0,0,0,0,0,0,3,0,0,1)	(0,5/2)	(0,0,0,0,0,0,3,1,0,1)	(0,3/2)⊕(0,5/2)
(0,0,0,0,0,0,3,2,0,1)	(0,1/2)⊕(0,3/2)⊕(0,5/2)	(0,0,0,0,0,0,4,0,0,1)	(0,7/2)
(0,0,0,0,0,0,4,1,0,1)	(0,5/2)⊕(0,7/2)	(0,0,0,0,0,0,5,0,0,1)	(0,9/2)
(0,0,0,0,0,1,0,0,0,1)	(0,3/2)	(0,0,0,0,0,1,0,0,1,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,0,0,2,1)	(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,0,0,3,1)	(0,3/2)⊕(0,5/2)
(0,0,0,0,0,1,0,0,4,1)	(0,5/2)⊕(0,7/2)	(0,0,0,0,0,1,1,0,0,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,1,0,1,1)	2(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,1,0,2,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,1,0,3,1)	(0,3/2)⊕(0,5/2)	(0,0,0,0,0,1,1,1,0,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,1,1,1,1)	2(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,1,1,2,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,2,0,0,1)	(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,2,0,1,1)	(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,2,1,0,1)	2(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,2,1,1,1)	2(0,1/2)⊕(0,3/2)
(0,0,0,0,0,1,2,2,0,1)	(0,1/2)⊕(0,3/2)	(0,0,0,0,0,1,3,0,0,1)	(0,3/2)⊕(0,5/2)
(0,0,0,0,0,1,3,0,1,1)	(0,3/2)⊕(0,5/2)	(0,0,0,0,0,1,3,1,0,1)	(0,1/2)⊕2(0,3/2)⊕(0,5/2)
(0,0,0,0,0,1,4,0,0,1)	(0,5/2)⊕(0,7/2)	(0,0,0,0,0,2,0,0,0,1)	(0,5/2)
(0,0,0,0,0,2,0,0,1,1)	(0,3/2)⊕(0,5/2)	(0,0,0,0,0,2,0,0,2,1)	(0,1/2)⊕(0,3/2)⊕(0,5/2)
(0,0,0,0,0,2,0,0,3,1)	(0,1/2)⊕(0,3/2)⊕(0,5/2)	(0,0,0,0,0,2,1,0,0,1)	(0,3/2)⊕(0,5/2)
(0,0,0,0,0,2,1,0,1,1)	(0,1/2)⊕2(0,3/2)⊕(0,5/2)	(0,0,0,0,0,2,1,0,2,1)	2(0,1/2)⊕2(0,3/2)⊕(0,5/2)
(0,0,0,0,0,2,1,1,0,1)	(0,3/2)⊕(0,5/2)	(0,0,0,0,0,2,1,1,1,1)	(0,1/2)⊕2(0,3/2)⊕(0,5/2)

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,0,2,2,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,2,2,0,1,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,2,2,1,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,2,3,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,3,0,0,0,1)	$(0,7/2)$	(0,0,0,0,0,3,0,0,1,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,3,0,0,2,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,3,1,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,3,1,0,1,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,3,1,1,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,3,2,0,0,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,4,0,0,0,1)	$(0,9/2)$
(0,0,0,0,0,4,0,0,1,1)	$(0,7/2) \oplus (0,9/2)$	(0,0,0,0,0,4,1,0,0,1)	$(0,7/2) \oplus (0,9/2)$
(0,0,0,0,0,5,0,0,0,1)	$(0,11/2)$	(0,0,0,0,1,0,0,0,0,1)	$(0,1/2)$
(0,0,0,0,1,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,0,1,1,0,0,2,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,0,0,3,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,1,0,1,1)	$3(0,1/2) \oplus (0,3/2)$
(0,0,0,0,1,1,1,0,2,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,0,1,1,1,1,1,1)	$3(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,2,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,0,1,1,2,0,1,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,1,1,2,1,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,0,1,1,3,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,2,0,0,1,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,0,0,2,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,2,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,1,0,1,1)	$4(0,1/2) \oplus 4(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,2,1,1,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,2,0,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,3,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,1,3,0,0,1,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,0,1,3,1,0,0,1)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,1,4,0,0,0,1)	$(0,7/2) \oplus (0,9/2)$
(0,0,0,0,2,0,0,0,0,1)	$(0,3/2)$	(0,0,0,0,2,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,0,2,1,0,0,1,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,2,1,1,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,0,2,1,1,0,1,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,0,2,1,1,1,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,0,2,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,0,2,2,0,0,1,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,2,2,1,0,0,1)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,2,3,0,0,0,1)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$
(0,0,0,0,3,0,0,0,0,1)	$(0,5/2)$	(0,0,0,0,3,1,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,3,1,0,0,1,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,3,1,1,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,3,2,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,0,4,0,0,0,0,1)	$(0,7/2)$
(0,0,0,0,4,1,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,5,0,0,0,0,1)	$(0,9/2)$
(0,0,0,1,1,0,0,0,0,1)	$(0,1/2)$	(0,0,0,1,1,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,1,1,1,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,0,1,1,1,0,0,2,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,0,1,1,1,1,0,1,1)	$3(0,1/2) \oplus (0,3/2)$
(0,0,0,1,1,1,1,1,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,0,1,1,1,2,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,1,1,2,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,1,1,2,0,0,1,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,1,1,2,1,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,1,3,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,1,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,1,2,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,1,2,1,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,0,1,2,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,0,1,2,2,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,3,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,1,3,1,0,0,0,1)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,4,0,0,0,0,1)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,2,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,0,0,2,2,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,0,2,3,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,3,2,0,0,0,0,1)	$(0,3/2)$
(0,0,1,1,1,0,0,0,0,1)	$(0,1/2)$	(0,0,1,1,1,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,1,0,0,1,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,1,1,1,1,0,0,1)	$2(0,1/2) \oplus (0,3/2)$
(0,0,1,1,1,2,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$	(0,0,1,1,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(0,0,1,1,2,1,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,0,1,1,3,0,0,0,0,1)	$(0,3/2) \oplus (0,5/2)$
(0,0,1,2,2,0,0,0,0,1)	$2(0,1/2) \oplus (0,3/2)$	(0,1,1,1,1,0,0,0,0,1)	$(0,1/2)$
(0,1,1,1,1,1,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$	(0,1,1,1,2,0,0,0,0,1)	$(0,1/2) \oplus (0,3/2)$
(1,1,1,1,1,0,0,0,0,1)	$(0,1/2)$	(0,0,0,0,0,0,0,0,3,2)	$(0,5/2)$
(0,0,0,0,0,0,0,0,4,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$	(0,0,0,0,0,0,3,0,0,2)	$(0,5/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,0,0,3,1,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,0,4,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$
(0,0,0,0,0,1,0,0,0,2)	$(0,5/2)$	(0,0,0,0,0,1,0,0,1,2)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,1,0,0,2,2)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,0,0,3,2)	$(0,1/2) \oplus 2(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,1,1,0,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,1,0,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,1,1,0,2,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,1,1,0,2)	$(0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,1,1,1,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,2,0,0,2)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,1,2,0,1,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,2,1,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,0,1,3,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,2,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$
(0,0,0,0,0,2,0,0,1,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 2(0,7/2) \oplus (1/2,3) \oplus (1/2,4)$	(0,0,0,0,0,2,0,0,2,2)	$(0,1/2) \oplus 3(0,3/2) \oplus 4(0,5/2) \oplus 2(0,7/2) \oplus (1/2,2) \oplus (1/2,3) \oplus (1/2,4)$
(0,0,0,0,0,2,1,0,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 2(0,7/2) \oplus (1/2,3) \oplus (1/2,4)$	(0,0,0,0,0,2,1,0,1,2)	$(0,1/2) \oplus 5(0,3/2) \oplus 7(0,5/2) \oplus 3(0,7/2) \oplus (1/2,2) \oplus 2(1/2,3) \oplus (1/2,4)$
(0,0,0,0,0,2,1,1,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 2(0,7/2) \oplus (1/2,3) \oplus (1/2,4)$	(0,0,0,0,0,2,2,0,0,2)	$(0,1/2) \oplus 3(0,3/2) \oplus 4(0,5/2) \oplus 2(0,7/2) \oplus (1/2,2) \oplus (1/2,3) \oplus (1/2,4)$
(0,0,0,0,0,3,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus 2(0,9/2) \oplus (1/2,4) \oplus (1/2,5) \oplus (1,11/2)$	(0,0,0,0,0,3,0,0,1,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$
(0,0,0,0,0,3,1,0,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$	(0,0,0,0,0,4,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus 2(0,9/2) \oplus 2(0,11/2) \oplus (1/2,4) \oplus (1/2,5) \oplus 2(1/2,6) \oplus (1,11/2) \oplus (1,13/2) \oplus (3/2,7)$
(0,0,0,0,1,1,0,0,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,1,0,0,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,1,0,0,2,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,1,1,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,1,1,0,1,2)	$3(0,1/2) \oplus 3(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,1,1,1,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,1,1,2,0,0,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,1,2,0,0,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 2(0,7/2) \oplus (1/2,3) \oplus (1/2,4)$
(0,0,0,0,1,2,0,0,1,2)	$(0,1/2) \oplus 5(0,3/2) \oplus 7(0,5/2) \oplus 3(0,7/2) \oplus (1/2,2) \oplus 2(1/2,3) \oplus (1/2,4)$	(0,0,0,0,1,2,1,0,0,2)	$(0,1/2) \oplus 5(0,3/2) \oplus 7(0,5/2) \oplus 3(0,7/2) \oplus (1/2,2) \oplus 2(1/2,3) \oplus (1/2,4)$
(0,0,0,0,1,3,0,0,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$	(0,0,0,0,2,1,0,0,0,2)	$(0,1/2) \oplus (0,3/2) \oplus (0,5/2)$
(0,0,0,0,2,1,0,0,1,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,0,2,1,1,0,0,2)	$2(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,0,2,2,0,0,0,2)	$(0,1/2) \oplus 3(0,3/2) \oplus 4(0,5/2) \oplus 2(0,7/2) \oplus (1/2,2) \oplus (1/2,3) \oplus (1/2,4)$	(0,0,0,0,3,0,0,0,0,2)	$(0,5/2)$
(0,0,0,0,3,1,0,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,4,0,0,0,0,2)	$(0,5/2) \oplus (0,7/2) \oplus (1/2,4)$
(0,0,0,1,1,1,0,0,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,1,1,1,0,0,1,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$
(0,0,0,1,1,1,1,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,1,2,0,0,0,2)	$(0,3/2) \oplus 3(0,5/2) \oplus 2(0,7/2) \oplus (1/2,3) \oplus (1/2,4)$
(0,0,0,1,2,1,0,0,0,2)	$(0,1/2) \oplus 2(0,3/2) \oplus (0,5/2)$	(0,0,0,1,3,0,0,0,0,2)	$(0,3/2) \oplus (0,5/2)$
(0,0,1,1,1,1,0,0,0,2)	$(0,3/2) \oplus (0,5/2)$	(0,0,0,0,0,1,0,0,0,3)	$(0,7/2)$
(0,0,0,0,0,1,0,0,1,3)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,1,0,0,2,3)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$

continued on next page

β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$	β	$\oplus N_{j_l, j_r}^\beta(j_l, j_r)$
(0,0,0,0,0,1,1,0,0,3)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,1,1,0,1,3)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,1,1,1,0,3)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,0,1,2,0,0,3)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,2,0,0,0,3)	$(0,5/2) \oplus (0,7/2) \oplus 2(0,9/2) \oplus (1/2,4) \oplus (1/2,5) \oplus (1,11/2)$	(0,0,0,0,0,2,0,0,1,3)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$
(0,0,0,0,0,2,1,0,0,3)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$	(0,0,0,0,0,3,0,0,0,3)	$(0,3/2) \oplus (0,5/2) \oplus 3(0,7/2) \oplus 3(0,9/2) \oplus 4(0,11/2) \oplus (1/2,3) \oplus 2(1/2,4) \oplus 3(1/2,5) \oplus 3(1/2,6) \oplus (1/2,7) \oplus (1,9/2) \oplus 2(1,11/2) \oplus 3(1,13/2) \oplus (3/2,6) \oplus (3/2,7) \oplus (2,15/2)$
(0,0,0,0,1,1,0,0,0,3)	$(0,5/2) \oplus (0,7/2)$	(0,0,0,0,1,1,0,0,1,3)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$
(0,0,0,0,1,1,1,0,0,3)	$(0,3/2) \oplus 2(0,5/2) \oplus (0,7/2)$	(0,0,0,0,1,2,0,0,0,3)	$(0,3/2) \oplus 3(0,5/2) \oplus 5(0,7/2) \oplus 3(0,9/2) \oplus (1/2,3) \oplus 3(1/2,4) \oplus 2(1/2,5) \oplus (1,9/2) \oplus (1,11/2)$
(0,0,0,0,2,1,0,0,0,3)	$(0,3/2) \oplus (0,5/2) \oplus (0,7/2)$	(0,0,0,1,1,1,0,0,0,3)	$(0,5/2) \oplus (0,7/2)$
(0,0,0,0,0,1,0,0,0,4)	$(0,9/2)$	(0,0,0,0,0,1,0,0,1,4)	$(0,7/2) \oplus (0,9/2)$
(0,0,0,0,0,1,1,0,0,4)	$(0,7/2) \oplus (0,9/2)$	(0,0,0,0,0,2,0,0,0,4)	$(0,5/2) \oplus (0,7/2) \oplus 2(0,9/2) \oplus 2(0,11/2) \oplus (1/2,4) \oplus (1/2,5) \oplus 2(1/2,6) \oplus (1,11/2) \oplus (1,13/2) \oplus (3/2,7)$
(0,0,0,0,1,1,0,0,0,4)	$(0,7/2) \oplus (0,9/2)$	(0,0,0,0,0,1,0,0,0,5)	$(0,11/2)$

Table 27. Refined BPS invariants of 6d E_8 minimal SCFT.

F_4

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34			
0	534	1076	1649	2209	2271	2920	3495	3828	3726	3081	2055	1051	398	102	12	1																						
1	244	773	1412	2209	3175	4247	5241	5848	5705	4599	2874	1326	431	88	8																							
2	248	593	1117	1871	1252	2866	4050	5232	6043	5997	4782	2841	1185	332	56	3																						
3	39	146	64	688	341	160	1285	2105	4293	3140	5278	4197	2379	902	218	28	1																					
4	21	64	64	160	355	710	1285	2105	3079	3919	4136	3306	1802	617	125	12																						
5	1	6	6	6	22	64	356	160	712	1276	2016	2728	2999	2418	1266	390	66	4																				
6	6	1	6	6	22	64	64	160	352	670	1058	1295	1079	522	123	12																						
7	6	6	6	6	22	64	64	160	64	159	338	588	772	660	308	62	4																					
8	6	6	6	6	22	64	64	160	64	64	155	302	433	384	172	28	1																					
9	6	6	6	6	22	64	64	160	64	64	22	63	141	224	207	89	11																					
10	6	6	6	6	22	64	64	160	64	64	6	6	59	107	105	44	4																					
11	6	6	6	6	22	64	64	160	64	64	1	6	21	46	49	20	8																					
12	6	6	6	6	22	64	64	160	64	64	1	6	6	1	6	17	2																					
13	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
14	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
15	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					

$$\beta = (1, 8, 1, 0, 0, 4)$$

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29								
0	630	1287	1456	2015	2793	3492	3828	3528	2572	1394	528	124	18																									
1	222	743	466	975	1783	2878	4064	4835	4553	3137	1455	414	62	9																								
2	177	79	216	504	1034	1854	2857	3625	3518	2402	1048	258	31	1																								
3	19	64	64	160	356	710	1276	2016	2728	2999	2418	1266	390	66	4																							
4	6	6	6	6	22	64	64	160	64	159	338	588	772	660	308	62	4																					
5	6	6	6	6	22	64	64	160	64	64	155	302	433	384	172	28	1																					
6	6	6	6	6	22	64	64	160	64	64	22	63	141	224	207	89	11																					
7	6	6	6	6	22	64	64	160	64	64	6	6	59	107	105	44	4																					
8	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
9	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
10	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
11	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					
12	6	6	6	6	22	64	64	160	64	64	6	6	6	6	6	6	6																					

$$\beta = (2, 7, 1, 0, 0, 4)$$

Table 28. Refined BPS invariants for selected degrees of 6d F_4 minimal SCFT.

630(6102)21dEHC

E_6

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34		
0	546	1101	1688	2327	2991	3578	3913	3807	3142	2077	1063	400	99	15																							
1	245	776	1420	2225	3204	4293	5306	5927	5788	4668	2906	1335	435	88	8																						
2	249	596	1126	1889	2900	4104	5310	6137	6095	4859	2868	1194	336	53	4																						
3	39	146	341	689	1255	2084	3159	4329	5234	5347	4260	2405	904	219	28	1																					
4	21	64	160	356	713	1294	2123	3113	3969	4198	3029	2451	1817	618	128	11																					
5	1	6	22	64	160	356	713	1279	2025	2746	1774	2055	1686	845	227	31	1																				
6	6	22	64	160	356	705	1219	2025	3252	671	1061	1303	1090	527	123	12	1																				
7	1	6	22	64	160	352	671	1061	159	339	591	780	1090	1638	272	34	1																				
8	6	22	64	160	352	705	1219	2025	3252	671	1061	1303	1090	527	123	12	1																				
9	1	6	22	64	160	352	671	1061	159	339	591	780	1090	1638	272	34	1																				
10	6	22	64	160	352	705	1219	2025	3252	671	1061	1303	1090	527	123	12	1																				
11	1	6	22	64	160	352	671	1061	159	339	591	780	1090	1638	272	34	1																				
12	6	22	64	160	352	705	1219	2025	3252	671	1061	1303	1090	527	123	12	1																				
13	1	6	22	64	160	352	671	1061	159	339	591	780	1090	1638	272	34	1																				
14	6	22	64	160	352	705	1219	2025	3252	671	1061	1303	1090	527	123	12	1																				
15	1	6	22	64	160	352	671	1061	159	339	591	780	1090	1638	272	34	1																				

$\beta = (0, 1, 0, 1, 4, 0, 0, 6)$

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
0	992	1980	3001	4100	5241	6249	6824	6654	5557	3777	2005	783	208	37	2																						
1	470	1483	2690	4158	5899	7804	9553	10592	10299	8357	5332	2551	872	193	24	1																					
2	527	1227	2253	3686	5532	7677	9776	11154	10971	8765	5308	2319	695	125	12																						
3	96	342	755	1458	2558	4112	6067	8136	9660	9729	7748	4493	1791	470	69	4																					
4	59	166	382	797	1522	2653	4203	5986	7449	7724	6157	3430	1251	285	31	1																					
5	5	22	65	168	387	807	1531	2624	4003	5262	5657	4543	2444	809	155	12																					
6	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
7	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																						
8	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
9	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																						
10	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
11	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																						
12	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
13	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																						
14	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
15	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																						
16	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					

$\beta = (0, 1, 0, 1, 5, 0, 0, 5)$

Table 29. Refined BPS invariants for selected degrees of 6d E_6 minimal SCFT.

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29				
0	647	1327	1466	2079	2891	3608	3957	3627	2627	1417	531	131	14	1																				
1	223	746	469	1806	3611	4754	5396	4980	3489	1728	558	108	9																					
2	178	469	2928	1806	1037	1865	4141	4945	4656	3199	1476	412	66	3																				
3	19	79	216	505	1037	2880	3670	3573	2444	1066	258	31	1																					
4	6	25	6	80	217	507	1027	1752	2408	2438	1665	688	141	13																				
5	6	1	6	25	80	216	496	1416	1504	1035	408	71	4																					
6	6	6	1	6	25	80	213	460	762	857	597	225	31	1																				
7	6	6	1	6	25	79	198	367	440	313	113	12	4																					
8	6	6	1	6	25	75	160	207	152	54	4																							
9	6	6	1	6	24	61	86	66	23	1																								
10	6	6	1	6	20	31	25	9																										
11	6	6	1	6	1	5	9	8	3																									
12	6	6	1	6	1	2	2	1																										

$$\beta = (0, 0, 0, 5, 1, 0, 0, 2, 4)$$

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
0	992	1483	1980	3001	4100	5241	6249	6824	6654	5557	3777	2005	783	208	37	2																					
1	470	527	1227	2253	3686	5532	7677	9776	11154	10971	8765	5308	2319	695	125	12																					
2	96	342	755	1458	2558	4112	6067	8136	9660	9729	7748	4493	1791	470	69	4																					
3	59	166	382	797	1522	2653	4203	5986	7449	7724	6157	3430	1251	285	31	1																					
4	5	22	65	168	387	807	1531	2624	4003	5262	5657	4543	2444	809	155	12																					
5	1	6	22	65	168	387	805	1505	2481	3472	3896	3164	1638	485	77	4																					
6	1	6	22	65	168	386	791	1420	2135	2518	2074	1035	272	34	1																						
7	1	6	22	65	167	368	661	889	768	356	69	4																									
8	1	6	22	65	163	328	480	429	191	30	1																										
9	1	6	22	64	148	241	225	97	12																												
10	1	6	22	60	112	112	47	4																													
11	1	6	21	46	49	20	8	1																													
12	1	6	17	6	1	5	7	3																													
13	1	6	1	5	2	2	1																														
14	1	6	1	5	2	1																															
15	1	6	1	5	2	1																															
16	1	6	1	5	2	1																															

$$\beta = (0, 0, 1, 5, 1, 0, 0, 0, 5)$$

Table 30. Refined BPS invariants for selected degrees of 6d E_7 minimal SCFT.

E_8

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
0		3	7	13		21		27		28		17		4		1				
1			1	4		10		19		28		31		19		4				
2					1		4		11		20		25		15		2			
3								1		4		10		15		9		1		
4											1		4		8		5			
5														1		3		2		
6																		1		1

$$\beta = (1, 1, 1, 1, 1, 3, 0, 0, 0, 4)$$

$2j_L \setminus 2j_R$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
0		17	44	79	91	63	23	2								
1	1		6	22	51	68	50	17	1							
2				1	6	20	31	25	9							
3						1	5	9	8	3						
4									1	2	2	1				

$$\beta = (0, 1, 1, 1, 1, 3, 0, 0, 2, 3)$$

Table 31. Refined BPS invariants for selected degrees of 6d E_8 minimal SCFT.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] W. Nahm, *Supersymmetries and their Representations*, *Nucl. Phys. B* **135** (1978) 149 [[INSPIRE](#)].
- [2] O.J. Ganor and A. Hanany, *Small E_8 instantons and tensionless noncritical strings*, *Nucl. Phys. B* **474** (1996) 122 [[hep-th/9602120](#)] [[INSPIRE](#)].
- [3] A. Klemm, P. Mayr and C. Vafa, *BPS states of exceptional noncritical strings*, *Nucl. Phys. Proc. Suppl.* **58** (1997) 177 [[hep-th/9607139](#)] [[INSPIRE](#)].
- [4] D.R. Morrison and C. Vafa, *Compactifications of F-theory on Calabi-Yau threefolds. 2.*, *Nucl. Phys. B* **476** (1996) 437 [[hep-th/9603161](#)] [[INSPIRE](#)].
- [5] J.J. Heckman, D.R. Morrison and C. Vafa, *On the Classification of 6D SCFTs and Generalized ADE Orbifolds*, *JHEP* **05** (2014) 028 [Erratum *ibid.* **06** (2015) 017] [[arXiv:1312.5746](#)] [[INSPIRE](#)].
- [6] J.J. Heckman, D.R. Morrison, T. Rudelius and C. Vafa, *Atomic Classification of 6D SCFTs*, *Fortsch. Phys.* **63** (2015) 468 [[arXiv:1502.05405](#)] [[INSPIRE](#)].

- [7] L. Bhardwaj, *Classification of 6d $\mathcal{N} = (1, 0)$ gauge theories*, *JHEP* **11** (2015) 002 [[arXiv:1502.06594](#)] [[INSPIRE](#)].
- [8] B. Haghighat, A. Klemm, G. Lockhart and C. Vafa, *Strings of Minimal 6d SCFTs*, *Fortsch. Phys.* **63** (2015) 294 [[arXiv:1412.3152](#)] [[INSPIRE](#)].
- [9] R. Pandharipande and R.P. Thomas, *Curve counting via stable pairs in the derived category*, *Invent. Math.* **178** (2009) 407.
- [10] R. Pandharipande and R.P. Thomas, *Stable pairs and BPS invariants*, *J. Am. Math. Soc.* **23** (2010) 267.
- [11] J. Choi, S. Katz and A. Klemm, *The refined BPS index from stable pair invariants*, *Commun. Math. Phys.* **328** (2014) 903 [[arXiv:1210.4403](#)] [[INSPIRE](#)].
- [12] N. Nekrasov and A. Okounkov, *Membranes and sheaves*, *Algebr. Geom.* **3** (2016) 320.
- [13] A. Iqbal, C. Kozçaz and C. Vafa, *The Refined topological vertex*, *JHEP* **10** (2009) 069 [[hep-th/0701156](#)] [[INSPIRE](#)].
- [14] M.-x. Huang and A. Klemm, *Direct integration for general Ω backgrounds*, *Adv. Theor. Math. Phys.* **16** (2012) 805 [[arXiv:1009.1126](#)] [[INSPIRE](#)].
- [15] M.-x. Huang, K. Sun and X. Wang, *Blowup Equations for Refined Topological Strings*, *JHEP* **10** (2018) 196 [[arXiv:1711.09884](#)] [[INSPIRE](#)].
- [16] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. II. K-theoretic partition function*, [math/0505553](#).
- [17] L. Gottsche, H. Nakajima and K. Yoshioka, *K-theoretic Donaldson invariants via instanton counting*, *Pure Appl. Math. Quart.* **5** (2009) 1029 [[math/0611945](#)] [[INSPIRE](#)].
- [18] H. Nakajima and K. Yoshioka, *Perverse coherent sheaves on blowup, III: Blow-up formula from wall-crossing*, *Kyoto J. Math.* **51** (2011) 263.
- [19] B. Haghighat, A. Iqbal, C. Kozçaz, G. Lockhart and C. Vafa, *M-Strings*, *Commun. Math. Phys.* **334** (2015) 779 [[arXiv:1305.6322](#)] [[INSPIRE](#)].
- [20] A. Gadde and S. Gukov, *2d Index and Surface operators*, *JHEP* **03** (2014) 080 [[arXiv:1305.0266](#)] [[INSPIRE](#)].
- [21] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of 2d $\mathcal{N} = 2$ Gauge Theories*, *Commun. Math. Phys.* **333** (2015) 1241 [[arXiv:1308.4896](#)] [[INSPIRE](#)].
- [22] M.-x. Huang, S. Katz and A. Klemm, *Topological String on elliptic CY 3-folds and the ring of Jacobi forms*, *JHEP* **10** (2015) 125 [[arXiv:1501.04891](#)] [[INSPIRE](#)].
- [23] J. Gu, M.-x. Huang, A.-K. Kashani-Poor and A. Klemm, *Refined BPS invariants of 6d SCFTs from anomalies and modularity*, *JHEP* **05** (2017) 130 [[arXiv:1701.00764](#)] [[INSPIRE](#)].
- [24] M. Del Zotto, J. Gu, M.-X. Huang, A.-K. Kashani-Poor, A. Klemm and G. Lockhart, *Topological Strings on Singular Elliptic Calabi-Yau 3-folds and Minimal 6d SCFTs*, *JHEP* **03** (2018) 156 [[arXiv:1712.07017](#)] [[INSPIRE](#)].
- [25] J. Gu, B. Haghighat, K. Sun and X. Wang, *Blowup Equations for 6d SCFTs. I*, *JHEP* **03** (2019) 002 [[arXiv:1811.02577](#)] [[INSPIRE](#)].
- [26] S.H. Katz, A. Klemm and C. Vafa, *Geometric engineering of quantum field theories*, *Nucl. Phys. B* **497** (1997) 173 [[hep-th/9609239](#)] [[INSPIRE](#)].

- [27] N.A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831 [[hep-th/0206161](#)] [[INSPIRE](#)].
- [28] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. 1.*, *Invent. Math.* **162** (2005) 313 [[math/0306198](#)].
- [29] A. Grassi and J. Gu, *BPS relations from spectral problems and blowup equations*, *Lett. Math. Phys.* **109** (2019) 1271 [[arXiv:1609.05914](#)] [[INSPIRE](#)].
- [30] C.A. Keller and J. Song, *Counting Exceptional Instantons*, *JHEP* **07** (2012) 085 [[arXiv:1205.4722](#)] [[INSPIRE](#)].
- [31] <http://www.th.physik.uni-bonn.de/klemm/data.php>.
- [32] J. Kim, S. Kim, K. Lee, J. Park and C. Vafa, *Elliptic Genus of E-strings*, *JHEP* **09** (2017) 098 [[arXiv:1411.2324](#)] [[INSPIRE](#)].
- [33] J. Kim, S. Kim and K. Lee, *Higgsing towards E-strings*, [arXiv:1510.03128](#) [[INSPIRE](#)].
- [34] H.-C. Kim, S. Kim and J. Park, *6d strings from new chiral gauge theories*, [arXiv:1608.03919](#) [[INSPIRE](#)].
- [35] M.-X. Huang, A. Klemm and M. Poretschkin, *Refined stable pair invariants for E-, M- and [p, q]-strings*, *JHEP* **11** (2013) 112 [[arXiv:1308.0619](#)] [[INSPIRE](#)].
- [36] Z. Duan, J. Gu and A.-K. Kashani-Poor, *Computing the elliptic genus of higher rank E-strings from genus 0 GW invariants*, *JHEP* **03** (2019) 078 [[arXiv:1810.01280](#)] [[INSPIRE](#)].
- [37] M. Del Zotto and G. Lockhart, *On Exceptional Instanton Strings*, *JHEP* **09** (2017) 081 [[arXiv:1609.00310](#)] [[INSPIRE](#)].
- [38] M. Del Zotto and G. Lockhart, *Universal Features of BPS Strings in Six-dimensional SCFTs*, *JHEP* **08** (2018) 173 [[arXiv:1804.09694](#)] [[INSPIRE](#)].
- [39] S.-S. Kim, M. Taki and F. Yagi, *Tao Probing the End of the World*, *PTEP* **2015** (2015) 083B02 [[arXiv:1504.03672](#)] [[INSPIRE](#)].
- [40] H. Hayashi and K. Ohmori, *5d/6d DE instantons from trivalent gluing of web diagrams*, *JHEP* **06** (2017) 078 [[arXiv:1702.07263](#)] [[INSPIRE](#)].
- [41] S. Benvenuti, A. Hanany and N. Mekareeya, *The Hilbert Series of the One Instanton Moduli Space*, *JHEP* **06** (2010) 100 [[arXiv:1005.3026](#)] [[INSPIRE](#)].
- [42] A. Hanany, N. Mekareeya and S.S. Razamat, *Hilbert Series for Moduli Spaces of Two Instantons*, *JHEP* **01** (2013) 070 [[arXiv:1205.4741](#)] [[INSPIRE](#)].
- [43] S. Cremonesi, G. Ferlito, A. Hanany and N. Mekareeya, *Coulomb Branch and The Moduli Space of Instantons*, *JHEP* **12** (2014) 103 [[arXiv:1408.6835](#)] [[INSPIRE](#)].
- [44] A. Gadde, L. Rastelli, S.S. Razamat and W. Yan, *Gauge Theories and Macdonald Polynomials*, *Commun. Math. Phys.* **319** (2013) 147 [[arXiv:1110.3740](#)] [[INSPIRE](#)].
- [45] D. Gaiotto and S.S. Razamat, *Exceptional Indices*, *JHEP* **05** (2012) 145 [[arXiv:1203.5517](#)] [[INSPIRE](#)].
- [46] P. Putrov, J. Song and W. Yan, *(0, 4) dualities*, *JHEP* **03** (2016) 185 [[arXiv:1505.07110](#)] [[INSPIRE](#)].
- [47] A. Gadde, S.S. Razamat and B. Willett, *“Lagrangian” for a Non-Lagrangian Field Theory with $\mathcal{N} = 2$ Supersymmetry*, *Phys. Rev. Lett.* **115** (2015) 171604 [[arXiv:1505.05834](#)] [[INSPIRE](#)].

- [48] P. Agarwal, K. Maruyoshi and J. Song, A “Lagrangian” for the E_7 superconformal theory, *JHEP* **05** (2018) 193 [[arXiv:1802.05268](#)] [[INSPIRE](#)].
- [49] B. Haghighat, G. Lockhart and C. Vafa, Fusing E -strings to heterotic strings: $E + E \rightarrow H$, *Phys. Rev. D* **90** (2014) 126012 [[arXiv:1406.0850](#)] [[INSPIRE](#)].
- [50] W. Cai, M.-x. Huang and K. Sun, On the Elliptic Genus of Three E -strings and Heterotic Strings, *JHEP* **01** (2015) 079 [[arXiv:1411.2801](#)] [[INSPIRE](#)].
- [51] J. Gu, B. Haghighat, A. Klemm, K. Sun and X. Wang, Elliptic Blowup Equations for 6d SCFTs. IV: Matters, to appear.
- [52] J. Gu, B. Haghighat, A. Klemm, K. Sun and X. Wang, Elliptic Blowup Equations for 6d SCFTs. III: E -strings, M -strings and Chains, to appear.
- [53] M. Del Zotto, J.J. Heckman and D.R. Morrison, 6D SCFTs and Phases of 5D Theories, *JHEP* **09** (2017) 147 [[arXiv:1703.02981](#)] [[INSPIRE](#)].
- [54] M.-x. Huang, A.-K. Kashani-Poor and A. Klemm, The Ω deformed B -model for rigid $\mathcal{N} = 2$ theories, *Annales Henri Poincaré* **14** (2013) 425 [[arXiv:1109.5728](#)] [[INSPIRE](#)].
- [55] M. Esole, P. Jefferson and M.J. Kang, The Geometry of F_4 -Models, [arXiv:1704.08251](#) [[INSPIRE](#)].
- [56] S. Hosono, A. Klemm and S. Theisen, Lectures on mirror symmetry, *Lect. Notes Phys.* **436** (1994) 235 [[hep-th/9403096](#)] [[INSPIRE](#)].
- [57] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, Mirror symmetry, mirror map and applications to Calabi-Yau hypersurfaces, *Commun. Math. Phys.* **167** (1995) 301 [[hep-th/9308122](#)] [[INSPIRE](#)].
- [58] T. Schimannek, Modularity from Monodromy, *JHEP* **05** (2019) 024 [[arXiv:1902.08215](#)] [[INSPIRE](#)].
- [59] C.A. Keller, N. Mekareeya, J. Song and Y. Tachikawa, The ABCDEFG of Instantons and W -algebras, *JHEP* **03** (2012) 045 [[arXiv:1111.5624](#)] [[INSPIRE](#)].
- [60] A. Hanany and R. Kalveks, Highest Weight Generating Functions for Hilbert Series, *JHEP* **10** (2014) 152 [[arXiv:1408.4690](#)] [[INSPIRE](#)].
- [61] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*, Graduate Texts in Contemporary Physics. Springer-Verlag, New York (1997) [[DOI:10.1007/978-1-4612-2256-9](#)].
- [62] P.C. Argyres, M.R. Plesser, N. Seiberg and E. Witten, New $N = 2$ superconformal field theories in four-dimensions, *Nucl. Phys. B* **461** (1996) 71 [[hep-th/9511154](#)] [[INSPIRE](#)].
- [63] T. Banks, M.R. Douglas and N. Seiberg, Probing F -theory with branes, *Phys. Lett. B* **387** (1996) 278 [[hep-th/9605199](#)] [[INSPIRE](#)].
- [64] M.R. Douglas, D.A. Lowe and J.H. Schwarz, Probing F -theory with multiple branes, *Phys. Lett. B* **394** (1997) 297 [[hep-th/9612062](#)] [[INSPIRE](#)].
- [65] J.A. Minahan and D. Nemeschansky, An $N = 2$ superconformal fixed point with E_6 global symmetry, *Nucl. Phys. B* **482** (1996) 142 [[hep-th/9608047](#)] [[INSPIRE](#)].
- [66] J.A. Minahan and D. Nemeschansky, Superconformal fixed points with $E(n)$ global symmetry, *Nucl. Phys. B* **489** (1997) 24 [[hep-th/9610076](#)] [[INSPIRE](#)].
- [67] P.C. Argyres and N. Seiberg, S -duality in $N = 2$ supersymmetric gauge theories, *JHEP* **12** (2007) 088 [[arXiv:0711.0054](#)] [[INSPIRE](#)].

- [68] A. Kapustin, *Holomorphic reduction of $N = 2$ gauge theories, Wilson-'t Hooft operators and S -duality*, [hep-th/0612119](#) [INSPIRE].
- [69] V.P. Spiridonov and S.O. Warnaar, *Inversions of integral operators and elliptic beta integrals on root systems*, *Adv. Math.* **207** (2006) 91.
- [70] F. Benini, S. Benvenuti and Y. Tachikawa, *Webs of five-branes and $N = 2$ superconformal field theories*, *JHEP* **09** (2009) 052 [[arXiv:0906.0359](#)] [INSPIRE].
- [71] D. Xie, *General Argyres-Douglas Theory*, *JHEP* **01** (2013) 100 [[arXiv:1204.2270](#)] [INSPIRE].
- [72] G. Bonelli, K. Maruyoshi and A. Tanzini, *Wild Quiver Gauge Theories*, *JHEP* **02** (2012) 031 [[arXiv:1112.1691](#)] [INSPIRE].
- [73] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli and B.C. van Rees, *Infinite Chiral Symmetry in Four Dimensions*, *Commun. Math. Phys.* **336** (2015) 1359 [[arXiv:1312.5344](#)] [INSPIRE].
- [74] C. Beem, W. Peelaers, L. Rastelli and B.C. van Rees, *Chiral algebras of class S* , *JHEP* **05** (2015) 020 [[arXiv:1408.6522](#)] [INSPIRE].
- [75] Y. Pan and W. Peelaers, *Chiral Algebras, Localization and Surface Defects*, *JHEP* **02** (2018) 138 [[arXiv:1710.04306](#)] [INSPIRE].
- [76] Y. Pan and W. Peelaers, *Schur correlation functions on $S^3 \times S^1$* , *JHEP* **07** (2019) 013 [[arXiv:1903.03623](#)] [INSPIRE].
- [77] J. Oh and J. Yagi, *Chiral algebras from Ω -deformation*, *JHEP* **08** (2019) 143 [[arXiv:1903.11123](#)] [INSPIRE].
- [78] M. Dedushenko and M. Fluder, *Chiral Algebra, Localization, Modularity, Surface defects, And All That*, [arXiv:1904.02704](#) [INSPIRE].
- [79] S. Jeong, *SCFT/VOA correspondence via Ω -deformation*, *JHEP* **10** (2019) 171 [[arXiv:1904.00927](#)] [INSPIRE].
- [80] R. Eager, G. Lockhart and E. Sharpe, *Hidden exceptional symmetry in the pure spinor superstring*, [arXiv:1902.09504](#) [INSPIRE].
- [81] J. Kinney, J.M. Maldacena, S. Minwalla and S. Raju, *An Index for 4 dimensional super conformal theories*, *Commun. Math. Phys.* **275** (2007) 209 [[hep-th/0510251](#)] [INSPIRE].
- [82] C. Romelsberger, *Counting chiral primaries in $N = 1$, $d = 4$ superconformal field theories*, *Nucl. Phys. B* **747** (2006) 329 [[hep-th/0510060](#)] [INSPIRE].
- [83] A. Gadde, L. Rastelli, S.S. Razamat and W. Yan, *The 4d Superconformal Index from q -deformed 2d Yang-Mills*, *Phys. Rev. Lett.* **106** (2011) 241602 [[arXiv:1104.3850](#)] [INSPIRE].
- [84] C. Cordova and S.-H. Shao, *Schur Indices, BPS Particles and Argyres-Douglas Theories*, *JHEP* **01** (2016) 040 [[arXiv:1506.00265](#)] [INSPIRE].
- [85] A. Gadde, L. Rastelli, S.S. Razamat and W. Yan, *The Superconformal Index of the E_6 SCFT*, *JHEP* **08** (2010) 107 [[arXiv:1003.4244](#)] [INSPIRE].
- [86] C. Cordova, D. Gaiotto and S.-H. Shao, *Infrared Computations of Defect Schur Indices*, *JHEP* **11** (2016) 106 [[arXiv:1606.08429](#)] [INSPIRE].
- [87] H.-C. Kim, J. Kim, S. Kim, K.-H. Lee and J. Park, *6d strings and exceptional instantons*, [arXiv:1801.03579](#) [INSPIRE].

- [88] B. Haghighat, S. Murthy, C. Vafa and S. Vandoren, *F-Theory, Spinning Black Holes and Multi-string Branches*, *JHEP* **01** (2016) 009 [[arXiv:1509.00455](#)] [[INSPIRE](#)].
- [89] H. Hayashi, P. Jefferson, H.-C. Kim, K. Ohmori and C. Vafa, *SCFTs, Holography and Topological Strings*, [arXiv:1905.00116](#) [[INSPIRE](#)].
- [90] V. Braun and D.R. Morrison, *F-theory on Genus-One Fibrations*, *JHEP* **08** (2014) 132 [[arXiv:1401.7844](#)] [[INSPIRE](#)].
- [91] D.R. Morrison and W. Taylor, *Sections, multisections and U(1) fields in F-theory*, [arXiv:1404.1527](#) [[INSPIRE](#)].
- [92] J.J. Heckman and T. Rudelius, *Top Down Approach to 6D SCFTs*, *J. Phys. A* **52** (2019) 093001 [[arXiv:1805.06467](#)] [[INSPIRE](#)].
- [93] A. Braverman, M. Finkelberg and H. Nakajima, *Instanton moduli spaces and \mathcal{W} -algebras*, [arXiv:1406.2381](#) [[INSPIRE](#)].
- [94] M. Taki, *On AGT-W Conjecture and q-Deformed W-Algebra*, [arXiv:1403.7016](#) [[INSPIRE](#)].
- [95] <https://www.tfc.tohoku.ac.jp/wp-content/uploads/2018/06/Fukuda.pdf>.
- [96] A. Klemm, B. Lian, S.S. Roan and S.-T. Yau, *Calabi-Yau fourfolds for M-theory and F-theory compactifications*, *Nucl. Phys. B* **518** (1998) 515 [[hep-th/9701023](#)] [[INSPIRE](#)].
- [97] J. Kim, K. Lee and J. Park, *On elliptic genera of 6d string theories*, *JHEP* **10** (2018) 100 [[arXiv:1801.01631](#)] [[INSPIRE](#)].