

REVISTA MATEMÁTICA de la
Universidad Complutense de Madrid
Volumen 8, número 2: 1995
http://dx.doi.org/10.5209/rev_REMA.1995.v8.n2.17642

Embedding of Real Varieties and their Subvarieties into Grassmannians

M.A. BUCHNER

ABSTRACT. Given a compact affine nonsingular real algebraic variety X and a nonsingular subvariety $Z \subset X$ belonging to a large class of subvarieties, we show how to embed X in a suitable Grassmannian so that Z becomes the transverse intersection of the zeros of a section of the tautological bundle on the Grassmannian.

In [2] Bochnak and Kucharz prove the following characterization of a compact nonsingular algebraic hypersurface Z in a compact affine nonsingular real algebraic variety X : There is an algebraic embedding $f : X \rightarrow RP^n$ (for some n) and a projective hyperplane $H \subset RP^n$ transverse to $f(X)$ such that $H \cap f(X) = f(Z)$. This fact (or rather a closely related statement about strongly algebraic real line bundles) plays a crucial role in their construction of algebraic models Y of a compact, connected, smooth manifold M of dimensions $m \geq 3$ such that

Supported by Nato Grant CRG 930238.

1991 Mathematics Subject Classification: 14P05

Servicio publicaciones Univ. Complutense. Madrid, 1995.

the algebraic homology elements in $H^1(Y, Z/2) = H^1(M, Z/2)$ form a prescribed subgroup $G \subset H^1(M, Z/2)$. If we wish to extend this result to subgroups of $H^k(M, Z/2)$ for $k > 1$ it seems desirable, as a first step, to extend the above characterization of hypersurfaces to subvarieties of higher codimension.

Let $G_{n,k}(R)$ denote the Grassmannian of k -planes in R^n . Let $\gamma_{n,k}$ denote the universal bundle over $G_{n,k}(R)$. For definitions and results concerning real varieties, strongly algebraic vector bundles etc. see [1].

Theorem 1. *Let X be a compact affine nonsingular real algebraic variety. Let ζ be a strongly algebraic real vector bundle over X of rank k . Let σ be a regular section of ζ transverse to the zero section. Let $Z = \sigma^{-1}(0)$. Then*

(i) *There exists a regular embedding $f : X \rightarrow G_{n,k}(R)$ for suitable n such that ζ and $f^*(\gamma_{n,k})$ are isomorphic.*

(ii) *There exists a regular section s of $\gamma_{n,k}$ such that s is transversal to the zero section and $s^{-1}(0) \cap f(X) = f(Z)$ (the intersection $s^{-1}(0) \cap f(X)$ being transverse intersection).*

Proof. We can assume that X is a subvariety of real projective q space RP^q for some q . By theorem 12.1.7 of [1] there is a regular map $g : X \rightarrow G_{\ell,k}(R)$ (for suitable ℓ) such that $g^*(\gamma_{\ell,k})$ and ζ are isomorphic. Let $G_{\ell,k}(C)$ denote the Grassmannian of complex k -planes in C^ℓ and $\gamma_{\ell,k}^C$ the corresponding universal complex bundle. Let X_C denote the complexification of X in CP^q . Then g extends to a regular map $\tilde{g} : U \rightarrow G_{\ell,k}(C)$ where $U \subset X_C$ is a Zariski open set containing X . We can assume U and \tilde{g} are defined over R . By resolution of singularities we can find a complex nonsingular subvariety Y of some complex projective space CP^m with Y defined over R and a regular map (defined over R) $\tau : Y \rightarrow X_C$ where τ is the composition of a sequence of blowings-up with real centers outside U such that $\tilde{g} \circ \tau$ extends to a regular map on Y . Denote this extension by h . To simplify notation we identify X with $\tau^{-1}(X)$. Then $h^*(\gamma_{\ell,k}^C)$ is a bundle defined over R and $h^*(\gamma_{\ell,k}^C)|_X$ is isomorphic to $\zeta \otimes C$.

Now, for $E \rightarrow M$ a holomorphic vector bundle of rank k over the compact complex manifold M , let $H^0(M, E)$ denote the space of holomorphic sections. Denote the dimension of $H^0(M, E)$ by n . Let

$i_E(x) = \{ \text{sections vanishing at } x \}$. Assume that each fiber of E is generated by global sections. Then identifying $H^0(M, E)$ with C^n we see that i_E maps M to $G_{n,n-k}(C) \simeq G_{n,k}(C)$. If $F \rightarrow M$ is a positive holomorphic line bundle then for p sufficiently large $i_{E \otimes F^p}$ is an embedding of M into $G_{n,k}(C)$ where, now, $n = \dim_C H^0(M, E \otimes F^p)$ and $i_{E \otimes F^p}^*(\gamma_{n,k}^C)$ is isomorphic to the bundle $E \otimes F^p \rightarrow M$. Apply this to $E \rightarrow M$ replaced by $h^*(\gamma_{\ell,k}^C)$ (so M is replaced by Y) and F replaced by $\gamma_{m,1}^C|Y$. In this case $i_{E \otimes F^p}$ is a regular map defined over R . Abbreviating $i_{E \otimes F^p}$ by i , we can write

$$i^*(\gamma_{n,k}^C) \simeq h^*(\gamma_{\ell,k}^C) \otimes (\gamma_{m,1}^C|Y)^p$$

(as complex bundles). We now restrict both sides to X and obtain

$$(i|X)^*(\gamma_{n,k}) \otimes C \simeq (\zeta \otimes C) \otimes ((\gamma_{m,1}|X) \otimes C)^p$$

and hence

$$(i|X)^*(\gamma_{n,k}) \simeq \zeta \otimes (\gamma_{m,1}|X)^p .$$

We can assume p is even. Then $(\gamma_{m,1}|X)^p$ is topologically trivial. Hence $(i|X)^*(\gamma_{n,k})$ is topologically and hence algebraically isomorphic to ζ . This completes the proof of (i) with $f = i|X$.

To simplify notation we now identify X with $f(X)$ and ζ with $\gamma_{n,k}|X$. Let s_1, \dots, s_n be sections of $\gamma_{n,k}$ (over $G_{n,k}(R)$) spanning the fiber at each point of $G_{n,k}(R)$. Write $\sigma = \sum \lambda_i (s_i|X)$ where λ_i are regular real-valued functions on X . Let $\tilde{\lambda}$ be a regular extension of λ_i to $G_{n,k}(R)$. Let ϕ be a regular real-valued function on $G_{n,k}(R)$ such that $\phi^{-1}(0) = Z (= \sigma^{-1}(0))$. For $t = (t_1, \dots, t_n)$, define $s_t = \sum_{i=1}^n (\tilde{\lambda}_i + t_i \phi^2) s_i$. We can find t (suitably small) so that s_t is transverse to the zero section, $s_t^{-1}(0)$ is transverse to X and $s_t^{-1}(0) \cap X = \sigma^{-1}(0) (= Z)$. This completes the proof of (ii).

References

[1] Bochnak, J., Coste, M. and Roy, M.-F., *Géométrie algébrique réelle*. Ergebnisse der Math. Vol. 12, Berlin, Heidelberg, New York: Springer 1987.

[2] Bochnak, J. and Kucharz, W., *Algebraic models of smooth manifolds*.
Invent. Math. Vol. **97**, pp. 585-611 (1989).

Department of Mathematics and Statistics
University of New Mexico
Albuquerque, New Mexico 87131

Recibido: 19 de Abril de 1994