

## Review

# Emerging roles of extracellular vesicles in neurodegenerative disorders: focus on HIV-associated neurological complications

Guoku Hu<sup>1,3</sup>, Lu Yang<sup>2,3</sup>, Yu Cai<sup>1,3</sup>, Fang Niu<sup>1</sup>, Frank Mezzacappa<sup>1</sup>, Shannon Callen<sup>1</sup>, Howard S Fox<sup>1</sup> and Shilpa Buch<sup>\*1</sup>

Exosomes are membrane-enriched extracellular vesicles with a proposed diameter in the range of 30–100 nm. They are released during both normal homeostasis as well as under pathological conditions by most cell types. In recent years, there has been robust interest in the study of these vesicles as conduits for the delivery of information between cells in both analogous as well as disparate tissues. Their ability to transport specialized cargo including signaling mediators, proteins, messenger RNA and miRNAs characterizes these vesicles as primary facilitators of cell-to-cell communication and regulation. Exosomes have also been demonstrated to have important roles in the field of cancer biology and metastasis. More recently, their role in several neurodegenerative disorders has been gaining increased momentum as these particles have been shown to promote the spread of toxic factors such as amyloid beta and prions, adding further validity to their role as important regulators of disease pathogenesis. This review briefly summarizes current findings and thoughts on exosome biology in the context of neurodegenerative disorders and the manipulation of these particles for the development of potential therapeutic strategies.

*Cell Death and Disease* (2016) 7, e2481; doi:10.1038/cddis.2016.336; published online 24 November 2016

## Facts

- Exosomes are globular, membrane-bound extracellular nanovesicles (30–100 nm in diameter) that are released by almost all types of cells.
- ARF6 and PLD2 have important roles in extracellular vesicle release through the regulation of the budding of ILVs into MVBs.
- Extracellular vesicular molecules (including ADAM17, TNF $\alpha$  and Nef) released from HIV-infected cells induce activation, apoptosis and HIV susceptibility in the recipient cells.
- Extracellular vesicles released from CD8<sup>+</sup> T cells contain antiviral membrane-bound factors that inhibit HIV-1 transcription.

- What is the role of EVs in propagation of pathogenic proteins in the neurodegenerative disorders?
- How can extracellular vesicle therapeutics be applied in the context of neurodegenerative diseases?

Cellular cross talk underlies most pathological conditions including those within the central nervous system (CNS). Although various factors have been identified as instigators of disease pathogenesis, it is now becoming clear that unrestrained neuroinflammation and, subsequent cellular toxicity are the key hallmark features of various neurological disorders. In this light, the notion that disease pathogenesis may be accelerated or mediated by exosomes and their associated cargos is recently gaining momentum. Exosomes are globular; membrane-bound extracellular nanovesicles (30–100 nm in diameter) that are released by almost all types of cells during normal cellular functioning and specifically, in response to cellular stressors. These small vesicles originally thought to contain 'junk' cellular debris were first described by Trams *et al.*<sup>1</sup> when they observed smaller membrane-bound vesicles within the larger endosomes (later termed multi-vesicular bodies (MVBs)). An electron micrographic study related the

## Open questions

- Are HIV proteins such as Tat /gp120 released in the extracellular vesicles and if so, do they disseminate CNS toxicity?

<sup>1</sup>Department of Pharmacology and Experimental Neuroscience, University of Nebraska Medical Center, Omaha, NE, USA and <sup>2</sup>School of Medicine, University of Electronic Science and Technology of China, Chengdu 610054, China

\*Corresponding author: S Buch, Department of Pharmacology and Experimental Neuroscience, University of Nebraska Medical Center, Omaha, NE 68198-5880, USA. Tel: 402 5593165; Fax: 402 5597495; E-mail: sbuch@unmc.edu

<sup>3</sup>These authors contributed equally to this work.

**Abbreviations:** CNS, central nervous system; MVBs, multi-vesicular bodies; EVs, extracellular vesicles; ILVs, intraluminal vesicles; EBV, Epstein–Barr virus; RME, receptor-mediated endocytosis; ESCRT, endosomal sorting complexes required for transport; ARF6, ADP ribosylation factor 6; PLD2, phospholipase D2; miRNAs, microRNAs; PS, phosphatidylserine; MDMs, monocyte-derived macrophages; AD, Alzheimer disease; A $\beta$ , amyloid beta; APP, amyloid precursor protein; APP-CTFs, C-terminal fragments of APP; PD, Parkinson's disease; ALP, autophagy-lysosomal pathway; ALS, amyotrophic lateral sclerosis; SOD1, superoxide dismutase-1 protein; TARDBP, TAR DNA-binding protein; vCJD, variant Creutzfeldt–Jakob disease; MDDCs, monocyte-derived dendritic cells

Received 11.7.16; revised 12.9.16; accepted 19.9.16; Edited by A Verkhratsky

exosomes to sheep reticulocytes.<sup>2</sup> The release of these small vesicles into the extracellular environment was proposed as a mechanism by which reticulocytes could secrete transferrin receptor. This proposed mechanism was further supported by *in vitro* analysis of sheep reticulocytes, which demonstrated the selective loss of certain proteins from maturing cells.<sup>3</sup> An understanding of the role of exosomes in various cell types has evolved greatly. They are no longer viewed as waste bags; instead, exosomes are thought to have an important role as cargo-carrying vesicles mediating communication among different cells and tissues including the CNS.<sup>4</sup> Exosomes are known to carry nucleic acids (RNA, microRNAs (miRNA) and DNA), functional proteins (including those of viral origin) and other cellular products. In the literature extracellular vesicle (EV) subtypes have often been given names such as exosomes, microvesicles, ectosomes or microparticles based on their biogenesis, physical characteristics (such as size), or function.

A growing body of evidence suggests the involvement of exosomes in many neuroinflammatory diseases. These small vesicles are important in CNS communication as most CNS cells secrete these particles.<sup>4</sup> Cell–cell communication via exosomes can be envisioned to have an important role in pathogenesis through their ability to transmit disease-causing agents from one cell to the other. Indeed, exosomes have been associated with numerous neuroinflammatory diseases including Parkinson's, Alzheimer's and Creutzfeldt–Jakob diseases. Further research into the role of these vesicles in disease progression is important for the development of effective preventative and therapeutic options. The focus of this review is to examine the role of exosomes in the progression of various neurodegenerative disorders.

### Exosome Cargo


Exosomes are generated via inward budding of the late endosomal membrane with the newly formed intraluminal vesicles (ILVs) destined for one of the two outcomes: either the late endosome merges with a lysosome, which degrades the ILVs along with their cytoplasmically derived cargo or the late endosome binds to the plasma membrane and releases the ILVs as exosomes with their cargo into the extracellular environment. Specifically, lysosome-directed vesicle formation occurs via the endosomal sorting complexes required for transport (ESCRT) machinery, whereas exosome budding is directed by the sphingolipid ceramide on the membrane of the endosome.<sup>5</sup> Recent studies have shown that the small GTPase ADP ribosylation factor 6 (ARF6) and its effector phospholipase D2 (PLD2) regulate exosome release.<sup>6</sup> ARF6 and PLD2 function together to regulate the budding of ILVs into MVBs.<sup>6</sup> After formation of ILVs, exosomes are released from the cell upon fusion of the late endosome with the plasma membrane. These vesicles are then free to carry their cargo throughout the surrounding environment and are associated with juxtacrine, paracrine and endocrine uptake in the host tissue (see the detailed reviews discussing exosome biogenesis:<sup>7,8</sup> (Figure 1a). The specific composition of an exosome largely depends upon the originating cell and can vary widely depending on the cellular and environmental

factors. Large-scale proteomic and phosphoproteomic studies of exosomes derived from various cell types suggest that these vesicles shuttle a wide array of biologically relevant molecules, including lipids, carbohydrates, RNAs and proteins.<sup>9</sup> For example, the study from Knepper's group identified 1132 proteins contained within exosomes isolated from urine. In addition, unique phosphorylation sites have also been identified on exosomal proteins.<sup>10</sup> Interestingly, miRNAs are abundantly present in the exosomes.<sup>11</sup> miRNAs regulate gene expression at the post-transcriptional level by binding to the 3'-UTR and/or the coding regions of their target mRNAs.<sup>12</sup> In fact, hundreds of miRNAs have been found in exosomes.<sup>13</sup> Many cell types, including reticulocytes, epithelial cells, neurons and tumor cells, have been reported to deliver exosomal miRNAs to recipient cells.<sup>14,15</sup> In addition, Epstein–Barr virus (EBV)-infected cells have been shown to secrete exosomes containing EBV-encoded miRNAs.<sup>16</sup> Importantly, exosomal miRNAs can repress mRNAs in target cells and subsequently influence target cell function. Furthermore, these exosomal miRNAs have been implicated in a number of cellular processes and human diseases including cell migration, cell differentiation, cell viability, aging, neurodegeneration, cancer and immune disorders.<sup>8</sup> These studies support the notion that exosomes obtained from body fluids have the potential to serve as biomarkers of disease development and/or progression. In this regard, Witwer *et al.*<sup>17</sup> have suggested the need for standardization of specimen handling, appropriate normative controls, and isolation and analysis techniques for EVs/exosomes to facilitate comparison of results.

### Mechanism(s) of Exosome Interactions with Recipient Cells

A primary function of exosomes is their ability to deposit their cargo inside a recipient cell. Although many roles of exosomes have been extensively reported in the literature, detailed interactions between target cells and exosomes remain to be elucidated. Currently, it is hypothesized that the uptake of exosomes into a target cell occurs by one of the three mechanisms: phagocytosis, receptor-mediated endocytosis (RME) or direct fusion of exosomes with the plasma membrane of the recipient cell<sup>15,18,19</sup> (Figure 1b). The latter mechanism involves the release of exosomal cargo directly into a cell following the fusion of the exosomes with the plasma membrane of the recipient cell. RME, on the other hand, consists of binding of exosomal surface proteins to proteins on the plasma membrane of recipient cells, thereby facilitating the targeting of exosomes to distinct cell types. The exosomes can then either fuse directly with the plasma membrane or follow a different endocytic pathway consisting of fusing with the delimiting membrane of an endocytic compartment (i.e., endosome, lysosome, etc.).<sup>20,21</sup>

The uptake of exosomes by RME in any given cell is largely dependent on the exosomal surface protein and lipid composition, which in turn, is primarily based on the type and condition of the secreting cell. The types of exosomal surface proteins and the kinds of receptors on the target cell determine the interaction between the exosome and the recipient cell. In addition, it has been hypothesized that lipid receptors could



**Figure 1** (a) Exosomes delivery via various signaling pathways. Exosomes can deliver molecules to distant cells (paracrine), adjacent cells (juxtacrine) or neighboring cells (autocrine). (b) The uptake of exosomes into a target cell occurs by one of the three mechanisms: phagocytosis, receptor-mediated endocytosis (RME) or direct fusion of exosomes with the plasma membrane of the recipient cell

also have a vital role in exosome recognition by the target cells. For example, phosphatidylserine (PS) on exosomal outer surface can interact with PS receptors, TIM1 and TIM4,<sup>22</sup> inducing accumulation of neutral lipids in the recipient cells.<sup>23</sup> In regard to the role of proteins in this uptake, an elegant study by Escrevente *et al.*<sup>24</sup> has shown that pretreatment of exosomes and recipient cells with a broad specificity protease K significantly decreased exosomal uptake efficiency. This study concluded that proteins from both exosomes and cells were required for uptake.<sup>24</sup> It has also been shown that heat-shock proteins such as Hsp90/Hsp70, which are chaperone proteins enriched in exosomes can interact with receptors such as low-density lipoprotein receptor-related protein 1 (LRP1), and that this interaction is critical for recognition of exosomes by target cells.<sup>25</sup> These findings further validated the notion that the interaction between exosomal membrane proteins and recipient cell-surface proteins is fundamental for the uptake of exosomes via RME. Upon further inquiry, it was shown that blocking CD9

and CD81 tetraspanins, which are commonly expressed on exosomes, resulted in a significant decrease in the exosomal uptake efficiency by dendritic cells.<sup>26</sup>

It has been demonstrated that phagocytic cells including RAW 264.7 macrophages and U937 monocyte-derived macrophages (MDM) internalize exosomes via phagocytosis.<sup>27</sup> More specifically, Feng *et al.*<sup>27</sup> demonstrated that macrophages internalized exosomes more efficiently than other non-phagocytic cells. Using electron microscopy, it was observed that exosomes remained localized on the surface of non-phagocytic cells, whereas exosomes in the vicinity of phagocytic cells were enveloped into large cellular extensions. Furthermore, the colocalization of fluorescent phagocytic tracers and the PKH26-dyed exosomes further confirmed that phagocytosis was the primary exosomal uptake mechanism employed by the macrophages. Furthermore, actin polymerization, which is necessary for phagocytosis, when inhibited by either cytochalasin D or latrunculin B, resulted in significantly reduced efficiency of exosomal uptake. This

further reinforced the idea that phagocytosis is an important mechanism for exosome internalization in the macrophages and likely also in other phagocytic cell types.

It has also been documented that oligodendroglia-derived exosomes can be taken up by microglia via macropinocytosis.<sup>28</sup> Furthermore, the authors also demonstrated that oligodendroglia-derived exosomes colocalized with Lamp1, a lysosome or late endosome marker, indicating thereby that an endocytic pathway was involved in the uptake of these exosomes. In order to deduce that macropinocytosis was indeed the mechanism in question, the microglia were treated with several reagents that interfered with specific steps in the macropinocytosis pathway. Following inhibition of macropinocytosis, microglial uptake of exosomes was inhibited, thereby underscoring the role of macropinocytosis in internalization of exosomes by microglia.<sup>28</sup>

### HIV Budding and Exosome Biogenesis

HIV acquires its envelope and propagates infection by budding through the limiting membranes of infected cells. HIV usurps a cellular pathway – that of formation of ILVs into MVB, to facilitate budding, indicating thereby that HIV budding and exosome release share a common mechanism. Indeed, published data on HIV-infected macrophages demonstrate the presence of HIV-containing vacuoles in macrophages that are reminiscent of MVBs.<sup>29</sup> Using human MDMs, the study by Nguyen *et al.*<sup>30</sup> revealed that the host protein profile of macrophage-derived exosomes and that of the HIV particles have a strong concordance, supporting the hypothesis that retroviral budding results from the exploitation of a pre-existing cellular pathway of intercellular vesicle trafficking. Proteomic analyses revealed that MDM-derived HIV virions contained 26 of 37 cellular proteins previously found in exosomes, consistent with the idea that HIV uses the late endosome/MVB pathway during virion budding from macrophages.<sup>21</sup> In addition, Bieniasz *et al.*<sup>31</sup> demonstrated that ESCRT components are recruited by HIV gag at the plasma membrane site of HIV budding. Nabhan *et al.*<sup>32</sup> revealed a role for ESCRT components in ectosomes budding at the plasma membrane. It is thus speculated that the budding of virus and EVs could utilize similar cellular pathways at the plasma membrane and inside the MVBs. An excellent review on the similarities between EVs and viral entry has been published and could provide further insights on this phenomenon.<sup>33</sup> Furthermore, it is also interesting to note that the HIV-1 virion has a similar size to that of the exosomes, an evidence supporting a similar origin of the two. This, however, also poses a potential confound for the isolation of exosomes from HIV-1-infected materials.<sup>34</sup> In addition, studies on T cells and exosomes have been inconsistent and confounding. For example, some studies implicate that HIV-1 budding does not involve either endosomes or exosomes,<sup>34–36</sup> whereas other reports have shown that HIV-1 budding from T cells is closely associated with exosomes.<sup>37</sup> In this regard, Park and He<sup>34</sup> have shown that high-speed centrifugation with 20% sucrose cushion during the last step can yield exosome-free HIV-1 virions compared with centrifugation only. Herein the authors provided a technical platform that could be employed to define the relationship between exosome biogenesis and budding of

HIV-1.<sup>34</sup> Though the small GTPase ARF6 and its effector PLD2 have an important role in exosome release, ARF6 is not involved in HIV-1 budding.<sup>6</sup> Early studies on HIV budding demonstrated that loss of the viral envelope Gag p6 domain caused a severe defect in virus budding.<sup>38</sup> However, recent studies showed that the N-terminal 433 amino acids of HIV Gag-Pol were sufficient to cause budding from cells, supporting the hypothesis that HIV budding was mediated by the exosome/microvesicle biogenesis pathway.<sup>39</sup> In other cell types such as the dendritic cells, exosomes can be internalized and transfer signaling molecules to the recipient cells. HIV-1 particles exploit this exosome-dissemination pathway to spread infection to the dendritic cells, thereby underscoring a potentially new viral dissemination pathway (Figure 2 and Table 1). Taken together, it remains an exciting question of whether HIV-1 budding and exosomal biogenesis are closely related processes and/or whether they mutually influence the respective processes.

### HIV Infection Alters Exosomal Release and Composition

Both exosome release as well as exosomal composition are regulated through cell signaling pathways that are activated by many factors, including but not limited to, HIV infection and subsequent immune activation. For example, it has been shown by Kadiu *et al.* that exosome numbers are increased in MDMs following HIV-1 infection.<sup>40</sup> Furthermore, HIV-1 was shown to accelerate infection and viral dissemination by surrounding itself with exosomes.<sup>40</sup> HIV infection not only affects exosome release, but also impacts exosomal cargo. Indeed, large-scale proteomics studies have revealed that compared with uninfected cells, exosomes released from HIV-1-infected cells harbor distinct regulatory molecules and are composed of a unique and quantitatively different protein signature.<sup>41,42</sup> Fourteen proteins out of 770 were identified to be differentially expressed in the exosomal fractions of HIV-1-infected cells compared with the uninfected cells. Three immunomodulatory molecules included ADP-ribosyl cyclase 1 (CD38), L-lactate dehydrogenase B chain and Annexin A5.<sup>41</sup> Recent studies have found that HIV-1 RNAs, such as HIV miRNA TAR, can also be incorporated into exosomes released from HIV-infected cells.<sup>43,44</sup> Furthermore, TAR RNA is sorted into exosomes in a chromosome region maintenance 1-dependent manner.<sup>44</sup> Importantly, exosomal TAR inhibited apoptosis by downregulating Bim and Cdk9 protein levels in recipient cells,<sup>44</sup> and stimulated proinflammatory cytokines, including IL-6 and TNF- $\beta$  in primary macrophages.<sup>45</sup> In a separate study, it was also shown that specific miRs such as miR-29b are transported via the EVs from HIV Tat and morphine-treated astrocytes to neurons and that, this transfer resulted in downregulation of PDGF-B (miR-29b target) in neurons, leading to neuronal apoptosis<sup>46</sup> (Figure 3). A recent study by Yelamanchili *et al.*<sup>47</sup> has also suggested increased expression of miR-21 in EVs derived from SIV-infected brains compared with the uninfected controls. Herein the authors demonstrated that in the brains of macaques with SIV-encephalitis, EV-miR-21 from donor macrophages/microglia resulted in neurotoxicity via activation of a TLR7-dependent downstream cell death pathway<sup>47</sup> (Figure 3). HIV-1 infection of MDMs resulted in significant upregulation of a






Table 1 Similarities and differences among HIV and EVs

	HIV	Exosome	Microvesicle, Ectosome	Apoptotic body
Size	90–160 nm	30–100 nm	100–1000 nm	50–5000 nm
Origin cell	Infected cell	Resting or activated cell	Activated or tumor cell	Apoptotic cell
Marker/Protein	Gag, Pol, gp120, Tat, Rev, Nef, Vpr, Vif and Vpu	CD63, CD9, Alix, Tsg101 and HSP70	Annexin V, flotillin-2, selectin, integrin and CD40 metalloproteinase	Annexin V, DNA and histones
Budding mechanism	Exocytosis from MVB	Exocytosis from MVB	Budding directly from a plasma membrane (outward budding)	Programmed cell death (blebbing)
Involved components	ESCRT components, Tsg101, Alix and HIV-1 Gag	nSMase2, ceramide, ESCRT components and ARDC1, Tsg101, Alix and tetraspanin	aSMase, ceramide, ESCRT components, Tsg101, Vps4 and tetraspanin	Caspase-3, rho-associated coiled-coil-forming kinase I and actin polymerization
Entry mechanism	HIV viral envelope, gp120 and gp41, binding CD4	Direct fusion, phagocytosis and RME (Figure 1).	Direct fusion, phagocytosis and RME (Figure 1).	Phagocytosis
Occur position	Plasma membrane and MVB	Plasma membrane and MVB	Plasma membrane	Plasma membrane

exosomal proteins such as Alix and flotillins have been reported to be localized within the amyloid plaques in the brains of the Tg2576 mice (AD model) as well as in the postmortem tissues of human AD patients.<sup>69,70</sup> In support of this is also another clinical report indicating upregulation of AD pathogenic proteins including P-T181-tau, P-S396-tau and A $\beta_{1-42}$  in the plasma exosomes isolated from AD patients, compared with cognitively normal-matched healthy individuals.<sup>67</sup> Furthermore, mechanistic studies have also implicated that both cleavage and endocytic transportation of amyloid precursor protein (APP) have cardinal roles in packaging A $\beta$  into exosomes for dispersion.<sup>70-72</sup> A $\beta$  is a cleaved product resulting from the cleavage of APP by the  $\beta$ - and  $\gamma$ -secretases.<sup>73</sup> A finding by Rajendran *et al.*<sup>70</sup> demonstrated that A $\beta$  was sorted into MVBs in both HeLa and N2a cells following  $\beta$ -cleavage in the early endosomes. Interestingly, Sharples *et al.*<sup>72</sup> also reported that inhibition of  $\gamma$ -secretase stimulated  $\alpha$ - and  $\beta$ -cleavage, leading in turn, to C-terminal fragments of APP (APP-CTFs) in the exosomes. Furthermore,  $\gamma$ -secretase has also been shown to be required for the clearance of APP-CTFs from the endocytic recycling compartments, which constitute a series of perinuclear tubular and vesicular membranes, that regulate recycling of APP-CTFs to the plasma membrane.<sup>74,75</sup> In addition, deficits in retromer, a multimeric complex that mediates retrograde protein transportation from endosome to the trans-Golgi network, has also been shown to promote amyloidogenic APP processing by enhancing interactions between APP and secretase enzymes in the late endosomes.<sup>71</sup> As endocytic trafficking is becoming increasingly recognized as a possible mechanism(s) of AD pathogenesis, examining the role of other endocytic trafficking regulators such as diacylglycerol kinase, Eps15 homology domain and molecules interacting with CasL-like1 (MICAL-L1) in both processing and release of APP via the exosomes can be developed as future areas of research that will provide valuable insights into the pathogenesis of AD.<sup>76-78</sup> Interestingly, aggregated tau protein has also been found to be present in the exosomes in both the *in vitro* taupathy models as well as in the cerebrospinal fluids of early Alzheimer's patients.<sup>68</sup> It is worth noting that microglia have an important role in spreading both A $\beta$  and tau through the EVs.<sup>79,80</sup> The advent and widespread application of new imaging techniques such as super-resolution microscopy and quantitative methodology are also avenues that continue to contribute in our understanding of the molecular mechanism (s) involved in the processing of APP and tau and their roles in the pathogenesis of AD.<sup>75,81</sup>

### Parkinson's Disease and EVs

Parkinson's disease (PD), clinically characterized by hypokinesia, rigidity and tremor, is the second most common neurodegenerative disorder. The pathological features of PD comprise of widespread degeneration of dopaminergic neurons and aggregation of Lewy bodies and cytoplasm inclusion bodies of  $\alpha$ -synuclein.<sup>82</sup> Interestingly, similar to the spread of pathogenic tau and A $\beta$  proteins, EVs also have crucial roles in the aggregation and the spread of  $\alpha$ -synuclein, thereby propagating disease pathogenesis in PD.<sup>83-85</sup>



**Figure 3** Exosome released from astrocytes treated with morphine and Tat carry miR-29b, which can be taken up by neurons, resulting in neuronal death. MiR-21 in EVs leads to neurotoxicity via the TLR7 signaling pathway

Increased levels of  $\alpha$ -synuclein have been reported in the exosomes isolated from both plasma and CSF of PD patients with a significant correlation observed between disease severity and plasma exosomal  $\alpha$ -synuclein levels.<sup>85</sup> It has been reported that spread of  $\alpha$ -synuclein between neurons via the exosome route confers cytotoxicity to the recipient cells, leading to increased accumulation of Lewy body throughout the various brain regions.<sup>84</sup> Although it is not well understood how this process is regulated, both Tsunemi *et al.*<sup>86</sup> and Kong *et al.*<sup>87</sup> suggest the role of P-type ATPase ion pump (PARK9/ATP13A2) in regulating both the biogenesis of exosomes, as well as secretion of exosomal  $\alpha$ -synuclein. In case of juvenile-onset PD, that is attributed to loss-of-function mutations in PARK9, it was demonstrated that knocking down of PARK9 resulted in the inhibition of exosomal secretion of  $\alpha$ -synuclein, and that, reciprocally, overexpression of PARK9 resulted in localization of PARK9 in MVBs and was associated with release of  $\alpha$ -synuclein from the exosomes.<sup>86,87</sup>

Another molecular mechanism regulating the exosomal release of  $\alpha$ -synuclein is the autophagy-lysosomal pathway (ALP) that degrades the enclosed cargo in the lysosome.<sup>88–90</sup> It can be speculated that exosomal release of  $\alpha$ -synuclein is likely an adaptive response to insufficient autophagic activity needed for the elimination of  $\alpha$ -synuclein.<sup>89</sup> Using bafilomycin A1, a pharmacological inhibitor that disrupts ALP by blocking fusion between autophagosomes and lysosomes, Alvarez-Erviti *et al.*<sup>91</sup> demonstrated that disruption of lysosomal functions resulted in enhanced exosomal release and uptake of  $\alpha$ -synuclein in SH-SY5Y cells. In line with this, Poehler *et al.*<sup>88</sup> also observed that ameliorating cytosolic accumulation of  $\alpha$ -synuclein in bafilomycin A1-treated  $\alpha$ -synuclein-transgenic mice, resulted in enhanced exosomal release of  $\alpha$ -synuclein, which in turn, resulted in neuroinflammation and


cellular damage. It is also worth noting that there is evidence suggesting that the gangliosides present in the exosomes can contribute to a catalytic environment leading to increased aggregation of  $\alpha$ -synuclein.<sup>83</sup>

The role of glial cells in mediating neuroinflammatory responses leading to neurodegenerative disorders has been well documented.<sup>92–94</sup> It is not surprising, therefore, that microglial/monocytes also have a role in the PD through the regulation of exosomal activities.<sup>95</sup> For example, Chang *et al.*<sup>95</sup> found that exosomes derived from  $\alpha$ -synuclein-exposed BV-2 microglia contained high levels of MHC class II and membrane TNF- $\alpha$  and, that treatment of rat cortical neurons with these activated exosomes resulted in the neuronal apoptosis.

### Amyotrophic Lateral Sclerosis and EVs

Amyotrophic lateral sclerosis (ALS) is a progressive neurodegenerative disorder often accompanied with frontotemporal dementia.<sup>96</sup> In the United States ALS affects ~3.9 cases per 100 000 individuals, with increased prevalence in persons aged 60–69 years.<sup>97</sup> Loss of spinal cord motor neurons is the prominent pathological feature of ALS that manifests as muscle weakness and respiratory failure.<sup>98,99</sup> Several genes have been found to be associated with ALS including superoxide dismutase-1 protein (SOD1), RNA-binding protein fused in sarcoma and TAR DNA-binding protein 43 (TARDBP).<sup>100,101</sup>

Similar to A $\beta$ , mutations of SOD1 in ALS result in aggregation of intracellular misfolded SOD1 protein and its spread.<sup>102</sup> Both mutant and wild-type SOD1 have been shown to be transmitted through both micropinocytosis of the released protein aggregates from dying cells as well as by uptake of SOD1 in the EVs by the recipient cells.<sup>102</sup> Secretion



**Figure 4** The exosome response to HIV infection. (a) Exosomes from HIV-infected cells contain various molecules including ADAM17, TNF $\alpha$ , Nef and so on. These molecules upon entering the recipient cells then induce the target cell activation, apoptosis and HIV susceptibility. (b) Exosomes released from CD8 $^+$  T cells contain an antiviral membrane-bound factor that inhibits HIV-1 transcription

of exosomal SOD1 has also been observed in a widely used *in vitro* model of ALS that of mouse motor neuron-like NSC-34 cells overexpressing the mutant human SOD1 (G93A).<sup>103,104</sup> Moreover, it has also been reported that mutant SOD1-containing exosomes released from the astrocytes exerted neurotoxicity.<sup>105</sup> Mechanisms by which SOD1 is packaged into the exosomes, however, remain elusive. Intriguingly, the expression of mutant SOD1 (G127X, G85R) has been reported on the surface of exosomes,<sup>104</sup> whereas that of the wild-type SOD1 is in the lumen of the exosomes.<sup>106,107</sup> Upon uptake by recipient cells, mutant SOD1 proteins function as templates for further misfolding and aggregation of the natively folded counterparts.<sup>102</sup> Studies by Grad *et al.*<sup>104</sup> reported inhibition of the conversion of SOD1 into its misfolded form in the recipient cells via the treatment of exosomes with antibodies specific for mutant SOD1, indicating thereby the therapeutic potential of drug delivery by exosomes.

Similarly, TDP-43 aggregates have also implicated to be released in the EVs and to initiate the intracellular aggregation of TDP-43 in the recipient cells.<sup>108</sup> Although TDP-43 aggregates are present in both microvesicles and exosomes, microvesicular TDP-43 has been shown to be the favored source for uptake by the recipient cells, leading in turn, to increased toxicity compared with the free TDP-43.<sup>109</sup> In the same study it has also been suggested that TDP-43 could also spread trans-synaptically in both the anterograde and retrograde manner. In addition, it was also shown that neurons exposed to either TDP-43-containing media (derived from cultured cells) or to brain lysates from ALS patients did take up TDP-43.

### Prion Disease and EVs

Prion diseases are a spectrum of transmissible lethal neurodegenerative diseases characterized by cognitive impairment, motor dysfunction, spongiosis, astrogliosis and cerebral deposition of insoluble PRNP (prion protein).<sup>110,111</sup>

In prion diseases, the native form of prion protein PRNP<sup>C</sup> is converted into PRNP<sup>SC</sup>, a pathogenic form that is protease-resistant and prone to aggregation.<sup>112</sup>

PRNP has been identified in the EVs derived from both CSF<sup>113</sup> and blood.<sup>114</sup> Accumulating evidence further suggests a key role of EVs in both the pathogenesis and propagation of prion diseases.<sup>115–117</sup> One of the initial studies has demonstrated that PRNP-expressing cell line (RK13) robustly secreted PRNP through exosomes.<sup>118</sup> Vella *et al.*<sup>119</sup> further showed that exosomes released from PRNP-infected neuronal cell line (GT1-7) also induced prion propagation in both neuronal as well as non-neuronal cells. Guo *et al.*<sup>120</sup> have shown that inhibiting exosome release reduces the intercellular transmission of PRNP in different PRNP-expressing cell lines *in vitro*. Taking advantage of pharmacological inhibitors and genetic approaches, the same group also demonstrated that neutral sphingomyelinase pathway has a crucial role in regulating the packaging of PRNP into exosomes.<sup>121</sup> Consistent with these *in vitro* findings, it has also been shown that EVs derived from plasma of mice infected with variant Creutzfeldt–Jakob disease (vCJD), one of the prion diseases, contains PRNP.<sup>122</sup> Due to the fact that transmission of vCJD has also been clinically reported to be associated with blood transfusion,<sup>123</sup> it is plausible to hypothesize that EVs might have a significant role in the transmission of prion diseases. Interestingly, upregulated miRNAs including let-7b, miR-146a, miR-103, miR-125a-5p and miR-342-3p were found to be present in the EVs isolated from human tissue samples affected with prion diseases,<sup>124</sup> suggesting thereby that PRNP-infected EVs also mediate the pathogenesis of prion disease through their contents in addition to spreading of PRNP.

However, much still remains to be discovered about the role of exosomes in prion diseases. It has been demonstrated that different strains of PRNP are secreted differentially from RK13 cell line possibly through various disparate cellular mechanisms.<sup>125</sup> A better understanding of the role of EVs in prion diseases is thus critical in dissecting mechanism(s)



underlying disease pathogenesis while also providing insights for other prion-like diseases such as AD and PD.<sup>126</sup>

### Exosomes as Therapeutic Conduits

Exosomes via their ability to deliver specific cargo are critical for cellular communication and physiology. This property of exosomes can also be exploited as a treatment strategy to deliver specific therapeutic molecules to the diseased tissues. An elegant study by Zhuang *et al.*<sup>127</sup> analyzed the potential for exosome therapeutics, through a noninvasive nasal delivery method, for the treatment of neuroinflammatory diseases. Herein the authors showed that exosome-mediated delivery of curcumin and an inhibitor of signal transducer and activator of transcription selectively and rapidly targeted the microglia, and significantly mitigated disease pathogenesis in three experimental models of neuroinflammation. This study highlights a novel and a noninvasive therapeutic approach for drug delivery into the CNS. A review by Andaloussi *et al.*<sup>128</sup> covers the potential for exosomes as a delivery system for transporting siRNAs across biological barriers such as the blood–brain barrier. Exosomes-containing siRNAs could provide a means of targeted therapeutic delivery into areas that are difficult to traverse, such as the brain.<sup>129,130</sup> The ability to non-invasively deliver therapeutic exosomes to the CNS, with cell-specific targeting, could provide a potential therapeutic approach for the treatment and eradication of HIV-associated neurocognitive disorders by targeting the latent reservoirs in the CNS. Indeed, the application of exosomal delivery of natural HIV-defense molecules to host cells has been shown to establish HIV resistance to uninfected cells of *vif*-deficient HIV infection.<sup>131</sup> Combining noninvasive therapies with siRNA targeting to HIV-1/HIV proteins could, thus, provide new therapeutic approaches to treat HIV-associated end-organ pathologies. Delivery of HIV gene-specific siRNA to infected cells could control virus infection and could be considered as adjunctive treatment(s) in combination with other therapeutic modalities. An elegant review by Vlassov *et al.*<sup>132</sup> covering a broader view of exosome therapeutics, including their use against tumors, sets the stage for future work. Further studies on specific applications of exosome-mediated therapeutic delivery to specific cells are warranted in the field.

Intriguingly, assessment of the specific, signature contents of disease-specific exosomes could provide useful information for future development of exosome-based therapeutics. Indeed, exosomes from milk, but not plasma, have been shown to contain inhibitory factors for HIV-1 infection of monocyte-derived dendritic cells (MDDCs) and for the subsequent viral transfer to CD4 T cells through binding of MDDCs via DC-SIGN.<sup>133</sup> In addition, there are several reports highlighting the potential for delivery of modified miRNAs and normal siRNAs to specific targets via the exosomes.<sup>129,130,134,135</sup> For more details please refer to the comprehensive review on the specific potential of targeting HIV and exosomal miRNAs as therapeutic approaches.<sup>8,136,137</sup>

Besides the targeting of exosomal contents, the exosome itself could be a potential therapeutic target. It is well documented that accumulation of reactive oxygen species underlies PD-associated neuroinflammation and neurotoxicity.<sup>138</sup> Exosomes loaded with either antioxidant enzyme catalase or a

plasmid DNA-encoding catalase have been used to reduce neuroinflammatory responses and exert neuroprotective effects.<sup>139,140</sup> Similarly, neurotropic factors that improve neuronal functioning can also be developed as potential therapeutic cargos for exosomal delivery. Along these lines Zhao *et al.* have demonstrated that delivery of macrophage exosomes containing glial cell line-derived neurotrophic factor ameliorated neurodegeneration and neuroinflammation in PD mice.<sup>141</sup>

### Conclusions

In summary the exponential growth in our understanding of exosome biogenesis, composition, function and their use continues to provide new insights into the normal physiology as well as disease processes. These small vesicles are secreted by many cell types, including all of the CNS cells, and are a key component for cell–cell communication. They have crucial roles in various diseases including cancer metastasis. This review focuses mainly on the possible link between exosomes, HIV-1 pathogenesis and HIV-associated CNS disease. The ability to target exosomes involved in HIV-1 pathogenesis could provide a new means of controlling infection, which in turn, could help to curtail many of the commonly associated complications of the CNS. Further studies are needed concerning the application of exosome therapeutics, involving the use of these vesicles as a drug delivery conduits and as therapeutic targets themselves, in the context of battling HIV-1 infection and its associated end-organ pathologies.

### Conflict of Interest

The authors declare no conflict of interest.

**Acknowledgements.** This work was supported by grants DA033150, DA035203, DA036157, DA040397, MH106425 and DA043138 (SB), and the pilot grant (GH) from CHAIN Center grant (P30 MH062261) (HSF) from the National Institutes of Health. The support by Nebraska Center for Substance Abuse Research is also acknowledged.

1. Trams EG, Lauter CJ, Salem N Jr., Heine U. Exfoliation of membrane ecto-enzymes in the form of micro-vesicles. *Biochim Biophys Acta* 1981; **645**: 63–70.
2. Pan BT, Teng K, Wu C, Adam M, Johnstone RM. Electron microscopic evidence for externalization of the transferrin receptor in vesicular form in sheep reticulocytes. *J Cell Biol* 1985; **101**: 942–948.
3. Johnstone RM, Adam M, Hammond JR, Orr L, Turbide C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* 1987; **262**: 9412–9420.
4. Gupta A, Pulliam L. Exosomes as mediators of neuroinflammation. *J Neuroinflammation* 2014; **11**: 68.
5. Trajkovic K, Hsu C, Chiantia S, Rajendran L, Wenzel D, Wieland F *et al.* Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science* 2008; **319**: 1244–1247.
6. Ghossoub R, Lembo F, Rubio A, Gaillard CB, Bouchet J, Vitale N *et al.* Syntenin-ALIX exosome biogenesis and budding into multivesicular bodies are controlled by ARF6 and PLD2. *Nat Commun* 2014; **5**: 3477.
7. Thery C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. *Nat Rev Immunol* 2002; **2**: 569–579.
8. Hu G, Drescher KM, Chen XM. Exosomal miRNAs: biological properties and therapeutic potential. *Front Genet* 2012; **3**: 56.
9. Schorey JS, Bhatnagar S. Exosome function: from tumor immunology to pathogen biology. *Traffic* 2008; **9**: 871–881.
10. Gonzales PA, Pisitkun T, Hoffert JD, Tchoprynjikov D, Star RA, Kleta R *et al.* Large-scale proteomics and phosphoproteomics of urinary exosomes. *J Am Soc Nephrol* 2009; **20**: 363–379.

11. Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 2007; **9**: 654–659.
12. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009; **136**: 215–233.
13. Mathivanan S, Fahner CJ, Reid GE, Simpson RJ. ExoCarta 2012: database of exosomal proteins, RNA and lipids. *Nucleic Acids Res* 2012; **40**: D1241–D1244.
14. Faure J, Lachenal G, Court M, Hirrlinger J, Chatellard-Causse C, Blot B et al. Exosomes are released by cultured cortical neurons. *Mol Cell Neurosci* 2006; **31**: 642–648.
15. Thery C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol* 2009; **9**: 581–593.
16. Meckes DG Jr, Shair KH, Marquitz AR, Kung CP, Edwards RH, Raab-Traub N. Human tumor virus utilizes exosomes for intercellular communication. *Proc Natl Acad Sci USA* 2010; **107**: 20370–20375.
17. Witwer KW, Buzas EI, Bemis LT, Bora A, Lasser C, Lotvall J et al. Standardization of sample collection, isolation and analysis methods in extracellular vesicle research. *J Extracell Vesicles* 2013; **2**.
18. Keller S, Konig AK, Marme F, Runz S, Wolterink S, Koensgen D et al. Systemic presence and tumor-growth promoting effect of ovarian carcinoma released exosomes. *Cancer Lett* 2009; **278**: 73–81.
19. Cocucci E, Racchetti G, Meldolesi J. Shedding microvesicles: artefacts no more. *Trends Cell Biol* 2009; **19**: 43–51.
20. Raposo G, Stoorvogel W. Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol* 2013; **200**: 373–383.
21. Chertova E, Chertov O, Coren LV, Roser JD, Trubey CM, Bess JW Jr et al. Proteomic and biochemical analysis of purified human immunodeficiency virus type 1 produced from infected monocyte-derived macrophages. *J Virol* 2006; **80**: 9039–9052.
22. Miyanishi M, Tada K, Koike M, Uchiyama Y, Kitamura T, Nagata S. Identification of Tim4 as a phosphatidylserine receptor. *Nature* 2007; **450**: 435–439.
23. Zakharova L, Svetlova M, Fomina AF. T cell exosomes induce cholesterol accumulation in human monocytes via phosphatidylserine receptor. *J Cell Physiol* 2007; **212**: 174–181.
24. Escrevente C, Keller S, Altevogt P, Costa J. Interaction and uptake of exosomes by ovarian cancer cells. *BMC Cancer* 2011; **11**: 108.
25. Skokos D, Botros HG, Demeure C, Morin J, Peronet R, Birkenmeier G et al. Mast cell-derived exosomes induce phenotypic and functional maturation of dendritic cells and elicit specific immune responses *in vivo*. *J Immunol* 2003; **170**: 3037–3045.
26. Kooijmans SA, Vader P, van Dommelen SM, van Solinge WW, Schiffelers RM. Exosome mimetics: a novel class of drug delivery systems. *Int J Nanomedicine* 2012; **7**: 1525–1541.
27. Feng D, Zhao WL, Ye YY, Bai XC, Liu RQ, Chang LF et al. Cellular internalization of exosomes occurs through phagocytosis. *Traffic* 2010; **11**: 675–687.
28. Fitzner D, Schnaars M, van Rossum D, Krishnamoorthy G, Dibaj P, Bakhti M et al. Selective transfer of exosomes from oligodendrocytes to microglia by macropinocytosis. *J Cell Sci* 2011; **124**: 447–458.
29. Stoorvogel W, Kleijmeer MJ, Geuze HJ, Raposo G. The biogenesis and functions of exosomes. *Traffic* 2002; **3**: 321–330.
30. Nguyen DG, Booth A, Gould SJ, Hildreth JE. Evidence that HIV budding in primary macrophages occurs through the exosome release pathway. *J Biol Chem* 2003; **278**: 52347–52354.
31. Bieniasz PD. The cell biology of HIV-1 virion genesis. *Cell Host Microbe* 2009; **5**: 550–558.
32. Nabhan JF, Hu R, Oh RS, Cohen SN, Lu Q. Formation and release of arrestin domain-containing protein 1-mediated microvesicles (ARMMs) at plasma membrane by recruitment of TSG101 protein. *Proc Natl Acad Sci USA* 2012; **109**: 4146–4151.
33. van Dongen HM, Masoumi N, Witwer KW, Pegtel DM. Extracellular vesicles exploit viral entry routes for cargo delivery. *Microbiol Mol Biol Rev* 2016; **80**: 369–386.
34. Park IW, He JJ. HIV-1 is budded from CD4+ T lymphocytes independently of exosomes. *Viral J* 2010; **7**: 234.
35. Garrus JE, von Schwedler UK, Pornillos OW, Morham SG, Zavitz KH, Wang HE et al. Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding. *Cell* 2001; **107**: 55–65.
36. Morita E, Sundquist W. Retrovirus budding. *Annu Rev Cell Dev Biol* 2004; **20**: 395–425.
37. Booth AM, Fang Y, Fallon JK, Yang JM, Hildreth JE, Gould SJ. Exosomes and HIV Gag bud from endosome-like domains of the T cell plasma membrane. *J Cell Biol* 2006; **172**: 923–935.
38. Gottlinger HG, Dorfman T, Sodroski JG, Haseltine WA. Effect of mutations affecting the p6 gag protein on human immunodeficiency virus particle release. *Proc Natl Acad Sci USA* 1991; **88**: 3195–3199.
39. Gan X, Gould SJ. HIV Pol inhibits HIV budding and mediates the severe budding defect of Gag-Pol. *PLoS One* 2012; **7**: e29421.
40. Kadiu I, Narayanasamy P, Dash PK, Zhang W, Gendelman HE. Biochemical and biologic characterization of exosomes and microvesicles as facilitators of HIV-1 infection in macrophages. *J Immunol* 2012; **189**: 744–754.
41. Li M, Aliotta JM, Asara JM, Tucker L, Quesenberry P, Lally M et al. Quantitative proteomic analysis of exosomes from HIV-1-infected lymphocytic cells. *Proteomics* 2012; **12**: 2203–2211.
42. Jaworski E, Narayanan A, Van Duyn R, Shabbeer-Meyering S, Iordanskiy S, Saifuddin M et al. Human T-lymphotropic virus type 1-infected cells secrete exosomes that contain Tax protein. *J Biol Chem* 2014; **289**: 22284–22305.
43. Columba Cabezas S, Federico M. Sequences within RNA coding for HIV-1 Gag p17 are efficiently targeted to exosomes. *Cell Microbiol* 2013; **15**: 412–429.
44. Narayanan A, Iordanskiy S, Das R, Van Duyn R, Santos S, Jaworski E et al. Exosomes derived from HIV-1-infected cells contain trans-activation response element RNA. *J Biol Chem* 2013; **288**: 20014–20033.
45. Sampey GC, Saifuddin M, Schwab A, Barclay R, Punya S, Chung MC et al. Exosomes from HIV-1-infected cells stimulate production of pro-inflammatory cytokines through trans-activating response (TAR) RNA. *J Biol Chem* 2016; **291**: 1251–1266.
46. Hu G, Yao H, Chaudhuri AD, Duan M, Yelamanchili SV, Wen H et al. Exosome-mediated shuttling of microRNA-29 regulates HIV Tat and morphine-mediated neuronal dysfunction. *Cell Death Dis* 2012; **3**: e381.
47. Yelamanchili SV, Lamberty BG, Rennard DA, Morse BM, Hochfelder CG, Meays BM et al. Correction: MIR-21 in extracellular vesicles leads to neurotoxicity via TLR7 signaling in SIV neurological disease. *PLoS Pathog* 2015; **11**: e1005131.
48. Roth WW, Huang MB, Adde Konadu K, Powell MD, Bond VC. Micro RNA in exosomes from HIV-infected macrophages. *Int J Environ Res Public Health* 2015; **13**: 32.
49. Ali SA, Huang MB, Campbell PE, Roth WW, Campbell T, Khan M et al. Genetic characterization of HIV type 1 Nef-induced vesicle secretion. *AIDS Res Hum Retroviruses* 2010; **26**: 173–192.
50. Campbell TD, Khan M, Huang MB, Bond VC, Powell MD. HIV-1 Nef protein is secreted into vesicles that can fuse with target cells and virions. *Ethn Dis* 2008; **18**: S2–14–19.
51. Lenassi M, Cagney G, Liao M, Vaupotic T, Bartholomeussen K, Cheng Y et al. HIV Nef is secreted in exosomes and triggers apoptosis in bystander CD4+ T cells. *Traffic* 2010; **11**: 110–122.
52. Muratori C, Cavallin LE, Kratzel K, Tinari A, De Milioto A, Fais S et al. Massive secretion by T cells is caused by HIV Nef in infected cells and by Nef transfer to bystander cells. *Cell Host Microbe* 2009; **6**: 218–230.
53. Raymond AD, Campbell-Sims TC, Khan M, Lang M, Huang MB, Bond VC et al. HIV Type 1 Nef is released from infected cells in CD45(+) microvesicles and is present in the plasma of HIV-infected individuals. *AIDS Res Hum Retroviruses* 2011; **27**: 167–178.
54. Shelton MN, Huang MB, Ali SA, Powell MD, Bond VC. Secretion modification region-derived peptide disrupts HIV-1 Nef's interaction with mortalin and blocks virus and Nef exosome release. *J Virol* 2012; **86**: 406–419.
55. Rahimian P, He JJ. Exosome-associated release, uptake, and neurotoxicity of HIV-1 Tat protein. *J Neurovirol* 2016 (e-pub ahead of print).
56. Konadu KA, Chu J, Huang MB, Amancha PK, Armstrong W, Powell MD et al. Association of cytokines with exosomes in the plasma of HIV-1-seropositive individuals. *J Infect Dis* 2015; **211**: 1712–1716.
57. Schwab A, Meyering SS, Lepene B, Iordanskiy S, van Hoek ML, Hakami RM et al. Extracellular vesicles from infected cells: potential for direct pathogenesis. *Front Microbiol* 2015; **6**: 1132.
58. Jaworski E, Saifuddin M, Sampey G, Shafagati N, Van Duyn R, Iordanskiy S et al. The use of Nanotrap particles technology in capturing HIV-1 virions and viral proteins from infected cells. *PLoS One* 2014; **9**: e96778.
59. Arenaccio C, Chiozzini C, Columba-Cabezas S, Manfredi F, Affabris E, Baur A et al. Exosomes from human immunodeficiency virus type-1 (HIV-1)-infected cells license quiescent CD4+ T lymphocytes to replicate HIV-1 through a Nef- and ADAM17-dependent mechanism. *J Virol* 2014; **88**: 11529–11539.
60. Arenaccio C, Chiozzini C, Columba-Cabezas S, Manfredi F, Federico M. Cell activation and HIV-1 replication in unstimulated CD4+ T lymphocytes ingesting exosomes from cells expressing defective HIV-1. *Retrovirology* 2014; **11**: 46.
61. Aqil M, Naqvi AR, Mallik S, Bandyopadhyay S, Maulik U, Jameel S. The HIV Nef protein modulates cellular and exosomal miRNA profiles in human monocyte cells. *J Extracell Vesicles* 2014; **3**.
62. Gulzar N, Copeland KF. CD8+ T-cells: function and response to HIV infection. *Curr HIV Res* 2004; **2**: 23–37.
63. Turne A, Prasad VS, Chen Y, Stolz DB, Saha K, Ratner DM et al. Noncytotoxic suppression of human immunodeficiency virus type 1 transcription by exosomes secreted from CD8+ T cells. *J Virol* 2009; **83**: 4354–4364.
64. Prince M, Wimo A, Guerchet M, Ali G, Wu Y, Prina M. World Alzheimer Report 2015. The global impact of dementia. An analysis of prevalence, incidence, cost and trends. *Alzheimers Dis Int* 2015.
65. Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 2001; **81**: 741–766.
66. Goetzl EJ, Boxer A, Schwartz JB, Abner EL, Petersen RC, Miller BL et al. Low neural exosomal levels of cellular survival factors in Alzheimer's disease. *Ann Clin Transl Neurol* 2015; **2**: 769–773.
67. Fiandaca MS, Kapogiannis D, Mapstone M, Boxer A, Eitan E, Schwartz JB et al. Identification of preclinical Alzheimer's disease by a profile of pathogenic proteins in neurally derived blood exosomes: a case-control study. *Alzheimers Dement* 2015; **11**: 600–607 e601.
68. Saman S, Kim W, Raya M, Visnick Y, Miro S, Saman S et al. Exosome-associated tau is secreted in tauopathy models and is selectively phosphorylated in cerebrospinal fluid in early Alzheimer disease. *J Biol Chem* 2012; **287**: 3842–3849.
69. Kokubo H, Saido TC, Iwata N, Helms JB, Shinohara R, Yamaguchi H. Part of membrane-bound Abeta exists in rafts within senile plaques in Tg2576 mouse brain. *Neurobiol Aging* 2005; **26**: 409–418.

70. Rajendran L, Honsho M, Zahn TR, Keller P, Geiger KD, Verkade P et al. Alzheimer's disease beta-amyloid peptides are released in association with exosomes. *Proc Natl Acad Sci USA* 2006; **103**: 11172–11177.
71. Sullivan CP, Jay AG, Stack EC, Pakaluk M, Wadlinger E, Fine RE et al. Retromer disruption promotes amyloidogenic APP processing. *Neurobiol Dis* 2011; **43**: 338–345.
72. Sharples RA, Vella LJ, Nisbet RM, Naylor R, Perez K, Barnham KJ et al. Inhibition of gamma-secretase causes increased secretion of amyloid precursor protein C-terminal fragments in association with exosomes. *FASEB J* 2008; **22**: 1469–1478.
73. Cai Y, Arikath J, Yang L, Guo ML, Periyasamy P, Buch S. Interplay of endoplasmic reticulum stress and autophagy in neurodegenerative disorders. *Autophagy* 2016; **12**: 225–244.
74. Zhang M, Haapasalo A, Kim DY, Ingano LA, Pettingell WH, Kovacs DM. Presenilin/gamma-secretase activity regulates protein clearance from the endocytic recycling compartment. *FASEB J* 2006; **20**: 1176–1178.
75. Xie S, Bahl K, Reinecke JB, Hammond GR, Naslavsky N, Caplan S. The endocytic recycling compartment maintains cargo segregation acquired upon exit from the sorting endosome. *Mol Biol Cell* 2016; **27**: 108–126.
76. Xie S, Naslavsky N, Caplan S. Diacylglycerol kinases in membrane trafficking. *Cell Logist* 2015; **5**: e1078431.
77. Xie S, Naslavsky N, Caplan S. Diacylglycerol kinase alpha regulates tubular recycling endosome biogenesis and major histocompatibility complex class I recycling. *J Biol Chem* 2014; **289**: 31914–31926.
78. Cai B, Xie S, Caplan S, Naslavsky N. GRAF1 forms a complex with MICAL-L1 and EHD1 to cooperate in tubular recycling endosome vesiculation. *Front Cell Dev Biol* 2014; **2**: 22.
79. Joshi P, Turolo E, Ruiz A, Bergami A, Libera DD, Benussi L et al. Microglia convert aggregated amyloid-beta into neurotoxic forms through the shedding of microvesicles. *Cell Death Differ* 2014; **21**: 582–593.
80. Asai H, Ikezu S, Tsunoda S, Medalla M, Luebke J, Haydar T et al. Depletion of microglia and inhibition of exosome synthesis halt tau propagation. *Nat Neurosci* 2015; **18**: 1584–1593.
81. Reinecke JB, Xie S, Naslavsky N, Caplan S. Qualitative and quantitative analysis of endocytic recycling. *Methods Cell Biol* 2015; **130**: 139–155.
82. Pollanen MS, Dickson DW, Bergeron C. Pathology and biology of the Lewy body. *J Neuropathol Exp Neurol* 1993; **52**: 183–191.
83. Grey M, Dunning CJ, Gaspar R, Grey C, Brundin P, Sparr E et al. Acceleration of alpha-synuclein aggregation by exosomes. *J Biol Chem* 2015; **290**: 2969–2982.
84. Gallegos S, Pacheco C, Peters C, Opazo CM, Aguayo LG. Features of alpha-synuclein that could explain the progression and irreversibility of Parkinson's disease. *Front Neurosci* 2015; **9**: 59.
85. Shi M, Liu C, Cook TJ, Bullock KM, Zhao Y, Ghingina C et al. Plasma exosomal alpha-synuclein is likely CNS-derived and increased in Parkinson's disease. *Acta Neuropathol* 2014; **128**: 639–650.
86. Tsunemi T, Hamada K, Krainc D. ATP13A2/PARK9 regulates secretion of exosomes and alpha-synuclein. *J Neurosci* 2014; **34**: 15281–15287.
87. Kong SM, Chan BK, Park JS, Hill KJ, Aitken JB, Cottle L et al. Parkinson's disease-linked human PARK9/ATP13A2 maintains zinc homeostasis and promotes alpha-synuclein externalization via exosomes. *Hum Mol Genet* 2014; **23**: 2816–2833.
88. Poehler AM, Xiang W, Spitzer P, May VE, Meixner H, Rockenstein E et al. Autophagy modulates SNCA/alpha-synuclein release, thereby generating a hostile microenvironment. *Autophagy* 2014; **10**: 2171–2192.
89. Danzer KM, Kranich LR, Ruf WP, Cagsal-Getkin O, Winslow AR, Zhu L et al. Exosomal cell-to-cell transmission of alpha synuclein oligomers. *Mol Neurodegener* 2012; **7**: 42.
90. Cai B, Xie S, Liu F, Simone LC, Caplan S, Qin X et al. Rapid degradation of the complement regulator, CD59, by a novel inhibitor. *J Biol Chem* 2014; **289**: 12109–12125.
91. Alvarez-Erviti L, Seow Y, Schapira AH, Gardiner C, Sargent IL, Wood MJ et al. Lysosomal dysfunction increases exosome-mediated alpha-synuclein release and transmission. *Neurobiol Dis* 2011; **42**: 360–367.
92. Yang L, Yao H, Chen X, Cai Y, Callen S, Buch S. Role of sigma receptor in cocaine-mediated induction of glial fibrillary acidic protein: implications for HAND. *Mol Neurobiol* 2016; **53**: 1329–1342.
93. Duan M, Yao H, Cai Y, Liao K, Seth P, Buch S. HIV-1 Tat disrupts CX3CL1-CX3CR1 axis in microglia via the NF-kappaB pathway. *Curr HIV Res* 2014; **12**: 189–200.
94. Yao H, Bethel-Brown C, Yang L, Cai Y, Kanmogne M, Mudgapalli V et al. Signal transduction in HIV protein-treated astrocytes. *Curr Sig Transduct Ther* 2012; **7**: 28–34.
95. Chang C, Lang H, Geng N, Wang J, Li N, Wang X. Exosomes of BV-2 cells induced by alpha-synuclein: important mediator of neurodegeneration in PD. *Neurosci Lett* 2013; **548**: 190–195.
96. Ferrari R, Kapogiannis D, Huey ED, Momeni P. FTD and ALS: a tale of two diseases. *Curr Alzheimer Res* 2011; **8**: 273–294.
97. Mehta P, Antao V, Kaye W, Sanchez M, Williamson D, Bryan L et al. Prevalence of amyotrophic lateral sclerosis - United States, 2010-2011. *MMWR Suppl* 2014; **63**: 1–14.
98. Boillee S, Vande Velde C, Cleveland DW. ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* 2006; **52**: 39–59.
99. Robberecht W, Philips T. The changing scene of amyotrophic lateral sclerosis. *Nat Rev Neurosci* 2013; **14**: 248–264.
100. Millicamps S, Salachas F, Cazeneuve C, Gordon P, Bricka B, Camuzat A et al. SOD1, ANG, VAPB, TARDBP, and FUS mutations in familial amyotrophic lateral sclerosis: genotype-phenotype correlations. *J Med Genet* 2010; **47**: 554–560.
101. Daoud H, Valdmanis PN, Kabashi E, Dion P, Dupre N, Camu W et al. Contribution of TARDBP mutations to sporadic amyotrophic lateral sclerosis. *J Med Genet* 2009; **46**: 112–114.
102. Silverman JM, Fernando SM, Grad LI, Hill AF, Turner BJ, Yerbury JJ et al. Disease Mechanisms in ALS: misfolded SOD1 transferred through exosome-dependent and exosome-independent pathways. *Cell Mol Neurobiol* 2016; **36**: 377–381.
103. Gomes C, Keller S, Altevogt P, Costa J. Evidence for secretion of Cu, Zn superoxide dismutase via exosomes from a cell model of amyotrophic lateral sclerosis. *Neurosci Lett* 2007; **428**: 43–46.
104. Grad LI, Yerbury JJ, Turner BJ, Guest WC, Pokrishevsky E, O'Neill MA et al. Intercellular propagated misfolding of wild-type Cu/Zn superoxide dismutase occurs via exosome-dependent and -independent mechanisms. *Proc Natl Acad Sci USA* 2014; **111**: 3620–3625.
105. Basso M, Pozzi S, Tortorolo M, Fiordaliso F, Bisighini C, Pasetto L et al. Mutant copper-zinc superoxide dismutase (SOD1) induces protein secretion pathway alterations and exosome release in astrocytes: implications for disease spreading and motor neuron pathology in amyotrophic lateral sclerosis. *J Biol Chem* 2013; **288**: 15699–15711.
106. Yuyama K, Igarashi Y. Physiological and pathological roles of exosomes in the nervous system. *Biomol Concepts* 2016; **7**: 53–68.
107. Kim DK, Kang B, Kim OY, Choi DS, Lee J, Kim SR et al. EVpedia: an integrated database of high-throughput data for systemic analyses of extracellular vesicles. *J Extracell Vesicles* 2013; **2**.
108. Nonaka T, Masuda-Suzukake M, Arai T, Hasegawa Y, Akatsu H, Obi T et al. Prion-like properties of pathological TDP-43 aggregates from diseased brains. *Cell Rep* 2013; **4**: 124–134.
109. Feiler MS, Strobel B, Freischmidt A, Heflerich AM, Kappel J, Brewer BM et al. TDP-43 is intercellularly transmitted across axon terminals. *J Cell Biol* 2015; **211**: 897–911.
110. Ugalde CL, Finkelstein DI, Lawson VA, Hill AF. Pathogenic mechanisms of prion protein, amyloid-beta and alpha-synuclein misfolding: the prion concept and neurotoxicity of protein oligomers. *J Neurochem* 2016; **139**: 162–180.
111. Tagliapietra M, Zanusso G, Fiorini M, Bonetto N, Zaranonello G, Zambon A et al. Accuracy of diagnostic criteria for sporadic Creutzfeldt-Jakob disease among rapidly progressive dementia. *J Alzheimers Dis* 2013; **34**: 231–238.
112. Hetz CA, Soto C. Stressing out the ER: a role of the unfolded protein response in prion-related disorders. *Curr Mol Med* 2006; **6**: 37–43.
113. Vella LJ, Greenwood DL, Cappai R, Scheerlinck JP, Hill AF. Enrichment of prion protein in exosomes derived from ovine cerebral spinal fluid. *Vet Immunol Immunopathol* 2008; **124**: 385–393.
114. Robertson C, Booth SA, Beniac DR, Coulthart MB, Booth TF, McNicol A. Cellular prion protein is released on exosomes from activated platelets. *Blood* 2006; **107**: 3907–3911.
115. Basso M, Bonetto V. Extracellular vesicles and a novel form of communication in the brain. *Front Neurosci* 2016; **10**: 127.
116. Vella LJ, Hill AF, Cheng L. Focus on extracellular vesicles: exosomes and their role in protein trafficking and biomarker potential in Alzheimer's and Parkinson's disease. *Int J Mol Sci* 2016; **17**: 173.
117. Coleman BM, Hill AF. Extracellular vesicles—their role in the packaging and spread of misfolded proteins associated with neurodegenerative diseases. *Semin Cell Dev Biol* 2015; **40**: 89–96.
118. Fevrier B, Vilette D, Archer F, Loew D, Faigle W, Vidal M et al. Cells release prions in association with exosomes. *Proc Natl Acad Sci USA* 2004; **101**: 9683–9688.
119. Vella LJ, Sharples RA, Lawson VA, Masters CL, Cappai R, Hill AF. Packaging of prions into exosomes is associated with a novel pathway of PrP processing. *J Pathol* 2007; **211**: 582–590.
120. Guo BB, Bellingham SA, Hill AF. Stimulating the release of exosomes increases the intercellular transfer of prions. *J Biol Chem* 2016; **291**: 5128–5137.
121. Guo BB, Bellingham SA, Hill AF. The neutral sphingomyelinase pathway regulates packaging of the prion protein into exosomes. *J Biol Chem* 2015; **290**: 3455–3467.
122. Saa P, Yakovleva O, de Castro J, Vasilyeva I, De Paoli SH, Simak J et al. First demonstration of transmissible spongiform encephalopathy-associated prion protein (PrP<sup>TSE</sup>) in extracellular vesicles from plasma of mice infected with mouse-adapted variant Creutzfeldt-Jakob disease by *in vitro* amplification. *J Biol Chem* 2014; **289**: 29247–29260.
123. Wroe SJ, Pal S, Siddique D, Hyare H, Macfarlane R, Joiner S et al. Clinical presentation and pre-mortem diagnosis of variant Creutzfeldt-Jakob disease associated with blood transfusion: a case report. *Lancet* 2006; **368**: 2061–2067.
124. Bellingham SA, Coleman BM, Hill AF. Small RNA deep sequencing reveals a distinct miRNA signature released in exosomes from prion-infected neuronal cells. *Nucleic Acids Res* 2012; **40**: 10937–10949.
125. Arellano-Anaya ZE, Huor A, Leblanc P, Lehmann S, Provansal M, Raposo G et al. Prion strains are differentially released through the exosomal pathway. *Cell Mol Life Sci* 2015; **72**: 1185–1196.

126. Polanco JC, Scicluna BJ, Hill AF, Gotz J. Extracellular vesicles isolated from the brains of rTg4510 mice seed Tau protein aggregation in a threshold-dependent manner. *J Biol Chem* 2016; **291**: 12445–12466.
127. Zhuang X, Xiang X, Grizzle W, Sun D, Zhang S, Axtell RC *et al*. Treatment of brain inflammatory diseases by delivering exosome encapsulated anti-inflammatory drugs from the nasal region to the brain. *Mol Ther* 2011; **19**: 1769–1779.
128. El Andaloussi S, Lakhai S, Mager I, Wood MJ. Exosomes for targeted siRNA delivery across biological barriers. *Adv Drug Deliv Rev* 2013; **65**: 391–397.
129. Alvarez-Erviti L, Seow Y, Yin H, Betts C, Lakhai S, Wood MJ. Delivery of siRNA to the mouse brain by systemic injection of targeted exosomes. *Nat Biotechnol* 2011; **29**: 341–345.
130. El-Andaloussi S, Lee Y, Lakhai-Littleton S, Li J, Seow Y, Gardiner C *et al*. Exosome-mediated delivery of siRNA *in vitro* and *in vivo*. *Nat Protoc* 2012; **7**: 2112–2126.
131. Khatua AK, Taylor HE, Hildreth JE, Popik W. Exosomes packaging APOBEC3G confer human immunodeficiency virus resistance to recipient cells. *J Virol* 2009; **83**: 512–521.
132. Vlassov AV, Magdaleno S, Setterquist R, Conrad R. Exosomes: current knowledge of their composition, biological functions, and diagnostic and therapeutic potentials. *Biochim Biophys Acta* 2012; **1820**: 940–948.
133. Naslund TI, Paquin-Proulx D, Paredes PT, Vallhov H, Sandberg JK, Gabrielsson S. Exosomes from breast milk inhibit HIV-1 infection of dendritic cells and subsequent viral transfer to CD4+ T cells. *AIDS* 2014; **28**: 171–180.
134. Akao Y, Iio A, Itoh T, Noguchi S, Itoh Y, Ohtsuki Y *et al*. Microvesicle-mediated RNA molecule delivery system using monocytes/macrophages. *Mol Ther* 2011; **19**: 395–399.
135. van den Boorn JG, Schlee M, Coch C, Hartmann G. siRNA delivery with exosome nanoparticles. *Nat Biotechnol* 2011; **29**: 325–326.
136. DeMarino C, Schwab A, Pleet M, Mathiesen A, Friedman J, El-Hage N *et al*. Biodegradable nanoparticles for delivery of therapeutics in CNS infection. *J Neuroimmune Pharmacol* 2016 (e-pub ahead of print).
137. Sampey GC, Meyering SS, Asad Zadeh M, Saifuddin M, Hakami RM, Kashanchi F. Exosomes and their role in CNS viral infections. *J Neurovirol* 2014; **20**: 199–208.
138. Dias V, Junn E, Mouradian MM. The role of oxidative stress in Parkinson's disease. *J Parkinsons Dis* 2013; **3**: 461–491.
139. Haney MJ, Klyachko NL, Zhao Y, Gupta R, Plotnikova EG, He Z *et al*. Exosomes as drug delivery vehicles for Parkinson's disease therapy. *J Control Release* 2015; **207**: 18–30.
140. Haney MJ, Zhao Y, Harrison EB, Mahajan V, Ahmed S, He Z *et al*. Specific transfection of inflamed brain by macrophages: a new therapeutic strategy for neurodegenerative diseases. *PLoS One* 2013; **8**: e61852.
141. Zhao Y, Haney MJ, Gupta R, Bohnsack JP, He Z, Kabanov AV *et al*. GDNF-transfected macrophages produce potent neuroprotective effects in Parkinson's disease mouse model. *PLoS One* 2014; **9**: e106867.



**Cell Death and Disease** is an open-access journal published by **Nature Publishing Group**. This work is licensed under a **Creative Commons Attribution 4.0 International License**. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016