

 Open access • Journal Article • DOI:10.1007/S11240-011-9975-X

Encapsulation of cauliflower (*Brassica oleracea* var *botrytis*) microshoots as artificial seeds and their conversion and growth in commercial substrates — Source link

Hail Z. Rihan, Mohammed Al-Issawi, S Burchett, Michael P. Fuller

Institutions: University of Plymouth

Published on: 28 May 2011 - Plant Cell Tissue and Organ Culture (Springer Netherlands)

Topics: Kinetin

Related papers:

- [A revised medium for rapid growth and bio assays with tobacco tissue cultures](#)
- [Assessment of genetic fidelity of encapsulated microshoots of *Picrorhiza kurroo*](#)
- [The encapsulation technology in fruit plants--a review.](#)
- [Alginate encapsulation of shoot tips and nodal segments for short-term storage and distribution of the eucalypt *Corymbia torelliana* × *C. citriodora*](#)
- [Synthetic seed: prospects and limitations.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/encapsulation-of-cauliflower-brassica-oleracea-var-botrytis-404nmv0g4k>

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle Encapsulation of cauliflower (*Brassica oleraceae* var *botrytis*) microshoots as artificial seeds and their conversion and growth in commercial substrates

Article Sub-Title

Article CopyRight Springer Science+Business Media B.V.
(This will be the copyright line in the final PDF)

Journal Name Plant Cell, Tissue and Organ Culture (PCTOC)

Corresponding Author

Family Name	Fuller
Particle	
Given Name	Michael P.
Suffix	
Division	School of Biomedical and Biological Sciences
Organization	Faculty of Science and Technology, University of Plymouth
Address	PL4 8AA, Plymouth, UK
Email	mfuller@plymouth.ac.uk

Author

Family Name	Rihan
Particle	
Given Name	Hail
Suffix	
Division	School of Biomedical and Biological Sciences
Organization	Faculty of Science and Technology, University of Plymouth
Address	PL4 8AA, Plymouth, UK
Email	

Author

Family Name	Al-Issawi
Particle	
Given Name	Mohammed
Suffix	
Division	School of Biomedical and Biological Sciences
Organization	Faculty of Science and Technology, University of Plymouth
Address	PL4 8AA, Plymouth, UK
Email	

Author

Family Name	Burchett
Particle	
Given Name	Stephen
Suffix	
Division	School of Biomedical and Biological Sciences
Organization	Faculty of Science and Technology, University of Plymouth
Address	PL4 8AA, Plymouth, UK
Email	

Schedule

Received	3 February 2011
Revised	

Abstract

An effective protocol for the mass production of cauliflower microshoots was refined using the meristematic layer of cauliflower curd. After the meristematic layer was surface sterilized and shaved off, a commercial blender was used for homogenization and several blending treatments were tested in the range 15–120 s and 30 s was found to be optimal in terms of the amount explants produced and their subsequent growth ability. Explants were cultivated in S23 liquid medium (4.4 g L^{-1} MS (Murashige and Skoog) and 3% v/w sucrose) supplemented with several combinations of plant growth regulators (PGRs) including 1 and 2 mg L^{-1} of kinetin in combination with three types of auxins (indole butyric acid (IBA), Naphthaleneacetic acid (NAA) and Indole-3-acetic acid (IAA)), each at 1 and 2 mg L^{-1} concentration. The use of 2 mg L^{-1} kinetin and 1 mg L^{-1} IBA gave the best results in terms of its effects on explant induction. Microshoots of different sizes were encapsulated in a sodium alginate matrix and the optimal stage suitable for the production of artificial seeds was assessed in terms of both subsequent conversion and plantlet viability. The feasibility of cultivating cauliflower artificial seeds in commercial substrates (compost, vermiculite, perlite and sand) irrigated with different solution mixtures including sterilized distilled water (SDW), PGRs-free S23 medium and S23 medium supplemented with kinetin (1 and 2 mg L^{-1}) and IBA or NAA at (1 and 2 mg L^{-1}) was investigated. The use of 2 mg L^{-1} kinetin and 2 mg L^{-1} NAA applied with S23 gave the optimal response with both perlite and compost. This study showed high growth capacity of cauliflower artificial seeds in commercial substrates which is considered a promising step for their direct use in vivo.

Keywords (separated by '-') Sodium alginate - Encapsulation - Curd - Meristems - Artificial seeds

Footnote Information

Journal: 11240
Article: 9975



Author Query Form

**Please ensure you fill out your response to the queries raised below
and return this form along with your corrections**

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details required	Author's response
1.	Please check and confirm that the authors and their respective affiliations have been correctly identified and amend if necessary.	
2.	Please confirm the section headings are correctly identified.	
3.	The superscript alphabets were given in the Table 1 and 2 and their significance is not provided. Please check and confirm.	

2 **Encapsulation of cauliflower (*Brassica oleraceae* var *botrytis*)**
3 **microshoots as artificial seeds and their conversion and growth**
4 **in commercial substrates**

5 **Hail Rihan · Mohammed Al-Issawi ·**
6 **Stephen Burchett · Michael P. Fuller**

7 Received: 3 February 2011 / Accepted: 14 May 2011
8 © Springer Science+Business Media B.V. 2011

9 **Abstract** An effective protocol for the mass production
10 of cauliflower microshoots was refined using the meriste-
11 matic layer of cauliflower curd. After the meristematic
12 layer was surface sterilized and shaved off, a commercial
13 blender was used for homogenization and several blending
14 treatments were tested in the range 15–120 s and 30 s was
15 found to be optimal in terms of the amount explants pro-
16 duced and their subsequent growth ability. Explants were
17 cultivated in S23 liquid medium (4.4 g L⁻¹ MS (Murashige
18 and Skoog) and 3% v/w sucrose) supplemented with sever-
19 al combinations of plant growth regulators (PGRs)
20 including 1 and 2 mg L⁻¹ of kinetin in combination with
21 three types of auxins (indole butyric acid (IBA), Naph-
22 thaleneacetic acid (NAA) and Indole-3-acetic acid (IAA)),
23 each at 1 and 2 mg L⁻¹ concentration. The use of 2 mg
24 L⁻¹ kinetin and 1 mg L⁻¹ IBA gave the best results in
25 terms of its effects on explant induction. Microshoots of
26 different sizes were encapsulated in a sodium alginate
27 matrix and the optimal stage suitable for the production of
28 artificial seeds was assessed in terms of both subsequent
29 conversion and plantlet viability. The feasibility of culti-
30 vating cauliflower artificial seeds in commercial substrates
31 (compost, vermiculite, perlite and sand) irrigated with
32 different solution mixtures including sterilized distilled
33 water (SDW), PGRs-free S23 medium and S23 medium
34 supplemented with kinetin (1 and 2 mg L⁻¹) and IBA or
35 NAA at (1 and 2 mg L⁻¹) was investigated. The use of
36 2 mg L⁻¹ kinetin and 2 mg L⁻¹ NAA applied with S23
37 gave the optimal response with both perlite and compost.

This study showed high growth capacity of cauliflower 38
artificial seeds in commercial substrates which is consid- 39
ered a promising step for their direct use in vivo. 40

Keywords Sodium alginate · Encapsulation · Curd · 42
Meristems · Artificial seeds 43

Introduction 44

The outermost layer of cauliflower curd is made of several 45
million meristems (Kieffer et al. 2001) and the use of curd 46
meristematic tissue for in vitro culture has been investi- 47
gated for micropropagation and the production of virus free 48
plants (Grout and Crisp 1977; Kumar et al. 1993). The use 49
of the meristematic layer of cauliflower curd is considered 50
to be superior compared with the use of conventional 51
protocols using seedling or leaf explants which have been 52
found to be labour intensive (Pow 1969; Kumar et al. 53
1993). Kieffer et al. (2001) designed an effective protocol 54
for the production of cauliflower propagules from frac- 55
tionated and graded curd and these propagules were sug- 56
gested to be suitable for encapsulation in sodium alginate 57
matrices for the production of artificial seeds. The tech- 58
nique of artificial seed has been widely studied and works 59
with various plant species including fruits, cereals, 60
medicinal plants, vegetables, ornamentals, forest trees and 61
orchids (Germanà et al. 2011, Sundararaj et al. 2010, Singh 62
et al. 2009, Rai et al. 2008, Pintos et al. 2008, Micheli et al. 63
2007, Antonietta et al. 2007, Naik and Chand 2006, Mal- 64
abadi and Staden 2005, Chand and Singh 2004, Nyende 65
et al. 2003, Mandal et al. 2000). The use of microshoots has 66
been widely reported for the production of artificial seeds 67
in different plant species such as *Cineraria maritime* 68
(Sandoval-Yugar et al. 2009), *Musa* sp. (Sandoval-Yugar 69

A1 H. Rihan · M. Al-Issawi · S. Burchett · M. P. Fuller (✉)
A2 School of Biomedical and Biological Sciences,
A3 Faculty of Science and Technology, University of Plymouth,
A4 Plymouth PL4 8AA, UK
A5 e-mail: mfuller@plymouth.ac.uk

70 et al. 2009), *Gerbera jamesonii* (Taha et al. 2009) and
71 *Picrorhiza kurrooa* (Mishra et al. 2010).

72 Cauliflower is an open pollinated species and there are
73 technical challenges to producing in-bred lines with reli-
74 able self-incompatibility or male sterility for producing F1
75 hybrids particularly amongst the winter-heading maturity
76 sub-group. The production of cauliflower clones multiplied
77 by tissue culture and distributed as artificial seeds could be
78 useful alternative to F1 hybrids varieties and could also be
79 useful in the context of the maintenance of elite clonal
80 germplasm in cauliflower breeding programmes. The cur-
81 rent study aimed to further optimize the production of
82 cauliflower propagules and to investigate the ability of
83 using cauliflower microshoots for the production of artifi-
84 cial seeds. Moreover, the investigation of cauliflower artifi-
85 cial seeds capacity to be grown on in commercial
86 substrates was one of the main aims of this study.

87 Materials and methods

88 Plant materials

89 Three F1 hybrid varieties of cauliflower Clemen, Mascaret
90 and Broden previously found to be equally responsive in
91 tissue culture were used. Plants were obtained from the
92 field in Cornwall, courtesy of Simmonds & Sons Ltd, and
93 replanted in raised beds at the University of Plymouth. The
94 plants were grown according to good commercial practice
95 and raised to maturity when the curds were harvested and
96 stored at 2–4°C until required. The use of 3 varieties gave a
97 continuous supply of cauliflower heads over the experi-
98 mental period.

99 Cauliflower microshoot production

100 Large pieces (1–5 cm) of cauliflower curds were sterilized
101 by immersion in 10% by volume un-thickened domestic
102 bleach (0.06% sodium hypochlorite) for 15 min followed
103 by a double wash with sterile distilled water. Explants were
104 produced mechanically by eliminating the mass of non-
105 responsive tissue (stem branches) and shaving off the upper
106 meristematic layer using a sterilized scalpel working in a
107 laminar flow cabinet. The meristematic clusters were then
108 homogenized using a commercial blender (Waring model
109 800) at approximately 1,700 rev min⁻¹ in maintenance S23
110 (4.4 g L⁻¹ SigmaTM MS (Murashige and Skoog 1962) and
111 3% w/v sucrose) liquid media. Eight blending durations in
112 the range 15–120 s were assessed in terms of their effects
113 on the amount and subsequent growth of the micro-
114 explants produced after sieving through precision sieves
115 (212, 300 and 600 µm) (Endecotts Ltd, London, UK). The
116 micro-explants were collected off the sieves, weighed and

converted to aliquots of explants using small precision 117
volumetric measures (74 or 240 ± 2 µL). Six containers 118
(150 mL plastic pots), each containing 30 mL liquid S23 119
culture medium supplemented with 2 mg L⁻¹ Kinetin and 120
1 mg L⁻¹ IBA, were cultured with a constant volume of 121
micro-explants (74 µL) and used with every blending 122
duration in order to assess the effects of blending treat- 123
ments. The pots were constantly shaken (150 rev min⁻¹) 124
during culture at 20°C and exposed to 16 h photoperiod. 125
The proportion of micro-explants having meristematic 126
domes was determined by taking 4 mL samples of cultures 127
at 8 days from each pot and observing them under a zoom 128
binocular microscope (Nikon SMZ-2T). The growth ability 129
of micro-explants during culture was determined after 130
20 days as the number and fresh weight of microshoots 131
produced. 132

The effects of several PGR combinations were evaluated 133
in terms of their ability to induce micro-explant develop- 134
ment. Nine PGR combinations consisting of various combi- 135
nation of the cytokinin (Kinetin) (1 and 2 mg L⁻¹) with 136
the auxins IBA (1 and 2 mg L⁻¹) or NAA (1 and 2 mg 137
L⁻¹) were evaluated in the first stage and another 4 combi- 138
nations consisting of various combinations of Kinetin 139
(1 and 2 mg L⁻¹) with IAA (1 and 2 mg L⁻¹) compared 140
with the use of Kinetin (2 mg L⁻¹) and IBA (1 mg L⁻¹) 141
were evaluated in the second stage. Four containers, each 142
with 30 mL of culture medium, were cultivated with a 143
constant volume of explants (74 µL) of the 300–425 µm 144
explant size class and used with every treatment. In order 145
to preserve culture sterility the culture media was supple- 146
mented 1 mL L⁻¹ PPMTM (Plant Preservative Mixture) 147
and used with all the treatments. 148

Microshoot development was assessed from the 2 mg 149
L⁻¹ Kinetin, 1 mg L⁻¹ IBA and 1 mL L⁻¹ PPM treatment 150
cultivated with 74 µL of micro-explants from each of the 151
two size class 212–300 µm and 300–600 µm. Random 152
microshoot samples (n = 15) were taken from each culture 153
every 3 days and their length measured under a zoom 154
binocular microscope. Measurements commenced when 155
the microshoots were 5 days old and continued until 156
20 days old. 157

Cauliflower artificial seed production 158

Micro-explants of the 212–300 µm size class were used for 159
artificial seed production. Microshoots were mixed with 160
sterilized (by tyndallisation) sodium alginate 2% (w/v) and 161
dropped into a sterilized (autoclaved) solution of calcium 162
chloride 15 g L⁻¹ using a sterilized pipette. Microshoots 163
were left in the calcium chloride for 30 min for full com- 164
plexion. The artificial seeds were then transferred to a S23 165
liquid media (without PGRs) for 30 min followed by a 166
quick wash with sterile distilled water. The optimal age 167

168 suitable for microshoot encapsulation was determined for
 169 9–15 day-old microshoots after which the microshoots were
 170 too big to be encapsulated. Five replications of 10 artificial
 171 seeds, were cultivated in plastic pots (10 × 10 × 8 cm)
 172 containing 75 mL of semi-solid S23 supplemented with
 173 2 mg L⁻¹ IBA. The conversion rate and fresh weight of
 174 plantlets produced was evaluated after 20 days of culture.

175 Five replications of 6 artificial seeds were placed onto
 176 different sterilized substrates (compost, vermiculite, perlite
 177 and sand) to assess their suitability for conversion and
 178 establishment using 11 day old microshoots produced from
 179 212 to 300 μm micro-explants and cultivated in S23 liquid
 180 medium supplemented with 2 mg L⁻¹ Kinetin and 1 mg
 181 L⁻¹ IBA. Ten different irrigation solutions were assessed
 182 with each substrate as follows, (1) SDW. (2) S23 PGRs
 183 free. (3) Eight combinations of S23 supplemented with
 184 various combinations of kinetin (1 and 2 mg L⁻¹) with
 185 IBA or NAA (1 and 2 mg L⁻¹). Each pot was irrigated with
 186 75 mL of the irrigation solution. Artificial seed conversion
 187 rate and the fresh weight of plantlets produced were eval-
 188 uated after 50 days of culture.

189 Statistical analysis

190 Results are presented as means ± standard error (SE). All
 191 data were subjected to analysis of variance (ANOVA)
 192 using Minitab software (version 15) and comparisons of
 193 means were made with least significant difference test
 194 (LSD) at 5% level of probability.

195 Results and discussion

196 There was an interaction between the number of micro-
 197 explants produced and their subsequent development in
 198 response to blending duration. In terms of the number of
 199 growing explants during subsequent culture, the use of the
 200 30 and 15 s treatments were found to be optimal for size
 201 classes 212–300 μm and 300–600 μm respectively (Fig. 1).
 202 However, while no significant difference was found
 203 between these two treatments at size class 300–600 μm,
 204 the number of developing microshoots was significantly higher
 205 using 30 s treatment at size class 212–300 μm (Fig. 1). In
 206 terms of microshoots fresh weight (fresh weight were
 207 considered as a good indicator of microshoots viability
 208 since it expresses the speed of growth), the use of 60 s for
 209 size class 300–600 μm was found to be optimal but the
 210 number of developing explants using this treatment was
 211 significantly lower than the use of 30 s treatment
 212 ($P < 0.001$) (Fig. 1). The use of the 30 s blending treat-
 213 ments is recommended as optimal.

214 It was observed that the proportion of micro-explants
 215 which had meristematic domes and those that were classed
 216 as debris increased with the duration of blending for both




Fig. 1 The effect of the blending duration treatments on the number of growing microshoots (Number of microshoots (LSD = 19.58 for 212–300 μm size class, LSD = 70.52 for 300–600 μm size class)) and their average weights (Average weights of microshoots (LSD = 0.022 for 212–300 μm size class, LSD = 0.0045 for 300–600 μm size class)) at two size classes, 212–300 μm and 300–600 μm

size classes (212–300 and 300–600 μm). The use of
 217 blenders has also been described for mass production of
 218 initial explants of fern (Knauss 1976; Cooke 1979; Jans-
 219 sens and Sepelie 1989; Teng and Teng 1997) and for
 220 separating meristamoid aggregates of several species (Ziv
 221 and Ariel 1991; Standardi and Piccioni 1998; Ziv et al.
 222 1998; Teng and Ngai 1999). The use of a blender for
 223 cauliflower explants production was also previously
 224 reported by Kieffer et al. (2001) and as shown in this work
 225 the blender is a crude but effective way of producing small
 226 micro-explants which remain viable and capable of pro-
 227 ducing microshoots. Explant growth capacity however
 228 quickly diminishes as the blending duration increases and
 229 30 s gives the optimal response rate under the conditions
 230 described here.

231 PGRs added to the liquid medium were found to have a
 232 crucial role in the induction of development of the explants
 233 since none of the explants developed in PGR-free S23.
 234 Although the use of 1 mg L⁻¹ Kinetin and 1 mg L⁻¹ IBA
 235 and 2 mg L⁻¹ Kinetin and 1 mg L⁻¹ IBA treatments gave
 236 the optimal results in terms of the number of growing
 237 microshoot ($P < 0.001$), the use of 2 mg L⁻¹ Kinetin and
 238 1 mg L⁻¹ IBA was found to be one of the best in terms of
 239 the microshoot fresh weight ($P < 0.001$) (Fig. 2). In the
 240 second stage of this investigation, the use of 2 mg L⁻¹
 241 Kinetin and 1 mg L⁻¹ IBA treatment was found to be
 242 better than all combination of Kinetin with IAA. Therefore,
 243 the use of 2 mg L⁻¹ Kinetin and 1 mg L⁻¹ IBA is rec-
 244 ommended (Fig. 2).
 245

246 Cytokinins are reported to have a crucial role in the
 247 organization of sink activity and nutrient partitioning
 248 (Kuiper 1988; Kuiper et al. 1989). Cytokinins are essen-
 249 tially made in the root apex (Komor et al. 1993) and
 250 because the cauliflower explants have no roots, the main




Fig. 2 The effect of the various combination of Kinetin (1, 2) mg L⁻¹ incorporated with IBA (1, 2) mg L⁻¹ or NAA (1, 2) mg L⁻¹ added to S23 liquid medium on the number (Number of microshoots (LSD = 28.85)) and average weight (Average weights of microshoots (LSD = 0.0083)) of developing microshoots

cytokinin source is provided by the culture medium. The current investigation also showed that the type of auxin was important and there were large differences between the number and the viability of microshoots depending on the type of auxin used and IBA was found to be best for the development of the explants. Despite these differences all of the auxin types used had a positive effect in the induction of cauliflower micro-explants. This is in contrast to Vandemoortele et al. (2001) who reported that it is difficult to associate endogenous auxin with a function in the induction of cauliflower curd explants. However, none of the growing microshoots showed the capacity for the development of roots irrespective of the combination of PGRs used. This limitation of microshoot rooting might be caused by an interaction with Kinetin since transferred microshoots to semi solid medium containing 2 mg L⁻¹ IBA displayed roots within a few days. The current observations are in contrast with those reported by Kieffer et al. (2001) who reported the capacity of NAA at low concentration to encourage early rooting of microshoots even in the presence of Kinetin in the culture medium. This may be a cauliflower varietal effect. It has been reported by others that the use of cytokinin decreases the number of lateral roots in other species (Hinchee and Rost 1986, Goodwin and Morris 1979, Böttger 1974). Eriksen (1974) working in peas mentioned that the presence of high concentrations of cytokinin could have negative effects on the initial step of rooting by deterring the activity of auxin. Rani Debi et al. (2005) indicated that cytokinin inhibits lateral root initiation in rice (*Oryza sativa*) and Nakashimada et al. (1995) also observed inhibition effects of kinetin presence in the culture media on the root elongation of horseradish hairy roots (*Armoracia rusticana*) plantlets. However, Hinchee and Rost (1986) reported that the

auxin:cytokinin ratio has an essential role in co-ordinating lateral root growth in pea seedlings. It is clear that there is no universal explanation for the variations in these observations indicating a strong genotypic effect.

Explant development stages were determined for explants from the 212 to 300 μm and 300 to 600 μm size classes. The explants started growing very slowly for the first 11 days in both size classes but after that the growth rate increased exponentially with time (Fig. 3). The best growth rate was observed with size class 212–300 μm compared to the 300–600 μm size class and might be because of competition for nutrient supply since explants produced at size class 212–300 μm bear only 1 meristematic dome, giving one microshoot while the explants at size class 300–600 μm bear 2–3 meristematic domes giving 2–3 microshoots and more localized competition for nutrients (Fig. 4).

We divided microshoot development into three main stages: (a) 0–11 day stage when the growth rate was very low. (b) 11–15 day stage of culture when an acceleration of microshoot growth was observed. (3) A stage after 15 days when the growth rate was rapid (Fig. 3).

The optimal age for microshoot encapsulation was observed to be 13–14 days (Fig. 5). The encapsulation of both younger and older microshoots had a negative effect on the subsequent artificial seed conversion rate and fresh weights of plantlets produced. The growth of microshoots younger than 13 day-old was observed to be very slow and encapsulation seemed to be an inhibitor of microshoot growth at this sensitive stage. The growth of microshoots older than 14 days old was observed to be rapid in culture but this fast growth brought about metabolic activity which seemed to negatively affect the subsequent of microshoot encapsulation. It was concluded that microshoots derived in an accelerating stage of growth (i.e. stage 2) were optimal for encapsulation.




Fig. 3 Cauliflower explants growth assessed by measured length changes over time




Fig. 4 Cauliflower microshoots at different developmental stages. **a, b, c, d, e** and **f** were taken from 212 to 300 μm size class and **g, h, i, j, k** and **l** from 300 to 600 μm. Photos were taken at 3, 6, 9, 12, 15, 17

and 20 days old respectively. **a, b, c, d, g, h, i, and j** were taken at 100 times magnification, **e** and **k** were taken at 40 times magnification and **f** and **l** without any magnification




Fig. 5 The effect of the optimal microshoot age suitable for encapsulation (LSD = 0.687 for the conversion rate and LSD = 0.0675 for artificial seeds viability)

321 For cauliflower artificial seeds to be a promising candidate for agriculture their capacity to be sown into commercial substrates such as compost, perlite, vermiculite or sand needs to be evaluated. It was clear from this investigation that the moistening/irrigation solution composition used during conversion was the key factor for the success of these substrates. No conversion of artificial seeds was observed when the culture substrates were irrigated with sterile distilled water and this is in common with other

species tested (Naik and Chand 2006) (Soneji et al. 2002). The use of S23 PGR-free as an irrigating solution had a positive effect on the conversion rate and viability of the artificial seeds (Tables 1 and 2). The use of PGRs combinations had a significant influence on the fresh weight of plantlets produced 50 days after conversion but they did not improve the conversion rate of the artificial seeds. Moreover, some of the PGR combinations such as the use of S23, 1 mg L⁻¹ Kinetin and 2 mg L⁻¹ IBA and the use of S23, 1 mg L⁻¹ Kinetin and 2 mg L⁻¹ NAA significantly reduced the conversion rate. However, the use of S23, 2 mg L⁻¹ Kinetin and 2 mg L⁻¹ NAA treatment was recommended resulting in the optimal artificial seed conversion rate and the best fresh weight of plantlets (Tables 1 and 2).

330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354

The type of auxin used with the culture substrates showed that 2 mg L⁻¹ NAA was better than those used with semi solid culture media supplemented with 2 mg L⁻¹ IBA (data not shown). This could be either due to the presence of Kinetin in the liquid media used for culture substrate moistening or because of the physical structure of culture substrates led to less transportation efficiency of PGRs to the cauliflower microshoots. However, although the optimal conversion rate was obtained using perlite, the viability of artificial seeds was negatively affected by this substrate and the growth of plantlets stopped at a certain

Table 1 The effect of irrigation composition and culture substrates on artificial seed conversion rate (%)

PGRs (mg L ⁻¹) ¹			Culture substrates				Treatment averages
Kin	IBA	NAA	Perlite	Sand	Compost	Vermiculite	
1	1	0	90 ^c	67 ^g	36 ¹	63 ^h	64 ^b
1	2	0	82 ^e	93 ^b	34 ^{lm}	33 ^m	60 ^{bc}
1	0	1	93 ^b	43 ^k	33 ^m	56 ⁱ	56 ^{bc}
1	0	2	83 ^e	63 ^h	26 ^o	30 ⁿ	50 ^c
2	1	0	93 ^b	97 ^a	73 ^f	30 ⁿ	73 ^{ab}
2	2	0	87 ^d	43 ^k	55 ⁱ	26 ^o	52 ^c
2	0	1	93 ^b	90 ^c	43 ^k	30 ⁿ	64 ^b
2	0	2	97 ^a	87 ^d	67 ^g	46 ^j	74 ^a
0	0	0	97 ^a	83 ^e	90 ^c	46 ^j	79 ^a
Average			91 ^a	74 ^b	50 ^c	40 ^d	

¹ S23 was supplemented with the described PGRs. (LSD = 10.528 for PGR combinations, LSD = 5.84 for culture substrates and LSD = 2.105 for PGR combinations × Culture substrates (interaction))

Table 2 The effect of irrigation composition and culture substrate on the plantlet fresh weight (g)

PGRs (mg L ⁻¹) ¹			Culture substrates				Average
Kin	IBA	NAA	Perlite	Sand	Compost	Vermiculite	
1	1	0	0.068 ^{efghij}	0.064 ^{fghij}	0.109 ^{bcdefghi}	0.050 ^{ghij}	0.072
1	2	0	0.051 ^{ghij}	0.075 ^{efghij}	0.178 ^{ab}	0.030 ^{ij}	0.083
1	0	1	0.071 ^{efghij}	0.025 ^j	0.234 ^a	0.050 ^{ghij}	0.095
1	0	2	0.073 ^{efghij}	0.12 ^{bcdefgh}	0.044 ^{hij}	0.033 ^{ij}	0.067
2	1	0	0.061 ^{ghij}	0.073 ^{efghij}	0.092 ^{cdefghij}	0.034 ^{ij}	0.065
2	2	0	0.087 ^{defghij}	0.054 ^{ghij}	0.167 ^{abc}	0.033 ^{ij}	0.085
2	0	1	0.061 ^{ghij}	0.141 ^{bcdef}	0.124 ^{bedefg}	0.031 ^{ij}	0.089
2	0	2	0.072 ^{efghij}	0.160 ^{abcd}	0.144 ^{bcde}	0.044 ^{hij}	0.105
0	0	0	0.052 ^{ghij}	0.072 ^{efghij}	0.059 ^{ghij}	0.027 ^{ij}	0.052
Average			0.066 ^b	0.087 ^b	0.128 ^a	0.037 ^c	

¹ S23 was supplemented with the described PGRs. (LSD = 0.022 for culture substrates and LSD = 0.080 for the interaction between the PGR combinations and culture substrates)

355 point. It was suggested that the cessation of growth could
356 be caused by a lack of moistened liquid mixture supplied
357 since the same volume of liquid mixture was used with the
358 four culture substrates and the signs of dehydration were
359 observed with perlite in comparison with other substrates.

360 The optimal fresh weight of plantlets produced was
361 obtained using compost as a culture substrate (Table 2). It
362 seemed that the conversion rate and viability of artificial
363 seed depends not only on the irrigation liquid mixture but
364 also on the physical structure of the culture substrates. It
365 might be good idea to investigate the result of using a
366 mixture of compost and perlite and examining the effects
367 on the conversion rate and viability of artificial seeds since
368 the optimal conversion rate and viability were obtained
369 using perlite and compost respectively. A high interaction
370 between the moistening/irrigation solution compositions
371 and the culture substrates was observed in terms of

artificial seeds conversion rate and fresh weights of plant- 372
373 lets produced. While the use of perlite moistened with S23
374 or S23 supplemented with 2 mg L⁻¹ Kinetin and 2 mg L⁻¹
375 NAA irrigation solutions gave the best artificial seed con-
376 version rate, the use of compost supplemented with S23,
377 1 mg L⁻¹ Kinetin and 1 mg L⁻¹ NAA was optimal in
378 terms of fresh weight of plantlets. However, the use of S23,
379 2 mg L⁻¹ Kinetin and 2 mg L⁻¹ NAA as an irrigating
380 solution was recommended with both perlite and compost
381 resulting in the optimal conversion rate and producing the
382 best plantlets fresh weights respectively since the only
383 significantly better plantlets fresh weights than this treat-
384 ment was obtained using S23, 1 mg L⁻¹ Kinetin and 1 mg
385 L⁻¹ NAA and since the conversion rate was observed to be
386 quite low when S23, 1 mg L⁻¹ Kinetin and 1 mg L⁻¹ NAA
387 was used as irrigating solution (Tables 1 and 2). Several
388 studies have investigated the possibilities of sowing

389 artificial seeds in soil or commercial substrates, for
390 example, on the use vermiculite, sand and soil for the
391 cultivation of mulberry artificial seeds (Machii and
392 Yamanouchi 1993), the use of soil for as alfalfa artificial
393 seed conversion substrate (Fujii et al. 1989), the use of
394 perlite for M.26 apple rootstock (Micheli et al. 2002) and
395 *Citrus reticulata* (Antonietta et al. 2007) and the use of
396 sand for *elite indica* rice (Roy and Mandal 2008) and it is
397 clear that the optimal conditions need to be derived
398 empirically for each species examined.

399 Conclusion

400 The production of microshoots from cauliflower curd was
401 optimized. This study showed that the use of a commercial
402 blender was a useful methodology for micro-explant gen-
403 eration and a 30 s blending duration treatment was found to
404 be the best in terms of the number and viability of sub-
405 sequent microshoots produced. The use of PGRs was
406 essential for microshoot development and the best PGR
407 combination was found to be 2 mg L⁻¹ Kinetin and 1 mg
408 L⁻¹ IBA. The use of the described protocol is considered a
409 cost effect methodology for cauliflower micropropagation
410 to produce huge number of microshoots per curd.

411 This study also demonstrated the ability of encapsulat-
412 ing cauliflower microshoots in a sodium alginate matrix for
413 artificial seed production and was the first which has
414 investigated the capacity of the cauliflower artificial seed
415 growing in commercial substrates (compost, vermiculite,
416 perlite and sand). Fully functional in vivo plantlets were
417 obtained using these commercial substrates and this opens
418 promising vistas for the direct use of cauliflower artificial
419 seeds in in vivo situations.

420 **Acknowledgments** The authors gratefully acknowledge the provi-
421 sion of a PhD grant from the University of Damascus, Syria.

422 References

- 423 Antonietta G, Ahmad H, Maurizio M, Alvaro S (2007) Preliminary
424 research on conversion of encapsulated somatic embryos of
425 *Citrus reticulata* Blanco, cv. Mandarino Tardivo di Ciaculli.
426 *Plant Cell Tiss Organ Cult* 88(1):117–120
427 Böttger M (1974) Apical dominance in roots of *Pisum sativum* L.
428 *Planta* (Berl) 121:253–261
429 Chand S, Singh AK (2004) Plant regeneration from encapsulated
430 nodal segments of *Dalbergia sissoo* Roxb. a timber-yielding
431 leguminous tree species. *J Plant Physiol* 161(2):237–243
432 Cooke RC (1979) Homogenization as an aid in tissue culture
433 propagation of *Platyserium* and *Davallia*. *Hortic Sci* 14:21–22
434 Eriksen EN (1974) Root Formation in Pea Cuttings. *Physiol Plant*
435 30(2):163–167
436 Fujii J, Slade D, Redenbaugh K (1989) Maturation and greenhouse
437 planting of alfalfa artificial seeds. *In Vitro Cell Dev Bio Plant*
438 25(12):1179–1182

- Germanà M, Micheli M, Chiancone B, Macaluso L, Standardi A 439
(2011) Organogenesis and encapsulation of in vitro-derived 440
propagules of *Carrizo citrange*. *Citrus sinensis* (L.); *Poncirus* 441
trifoliata (L.). *Plant Cell Tiss Organ Cult* 78:1–9 442
Goodwin PB, Morris SC (1979) Application of phytohormones to pea 443
roots after removal of the apex: effect on lateral root production. 444
Aust J Plant Physiol 6:195–200 445
Grout BWW, Crisp P (1977) Practical aspects of the propagation of 446
cauliflower by meristem culture. *Acta Hort* 78:289–296 447
Hinchee MAW, Rost TL (1986) The Control of Lateral Root 448
Development in Cultured Pea Seedlings. I. The Role of Seedling 449
Organs and Plant Growth Regulators. *Bot Gaz* 147(2):137 450
Janssens J, Sepelie M (1989) Rapid multiplication of ferns by 451
homogenization. *Sci Hortic* 38:161–164 452
Kieffer M, Simkins N, Fuller MP, Jellings AJ (2001) A cost effective 453
protocol for in vitro mass propagation of cauliflower. *Plant Sci* 454
160(5):1015–1024 455
Knauss JF (1976) A partial tissue culture method for pathogen-free 456
propagation of selected ferns from spores. *Proc Fla State Hort* 457
Soc 89:63–65 458
Komor E, Liegl I, Schobert C (1993) Loading and translocation of 459
various cytokinins in phloem and xylem of the seedlings of 460
Ricinus communis L. *Planta* 191(2):252–255 461
Kuiper D (1988) Growth Responses of *Plantago major* L. ssp. 462
pleiosperma (Pilger) to Changes in Mineral Supply: Evidence for 463
Regulation by Cytokinins. *Plant Physiol* 87(3):555–557 464
Kuiper D, Kuiper PJC, Lambers H, Schuit J, Staal M (1989) 465
Cytokinin concentration in relation to mineral nutrition and 466
benzyladenine treatment in *Plantago major* ssp. *pleiosperma*. 467
Physiol Plant 75(4):511–517 468
Kumar A, Kumar VA, Kumar J (1993) Rapid in vitro propagation of 469
cauliflower. *Plant Sci* 90:175–178 470
Machii H, Yamanouchi H (1993) Growth of mulberry synthetic seeds 471
on vermiculite, sand and soil media. *J SericSci Jpn* 62:85–87 472
Malabadi RB, Staden Jv (2005) Storability and germination of sodium 473
alginate encapsulated somatic embryos derived from the vege- 474
tative shoot apices of mature *Pinus patula* trees. *Plant Cell Tiss* 475
Organ Cult 82(3):259–265 476
Mandal J, Pattnaik S, Chand P (2000) Alginate encapsulation of 477
axillary buds of *Ocimum americanum* L. (hoary basil), *Basili-* 478
cum L. (sweet basil), *Gratissimum* L. (shrubby basil), and 479
Sanctum L. *In Vitro Cell Dev Biol Plant* 36(4):287–292 480
Micheli M, Pellegrino S, Piccioni E, Standardi A (2002) Effects of 481
double encapsulation and coating on synthetic seed conversion 482
in M.26 apple rootstock. *J Microencapsul* 19:347–356 483
Micheli M, Hafiz IA, Standardi A (2007) Encapsulation of in vitro- 484
derived explants of olive (*Olea europaea* L. cv. Moraiolo): II. 485
Effects of storage on capsule and derived shoots performance. 486
Sci Hortic 113(3):286–292 487
Mishra J, Singh M, Palni L, Nandi S (2010) Assessment of genetic 488
fidelity of encapsulated microshoots of *Picrorhiza kurrooa*. *Plant* 489
Cell Tiss Organ Cult :1–6 490
Murashige T, Skoog F (1962) A revised medium for rapid growth and 491
bioassays with tobacco tissue cultures. *Physiol Plant* 15(3): 492
473–497 493
Naik SK, Chand PK (2006) Nutrient-alginate encapsulation of in vitro 494
nodal segments of pomegranate (*Punica granatum* L.) for 495
germplasm distribution and exchange. *Sci Hortic* 108(3): 496
247–252 497
Nakashimada Y, Uozumi N, Kobayashi T (1995) Production of 498
plantlets for use as artificial seeds from horseradish hairy roots 499
fragmented in a blender. *J Ferment* 79(5):458–464 500
Nyende A, Schittenhelm S, Mix-Wagner G, Greef J-M (2003) 501
Production, storability, and regeneration of shoot tips of potato 502
(*Solanum tuberosum* L.) encapsulated in calcium alginate hollow 503
beads. *In Vitro Cell Dev Biol Plant* 39(5):540–544 504

- 505 Pintos B, Bueno M, Cuenca B, Manzanera J (2008) Synthetic seed
506 production from encapsulated somatic embryos of cork oak
507 (*Quercus suber* L.) and automated growth monitoring. *Plant Cell*
508 *Tiss Organ Cult* 95(2):217–225
- 509 Pow JJ (1969) Clonal propagation in vitro from cauliflower curd.
510 *Hortic Sci* 9:151–152
- 511 Rai MK, Jaiswal VS, Jaiswal U (2008) Encapsulation of shoot tips of
512 guava (*Psidium guajava* L.) for short-term storage and germ-
513 plasm exchange. *Sci Hortic* 118(1):33–38
- 514 Rani Debi B, Taketa S, Ichii M (2005) Cytokinin inhibits lateral root
515 initiation but stimulates lateral root elongation in rice (*Oryza*
516 *sativa*). *J Plant Physiol* 162(5):507–515
- 517 Roy B, Mandal AB (2008) Development of synthetic seeds involving
518 androgenic and pro-embryos in *elite indica* rice. *Ind J Biotechnol*
519 7(4):515–519
- 520 Sandoval-Yugar EW, Dal Vesco LL, Steinmacher DA, Stolf EC,
521 Guerra MP (2009) Microshoots encapsulation and plant conver-
522 sion of *Musa* sp. cv. 'Grand Naine'. *Ciencia Rural* 39(4)
- 523 Singh S, Rai M, Asthana P, Pandey S, Jaiswal V, Jaiswal U (2009)
524 Plant regeneration from alginate-encapsulated shoot tips of
525 *Spilanthes acmella* L. Murr., a medicinally important and herbal
526 pesticidal plant species. *Acta Physiologiae Plantarum* 31(3):
527 649–653
- 528 Soneji JR, Rao PS, Mhatre M (2002) Germination of synthetic seeds
529 of pineapple (*Ananas comosus* L. Merr.). *Plant Cell Rep* 20:
530 891–894
- 531 Standardi A, Piccioni E (1998) Recent Perspectives on Synthetic Seed
532 Technology Using Nonembryogenic In vitro-Derived Explants.
533 *Plant Sci* 159:968–978
- Sundararaj SG, Agrawal A, Tyagi RK (2010) Encapsulation for in vitro short-term storage and exchange of ginger (*Zingiber officinale* Rosc.) germplasm. *Sci Hortic* 125(4):761–766
- Taha RM, Hasbullah NA, Awal A (2009) Production of synthetic seeds from micro shoots and somatic embryos of *Gerbera jamesoni* bolus ex. hook f. *Acta Hort* 829:91–98
- Teng WL, Ngai YW (1999) Regeneration of *Oxalis triangularis* ssp. *triangularis* from suspension cells cultured in three different systems (solid, liquid-flask and bioreactor cultures). *Plant Cell Rep* 18(9):701–706
- Teng WL, Teng MC (1997) In vitro regeneration patterns of *Platycerium bifurcatum* leaf cell suspension culture. *Plant Cell Rep* 16(12):820–824
- Vandemoortele J-L, Kevers C, Billard J-P, Gaspar T (2001) Osmotic pretreatment promotes axillary shooting from cauliflower curd pieces by acting through internal cytokinin level modifications. *J Plant Physiol* 158(2):221–225
- Ziv M, Ariel T (1991) Bud proliferation and plant regeneration in liquid-cultured philodendron treated with ancymidol and paclobutrazol. *J Plant Growth Regul* 10(1):53–57
- Ziv M, Ronen G, Raviv M (1998) Proliferation of meristematic clusters in disposable presterilized plastic bioreactors for the large-scale micropropagation of plants. *In Vitro Cell and Dev Biol Plant* 34:152–158