


## Electronic Supplementary Information (ESI)


# End-group Effects of Piezofluorochromic Aggregation-induced Enhanced Emission Compounds Containing Distyrylanthracene

Xiqi Zhang,<sup>a</sup> Zhenguo Chi,<sup>\*a</sup> Bingjia Xu,<sup>a</sup> Chengjian Chen,<sup>a</sup> Xie Zhou,<sup>b</sup> Yi Zhang,<sup>a</sup> Siwei Liu<sup>a</sup> and Jiarui Xu<sup>a</sup>

<sup>a</sup> PCFM Lab, DSAPM Lab, KLGHEI of Environment and Energy Chemistry, State Key Laboratory of Optoelectronic Materials and Technologies, School of Chemistry and Chemical Engineering, Sun Yat-sen University, Guangzhou 510275, P. R. China. Fax: +86 20 84112222; Tel: +86 20 84112712; E-mail: chizhg@mail.sysu.edu.cn. <sup>b</sup>School of Pharmaceutical Sciences, Sun Yat-sen University, Guangzhou 510275, China.



**Fig. S1** Calculated spatial electron distributions of LUMOs and HOMOs (left); and the optimized structure of the compounds (right).



**Fig. S2** Time-resolved emission decay curves of the annealed and pressed of AnP<sub>3</sub>, AnP<sub>3</sub>P, AnP<sub>4</sub> and AnP<sub>4</sub>P samples.