

1 **Endocrine Disruption in Male Mosquitofish (*Gambusia holbrooki*)**
2 **Inhabiting Wetlands in Western Australia**

3

4 Running Title: Endocrine disruption in mosquitofish, West Australia

5

6 Clare Game¹, M Monique Gagnon^{1*}, Diane Webb¹, and Richard Lim²

7

8 ¹Department of Environmental Biology, G.P.O. Box U1987, Curtin University of
9 Technology, Perth, Western Australia, 6845.

10 ²Institute for Water and Environmental Resource Management, and Department of
11 Environmental Sciences, University of Technology, Sydney, PO Box 123, Broadway,
12 NSW 2007, Australia.

13

14 * Corr: M. Monique Gagnon, PhD
15 Department of Environmental Biology
16 Curtin University of Technology
17 GPO Box U 1987
18 Perth, Western Australia, 6845
19 Voice: + 61 8 9266 3723
20 Fax: +61 8 9266 2495
21 Email: m.gagnon@curtin.edu.au

22

23

24

25 **Abstract**

26 The use of gonopodial indices as potential indicators of endocrine disruption in the
27 mosquitofish *Gambusia holbrooki* inhabiting south west Australian wetlands was
28 investigated. A minimum of fifty mature males was collected from each of five water-
29 bodies in the Swan Coastal Plain, Western Australia, in order to measure morphological
30 features related to reproduction. A set of morphological measurements were used to
31 derive the following indices: gonopodium length/standard body length, pre-anal
32 length/standard body length, the index of elongation and the percentage of male fish with
33 hooks on the distal end of the gonopodium. Indices of male mosquitofish collected from
34 Jack Finney Lake, located in the Curtin University campus, suggest the presence of
35 endocrine disrupting chemicals (EDCs) in this water-body, while those from Lake
36 Kulinup suggest this is a site of concern. Indices of male fish from the Wagerup wetland,
37 Lake Monger and Loch McNess indicate that fish inhabiting these wetlands are not
38 affected by EDCs. This preliminary study suggests that EDCs may be present in a number
39 of wetlands of the Swan Coastal Plain. Further study using EDC specific markers such as
40 vitellogenin induction in male mosquitofish is required to confirm whether EDCs are
41 present in these water-bodies.

42

43

44 **Keywords** – biomonitoring, endocrine disruption, morphological indices, *Gambusia*

45

46

47

48

49 **Introduction**

50 In recent years concern has increased over the effects of anthropogenic chemicals on the
51 endocrine systems of wildlife. The hypothesis that widespread low-level exposure to
52 endocrine disrupting chemicals (EDCs) can potentially affect an organism's development
53 and a population's viability is well documented. Studies have indicated that chemicals
54 with endocrine disrupting properties can alter the development of secondary sexual
55 characteristics and may impair the reproductive functions of fish (Jobling *et al.*, 1998;
56 Batty and Lim, 1999; Doyle and Lim, 2002; Toft *et al.*, 2004; 2005).

57

58 The environmental effects and risk posed by EDCs in Western Australia (WA) are poorly
59 understood, due to a paucity of recorded studies. It is recognised that EDCs have the
60 potential to enter WA's environment from local mining and manufacturing processes,
61 sewage discharges, and waste management practices (Rose *et al.*, 2003). A few studies
62 have been completed on the effects of EDCs in WA's marine environment such as
63 imposex in intertidal whelks as related to tributyltin (TBT) exposure (Reitsema *et al.*,
64 2003). To date, no studies have investigated the presence of EDCs in WA's freshwater
65 environments, although a field study on the east coast of Australia have described altered
66 morphological characteristics in freshwater fish chronically exposed to EDCs (Batty and
67 Lim, 1999).

68

69 This preliminary study aimed to investigate the possible presence of EDCs in freshwater
70 wetlands of the Swan Coastal Plain, by comparing morphological characteristics in male
71 mosquitofish (*Gambusia holbrooki*) living in potentially contaminated wetlands with
72 those from a reference lake.

73

74 *G. holbrooki* belong to the family of live bearing freshwater fish, the Poeciliidae (Batty
75 and Lim, 1999). They are native to Southern and Central America and were introduced
76 into Western Australia in the 1930s for mosquito control. The fish are easily collected in
77 Perth's metropolitan water-bodies due to their wide distribution, large population sizes
78 and, as they are strongly tolerant to a wide range of environmental conditions, and survive
79 in heavily contaminated water-bodies (Morgan *et al.*, 2004).

80

81 Male *G. holbrooki* are excellent biological indicators of endocrine disruption, due to the
82 fact that they have distinguishable hormone dependent sexual dimorphism under
83 androgenic or oestrogenic control (Batty and Lim, 1999; Toft *et al.*, 2003). On reaching
84 sexual maturity, the anal fin and spinal column of the male undergoes a series of hormone
85 dependent changes that result in the development of a gonopodium, which is used for the
86 transfer of spermatozeugmata during copulation. There are 10 rays in the anal fin of *G.*
87 *holbrooki* (Doyle and Lim, 2002). In sexually immature males, the anal fin rays numbered
88 3 through to 6 are all approximately the same length. However, during the development of
89 the male gonopodium, anal rays 3, 4, and 5 elongate and are modified becoming
90 approximately twice as long as the other rays in the fin. In the female both the anal fin and
91 the body increase in size proportionally, and growth of both continues after reaching
92 sexual maturity (Angus *et al.*, 2001).

93

94 The ratio of the length of ray 4 (which is in the centre of the gonopodium) to ray 6 can be
95 considered as an index of maturity, known as the index of elongation (Angus *et al.*, 2001).
96 During sexual maturation hooks and serrae develop on the distal portion of the
97 gonopodium. The hooks and serrae act as holdfast devices during the transfer of

98 spermatozeugmata, and their presence indicates that the gonopodium is fully developed
99 and the fish sexually mature (Doyle and Lim, 2002).

100

101 **Materials and Methods**

102 *Sampling sites*

103 *G. holbrooki* were randomly sampled at five water bodies located around Perth, Western
104 Australia. These included wetlands located in Alcoa Parklands surrounding bauxite
105 refinery sites at Wagerup and Pinjarra, as well as the metropolitan lakes, Lake Monger,
106 Jack Finney Lake and Loch McNess. The animals were treated humanely according to the
107 Curtin University Animal Ethics Approval number N-53-03.

108

109 The Wagerup Alcoa wetland is located on the land surrounding Alcoa's Wagerup Bauxite
110 Refinery approximately 125 km south east of the Perth central business district (CBD)
111 (Figure 1). Alcoa uses the land surrounding the refinery as experimental farmland for
112 cropping and livestock rearing. No drains enter or discharge from this wetland. Its main
113 water sources are rainfall, groundwater seepage and surface water run-off from the
114 surrounding area. The wetland may be contaminated by polycyclic aromatic hydrocarbons
115 (PAH's) associated with the use of farmland vehicles. Excess nutrient inputs may also
116 occur from fertilizers used in farming practices and from livestock waste. These
117 contaminants have the potential to be carried into the wetland with surface water runoff
118 and groundwater seepage.

119

120 The Pinjarra Alcoa wetland is situated in farmland surrounding Alcoa's Pinjarra Refinery
121 approximately 90km south east of the Perth CBD. This wetland is known as Lake Kulinup
122 and is an important regional wetland formed from a former clay "borrow" pit at the

123 refinery (Alcoa, n.d.). This wetland is relatively closer to the refinery than the wetland at
124 the Wagerup Refinery, which could increase the chance of contamination from by-
125 products of the refinery. However this is considered unlikely as all process fluids are
126 contained and there is no avenue for contamination via surface water runoff or ground
127 water seepages (Simon Sandover, Alcoa, pers comm.). The lake has the potential to
128 receive excess nutrient inputs with surface water runoff carrying fertilizers that are used in
129 the surrounding farmland areas.

130

131 Lake Monger is a large urban water-body with high recreational values located 5 km from
132 the Perth CBD. The lake has been used as a stormwater drainage basin since the early
133 1900s, being fed by underground springs, groundwater and 23 stormwater drains. It
134 eventually discharges into the Swan River via the Mounts Bay Drain. Lake Monger is
135 surrounded by urban land uses with rapid urbanization causing a decline in the health of
136 this lake. The clearing of riparian vegetation has led to nutrient enrichment in the
137 waterway (Lund and Davis, 2000).

138

139 For 59 years, Lake Monger was used as a site for rubbish dumping for the reclamation of
140 shallow swamplands. This practice ceased in 1966, however in some reclaimed areas the
141 rubbish remains 2 to 3 metres below the parkland surfaces. Studies conducted in the early
142 1990's found levels of arsenic, zinc copper, lead, phenols and fenitrophion in the water
143 column exceeded recommended levels for aquatic life. PAHs and high levels total
144 phosphorous are also present in the lake (City of Perth, 1992). The main sources of these
145 contaminants include surface water runoff from surrounding urban areas and extensive
146 drainage from surrounding roads including an adjacent freeway. The sandy soils
147 surrounding the lake lack any capacity to retain nutrients (Swan River Trust, 1999) and
148 fertilizers and herbicides used in lawn maintenance contribute to the lake's contamination

149 (City of Perth, 1992; Lund and Davis, 2000).

150

151 Jack Finney Lake is located on the Curtin University Bentley campus. It is an artificial
152 lake formed by lining the bottom of a hollow with clay. Two drains enter the lake, one
153 from the surrounding parking area and lawn, and the other from an adjacent roadway. The
154 water flowing in from the drains and surface water runoff from its surrounds are suspected
155 to contribute contamination. Consequently, the lake has the potential to receive PAHs via
156 surface water runoff and groundwater seepage. In addition, the lake receives inputs from
157 the surrounding grassed area that is regularly treated with fertilizers and herbicides.

158

159 Loch McNess is located in the northeast section of Yanchep National Park, approximately
160 50 km north of Perth. Compared to other metropolitan wetlands, the conductivity and
161 nutrient levels in Loch McNess are very low, which is attributed to the constant inflow of
162 fresh groundwater and minimal anthropogenic inputs (Gordon *et al.*, 1981), and as such
163 the lake is considered to have high environmental values. For these reasons, Loch McNess
164 was used as the reference site in this study. Yanchep National Park is used for
165 recreational purposes, and fertilizers are used to maintain the lawns, ovals and a golf
166 course in the area surrounding Loch McNess. Surface water runoff and groundwater
167 seepage have the potential to carry these fertilizers into the lake. Nutrients leaching into
168 surrounding ground from underground septic tanks, used in the park since the 1930s, enter
169 the groundwater. However, based on the low recorded nitrogen levels in the lake, the
170 tanks do not appear to have had a significant effect on the water quality of the lake
171 (CALM, 1989).

172

173

174 *Sampling and measurement of mosquitofish characteristics*

175 A minimum of fifty male *G. holbrooki* was captured at each site, using a dip net on a rod.
176 Fish were placed in sealed plastic bags immediately upon capture and put on ice, until
177 returned to the laboratory where they were frozen in a -20°C freezer. The fish were
178 thawed before they were sorted by sex and maturity levels. Only mature males from each
179 site were used for measurement. The presences of hooks on the gonopodium was
180 considered an indication of sexual maturity; however males that lacked hooks but had
181 standard lengths equal to or greater than the smallest individual with hooks from the
182 respective site were also measured as they were considered to be of similar age. Sampling
183 occurred in October 2004.

184

185 The measurements recorded for each mature male included standard body length,
186 gonopodium to snout length (pre-anal length), 4th ray length (gonopodium length), 6th ray
187 length, and the presence or absence of hooks on the distal portion of the gonopodium
188 (Figure 2). Measurements were taken under a dissecting microscope using a ruler with
189 0.1mm divisions. The following indices were calculated: (1) the gonopodium
190 length/standard body length ratio, (2) the pre-anal length/standard body length ratio, (3)
191 the 4th ray/6th ray ratio, and (4) percentage of fish with hooks. As no single index can
192 reveal the severity and extent of endocrine disruption, this suite of complementary ratios
193 were used in order to provide a better overview of endocrine disruption effects.

194

195 *Statistical Analysis*

196 For each index, the data were tested for normality and homoscedasticity. Where these
197 conditions were not met the data was log-transformed. Statistical analysis was undertaken
198 using the statistical package, SPSS 11.5 for Windows. Data for the morphological indices

199 of the gonopodium were analysed using one-way analysis of variance (ANOVA). Where
200 significant differences between sites were found ($p \leq 0.05$), Tukey's Multiple Comparison
201 Test was used to identify differences between the means. The presence/absence of hooks
202 on the gonopodium was tested using the Kruskal-Wallis Test with "site" as the grouping
203 variable. Data are presented as mean \pm standard error (SEM).

204

205 **Results**

206 The mean standard lengths of male fish were significantly different between sites ($p <$
207 0.001 ; Table 1). Fish at Jack Finney Lake and Lake Kulinup were significantly shorter
208 than those from the reference site, Loch McNess. The largest mean standard length was
209 recorded at the reference site.

210

211 Significant differences occurred in the mean gonopodium lengths of male fish between
212 sites ($p < 0.001$; Table 1). Male fish from Jack Finney Lake and Lake Kulinup had
213 significantly smaller mean gonopodium lengths relative to those from the reference site.
214 However, the largest mean gonopodium length was recorded at the Wagerup wetland.

215

216 The mean ratios of gonopodium length/standard body length of the male fish also
217 significantly differed between sites ($p < 0.001$; Figure 2A). Male fish from Jack Finney
218 Lake had a significantly smaller mean ratio than those from the other sample sites except
219 for Lake Kulinup. Male fish captured at the Wagerup site recorded the largest
220 gonopodium length /standard body length ratio.

221

222 The mean pre-anal length/standard body length ratio of male fish showed a significant
223 difference between the sites ($p < 0.001$; Figure 2B). Male fish from the Wagerup site had

224 a lower ratio than that of all other sample sites, while the ratios for male fish from Jack
225 Finney Lake were significantly higher than fish from all the other sites.

226

227 The mean index of elongation (the ratio of the 4th ray length/6th ray length) of male fish
228 showed a significant difference between the sites ($p < 0.001$; Figure 2C). Male fish from
229 Jack Finney Lake were shown to have a significantly smaller ratio than individuals from
230 all other sample sites, while fish from Lake Monger were found to have the largest ratio.

231

232 The percentage of male fish with gonopodial hooks was significantly different between
233 populations of the wetland sites sampled ($p < 0.001$; Figure 2D). All male fish sampled
234 from Loch McNess and Wagerup Alcoa wetlands, had hooks on their gonopodia, the
235 percentage of which were significantly higher than those from the other sites. The
236 percentage of male fish with hooks on their gonopodia from the latter sites (Jack Finney,
237 Lake Kulinup and Lake Monger) differed significantly from one another. Male fish from
238 Jack Finney Lake had the lowest percentage of gonopodia with hooks.

239

240 **Discussion**

241 The results from this preliminary survey (summarised in Table 2) suggest that endocrine
242 disruption is occurring in some wetlands around Perth, as there were significant
243 differences in the morphological indices between populations inhabiting the selected
244 sample sites. Doyle and Lim (2002) showed significant effects on the gonopodial
245 characteristics (e.g. gonopodium length, 4th to 6th ray ratio) when exposed to varying
246 dilutions of oestradiol under laboratory flow through conditions, which confirms that
247 gonopodial characteristics can be affected by EDCs.

248

249 The variable standard lengths recorded between the sites indicate that the fish are
250 exhibiting differential growth patterns. Since all the fish measured were mature males, a
251 shorter standard length might reflect reduced growth related to exposure to contaminants,
252 or may simply reflect the trophic status (i.e. food availability) of the environment in which
253 they were collected. The mean gonopodial length in fish from Jack Finney Lake was
254 significantly shorter than those at the other sites except Lake Kulinup. Batty and Lim
255 (1999) found that a reduced gonopodial length suggests the presence of endocrine
256 disruptors in the water, when they compared male *Gambusia* collected from up and
257 downstream of a sewage plant. In our study however, the standard fish lengths of male
258 fish from Jack Finney and Kulinup Lakes were the shortest recorded; correspondingly the
259 sites that recorded the longest gonopodial lengths also had the largest standard lengths.

260

261 In order to remove the confounding effect of age and the size of the fish, the ratio of
262 gonopodium length to standard length was investigated. As the fish are of different sizes,
263 and thus the gonopodium lengths vary, the standard length is used in the ratio to normalise
264 the data. The development of the gonopodium is under androgenic control and normally
265 occurs as the testes begin to produce androgens at the time of sexual maturation. Exposure
266 of male fish to EDCs could potentially interfere with the normal development of this
267 secondary sexual characteristic (Angus *et al.*, 2001). Results indicate that male *G.*
268 *holbrooki* inhabiting Jack Finney Lake and Lake Kulinup had the lowest gonopodium
269 length to standard length ratios relative to fish inhabiting the other sites, suggesting that
270 EDCs might be present at these sites. The Wagerup wetland had individual males with the
271 highest gonopodium length/standard length ratio, followed by fish from Lake Monger and
272 Loch McNess indicating that sexual development has progressed normally at these three
273 sites.

274

275 The pre-anal length/ standard body length ratio relates to the forward translocation of the
276 gonopodium as the male fish matures (Rosa-Molinar *et al.*, 1998), which is essential for
277 its proper function (Rosa-Molinar *et al.*, 1996). This forward translocation can be affected
278 by EDCs while the gonopodium is developing therefore; a large pre-anal length to
279 standard body length ratio indicates possible endocrine disruption. Male fish from Jack
280 Finney Lake were found to have the highest ratio; this was significantly greater than those
281 from all other sites. This again suggests the presence of EDCs in Jack Finney Lake.

282

283 As the fish matures the 4th ray elongates while the 6th ray does not, therefore this ratio
284 provides an index of gonopodium elongation related to proper sexual maturation (Angus
285 *et al.*, 2001). The ultimate length of the 4th ray is shorter when fish are exposed to EDCs
286 (Doyle and Lim, 2002) and therefore, a lower ratio suggests endocrine disruption has
287 occurred. It was observed that fish collected at Jack Finney Lake had the lowest 4th ray to
288 6th ray ratio pointing to a level of endocrine disruption; this concurs with other endpoints
289 such as the shortest gonopodium length and the lowest gonopodium/standard length ratio.
290 Interestingly, male fish in Loch McNess were found to have the next lowest 4th to 6th ray
291 ratio, although it was significantly higher than that in fish from Jack Finney Lake. Fish
292 collected from Loch McNess were not expected to be exposed to EDCs, as it is located in
293 a National Park. The low value of this index at a site which is theoretically pristine calls
294 for chemical analysis of water and sediments for the presence of EDCs. Alternatively, this
295 result might indicate that the 4th to 6th ray ration is too variable between fish populations,
296 to be used as an indicator of endocrine disruption. Male fish from Lake Monger and
297 Wagerup Alcoa wetlands had the highest elongation ratios, supporting the notion that
298 these sites are not affected by EDCs.

299

300 Significant differences were found in the presence/absence of anal hooks on the distal end
301 of the gonopodia of the fish sampled from the various sites. This characteristic is an
302 indicator of sexual maturity in males and was found to be lowest at Jack Finney Lake,
303 while hooks were present on the gonopodia of 100% of the fish sampled at the Wagerup
304 Alcoa wetlands and Loch McNess indicating they were all sexually mature males. In other
305 studies, fish that were exposed to high levels of EDCs had a lower proportion of
306 gonopodial with hooks compared with that of unexposed fish of the same age and size
307 (Doyle and Lim, 2002). The low percentage of fish with hooks on their gonopodia at Jack
308 Finney Lake once again suggests that endocrine disruption could be occurring in the *G.*
309 *holbrooki* inhabiting this site.

310

311 In summary, there were significant differences in gonopodial characteristics of male
312 mosquitofish from the sites sampled. Jack Finney Lake located on Curtin University
313 grounds had gonopodial characteristics that indicated the highest level of endocrine
314 disruption compared with that of fish from the other sample sites. This may be due to the
315 input of PAHs from surrounding roads and car parks and from the fertilizers and
316 herbicides used on the surrounding grassland. Added to this, Jack Finney Lake has no
317 form of flushing and does not allow for contaminants to seep out of the lake.

318

319 The sites least affected by EDCs appear to be the Wagerup Alcoa Wetland, Lake Monger
320 and Loch McNess. Lake Kulinup represented a site of concern that would require further
321 study to determine the extent of endocrine disruption occurring in the mosquitofish that
322 inhabit the lake. Lake Kulinup is located closer to the refinery than the wetland on the
323 Wagerup Refinery site. The refinery is a possible source of EDCs due to atmospheric

324 fallout; however this would require additional study that incorporated water quality
325 sampling. The lack of any indication of endocrine disruption in Lake Monger was
326 unexpected as this water body is highly impacted by surrounding anthropogenic activities.
327 It is possible that the efforts to revegetate the banks of Lake Monger with riparian
328 vegetation is contributing to filtering excess contaminants or the lake may is polluted with
329 chemicals that do not induce endocrine disruption.

330

331 As no single index alone can reliably detect endocrine disruption in the mosquitofish, it is
332 recommended that a number of indices be used to determine the presence of EDCs in the
333 environment. Coupled with this further studies should include more EDC specific
334 endpoints such as vitellogenin induction in exposed male fish as well as chemical analysis
335 for putatively identified EDCs in water and sediment samples. This will allow better
336 assessment of endocrine disrupting effects of water bodies based on a weight of evidence
337 approach. The age of the fish collected should also be determined by otolith analysis to
338 ensure all the fish sampled are of similar age and maturity level.

339

340 **Acknowledgements**

341 The authors wish to thank Simon Sandover for facilitating access to the Alcoa wetlands.

342 **References**

- 343 Alcoa. (n.d.). Water use in Western Australia: the facts. Accessed 23 February, 2006.
344 http://www.alcoa.com/australia/en/info_page/water_use.asp.
- 345 Angus R.A., McNatt H.B., Howell W.M., Peoples S.D. 2001. Gonopodium development
346 in normal male and 1-ketotestosterone-treated female mosquitofish (*Gambusia*
347 *affinis*): a quantitative study using computer image analysis. Gen. Comp.
348 Endocrinol. 123:222-234.
- 349 Batty J., Lim R. 1999. Morphological and reproductive characteristics of male
350 mosquitofish (*Gambusia affinis holbrooki*) inhabiting sewage-contaminated
351 waters in New South Wales, Australia. Arch. Environ. Contam. Toxicol. 36:301-
352 307.
- 353 CALM. 1989. Yanchep National Park Management Plan 1989-1999. Perth, Western
354 Australia: Department of Conservation and Land Management. Management Plan
355 No. 14. 178 p.
- 356 City of Perth. 1992. Lake Monger Management Plan. Perth, Western Australia: City of
357 Perth. 79 p.
- 358 Doyle C.J., Lim R.P. 2002. The effect of 17 β -estradiol on the gonopodial development
359 and sexual activity of *Gambusia holbrooki*. Environ. Toxicol. Chem. 21:2719-
360 2724.
- 361 Gordon D.M., Finlayson C.M., McComb A.J. 1981. Nutrients and phytoplankton in three
362 shallow, freshwater lakes of different trophic status in Western Australia. Aust. J.
363 Mar. Freshwater Res. 32:541-553.
- 364 Jobling S., Nolan M., Tyler C.R., Brighty G., Sumpter J.P. 1998. Widespread sexual
365 disruption in wild fish. Environ. Sci. Technol. 32:2498-2506.

- 366 Lund M.A., Davis J.A. 2000. Seasonal dynamics of plankton communities and water
367 chemistry in a eutrophic wetland (Lake Monger, Western Australia): implications
368 for biomanipulation. *Mar. Freshwater Res.* 51:321-332.
- 369 Morgan D.L., Gill H.S., Maddern M.G., Beatty S.K. 2004. Distribution and impacts of
370 introduced freshwater fishes in Western Australia. *NZJ Mar. Fresh. Res.* 38:511-
371 523.
- 372 Reitsema T.J., Field S., Spickett J.T. 2003. Surveying imposex in the coastal waters of
373 Perth, Western Australia, to monitor trends in TBT contamination. *Aust. J Ecotox.*
374 9:87-92.
- 375 Rosa-Molinar E., Fritsch B., Hendricks S.E. 1996. Organizational-activational concept
376 revisited: sexual differentiation in an atherinomorph teleost. *Horm. Behav.*
377 30:563-575.
- 378 Rosa-Molinar E., Proskocil B.J., Hendricks S.E., Fritsch B. 1998. A mechanism for
379 anterior transposition of the anal fin and its appendicular support in the western
380 mosquitofish, *Gambusia affinis affinis* (Baird and Girard, 1854). *Acta anat (Basel)*
381 163:75-91.
- 382 Rose T., McTaggart A., Hillman K. 2003. A review of EDC's on WA water: can any
383 meaningful assessment be made? Chemicals of Concern in Water Conference;
384 2003; Sydney, NSW, Australia.
- 385 Swan River Trust. 1999. Swan-Canning Cleanup Program: an Action Plan to clean up the
386 Swan-Canning Rivers and Estuary. Accessed 9 November 2004.
387 <<http://www.wrc.wa.gov.au/srt/sccp/index.html>>.
- 388 Toft G., Baatrup E., Guillete Jr L.J. 2004. Altered social behaviour and sexual
389 characteristics in mosquitofish (*Gambusia holbrooki*) living downstream of a
390 paper mill. *Aquat. Toxicol.* 70:213-222.

- 391 Toft G., Edwards T.M., Baatrup E., Guillete Jr L.J. 2003. Disturbed sexual characteristics
392 in male mosquitofish. *Environ. Health Perspect* 111:695-701.
- 393 Toft G., Guillete Jr L.J. 2005. Decreased sperm count and sexual behavior in
394 mosquitofish exposed to water from a pesticide-contaminated lake. *Ecotoxicol.*
395 *Environ. Saf.* 60:15-20.
- 396

397 **Table 1:** Morphological measurements (Mean \pm SEM) recorded on male *Gambusia*
 398 *holbrooki* collected from selected wetlands. Within each parameter, groups with similar
 399 letters are not statistically different ($p > 0.05$).

400

Site	Loch McNess	Jack Finney Lake	Kulinup	Wagerup	Lake Monger
Parameter					
N	60	79	73	55	62
Standard Length (mm)	21.9 \pm 0.2 ^b	20.6 \pm 0.2 ^a	20.8 \pm 0.2 ^a	21.7 \pm 0.2 ^b	21.3 \pm 0.2 ^{ab}
Gonopodium length (mm)	7.0 \pm 0.1 ^b	6.2 \pm 0.1 ^a	6.4 \pm 0.1 ^a	7.2 \pm 0.1 ^b	6.9 \pm 0.1 ^b

401

402

403 **Table 2:** Summary of results obtained with the morphological ratios measured on
 404 *Gambusia holbrooki* collected from selected wetlands. A '+' indicates that the indices
 405 suggests the presence of EDC at this site, while a 'O' indicates that no effects were
 406 observed.

407

Site	Loch McNess	Jack Finney Lake	Kulinup	Wagerup	Lake Monger
Index					
Gonopodium length/ Standard body length	O	+	+	O	O
Pre-anal Length/ Standard body length	O	+	+	O	O
Index of Elongation (4th ray length/6th ray length)	O	+	O	O	O
Hook Presence	O	+	+	O	O

408

409

410 **List of Figures**

411 **Figure 1.** Location of fish sampling sites in the Swan Coastal Plain.


412

413 **Figure 2.** Camera lucida drawings of: (A) the modified anal fin of a sexually mature male
414 *Gambusia holbrooki*. Fin rays 3 to 5 elongate to form the gonopodium, which is used for
415 sperm transfer during copulation; and (B) the distal tip of the gonopodium of a sexually
416 mature male *Gambusia holbrooki*. The presence of the hooks (H) and serrae (S) indicates
417 that the gonopodium is fully developed acting as holdfast mechanisms during sperm
418 transfer (modified from Doyle and Lim, 2002).

419

420 **Figure 3.** Mean indices values (\pm SEM) for the five wetlands sampled in the Swan
421 Coastal Plain (A) Gonopodium length/standard length ratio (B) Pre-anal length/standard
422 length ratio (C) Index of elongation (D) Hook presence (%). Bars with same letter have no
423 significant site differences ($p \geq 0.05$). N as in Table 1.

424



425

426

427


428

429

430

431

432



433

434

435

436

437


438

439


440

441


442




443



444



445



446