

# Energy in an Expanding Universe in the Teleparallel Geometry

A. A. Sousa,\* J. S. Moura, and R. B. Pereira

*Instituto de Ciências Exatas e da Terra Campus Universitário do Araguaia  
Universidade Federal de Mato Grosso 78698-000 Pontal do Araguaia, MT, Brazil  
(Received on 7 March, 2008)*

The main purpose of this paper is to explicitly verify the consistency of the energy-momentum and angular momentum tensor of the gravitational field established in the Hamiltonian structure of the Teleparallel Equivalent of General Relativity (TEGR). In order to reach these objectives, we obtained the total energy and angular momentum (matter plus gravitational field) of the closed universe of the Friedmann-Lemaître-Robertson-Walker (FLRW). The result is compared with those obtained from the pseudotensors of Einstein and Landau-Lifshitz. We also applied the field equations (TEGR) in an expanding FLRW universe. Considering the stress energy-momentum tensor for a perfect fluid, we found a teleparallel equivalent of Friedmann equations of General Relativity (GR).

Keywords: Gravitation; teleparallelism; gravitational energy- momentum tensor.

## I. INTRODUCTION

It is generally believed that the energy of a gravitational field is not localizable, that is, defined in a finite region of space. An example of this interpretation can be found in the works of Landau and Lifshitz [1], where they present a pseudotensor of the gravitational field that is dependent of the second derivative of the metric tensor. This quantity can be annulled by an adequate transformation of coordinates. The results would be consistent with Einstein's principle of equivalence. According to this principle, you can always find a small region of space-time that prevails in the space-time of Minkowski. In such space-time, the gravitational field is null. Therefore, it is only possible to define the energy of a gravitational field in a whole space-time region and not in a small region. To avoid this difficulty, alternative geometric models to GR were constructed out of the torsion tensor.

The notion of torsion in the space-time was introduced by Cartan [2][3], who also gave a geometric interpretation for this tensor. Consider a vector in some space-time point and transport it simultaneously along a closed infinitesimal curve projected in the space tangent. If the connection used to accomplish the transport parallel has torsion, we will obtain a "gap" among curve extremities in the space tangent. In other words, infinitesimal geodesic parallelograms do not close in the presence of torsion [4]. The curvature effect already produces a change in the vector direction when it returns to the starting point. This way, while the torsion appears directly related to translations, the curvature appears directly related to rotations in the space-time.

It is important to mention that Weitzenböck [5] independently introduced a space-time that presents torsion with null curvature during the 1920s. This space-time possesses a pseudo-Riemannian metric, based on tetrads, known as the Weitzenböck space-time. A tetrad is a set of four linearly independent vectors defined at every point in a space-time. The condition that we have null curvature in Weitzenböck space-time leads to an absolute parallelism or teleparallelism of a tetrads field. The first proposal of using tetrads for the description of the gravitational field was made by Einstein [6] in

1928 in the attempt to unify the gravitational and electromagnetic fields. However, his attempt failed when it did not find a Schwarzschild's solution for the simplified form of its field equation. The description of gravitation in terms of absolute parallelism and the tetrads field were forgotten for some time. Later, Møller [7] rescued Einstein's idea by showing that only in terms of tetrads we can obtain a Lagrangian density that leads to a tensor of gravitational energy-momentum. This tensor, constructed from the first derivatives of the tetrads, does not vanishes in any coordinates transformation. An alternative teleparallel geometric description to GR is the formulation of the Teleparallel Equivalent of General Relativity (TEGR). In this formalism, the Lagrangian density contains quadratic torsion terms and is invariant under global Lorentz transformation, general coordinate and parity transformation [8].

In 1994, Maluf [9] established the Hamiltonian formulation of the TEGR in Schwinger's time gauge [10]. An essential feature of the Hamiltonian formulation shows that we can define the energy of a gravitational field by means of an adequate interpretation of the Hamiltonian constraint. Several configurations of gravitational energies were investigated with success, such as in the space-time configurations of de Sitter [11], conical defects [12], static Bondi [13], disclination defects [14], Kerr black hole [15], Banãdos, Teitelboim and Zanelli (BTZ) black hole [8] and Kerr anti-de Sitter [16]. For Andrade and Pereira [17], the TEGR can indeed be understood as a gauge theory for the translation group. In this approach, the gravitational interaction is described by a force similar to the Lorentz force equation of electrodynamics, with torsion playing the role of force.

In 2000, Sousa and Maluf [18][19] established the Hamiltonian formulation of arbitrary teleparallel theories using Schwinger's time gauge. In this approach, they showed that the TEGR is the only viable consistent teleparallel gravity theory.

In 2001, Maluf and Rocha [20] established a theory in which Schwinger's time gauge was excluded from the geometry of absolute parallelism. In this formulation, the definition of the gravitational angular momentum arises by suitably interpreting the integral form of the constraint equation  $\Gamma^{ab} = 0$ . This definition was applied satisfactorily for the gravitational field of a thin, slowly-rotating mass shell [21] and the three-dimensional BTZ black hole [22].

---

\*Electronic address: adellane@ufmt.br

In GR, the problem of energy-momentum and angular momentum is generally addressed by the energy-momentum (angular momentum) complex. It is calculated as the sum of the energy-momentum (angular momentum) pseudotensor of the gravitational field and the energy-momentum (angular momentum) tensor of the matter. In the literature [23] these complexes appear with several names, such as Landau-Lifshitz, Bergman-Thompson, Einstein and others. They differ from each other in the way they are constructed. These complexes have been applied to several configurations of the gravitational field, such as the universe of Friedmann-Lemaître-Robertson-Walker (FLRW). In these works, Rosen [24], Cooperstock [25], Garecki [26], Johri *et al.* [27] and Vargas [28] show that for the spherical universe, the total energy is zero.

Although the TEGR approach produced consistent results for the energy of several configurations of space-time, the TEGR is not a different geometric structure of GR, but equivalent to it. It is found that the field equations of TEGR are equivalent to the equations of Einstein in the tetrads form [9].

In this work, we first explicitly verify the equivalence between GR and TEGR. More specifically, we consider the solution for an isotropic and homogeneous universe described by the FLRW metric in Cartesian coordinates. The main reason for using these coordinates is for subsequent comparison of our work with other literature results. We find an identical equation to the cosmological equation of Friedmann. We also verify the consistency of the tensorial expressions of the total energy-momentum and angular momentum from the Hamiltonian formalism of the TEGR. For this, we apply the Hamiltonian formulation implemented by Maluf [21][29] to find the total energy-momentum (gravitational field plus matter) and gravitational angular momentum values in the FLRW universe. It is shown that all these quantities vanish for flat and spherical geometries.

The article is organized as follows. In section 1, we review the Lagrangian and Hamiltonian formulation of the TEGR. In section 2, using the field equations of the TEGR, we find the teleparallel version of Friedmann equations. In section 3, we calculate the total energy of the FLRW universe and compare it with those obtained from the pseudotensors. In section 4, we find the total three-momentum of the universe. In section 5, we obtain the gravitational angular momentum of the FLRW universe. Finally, in section 6, we present our conclusions.

The notation is the following: space-time indices  $\mu, \nu, \dots$  and global  $SO(3, 1)$  indices  $a, b, \dots$  run from 0 to 3. Time and space indices are indicated according to  $\mu = 0, a = (0), (i)$ . The tetrad field is denoted by  $e^a{}_{\mu}$  and the torsion tensor reads  $T_{a\mu\nu} = \partial_{\mu}e_{a\nu} - \partial_{\nu}e_{a\mu}$ . The flat, Minkowski space-time metric tensor raises and lowers tetrad indices and is fixed by  $\eta_{ab} = e_{a\mu}e_{b\nu}g^{\mu\nu} = (-+++)$ . The determinant of the tetrad field is represented by  $e = \det(e^a{}_{\mu})$ . We use units **in which**  $c = 1$ , where  $c$  is the light speed.

## II. THE HAMILTONIAN CONSTRAINTS EQUATIONS AS AN ENERGY AND GRAVITATIONAL ANGULAR MOMENTUM EQUATIONS

We will briefly recall both the Lagrangian and Hamiltonian formulations of the TEGR. The Lagrangian density for the gravitational field in the TEGR [10] with the cosmological constant  $\Lambda$  is given by

$$L(e_{a\mu}) = -k' e \left( \frac{1}{4} T^{abc} T_{abc} + \frac{1}{2} T^{abc} T_{bac} - T^a T_a \right) - L_M - 2ek' \Lambda \equiv -k' e \Sigma^{abc} T_{abc} - L_M - 2ek' \Lambda, \quad (1)$$

where  $k' = 1/(16\pi G)$ ,  $G$  is the Newtonian gravitational constant and  $L_M$  stands for the Lagrangian density for the matter fields. As usual, tetrad fields convert space-time into Lorentz indices and vice versa. The tensor  $\Sigma^{abc}$  is defined by

$$\Sigma^{abc} = \frac{1}{4} (T^{abc} + T^{bac} - T^{cab}) + \frac{1}{2} (\eta^{ac} T^b - \eta^{ab} T^c), \quad (2)$$

and  $T^b = T^b{}_{b^a}$ . The quadratic combination  $\Sigma^{abc} T_{abc}$  is proportional to the scalar curvature  $R(e)$ , except for a total divergence. The field equations for the tetrad field read

$$e_{a\lambda} e_{b\mu} \partial_{\nu} (e \Sigma^{b\lambda\nu}) - e \left( \Sigma^{b\nu}{}_{\mu} T_{b\nu\mu} - \frac{1}{4} e_{a\mu} T_{bcd} \Sigma^{bcd} \right) + \frac{1}{2} e e_{a\mu} \Lambda = \frac{1}{4k'} e T_{a\mu}. \quad (3)$$

where  $e T_{a\mu} = \delta L_M / \delta e^{a\mu}$ . It is possible to prove by explicit calculations that the left-hand side of Eq. (3) is exactly given by

$$\frac{1}{2} e \left\{ R_{a\mu}(e) - \frac{1}{2} e_{a\mu} R(e) + e_{a\mu} \Lambda \right\}, \quad (4)$$

and thus, it follows that the field equations arising from the variation of  $L$  with respect to  $e^a{}_{\mu}$  are strictly equivalent to Einstein's equations in tetrad form.

The field equations (3) may be rewritten in the form

$$\partial_{\nu} (e \Sigma^{a\lambda\nu}) = \frac{1}{4k'} e e^a{}_{\mu} (t^{\lambda\mu} + \tilde{T}^{\lambda\mu}), \quad (5)$$

where

$$t^{\lambda\mu} = k' \left( 4 \Sigma^{bc\lambda} T_{bc}{}^{\mu} - g^{\lambda\mu} \Sigma^{bcd} T_{bcd} \right), \quad (6)$$

and

$$\tilde{T}^{\lambda\mu} = T^{\lambda\mu} - 2k' g^{\lambda\mu} \Lambda, \quad (7)$$

are interpreted as the gravitational energy-momentum tensor [29][30] and the matter energy-momentum tensor respectively.

The Hamiltonian formulation of the TEGR is obtained by first establishing the phase space variables. The Lagrangian density does not contain the time derivative of the tetrad component  $e_{a0}$ . Therefore this quantity will arise as a Lagrange multiplier [31]. The momentum canonically conjugated to  $e_{ai}$  is given by  $\Pi^{ai} = \delta L / \delta \dot{e}_{ai}$ . The Hamiltonian formulation is obtained by rewriting the Lagrangian density in the form  $L = p\dot{q} - H$ , in terms of  $e_{ai}$ ,  $\Pi^{ai}$  and Lagrange multipliers. The Legendre transform can be successfully carried out and the final form of the Hamiltonian density reads [20]

$$H = e_{a0}C^a + \alpha_{ik}\Gamma^{ik} + \beta_k\Gamma^k, \quad (8)$$

plus a surface term. Here  $\alpha_{ik}$  and  $\beta_k$  are Lagrange multipliers that (after solving the field equations) are identified as  $\alpha_{ik} = 1/2(T_{i0k} + T_{k0i})$  and  $\beta_k = T_{00k}$ .  $C^a$ ,  $\Gamma^{ik}$  and  $\Gamma^k$  are first class constraints. The Poisson brackets between any two field quantities F and G is given by

$$\{F, G\} = \int d^3x \left( \frac{\delta F}{\delta e_{ai}(x)} \frac{\delta G}{\delta \Pi^{ai}(x)} - \frac{\delta F}{\delta \Pi^{ai}(x)} \frac{\delta G}{\delta e_{ai}(x)} \right). \quad (9)$$

The constraint  $C^a$  is written as  $C^a = -\partial_i \Pi^{ai} + h^a$ , where  $h^a$  is an intricate expression of the field variables. The integral form of the constraint equation  $C^a = 0$  motivates the definition of the energy-momentum  $P^a$  four-vector [15]

$$P^a = - \int_V d^3x \partial_i \Pi^{ai}, \quad (10)$$

where  $V$  is an arbitrary volume of the three-dimensional space. In the configuration space we have

$$\Pi^{ai} = -4k' e \Sigma^{a0i}. \quad (11)$$

The emergence of total divergences in the form of scalar or vector densities is possible in the framework of theories constructed out of the torsion tensor. Metric theories of gravity do not share this feature.

By making  $\lambda = 0$  in equation (5) and identifying  $\Pi^{ai}$  in the left-hand side of the latter, the integral form of Eq. (5) is written as

$$P^a = \int_V d^3x e e^a{}_{\mu} (t^{0\mu} + \tilde{T}^{0\mu}). \quad (12)$$

This equation suggests that  $P^a$  is now understood as the total, gravitational and matter fields (plus a cosmological constant fluid) energy-momentum [29]. The spatial components  $P^{(i)}$  form a total three-momentum, while temporal component  $P^{(0)}$  is the total energy (gravitational field plus matter) [1].

It is important to rewrite the Hamiltonian density  $H$  in the simplest form. It is possible to simplify the constraints into a single constraint  $\Gamma^{ab}$ . It is then simple to verify that the Hamiltonian density (8) may be written in the equivalent form [21]

$$H = e_{a0}C^a + \frac{1}{2}\lambda_{ab}\Gamma^{ab}, \quad (13)$$

where  $\lambda_{ab} = -\lambda_{ba}$  are Lagrange multipliers that are identified as  $\lambda_{ik} = \alpha_{ik}$  and  $\lambda_{0k} = -\lambda_{k0} = \beta_k$ . The constraints  $\Gamma^{ab} = -\Gamma^{ba}$  embodies both constraints  $\Gamma^{ik}$  and  $\Gamma^k$  by means of relation

$$\Gamma^{ik} = e_a{}^i e_b{}^k \Gamma^{ab}, \quad (14)$$

and

$$\Gamma^k \equiv \Gamma^{0k} = \Gamma^{ik} = \Gamma^{ik} = e_a{}^0 e_b{}^k \Gamma^{ab}. \quad (15)$$

It reads

$$\Gamma^{ab} = M^{ab} + 4k' e \left( \Sigma^{a0b} - \Sigma^{b0a} \right). \quad (16)$$

Similar to the definition of  $P^a$ , the integral form of the constraint equation  $\Gamma^{ab} = 0$  motivates the new definition of the space-time angular momentum. The equation  $\Gamma^{ab} = 0$  implies

$$M^{ab} = -4k' e \left( \Sigma^{a0b} - \Sigma^{b0a} \right). \quad (17)$$

Therefore Maluf [21] defines

$$L^{ab} = \int_V d^3x e^a{}_{\mu} e^b{}_{\nu} M^{\mu\nu}, \quad (18)$$

where

$$M^{ab} = e^a{}_{\mu} e^b{}_{\nu} M^{\mu\nu} = -M^{ba}. \quad (19)$$

as the four-angular momentum of the gravitational field.

The quantities  $P^a$  and  $L^{ab}$  are separately invariant under general coordinate transformations of the three-dimensional space and under time reparametrizations, which is an expected feature since these definitions arise in the Hamiltonian formulation of the theory. Moreover, these quantities transform covariantly under global SO(3,1) transformations.

### III. TELEPARALLEL VERSION OF FRIEDMANN EQUATIONS

In this section, we will derive several TEGR expressions capable of calculating the total momentum and angular momentum of the gravitational field in the FLRW universe model. Indeed, we will explicitly demonstrate the equivalence between GR and TEGR in this specific model.

In order to solve the field equations (3) of TEGR, it is necessary to determine the tetrads field. In the Cartesian coordinate system, the line element of the FLRW space-time [32] is given by

$$ds^2 = -dt^2 + \frac{R(t)^2}{\left(1 + \frac{kr^2}{4}\right)^2} (dx^2 + dy^2 + dz^2), \quad (20)$$

where  $r^2 = x^2 + y^2 + z^2$ ,  $R(t)$  is the time-dependent cosmological scale factor, and  $k$  is the curvature parameter, which can assume the values  $k = 0$  (flat FLRW universe),  $k = +1$

(spherical FLRW universe) and  $k = -1$  (hyperbolic FLRW universe).

Using the relations

$$g_{\mu\nu} = e^a{}_{\mu} e_{a\nu}, \quad (21)$$

and

$$e_{a\mu} = \eta_{ab} e^b{}_{\mu}, \quad (22)$$

a set of tetrads fields that satisfy the metric is given by

$$e_{a\mu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & \frac{R(t)}{(1+\frac{kr^2}{4})} & 0 & 0 \\ 0 & 0 & \frac{R(t)}{(1+\frac{kr^2}{4})} & 0 \\ 0 & 0 & 0 & \frac{R(t)}{(1+\frac{kr^2}{4})} \end{pmatrix}. \quad (23)$$

We remember here that we use two simplifications to choose this tetrads field. The first simplification is the Schwinger's time gauge condition [10]

$$e^{(k)}{}_{0} = 0. \quad (24)$$

which implies

$$e^{(0)}{}_{k} = 0. \quad (25)$$

The second simplification is the symmetry valid in Cartesian coordinates for space tetrads components

$$e_{(i)j} = e_{(j)i}, \quad (26)$$

that establish a unique reference space-time that is neither related by a boost transformation nor rotating with respect to the physical space-time [15].

Now, with the help of the inverse metric tensor  $g^{\mu\nu}$ , we can write the inverse tetrads

$$e_a{}^{\mu} = g^{\mu\nu} e_{a\nu}, \quad (27)$$

as

$$e_a{}^{\mu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{(1+\frac{kr^2}{4})}{R(t)} & 0 & 0 \\ 0 & 0 & \frac{(1+\frac{kr^2}{4})}{R(t)} & 0 \\ 0 & 0 & 0 & \frac{(1+\frac{kr^2}{4})}{R(t)} \end{pmatrix}, \quad (28)$$

where the determinant of  $e^a{}_{\mu}$  is

$$e = \frac{R^3(t)}{(1+\frac{kr^2}{4})^3}. \quad (29)$$

Before solving the field equations, it is necessary to consider the material content of the universe. We restrict our

consideration here to the stress-energy-momentum tensor of a perfect fluid [32] given by

$$T^{\mu}{}_{\nu} = \begin{pmatrix} \rho & 0 & 0 & 0 \\ 0 & -p & 0 & 0 \\ 0 & 0 & -p & 0 \\ 0 & 0 & 0 & -p \end{pmatrix}, \quad (30)$$

where  $\rho = \rho(x)$  is the matter energy density and  $p$  is the matter pressure. It is convenient rewrite the field equations of the TEGR (3) as

$$\begin{aligned} & e_{a\lambda} e_{b\mu} \partial_{\nu} (e e_d{}^{\lambda} e_c{}^{\nu} \Sigma^{bcd}) \\ & - e \left( \eta_{ad} e_c{}^{\nu} \Sigma^{bcd} T_{b\nu\mu} - \frac{1}{4} e_{a\mu} e_c{}^{\nu} e_d{}^{\nu} T_{b\nu\lambda} \Sigma^{bcd} \right) \\ & + \frac{1}{2} e e_{a\mu} \Lambda = \frac{1}{4k} e e_a{}^{\nu} T_{\nu\mu}, \end{aligned} \quad (31)$$

where:

$$\begin{aligned} \Sigma^{abc} &= \frac{1}{4} \left( \eta^{ad} e^{b\mu} e^{c\nu} T_{d\mu\nu} + \eta^{bd} e^{a\mu} e^{c\nu} T_{d\mu\nu} - \eta^{cd} e^{a\mu} e^{b\nu} T_{d\mu\nu} \right) \\ & + \frac{1}{2} \left( \eta^{ac} e^{b\nu} e^{d\mu} T_{d\mu\nu} - \eta^{ab} e^{c\nu} e^{d\mu} T_{d\mu\nu} \right). \end{aligned} \quad (32)$$

These equations were obtained using the transformations given by

$$T^{abc} = T_{d\mu\nu} e^{b\mu} e^{c\nu} \eta^{ad}, \quad (33)$$

$$T_{abc} = T_{a\mu\nu} e_b{}^{\mu} e_c{}^{\nu}, \quad (34)$$

$$T^b = T_{d\mu\nu} e^{b\nu} e^{d\mu}, \quad (35)$$

$$\Sigma^{a\nu}{}_c = \Sigma^{abd} e_b{}^{\nu} \eta_{dc}, \quad (36)$$

$$\Sigma^{a\mu\nu} = \Sigma^{abc} e_b{}^{\mu} e_c{}^{\nu}, \quad (37)$$

$$T_{a\mu} = T_{\nu\mu} e_a{}^{\nu}. \quad (38)$$

The non-zero components of the torsion tensor  $T_{a\mu\nu}$  are given by

$$T_{(1)01} = T_{(2)02} = T_{(3)03} = \frac{\dot{R}(t)}{1+\frac{kr^2}{4}}, \quad (39)$$

$$T_{(1)12} = T_{(3)32} = \frac{R(t)ky}{2\left(1+\frac{kr^2}{4}\right)^2}, \quad (40)$$

$$T_{(1)13} = T_{(2)23} = \frac{R(t)kz}{2\left(1+\frac{kr^2}{4}\right)^2}, \quad (41)$$

$$T_{(2)21} = T_{(3)31} = \frac{R(t)kx}{2\left(1+\frac{kr^2}{4}\right)^2}, \quad (42)$$

remembering that the torsion components are anti-symmetrical under the exchange of the two last indexes.

After tedious but straightforward calculations, we obtain

the non-zero components of the tensor  $\Sigma^{abc}$

$$\Sigma^{(0)(0)(1)} = \frac{kx}{2R(t)}, \quad (43)$$

$$\Sigma^{(0)(0)(2)} = \frac{ky}{2R(t)}, \quad (44)$$

$$\Sigma^{(0)(0)(3)} = \frac{kz}{2R(t)}, \quad (45)$$

$$\Sigma^{(1)(0)(1)} = \Sigma^{(2)(0)(2)} = \Sigma^{(3)(0)(3)} = \frac{\dot{R}(t)}{R(t)}, \quad (46)$$

$$\Sigma^{(1)(1)(2)} = \Sigma^{(3)(3)(2)} = -\frac{ky}{4R(t)}, \quad (47)$$

$$\Sigma^{(1)(1)(3)} = \Sigma^{(2)(2)(3)} = -\frac{kz}{4R(t)}, \quad (48)$$

$$\Sigma^{(2)(1)(2)} = \Sigma^{(3)(1)(3)} = \frac{kx}{4R(t)}. \quad (49)$$

Next, we proceed to obtain the components  $\{a = (0), \mu = 0\}$ ,  $\{a = (1), \mu = 1\}$ ,  $\{a = (2), \mu = 2\}$ , and  $\{a = (3), \mu = 3\}$  of the field equations. The other components of field equations are identically zero. This is carried out in two steps. First, we calculate the component  $\{a = (0), \mu = 0\}$ . It is not difficult to obtain

$$\begin{aligned} & e_{(0)0}e_{(0)0} \left[ \partial_1 \left( ee_{(0)}^0 e_{(1)}^1 \Sigma^{(0)(0)(1)} \right) \right. \\ & + \partial_2 \left( ee_{(0)}^0 e_{(2)}^2 \Sigma^{(0)(0)(2)} \right) \\ & + \partial_3 \left( ee_{(0)}^0 e_{(3)}^3 \Sigma^{(0)(0)(3)} \right) \left. \right] \\ & - e \left( 3\eta_{(0)(0)} e_{(1)}^1 \Sigma^{(1)(1)(0)} T_{(1)10} \right. \\ & - \frac{3}{2} e_{(0)0} e_{(0)}^0 e_{(1)}^1 T_{(1)01} \Sigma^{(1)(0)(1)} \\ & - e_{(0)0} e_{(1)}^1 e_{(2)}^2 T_{(1)12} \Sigma^{(1)(1)(2)} \\ & - e_{(0)0} e_{(1)}^1 e_{(2)}^2 T_{(2)12} \Sigma^{(2)(1)(2)} \\ & \left. - e_{(0)0} e_{(1)}^1 e_{(3)}^3 T_{(1)13} \Sigma^{(1)(1)(3)} \right) \\ & + \frac{1}{2} ee_{(0)0} \Lambda = \frac{1}{4k'} ee_{(0)}^0 T_{00}. \end{aligned} \quad (50)$$

By substituting (23), (28), (29), (30) and (40)-(48) into the above equation, we arrive at

$$3 \frac{\dot{R}^2(t) + k}{R^2(t)} - \Lambda = 8\pi G\rho. \quad (51)$$

The second step consists of calculating the component  $\{a = (1), \mu = 1\}$  of the field equation. Eliminating the null

terms, we find

$$\begin{aligned} & e_{(1)1}e_{(1)1} \left[ \partial_0 \left( ee_{(1)}^1 e_{(0)}^0 \Sigma^{(1)(1)(0)} \right) \right. \\ & + \partial_2 \left( ee_{(1)}^1 e_{(2)}^2 \Sigma^{(1)(1)(2)} \right) \\ & + \partial_3 \left( ee_{(1)}^1 e_{(3)}^3 \Sigma^{(1)(1)(3)} \right) \left. \right] \\ & - e \left( \eta_{(1)(1)} e_{(0)}^0 \Sigma^{(1)(0)(1)} T_{(1)01} \right. \\ & + \eta_{(1)(1)} e_{(2)}^2 \Sigma^{(1)(2)(1)} T_{(1)21} \\ & + 2\eta_{(1)(1)} e_{(2)}^2 \Sigma^{(2)(2)(1)} T_{(2)21} \\ & + \eta_{(1)(1)} e_{(3)}^3 \Sigma^{(1)(3)(1)} T_{(1)31} \\ & - \frac{3}{2} e_{(1)1} e_{(0)}^0 e_{(1)}^1 T_{(1)01} \Sigma^{(1)(0)(1)} \\ & - e_{(1)1} e_{(1)}^1 e_{(2)}^2 T_{(1)12} \Sigma^{(1)(1)(2)} \\ & - e_{(1)1} e_{(1)}^1 e_{(2)}^2 T_{(2)12} \Sigma^{(2)(1)(2)} \\ & \left. - e_{(1)1} e_{(1)}^1 e_{(3)}^3 T_{(1)13} \Sigma^{(1)(1)(3)} \right) \\ & + \frac{1}{2} ee_{(1)1} \Lambda = \frac{1}{4k'} ee_{(1)}^1 T_{11}. \end{aligned} \quad (52)$$

By replacing (23), (28)-(30), (39)-(42) and (47)-(49) into the above equation, we arrive at

$$\frac{2\dot{R}(t)R(t) + \dot{R}^2(t) + k}{R^2(t)} - \Lambda = -8\pi G\rho. \quad (53)$$

The differential equations for the components  $\{a = (2), \mu = 2\}$  and  $\{a = (3), \mu = 3\}$  are the same as equation (53). The field equations' solutions of the TEGR reduce to the system of equations (51) e (55).

$$\begin{aligned} & 3 \frac{\dot{R}^2(t) + k}{R^2(t)} - \Lambda = 8\pi G\rho, \\ & \frac{\dot{R}^2(t) + 2\dot{R}(t)R(t) + k}{R^2(t)} - \Lambda = -8\pi G\rho, \end{aligned}$$

which is equivalent to the Friedmann equations of General Relativity [33].

#### IV. TOTAL ENERGY OF THE FLRW UNIVERSE

Let us now calculate the total energy of the FLRW universe using the equations shown in section 2. By replacing the equation (5) in (12) and using that

$$\Sigma^{a\lambda\nu} = \Sigma^{abc} e_b^\lambda e_c^\nu, \quad (54)$$

we have

$$P^a = \int_V d^3x 4k' \partial_\nu \left( e \Sigma^{a(0)c} e_{(0)}^0 e_c^\nu \right). \quad (55)$$

As previously observed, the temporal component represents the system energy. Therefore, the energy will be given by

$$P^{(0)} = \int_V d^3x 4k' \partial_\nu \left( e \Sigma^{(0)(0)c} e_0^0 e_c^\nu \right). \quad (56)$$

Such quantity can be written as

$$P^{(0)} = \int_V d^3x 4k' \left[ \partial_1 \left( e \Sigma^{(0)(0)(1)} e_0^0 e_{(1)}^1 \right) + \partial_2 \left( e \Sigma^{(0)(0)(2)} e_0^0 e_{(2)}^2 \right) + \partial_3 \left( e \Sigma^{(0)(0)(3)} e_0^0 e_{(3)}^3 \right) \right]. \quad (57)$$

By replacing (28), (29), (43)-(45) in (57), we obtain

$$P^{(0)} = \frac{3R}{8\pi G} \int_V d^3x \frac{k}{\left(1 + \frac{kr^2}{4}\right)^2} - \frac{R}{8\pi G} \int_V d^3x \frac{k^2 r^2}{\left(1 + \frac{kr^2}{4}\right)^3}. \quad (58)$$

It is convenient to perform the integration in spherical coordinates. By solving the integral in  $\theta$  and  $\phi$ , we find

$$P^{(0)} = \frac{3R}{2G} \int_0^\infty \frac{kr^2}{\left(1 + \frac{kr^2}{4}\right)^2} dr - \frac{R}{2G} \int_0^\infty \frac{k^2 r^4}{\left(1 + \frac{kr^2}{4}\right)^3} dr. \quad (59)$$

We can now obtain the total energy  $P^{(0)}$  of the spherical FLRW universe. By making  $k = 1$ , the component  $P^{(0)}$  results

$$P^{(0)} = \frac{R}{2G} [6\pi - 3(2\pi)] = 0. \quad (60)$$

We found that the total energy of a FLRW spatially-spherical universe is zero at all times, irrespective of the equations of the state of the cosmic fluid.

We remark that by making  $k = 0$  in equation (59), it follows that the total energy in the expanding FLRW flat universe is also zero.

It is important to note that by fixing  $k = -1$ , we obtain an infinite energy  $P^{(0)}$  in the integration interval  $[0, 2]$  in accordance with Vargas [28].

In the past, many researchers used the energy-momentum complexes of general relativity to obtain the energy and momentum of the FLRW universe. Rosen [24] and Cooperstock [25] calculated the energy of the universe, including matter and gravitational field. They used the Einstein pseudotensor of energy-momentum to represent the gravitational energy. The result revealed that the total energy of a FLRW spherical universe is zero. Garecki [26] and Johri *et al.* [27] used the energy complex of Landau-Lifshitz and found the same result. We stress that Vargas [28], using the teleparallel version of Einstein and Landau-Lifshitz pseudotensors, has also obtained zero total energy in a FLRW spherical universe. Our result is compatible with these results.

## V. THE TOTAL MOMENTUM OF THE FLRW UNIVERSE

Clearly the total three-momentum (matter plus gravitational field) of the FLRW universe vanishes according to the physical principle by homogeneity. Let us verify the consistency of our formalism. As seen in section 2, it is noted that the total three-momentum is given by space components  $a = \{1\}$ ,  $\{2\}$  and  $\{3\}$  of the equation (55).

In order to obtain the space component  $a = \{1\}$  of the total momentum, we can write the quantity  $P^{(1)}$  as

$$P^{(1)} = \int_V d^3x 4k' \partial_1 \left( e \Sigma^{(1)(0)(1)} e_{(0)}^0 e_{(1)}^1 \right). \quad (61)$$

By substituting (28), (29) and (46) in the previous equation, we obtain

$$P^{(1)} = -\frac{R\dot{R}}{4\pi G} \int_V d^3x \left[ \frac{kx}{\left(1 + \frac{kr^2}{4}\right)^3} \right]. \quad (62)$$

In order to calculate this integral, we observe that the integrand in (62) is an odd function. Thus

$$P^{(1)} = 0. \quad (63)$$

The calculations to obtain the other two components of the total three-momentum are analogous. We found  $P^{(2)} = P^{(3)} = 0$ . All three components of the total momentum are zero regardless of the curvature parameter in the expanding universe.

## VI. GRAVITATIONAL ANGULAR MOMENTUM

According to the physical principle by isotropy, the angular momentum must vanish. Let us verify the consistency of the expression of gravitational angular momentum (18). By making use of (19) and (17) we can write (18) in the form

$$L^{ab} = -\int_V d^3x 4k' e \left( \Sigma^{a0b} - \Sigma^{b0a} \right). \quad (64)$$

By making use  $\Sigma^{a0b} = e_c^0 \Sigma^{acb}$  and reminding that the tetrad field matrix (28) is diagonal, then the equation (64) can be rewritten as

$$L^{ab} = -\int_V d^3x 4k' e e_{(0)}^0 \left( \Sigma^{a(0)b} - \Sigma^{b(0)a} \right). \quad (65)$$

Consider the non-zero components of the tensor  $\Sigma^{abc}$  given by equation (43) up to (49). It is clear that the non-zero components of the angular momentum tensor are  $L^{(0)(1)}$ ,  $L^{(0)(2)}$  and  $L^{(0)(3)}$ . Therefore, it is simple to obtain

$$L^{(0)(1)} = -\int_V d^3x 4k' e e_{(0)}^0 \Sigma^{(0)(0)(1)}. \quad (66)$$

By replacing (28), (29) and (43), we have

$$L^{(0)(1)} = -\frac{R^2}{8\pi G} \int_V d^3x \frac{kx}{\left(1 + \frac{kr^2}{4}\right)^3}. \quad (67)$$

Again, we observe that the integrand is an odd function, thus

$$L^{(0)(1)} = 0. \quad (68)$$

By analogous calculations we found the components  $L^{(0)(2)} = L^{(0)(3)} = 0$ .

## VII. CONCLUSIONS

In the first part of this work, we analyzed the equivalence between General Relativity and TEGR. According to this equivalence, while the GR describes the gravitation through the curvature, teleparallelism describes the same gravitation, but using torsion. Starting with a Lagrangian density composed by a quadratic combination of terms in the torsion, and that contains the condition of null curvature as a constraint, it is obtained that the dynamic equation of tetrads is equivalent to Einstein's equations. This characterizes the Teleparallelism Equivalent of General Relativity. In order to show explicitly the equivalence between GR and TEGR, we found a tetrads field in FLRW space-time that describes the cosmological model standard (isotropic and homogeneous). In this cosmological model, we concluded that field equations of the TEGR are equivalent to the Friedmann equations of GR.

In the second part of this work, using the gravitational tensor in the context of the TEGR, we calculated the total energy of the FLRW universe. For spherical universe ( $k = 1$ ), the total energy is zero, irrespective of the equations of the state of the cosmic fluid, agreeing with the results using the pseudotensors of Rosen, Cooperstock, Garecki, Johri *et al.* and Vargas. This result is in accord with the arguments presented by Tryon [34]. He proposed that our universe may have arisen as a quantum fluctuation of the vacuum and mentioned that no conservation law of physics needed to have been violated at the time of its creation. He showed that in the early spherical universe, the gravitational energy cancels out the energy of the created matter. For the flat universe ( $k = 0$ ), the energy vanishes, as expected and in accordance with the previously

cited papers. Finally, for the hyperbolic universe ( $k = -1$ ), the energy diverges in the interval of integration  $[0, 2]$ . Any possible infinity energy in the universe could lead to "big rip" type singularities [35].

We showed by consistency of formalism that the total three-momentum of the FLRW universe vanishes according to the physical principle by homogeneity. Finally, we showed by consistency of formalism that the components of the angular momentum of the FLRW universe is zero. This result was also expected since there are no privileged directions in this expanding space-time.

Although, some results about total energy-momentum and angular momentum found in this paper were expected and previously obtained in the literature, we concluded from this work that the TEGR obtained equivalent results to the GR with the great advantage of addressing covariantly the definitions of quantities like energy-momentum and angular momentum tensors of the gravitational field.

In order to continue testing the gravitational energy-momentum tensor of the TEGR, we intend to calculate the total energy of the closed Bianchi type I and II universes and Gödel-type metric, among other configurations. These metrics are important mainly in the investigation of the possible anisotropic early universe models. In particular, we expected to find specific contributions of the anisotropy universe model to the components of the total energy-momentum and angular momentum tensors. Efforts in this respect will be carried out.

### Acknowledgements

One of us (J. S. M.) would like to thank the Brazilian agency CAPES for their financial support.

- 
- [1] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, Oxford, 1980).
- [2] E. Cartan, *C. R. Acad. Sci.* **174**, 734 (1922).
- [3] E. Cartan, *C. R. Acad. Sci.* **174**, 593 (1922).
- [4] J. G. Pereira and T. Vargas, *Class. Quantum Grav.* **19**, 4807 (2002).
- [5] R. Weitzenböck, *Invarianten Theorie* (Nordhoff, Groningen, 1923).
- [6] A. Einstein, *Sitzungsber. Preuss. Akad. Wiss.* **217**, 224 (1928); A. Unzicker and T. Case, *Translation of Einstein's Attempt of a Unified Field Theory with Teleparallelism*, e-print:physics/0503046, (2005).
- [7] C. Möller, *Ann. Phys.* **12**, 118 (1961).
- [8] A. A. Sousa and J. W. Maluf, *Prog. Theor. Phys.* **108**, 457 (2002).
- [9] J. W. Maluf, *J. Math. Phys.* **35**, 335 (1994).
- [10] J. Schwinger, *Phys. Rev.* **130**, 1253 (1963).
- [11] J. W. Maluf, *J. Math. Phys.* **37**, 6293 (1996).
- [12] J. W. Maluf and A. Kneip, *J. Math. Phys.* **38**, 458 (1997).
- [13] J. W. Maluf and J.F. da Rocha-Neto, *J. Math. Phys.* **40**, 1490 (1999).
- [14] J. W. Maluf and A. Goya, *Class. Quant. Grav.* **18**, 5143 (2001).
- [15] J. W. Maluf, J.F. da Rocha-Neto, T.M.L. Toribio and K.H. Castello-Branco, *Phys. Rev. D* **65**, 124001 (2002).
- [16] J. F. da Rocha-Neto and K. H. Castello-Branco, *JHEP* **0311**, 002 (2003).
- [17] V. C. Andrade and J. G. Pereira, *Phys. Rev. D* **56**, 4689 (1997).
- [18] J. W. Maluf and A. A. Sousa, *Hamiltonian formulation of teleparallel theories of gravity in the time gauge*, e-print:gr-qc/0002060, (2000).
- [19] A. A. Sousa and J. W. Maluf, *Prog. Theor. Phys.* **104**, 531 (2000).
- [20] J. W. Maluf and J. F. da Rocha-Neto, *Phys. Rev. D* **64**, 084014 (2001).
- [21] J. W. Maluf, S. C. Ulhoa, F. F. Faria and J. F. da Rocha-Neto, *Class. Quantum Grav.* **23**, 6245 (2006).
- [22] A. A. Sousa, R. B. Pereira and J. F. Rocha-Neto, *Prog. Theor. Phys.*, **114**, 1179 (2005).
- [23] J. M. Aguirregabiria, A. Chamorro, K. S. Virbhadra, *Gen. Rel. Grav.* **28**, 1393 (1996).
- [24] N. Rosen, *Gen. Rel. Grav.* **26**, 319 (1994).
- [25] F. I. Cooperstock, *Gen. Rel. Grav.* **26**, 323 (1994).
- [26] J. Garecki, *Gen. Rel. Grav.* **27**, 55 (1995).
- [27] V. B. Johri, D. Kalligas, G. P. Singh and C. W. F. Everitt, *Gen. Rel. Grav.* **27**, 313 (1995).
- [28] T. Vargas, *Gen. Rel. Grav.* **36**, 1255 (2004).
- [29] J. W. Maluf, *Annalen Phys.* **14**, 723 (2005).
- [30] J. W. Maluf and F. F. Faria, *Class. Quantum Grav.* **20**, 4683 (2003).
- [31] P. A. M. Dirac, *Lectures on Quantum Mechanics* (Belfer Graduate School of Science (Monographs Series No 2), Yeshiva University, New York, 1964).
- [32] R. A. d'Inverno, *Introducing Einstein's Relativity* (Clarendon Press, Oxford, 1992).
- [33] M. Dalarsson and N. Dalarsson, *Tensors, Relativity and Cosmology* (Elsevier Academic Press, 2005).

[34] E. P. Tryon, *Nature* **246**, 396 (1973).

[35] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, *Phys.*

*Rev. Lett.* **91**, 071301 (2003).