

Enhancement of cardiac function after adenoviral-mediated in vivo intracoronary β_2 -adrenergic receptor gene delivery

John P. Maurice,¹ Jonathan A. Hata,¹ Ashish S. Shah,¹ David C. White,¹ Patricia H. McDonald,² Paul C. Dolber,¹ Katrina H. Wilson,² Robert J. Lefkowitz,² Donald D. Glower,¹ and Walter J. Koch¹

¹Department of Surgery, and

²Departments of Medicine and Biochemistry, and The Howard Hughes Medical Institute, Duke University Medical Center, Durham, North Carolina 27710, USA

Address correspondence to: Walter J. Koch, Department of Surgery, Duke University Medical Center, Room 472, Medical Sciences Research Building, Box 2606, Research Drive, Durham, North Carolina 27710, USA. Phone: (919) 684-3007; Fax: (919) 684-5714; E-mail: koch0002@mc.duke.edu.

John P. Maurice and Jonathan A. Hata contributed equally to this work.

Received for publication December 10, 1998, and accepted in revised form May 18, 1999.

Exogenous gene delivery to alter the function of the heart is a potential novel therapeutic strategy for treatment of cardiovascular diseases such as heart failure (HF). Before gene therapy approaches to alter cardiac function can be realized, efficient and reproducible in vivo gene techniques must be established to efficiently transfer transgenes globally to the myocardium. We have been testing the hypothesis that genetic manipulation of the myocardial β -adrenergic receptor (β -AR) system, which is impaired in HF, can enhance cardiac function. We have delivered adenoviral transgenes, including the human β_2 -AR (*Adeno- β_2 AR*), to the myocardium of rabbits using an intracoronary approach. Catheter-mediated *Adeno- β_2 AR* delivery produced diffuse multichamber myocardial expression, peaking 1 week after gene transfer. A total of 5×10^{11} viral particles of *Adeno- β_2 AR* reproducibly produced 5- to 10-fold β -AR overexpression in the heart, which, at 7 and 21 days after delivery, resulted in increased in vivo hemodynamic function compared with control rabbits that received an empty adenovirus. Several physiological parameters, including dp/dt_{max} as a measure of contractility, were significantly enhanced basally and showed increased responsiveness to the β -agonist isoproterenol. Our results demonstrate that global myocardial in vivo gene delivery is possible and that genetic manipulation of β -AR density can result in enhanced cardiac performance. Thus, replacement of lost receptors seen in HF may represent novel inotropic therapy.

J. Clin. Invest. 104:21–29 (1999).

Introduction

Cardiovascular disease accounts for nearly 40% of all deaths annually in the United States. Advances in medical treatments have dramatically reduced the overall mortality rate due to heart disease; however, death due to chronic heart failure (HF) continues to rise, and effective therapy has been elusive (1). Therefore, novel therapeutic approaches for treating the failing heart are of interest. The β -adrenergic receptor (β -AR) signaling system, which mediates the inotropic and chronotropic effects of the sympathetic transmitters epinephrine and norepinephrine, plays a critical role in the regulation of cardiac performance, which is particularly important under pathological circumstances. Multiple lines of experimental evidence indicate fundamental alterations in the myocardial β -AR system that both precede and accompany the development of HF (2–4). In failing human myocardium, there is a 50% decrease in β -ARs that is specific for the β_1 -subtype, whereas remaining receptors (β_1 and β_2) are functionally uncou-

pled (5, 6). Functional uncoupling of myocardial β -ARs in human HF may be due to elevated levels of the β -AR kinase (β ARK1) that has been shown to be increased in human HF (6). Drugs targeting adrenergic signaling, including β -agonists for acute situations and β -blockers for chronic therapy, have been at the forefront of conventional therapy for HF; however, administration of β -AR agonists to stimulate inotropy in the failing heart have inherently limited efficacy, partly because of the specific reduction in β -AR targets (7).

Gene transfer in vivo to the heart has significant therapeutic potential for several cardiovascular disorders. The delivery of a transgene encoding a β -AR to the myocardium is an attractive strategy for improving cardiac function in HF by replacing lost receptors. In fact, transgenic mouse technology has already demonstrated that myocardial overexpression of β_2 -ARs results in mice with significantly enhanced cardiac function without significant negative effects (8, 9). Moreover, we have recently shown that adenoviral-mediated β_2 -AR




Figure 1

Expression of β -galactosidase in rabbit hearts after intracoronary in vivo delivery of a recombinant adenovirus. (a) X-Gal staining of a cross-section of a heart taken at midventricular level 6 days after intracoronary delivery of 1×10^{12} tvp of *Adeno- β Gal*. $\times 3$. (b) Representative higher magnification of *Adeno- β Gal*-treated hearts showing myocyte-specific β -Gal expression in cross (left) and longitudinal (right) sections. $\times 40$. (c) Representative sections from hearts injected with EV (left) or *Adeno- β_2 AR* (right). $\times 40$.

gene transfer to cultured ventricular myocytes isolated from rabbit hearts paced into HF restores β -AR signaling to normal levels (10).

Before gene therapy approaches to alter cardiac function can be realized, efficient and reproducible in vivo gene delivery to the heart must be established and optimized. Recombinant adenoviruses appear to offer the most advantages for myocardial gene transfer. Numerous reports have described attempts at direct injection of adenoviral constructs containing marker genes into the myocardium (11–13), but limitations of this technique include low efficiency of infection and insufficient volume of myocardial expression to alter cardiac function. In addition, there have been attempts to deliver adenoviral transgenes to the myocardium through the coronary arteries (14–16). The purpose of this study was to develop an intracoronary artery delivery method for efficient and reproducible global gene transfer to the rabbit myocardium. In addition, we wanted to test the hypothesis that overexpression of β_2 -ARs can alter both biochemical and in vivo cardiac function in the rabbit.

Methods

Adenoviral transgenes. The adenoviral backbone used for the construction of the vectors containing the human β_2 -AR transgene (*Adeno- β_2 AR*) and a cytoplasmic-expressing β -galactosidase transgene (*Adeno- β Gal*) was a replication-deficient first-generation type V adenovirus with deletions of the E1 and E3 genes as we have described previously (10, 17). Large-scale preparations of these adenoviruses were purified from infected EBNA-transfected 293 cells (Invitrogen Corp., Carlsbad, California, USA) as described (10, 17).

In vivo intracoronary delivery of adenoviruses. Animals used in this study were adult male New Zealand white rabbits (1–2 kg). Animals were housed under standard conditions and allowed to feed ad libitum. The Animal

Care and Use Committee of Duke University Medical Center approved all procedures performed in accordance with the regulations adopted by the National Institutes of Health. Rabbits were anesthetized with a mixture of ketamine (80 mg/kg) and acepromazine (0.1 mg/kg), intubated, and mechanically ventilated. Additional ketamine was administered intravenously if necessary during the procedure. A right thoracotomy was performed through the third or fourth intercostal space, and the aortic root was exposed by blunt dissection. A bolus of adenosine (0.75 mg/kg) was delivered to the right ventricular (RV) chamber by a 30-gauge syringe. Immediately, a 25-gauge catheter was inserted through the apical myocardium; the proximal aorta was cross-clamped; and a 1.5-mL bolus of either saline or adenoviral solution was rapidly injected into the left ventricular (LV) chamber. After a clamp time of 40 seconds, aortic flow was restored. When hemodynamic stability was achieved, anatomic closure was performed; the chest was evacuated of residual air using a 16-gauge angiocatheter attached to a syringe; and the animal was extubated when able to breathe spontaneously. The operative mortality associated with the gene delivery procedure was ~10%. In the different treatment groups, mortality after surgery was ~20%, with the majority of deaths occurring around postoperative day 2. There were no deaths after postoperative day 7.

Histology. Hearts were excised, rinsed in PBS, frozen in embedding compound, and stored at -80°C . Specimens were mounted on a freezing microtome, and 10- μm sections were transferred to glass slides pretreated with aminoalkylsilane (Sigma Chemical Co., St. Louis, Missouri, USA). Sections were fixed in 10% formalin for 2 minutes at room temperature and washed in PBS. β -Galactosidase staining was carried out in 2 mmol/L $\text{K}_4\text{Fe}(\text{CN})_6$, 2 mmol/L $\text{K}_3\text{Fe}(\text{CN})_6$, 2 mmol/L MgCl_2 , and 0.5 mg/mL X-Gal (5-bromo-4-chloro-3-indoyl- β -D-galactopyranoside) in PBS (pH 7.4) as

described (18). After being stained (30–60 minutes at 37°C), the sections were rinsed in PBS solution and counterstained with eosin. Additional sections were prepared in the same manner and stained with hematoxylin and eosin (H&E) for assessment of myocardial cellular infiltrates and inflammation.

β_2 -AR immunohistochemistry. Frozen myocardial sections were cut at 10 μ m for indirect immunofluorescence studies. Sections were rinsed in PBS and then in PBS with 0.05% Triton X-100 (Triton-PBS), blocked with serum diluent (10% goat serum in PBS with 0.1% BSA and 0.1% sodium azide), and then rinsed for 15 minutes in Triton-PBS before overnight incubation at 4°C with a primary rabbit antihuman β_2 -AR antiserum (1:500 dilution in serum diluent) as we have described (18). The sections were then washed 4 times for 10 minutes in Triton-PBS at room temperature and incubated for 1 hour in FITC-conjugated goat anti-rabbit IgG (1:50 dilution in serum diluent). After rinsing with PBS, the sections were mounted and photographed (18).

β -AR radioligand binding. Myocardial membranes were prepared by homogenization of excised hearts in ice-cold lysis buffer (5 mM Tris-HCl [pH 7.4], 5 mM EDTA) as described previously (18, 19). Final purified cardiac membranes were suspended at a concentration of 1–2 mg/mL in ice-cold β -AR binding buffer (75 mM Tris-HCl [pH 7.4], 12.5 mM MgCl₂, and 2 mM EDTA),

and receptor binding was performed using the nonselective β -AR ligand [¹²⁵I]cyanopindolol (18, 19). Non-specific binding was determined in the presence of 20 μ M alprenolol. All assays were performed in triplicate, and receptor density (measured in femtomoles) was normalized to milligrams of membrane protein.

Membrane adenylyl cyclase activity. Myocardial membranes were prepared as already described here and incubated (20–30 μ g of protein) for 15 minutes at 37°C with [α -³²P]ATP under basal conditions or in the presence of either 100 μ M isoproterenol or 10 mM NaF, and cAMP was quantitated by standard methods that have been described (19).

In vivo hemodynamic measurements. To evaluate in vivo physiology and contractile function, all animals were sedated with ketamine (80 mg/kg) and acepromazine (0.1 mg/kg). After local infiltration of 1% lidocaine, venous access was obtained by way of the external jugular vein, and a 2.5 French micromanometer was placed into the LV through the carotid artery. LV pressure was then obtained using custom data acquisition software at a sampling rate of 200 Hz (Physiologic Data Systems Inc., Durham, North Carolina, USA). Data were acquired at baseline and after infusion of isoproterenol at 0.1, 0.5, and 1.0 μ g/kg/min as we have described (20). The LV pressure data were then analyzed on a VAX/Microsystem II (Digital Equipment Corp.,




Figure 2

β -AR overexpression in rabbit hearts after intracoronary delivery of Adeno- β_2 AR. (a) Total β -AR density in different chambers of rabbit myocardium 6 days after 1×10^{12} tvp of Adeno- β_2 AR was delivered via intracoronary perfusion. β -AR density after Adeno- β_2 AR treatment is compared with rabbit hearts that received the same dose of the EV. (b) Representative immunohistochemical detection of expressed human β_2 -ARs in rabbit LV 6 days after intracoronary delivery of Adeno- β_2 AR. (c) Dose-dependent myocardial β_2 -AR overexpression after delivery of increasing doses of Adeno- β_2 AR. Total β -AR density is shown in the RV and LV compared with myocardial β -AR density after delivery of EV, which did not change from endogenous β -AR levels. (d) Time course of β -AR overexpression in rabbit hearts after delivery of 5×10^{11} tvp Adeno- β_2 AR. All binding values represent the mean \pm SEM.

Table 1In vivo hemodynamic measurements in rabbits with 5×10^{11} tvp of *Adeno- β_2 AR* compared with saline-treated controls

	Saline (n = 11)				<i>Adeno-β_2AR</i> (n = 11)			
	Basal	ISO 0.1	ISO 0.5	ISO 1.0	Basal	ISO 0.1	ISO 0.5	ISO 1.0
LV dP/dt _{max} (mmHg/s)	2,515 ± 182	4,085 ± 132	5,335 ± 134	5,589 ± 112	2,988 ± 185 ^A	4,584 ± 173 ^A	6,056 ± 145 ^B	5,841 ± 192
LV dP/dt _{min} (mmHg/s)	-2,203 ± 67	-1,948 ± 479	-2,557 ± 80	-2,635 ± 96	-2,302 ± 127	-2,758 ± 138 ^A	-3,119 ± 171 ^B	-2840 ± 123
MEP (mmHg)	51.9 ± 1.3	52.2 ± 1.8	54.8 ± 1.8	56.7 ± 1.9	53.1 ± 2.8	54.5 ± 2.6	62.4 ± 1.5 ^B	55.7 ± 2.4
LVSP (mmHg)	58 ± 1.3	60 ± 1.8	65 ± 1.7	66 ± 1.6	61.3 ± 2.2	65.2 ± 2.4 ^A	73.3 ± 2.4 ^B	70.4 ± 3.6
HR (bpm)	203 ± 9	268 ± 4.9	291 ± 4.7	293 ± 4.7	241 ± 8 ^B	285 ± 9	307 ± 8.2	328 ± 9.9 ^B

ISO, isoproterenol; MEP, left ventricular mean ejection pressure; LVSP, left ventricular systolic pressure; HR, heart rate. *P* value determined by Student's *t*-test. ^A*P* < 0.05 vs. saline. ^B*P* < 0.005 vs. saline.


Woburn, Massachusetts, USA) with custom software to derive the maximal and minimal first derivative of the pressure rise (LV dP/dt_{max} and LV dP/dt_{min}, respectively), as well as heart rate (HR), peak systolic pressure (SP), and mean ejection pressure. To determine the effect of HR on ventricular function, 6 animals underwent placement of epicardial pacing wires on the right atrium at the time of virus injection.

Myocyte studies. To determine whether β_2 -AR overexpression might benefit the failing myocyte, we carried out a study in cultured ventricular myocytes isolated from normal rabbits or rabbits in HF induced by left circumflex coronary artery ligation (21). This model of HF in the rabbit has recently been described by our laboratory to recapitulate several features of the human disease (21). Myocytes were isolated from rabbit hearts as described by us using a Langendorff perfusion technique (10, 17). To assess the effect of β_2 -AR overexpression on intracellular signaling, cultured myocytes were plated at a density of 1×10^5 per 35-mm well that were precoated with 20 μ g/mL of mouse laminin for 1 hour. After 5 hours, the myocytes adhered to the tissue culture plates, and cells were infected with an moi of 300:1 of the adenovirus in 0.5 mL of solution (10, 17). The cells were incubated with the virus for 15 minutes, after which culture media were added back to the plates. Typically, infection of rabbit ventricular myocytes with *Adeno- β_2 AR* results in β -AR levels > 1 pmol/mg membrane protein (10, 17). Thirty-six hours later, myocytes were labeled overnight in 1.5 μ Ci/mL [³H]adenine in Medium 199 and then preincubated in MEM with 10 mM HEPES and 1 mM 3-isobutyl-1-methylxanthine (IBMX) for 30 minutes. For dose-response studies, different concentrations of isoproterenol were added for 15 minutes, and intracellular cAMP was determined by standard chromatography methods as described (10, 17).

Statistical analysis. Quantitative data, such as myocardial β -AR density after adenoviral transgene delivery and physiological parameters, are expressed as the mean \pm SEM. The difference in the level of myocardial transgene expression and cardiac function between control and adenovirus-treated rabbits was evaluated using a Student's *t* test. For physiological and myocyte isoproterenol responses, differences between groups were analyzed using single-factor ANOVA. For all analyses, *P* < 0.05 was considered statistically significant.

Results

In vivo myocardial adenoviral transgene delivery. Intracoronary delivery is the most efficient way to deliver adenoviral transgenes globally to the myocardium. To achieve this, we developed a surgical technique in rabbits, whereby a catheter is placed into the LV chamber through the apex of the heart by way of a right thoracotomy. The adenovirus solution in PBS (1.5 mL) is rapidly injected while the aorta is cross-clamped for 40 seconds. This enables all coronary beds to be perfused by the adenovirus. During clamping, the heart visibly blanches, demonstrating a washout of blood and replacement by the viral solution. Before adenovirus delivery, adenosine (0.75 mg/kg) is injected, which serves to slow the heart. To assess myocardial expression of adenoviral transgenes, we delivered either an adenovirus containing a cytoplasmic β -galactosidase marker gene (*Adeno- β Gal*), the cDNA for the human β_2 -AR (*Adeno- β_2 AR*), or no transgene (empty adenovirus [EV]). As shown in Figure 1a, 6 days after *Adeno- β Gal* delivery (1×10^{12} total viral particles [tvp]), glob-

**Figure 3**

Membrane adenylyl cyclase activity from hearts of *Adeno- β_2 AR*-treated rabbits. Membranes were purified from rabbits 6 days after intracoronary delivery of 5×10^{11} tvp of *Adeno- β_2 AR* or EV (*n* = 6 each), and adenylyl cyclase activity was measured under basal conditions and after isoproterenol (10^{-4} M) or NaF (10 mM). Values shown are the mean \pm SEM normalized to the value seen with NaF. Activity in the 2 groups with NaF was not significantly different (data not shown), indicating no postreceptor defects in signaling. **P* < 0.05 vs. EV (Student's *t* test).

al myocardial β -Gal expression is evident by the robust X-Gal staining of both the LV and RV as well as throughout the septum. Although there is an abundance of epicardial expression, transmural penetration of the β -Gal transgene is clearly evident. With higher magnification, *Adeno- β Gal*-treated hearts reveal that the expression is apparent in individual myocytes (Figure 1b). Control sections of hearts treated with equivalent doses of *Adeno- β_2 AR* or EV revealed no specific X-Gal staining (Figure 1c).

Adenoviral-mediated myocardial β_2 -AR overexpression. After the demonstration of in vivo global myocardial transgene delivery with *Adeno- β Gal*, we assessed the delivery of the therapeutic transgene, *Adeno- β_2 AR*. Similar to *Adeno- β Gal* delivery, β_2 -AR overexpression, determined by radioligand binding 1 week after *Adeno- β_2 AR* intracoronary administration, is global in nature, as demonstrated by robust overexpression in all cardiac chambers – including the septum (Figure 2a). Diffuse β_2 -AR overexpression is evident throughout the heart in the sarcolemmal membranes of individual myocytes, as demonstrated by immunohistochemical staining (Figure 2b). We also found measurable β -AR overexpression in other tissues, but appreciable levels were limited to the liver and lungs (data not shown), consistent with previous studies (14, 16).

To determine whether β_2 -AR overexpression depends on the adenoviral dose injected into the rabbit heart, we delivered varying doses of *Adeno- β_2 AR*. As shown in Figure 2c, β_2 -AR overexpression in the ventricles shows a dose dependence, as assessed at 1 week after in vivo gene delivery of *Adeno- β_2 AR*. Normal, endogenous β -AR density in rabbit myocardium is ~ 75 fmol/mg membrane protein, which was not altered by delivery of the EV (Figure 2c). Significant β_2 -AR overexpression was seen beginning with 5×10^{11} tvp of *Adeno- β_2 AR*, and expression approached 1 pmol/mg membrane protein with 20×10^{11} tvp (Figure 2c). However, there was not a significant increase in β_2 -AR overexpression between 10×10^{11} and 20×10^{11} tvp of *Adeno- β_2 AR*. Furthermore, in preliminary physiological experiments, β_2 -AR overexpression above 800 fmol/mg membrane protein produced diminished cardiac function, which is probably the result of enhanced inflammation observed with higher doses of adenovirus (see later here). Thus, 5×10^{11} tvp of *Adeno- β_2 AR* was chosen as the “therapeutic dose.” Injection of this amount of virus reproducibly results in 5- to 10-fold global myocardial β_2 -AR overexpression.

A property of adenovirus gene transfer in vivo is the loss of transgene expression, due to several factors inherent in the virus construct itself or in the host animal. To test how long β_2 -AR transgene overexpression is supported in the heart after in vivo intracoronary delivery, we injected 5×10^{11} tvp of *Adeno- β_2 AR* and measured β -AR density at varying time points. As shown in Figure 2d, β_2 -AR overexpression peaked ~ 1 week after gene delivery but was still significantly increased at 12 days. Three weeks after gene delivery, β_2 -




Figure 4

In vivo assessment of LV contractile function in rabbits treated with *Adeno- β_2 AR*. Cardiac catheterization was performed in conscious sedated rabbits 6 days after intracoronary delivery (1.5 mL in PBS) of either 5×10^{11} tvp *Adeno- β_2 AR* ($n = 11$), EV ($n = 7$), or saline ($n = 11$) using a high-fidelity micromanometer. Parameters measured include (a) LV dP/dt_{max} and (b) LVSP. Shown are the baseline values (as mean \pm SEM) and responses to progressive doses of isoproterenol. * $P < 0.05$; # $P < 0.005$ vs. saline (Student's t test). For LV dP/dt_{max} and LVSP, isoproterenol responses in the *Adeno- β_2 AR* group are significantly different than those in the EV-treated group ($P < 0.01$, ANOVA).

AR overexpression is minimal, which is consistent with the duration of in vivo expression seen by others (14). However, significant overexpression at ~ 2 weeks is quite a bit longer than the duration of expression observed in a rat heart transplant model of ex vivo *Adeno- β_2 AR* gene transfer (18). Because the major reason for loss of adenoviral transgene expression in vivo is believed to be host immunological responses due to the virus (22, 23), we assessed the degree of inflammation in our hearts histologically. Six days after 5×10^{11} tvp *Adeno- β_2 AR* or EV delivery by coronary perfusion, there is minor inflammation signified by cellular infil-




Figure 5
Relationship between myocardial β_2 -AR overexpression and in vivo contractility. Total cardiac β -AR density was plotted against maximal isoproterenol-stimulated LV $+dP/dt_{max}$ of rabbits treated with *Adeno- β_2 AR*. A significant linear relationship was found (*t* test).

tration in the myocardium, as assessed by H&E staining, compared with saline-injected control rabbits (data not shown); this finding is consistent with previous studies using adenovirus (12, 14).

Effect of β_2 -AR overexpression on myocardial signaling and function. To assess whether the β_2 -ARs expressed in the myocardium after in vivo intracoronary delivery were functionally coupled, we examined adenylyl cyclase activity in membranes prepared from rabbit hearts injected with either the EV or *Adeno- β_2 AR*. Basal adenylyl cyclase activity was measured, as well as activity in response to the β -agonist isoproterenol. As shown in Figure 3, both basal and isoproterenol-stimulated adenylyl cyclase activities were significantly enhanced in hearts overexpressing β_2 -ARs compared with myocardial membranes prepared from EV-treated rabbits.

The ultimate goal of this study was to test whether myocardial β_2 -AR overexpression after in vivo adenovirus delivery by coronary perfusion could alter global heart function. To do this, we evaluated in vivo hemodynamic parameters of cardiac function by way of catheterization of sedated but conscious rabbits. We initially assessed cardiac physiology 6 days after adenovirus administration, which was the time of peak expression (Figure 2d). Rabbits were either injected with saline, EV (5×10^{11} tvp), or *Adeno- β_2 AR* (5×10^{11} tvp) as already described here. Measurements were recorded at baseline and after infusion of 0.1, 0.5, and 1.0 $\mu\text{g}/\text{kg}/\text{min}$ of isoproterenol. Physiological measurements of the saline-injected and *Adeno- β_2 AR*-treated rabbits are listed in Table 1. Values for LV $+dP/dt_{max}$ and LV $-dP/dt_{min}$ as measurements of cardiac contractility and relaxation, respectively, are shown, as are values for the LV mean ejection pressure (MEP), LV systolic pressure (LVSP), and HR. β_2 -AR overexpression as a result of the intracoro-

nary delivery of *Adeno- β_2 AR* leads to significant enhancement of several key hemodynamic measurements (Table 1). At baseline, there was a significant improvement in baseline LV $+dP/dt_{max}$ in the *Adeno- β_2 AR*-treated rabbits that was accompanied by a significant increase in HR (Table 1). After administration of 0.1 and 0.5 $\mu\text{g}/\text{kg}/\text{min}$ isoproterenol, HR was not significantly increased in *Adeno- β_2 AR*-treated rabbits; however, LV $+dP/dt_{max}$ and LV SP in β_2 -AR-overexpressing rabbits was significantly enhanced compared with saline-injected controls. LV contractility responses in *Adeno- β_2 AR*-, saline-, and EV-treated rabbits are shown graphically in Figure 4a. Figure 4b shows LVSP responses in the same groups. The isoproterenol dose-responses for both of these parameters were significantly enhanced in the *Adeno- β_2 AR* group compared with EV-treated rabbits. EV treatment was not significantly different than saline; however, values tended to be lower, as seen in Figure 4, a and b. This could be due to the small level of inflammation that was observed histologically after adenovirus administration. After physiological assessment, all animals were sacrificed to measure their level of β_2 -AR overexpression. In animals that received 5×10^{11} tvp *Adeno- β_2 AR*, receptor density ranged from 400 to 700 fmol/mg membrane protein. Within this window of overexpression, there was a significant direct linear relationship between maximum isoproterenol-induced LV $+dP/dt_{max}$ and β -AR density (Figure 5). This demonstrates that the enhanced cardiac function seen in vivo in these rabbits is the direct result of β_2 -AR overexpression.

In an additional set of rabbits, we delivered 5×10^{11} tvp *Adeno- β_2 AR* ($n = 9$) or EV ($n = 5$) and assessed in vivo hemodynamics 21 days after gene delivery. Although baseline LV $+dP/dt_{max}$ was not elevated, responses to




Figure 6
In vivo assessment of LV contractile function (LV $+dP/dt_{max}$) in rabbits 21 days after treatment with *Adeno- β_2 AR*. Cardiac catheterization was performed in conscious sedated rabbits after intracoronary delivery (1.5 mL in PBS) of either 5×10^{11} tvp *Adeno- β_2 AR* ($n = 9$) or EV ($n = 5$) using a high-fidelity micromanometer. Shown are the baseline values (mean \pm SEM) and responses to progressive doses of isoproterenol. $*P < 0.05$ vs. EV (Student's *t* test). Total β -AR density after *Adeno- β_2 AR* treatment was 115 ± 12 fmol/mg membrane protein vs. 79 ± 7 in hearts that received the same dose of EV ($P < 0.05$, Student's *t* test).

isoproterenol were significantly enhanced in the *Adeno-β₂AR*-treated animals compared with the control rabbits that received EV (Figure 6). Interestingly, the levels of β₂-AR overexpression seen in these rabbit hearts were minimal compared with what was seen in the animals tested 1 week after gene delivery; however, β-AR density was still significantly higher than endogenous levels in the hearts of EV-treated rabbits (115 ± 12 vs. 79 ± 7 fmol/mg membrane protein; *P* < 0.05).

The results of the physiology measurements indicate that β₂-AR overexpression does positively effect global in vivo cardiac performance. However, as HR can affect certain physiological parameters, especially LV +dP/dt_{max}, we studied 3 additional saline- and *Adeno-β₂AR*-treated rabbits that had atrial pacing wires surgically implanted at the time of gene delivery. LV +dP/dt_{max} was then assessed in these animals after 6 days, as already described here, at differently paced HRs; importantly, no significant change in LV +dP/dt_{max} was seen with HRs between 200 and 300 bpm in *Adeno-β₂AR*-treated rabbits.

β₂-AR overexpression in failing rabbit myocytes. These results suggest that replacing lost β-ARs in HF may be a novel therapeutic strategy to treat the failing heart. To examine whether β₂-AR overexpression in failing myocytes might enhance β-AR signaling, we carried out studies using *Adeno-β₂AR* treatment in failing rabbit myocytes. These failing myocytes were isolated from rabbits 21 days after myocardial infarction induced by left circumflex coronary artery ligation (21). In this model of rabbit HF, several features of human HF are recapitulated, including abnormal β-AR signaling (21). Myocytes isolated from infarcted rabbits have abnormal cAMP accumulation in response to β-AR stimulation (21). In Figure 7, we demonstrate that EV-treated ventricular myocytes from infarcted rabbits have attenuated cAMP responses compared with EV-treated myocytes from control (sham-operated) rabbits. However, *Adeno-β₂AR* treatment of myocytes isolated from infarcted rabbits enhances β-AR signaling to levels even greater than control, demonstrating a β-AR signaling rescue in these failing myocytes (Figure 7).

Discussion

In this study, we present the novel findings that acute global myocardial overexpression of the human β₂-AR in rabbits can lead to enhancement of cardiac physiology assessed by conscious in vivo hemodynamics after intracoronary administration of recombinant adenovirus. We have developed a technique for global myocardial transgene delivery in a larger animal species that provides an important in vivo model to study the functional significance of acute transgenesis of the heart. Given the well-characterized deficiencies in myocardial β-AR signaling in chronic human HF (2, 3, 6), our present findings suggest that gene transfer to replace lost β-ARs in failing hearts may have potential for producing therapeutic cardiac inotropy.

Our model of in vivo myocardial gene transfer demonstrates efficient and reproducible global transgene delivery to all chambers of the rabbit heart. This method was developed to ensure that all coronary beds are perfused by the adenoviral solution under pressure so that the myocardium may be globally infected. The method of in vivo global myocardial gene delivery described here has advantages over methods of direct coronary artery catheterization and injection. Percutaneous coronary artery delivery involves administering the virus to a single coronary artery, and transgene expression is limited to the area of the heart supplied by that coronary (14, 15). In contrast, by injection of the adenovirus into the LV cavity while the aorta is cross-clamped, we ensure that all coronary vessels are perfused, which results in transgene expression globally in the heart. While we were completing this study, a similar technique was used in rats, whereby adenovirus was injected into the LV cavity while briefly clamping both the aorta and pulmonary artery (16). However, our rabbit model is unique, as we do not clamp the pulmonary artery and aortic clamp time is considerably longer, which probably contributes to enhanced viral uptake. In the rat study, myocardial overexpression of a phospholamban transgene decreased certain hemodynamic parameters and, thus, did not represent a potential therapy (16). Overexpression of the phospholamban transgene was modest (2- to 3-fold) compared with the 5- to 10-fold overexpression of the β₂-AR transgene in the present




Figure 7

Rescue of attenuated signaling in failing ventricular myocytes by treatment with *Adeno-β₂AR*. Shown is the isoproterenol-induced cAMP production in sham and failing isolated ventricular cardiomyocytes treated with EV or *Adeno-β₂AR*. Failing ventricular myocytes were isolated from rabbit hearts 21 days after left circumflex coronary artery ligation (21). Experiments were performed 36 hours after adenoviral infection and after overnight labeling with 1.5 μCi/mL [³H]adenine (see Methods). The accumulation of intracellular cAMP is expressed as percent conversion from total ³H uptake. The data represent the mean ± SEM of 4–9 independent experiments, each performed in triplicate. Total responses of the failing/EV group were significantly attenuated compared with sham/EV-treated myocytes (*P* < 0.05, single-factor ANOVA). Total responses of the failing *Adeno-β₂AR*-treated myocytes were significantly enhanced compared with the failing/EV-treated cells (**P* < 0.05, ANOVA).

study, which could possibly be due to the prolonged aortic clamp time and higher adenoviral doses used. In addition, rabbits have slower HRs than rats; these already slow HRs were further reduced by adenosine administration, allowing more adenovirus contact time with the myocardium. Thus, all of these factors probably contribute to the robust global myocardial transgene expression seen in this rabbit model of intracoronary gene delivery, making it possible to examine the potential usefulness of this approach to enhance therapeutically the function of the heart in vivo.

The demonstration that overexpression of β_2 -ARs between 400 and 700 fmol/mg membrane protein enhances adenylyl cyclase activity in vitro, and several hemodynamic parameters in vivo, is of significance. In this study, both baseline and isoproterenol-stimulated $LV +dP/dt_{max}$ were enhanced after β_2 -AR overexpression compared with saline- and EV-treated control rabbits, which were not influenced by HR. Moreover, there was a significant direct relationship between β_2 -AR overexpression and contractility at 6 days after gene delivery. Significant enhancement of isoproterenol-stimulated in vivo cardiac function after *Adeno- β_2 AR* treatment was also seen 21 days after gene delivery when compared with EV-treated rabbits, even though β_2 -AR overexpression was drastically reduced compared with overexpression seen at 1 week. This minimal overexpression at 3 weeks was statistically significant compared with endogenous β -AR density, and although this level of overexpression no longer resulted in enhanced basal $LV +dP/dt_{max}$ compared with EV-treated rabbits, β -AR responsiveness was still significantly elevated. Thus, prolonged β_2 -AR overexpression appears to be responsible for maintained elevated in vivo contractile responses at 3 weeks.

Interestingly, there appears to be a “therapeutic window” of β_2 -AR overexpression, as levels >800 fmol/mg membrane protein, produced by increasing the viral dose, resulted in a decline of in vivo function (data not shown). This could be due to viral toxicity and immunological responses of the host animal at these higher doses. Attenuated function was seen when higher ($>10^{12}$ tvp) doses of EV were also administered (data not shown), indicating that it is a general feature of adenovirus delivery and not specifically due to higher levels of β_2 -AR overexpression.

Our study demonstrates that global myocardial expression is possible in vivo through intracoronary adenovirus delivery. Previous ex vivo models in the heart have demonstrated that intracoronary infusions of adenovirus can lead to global transgene expression (18, 24, 25). We have used a heterotopic rat heart transplant model to deliver *Adeno- β_2 AR* to the transplanted rat heart, and this has led to both basal and β -agonist-mediated increases in cardiac contractility measured in the isolated heart by Langendorff perfusion (25). Thus, β_2 -AR overexpression in both the transplanted rat heart and the rabbit heart in vivo can enhance global myocardial function.

These adenoviral-mediated gene transfer experiments were done to determine whether the hearts of larger animals could be manipulated similarly to those of transgenic mice. Previous work by our laboratory demonstrated that myocardial-targeted overexpression of the human β_2 -AR transgene in transgenic mice at extraordinary levels (>200 -fold over endogenous levels) led to dramatic increases in cardiac function (8). It is obvious that we will not approach this level of overexpression of β_2 -ARs using adenoviruses; however, Liggett and colleagues have generated additional transgenic mice overexpressing β_2 -ARs in the heart at levels in the range of the overexpression seen in our adenoviral-treated rabbit hearts, and cardiac in vivo contractility was still enhanced (9). Thus, our results in the rabbit demonstrate that ~ 10 -fold overexpression of myocardial β_2 -ARs can lead to enhanced contractility in vivo.

These studies demonstrate that β_2 -AR overexpression in the context of normal myocardium can lead to increases in cardiac contractility; however, potential clinical use would be in a heart in which function is compromised. Multiple lines of evidence, including experiments presented in this study, suggest that β_2 -AR overexpression in the failing heart might be of clinical utility. We have shown that *Adeno- β_2 AR* treatment of cardiac myocytes isolated and cultured from post-myocardial infarction rabbits that are in congestive HF can restore dysfunctional β -AR signaling, as measured by intracellular cAMP accumulation (Figure 7). Thus, alterations in the β -AR system in the failing heart can be restored after acute adenoviral-mediated gene transfer, demonstrating the feasibility of genetically replacing lost β -ARs in the failing heart.

In addition, hybrid transgenic mouse studies have shown that modest (30-fold) myocardial overexpression of β_2 -ARs can rescue signaling, hypertrophy, and baseline ventricular function in a $G\alpha_q$ overexpression model of hypertrophic dysfunction (26). More importantly, data supporting the potential clinical significance of β_2 -AR gene therapy for HF include the recent findings that HF patients possessing a β_2 -AR gene polymorphism resulting in a β_2 -AR that couples to adenylyl cyclase less efficiently have significantly lower survival rates (27). Thus, gene therapy using the wild-type β_2 -AR, which couples efficiently to adenylyl cyclase, may be potentially useful in human HF especially in this population of patients. Furthermore, 2 recent studies of patients with severe HF have demonstrated that there is contractile reserve in failing myocardium. It has been shown that failing myocardium retains a certain capacity to recover functionally; it also undergoes “reverse remodeling” and regression of hypertrophy after use of a LV mechanical assistance device (28, 29). These data suggest that it might be possible to treat failing myocardium by replacing lost receptors and that failing myocytes will support enhanced adrenergic signaling through β_2 -ARs.

Efficient gene transfer in vivo to the rabbit heart, as described here, now makes it possible to study potential molecular therapies, as there are reliable in vivo HF models available. It has yet to be determined whether our method of in vivo transgene delivery will rescue the failing rabbit heart, but this will be a focus of future studies. Nevertheless, our current results demonstrate the feasibility of acute in vivo global myocardial transgenesis in investigating important molecular targets for the treatment of cardiovascular disease.

Acknowledgments

The authors thank K. Shotwell, K. Campbell, and A. Phippen for technical assistance, and A. Kypson and R.E. Lilly for critical contributions throughout these studies. We also thank the Genzyme Corporation (Framingham, Massachusetts, USA) for preparation and purification of Adeno- β_2AR . This work was supported in part by National Institutes of Health grants HL-16037 (to R.J. Lefkowitz), HL-59533 (to W.J. Koch), and HL-56205 (to W.J. Koch), and a Grant-in-Aid from the American Heart Association (to W.J. Koch).

- Cohn, J.N., et al. 1997. Report of the National Heart, Lung, and Blood Institute Special Emphasis Panel on Heart Failure Research. *Circulation*. **95**:766–770.
- Brodde, O.E. 1993. Beta-adrenoceptors in cardiac disease. *Pharmacol. Ther.* **60**:405–430.
- Bristow, M.R., et al. 1982. Decreased catecholamine sensitivity and β -adrenergic receptor density in failing human hearts. *N. Engl. J. Med.* **307**:205–211.
- Ping, P., Anzai, T., Gao, M., and Hammond, H.K. 1997. Adenylyl cyclase and G protein receptor kinase expression during development of heart failure. *Am. J. Physiol.* **273**:H707–H717.
- Bristow, M.R., et al. 1993. Reduced β_1 receptor messenger RNA abundance in the failing human heart. *J. Clin. Invest.* **92**:2737–2745.
- Ungerer, M., Bohm, M., Elce, J.S., Erdmann, E., and Lohse, M.L. 1993. Altered expression of β -adrenergic receptor kinase and β_1 -adrenergic receptors in the failing heart. *Circulation*. **87**:454–463.
- Bristow, M.R., and Lowes, B.D. 1994. Low-dose inotropic therapy for ambulatory heart failure. *Coron. Artery Dis.* **5**:112–118.
- Milano, C.A., et al. 1994. Enhanced myocardial function in transgenic mice overexpressing the β_2 -adrenergic receptor. *Science*. **264**:582–586.
- Turki, J., et al. 1996. Myocardial signaling defects and impaired cardiac function of a human β_2 -adrenergic receptor polymorphism expressed in transgenic mice. *Proc. Natl. Acad. Sci. USA*. **93**:10483–10488.
- Akhter, S.A., et al. 1997. Restoration of β -adrenergic signaling in failing cardiac ventricular myocytes via adenoviral-mediated gene transfer. *Proc. Natl. Acad. Sci. USA*. **94**:12100–12105.
- Kass-Eisler, A., et al. 1993. Quantitative determination of adenovirus-mediated gene delivery to rat cardiac myocytes in vitro and in vivo. *Proc. Natl. Acad. Sci. USA*. **90**:11498–11502.
- French, B.A., Mazur, W., Geske, R.S., and Bolli, R. 1994. Direct in vivo gene transfer into porcine myocardium using replication-deficient adenoviral vectors. *Circulation*. **90**:2414–2424.
- Magovern, C.J., et al. 1996. Direct in vivo gene transfer to canine myocardium using a replication-deficient adenovirus vector. *Ann. Thorac. Surg.* **62**:425–433.
- Barr, E., et al. 1994. Efficient catheter-mediated gene transfer into the heart using replication-defective adenovirus. *Gene Ther.* **1**:51–58.
- Giordano, F.J., et al. 1996. Intracoronary gene transfer of fibroblast growth factor-5 increases blood flow and contractile function in an ischemic region of the heart. *Nat. Med.* **2**:534–539.
- Hajjar, R.J., et al. 1998. Modulation of ventricular function through gene transfer in vivo. *Proc. Natl. Acad. Sci. USA*. **95**:5251–5256.
- Drazner, M.H., et al. 1997. Potentiation of β -adrenergic signaling by adenoviral-mediated gene transfer in adult rabbit ventricular myocytes. *J. Clin. Invest.* **99**:288–296.
- Kypson, A.P., et al. 1998. Ex vivo adenoviral-mediated gene transfer to the transplanted adult rat heart. *J. Thorac. Cardiovasc. Surg.* **115**:623–630.
- Koch, W.J., et al. 1995. Cardiac function in mice overexpressing the β -adrenergic receptor kinase or a β ARK inhibitor. *Science*. **268**:1350–1353.
- Silvestry, S.C., et al. 1996. The in vivo quantification of myocardial performance in rabbits: a model for evaluation of cardiac gene therapy. *J. Mol. Cell. Cardiol.* **28**:815–823.
- Maurice, J.P., et al. 1999. Molecular β -adrenergic signaling abnormalities in failing rabbit hearts after infarction. *Am. J. Physiol.* **276**:H1853–1860.
- Engelhardt, J.F., Ye, X., Doranz, B., and Wilson, J.M. 1994. Ablation of E2A in recombinant adenoviruses improves transgene persistence and decreases inflammatory response in mouse liver. *Proc. Natl. Acad. Sci. USA*. **91**:6196–6200.
- Yang, Y., Jooss, K.U., Su, Q., Ertl, H.C.J., and Wilson, J.M. 1996. Immune responses to viral antigens versus transgene product in the elimination of recombinant adenovirus-infected hepatocytes in vivo. *Gene Ther.* **3**:137–144.
- Donahue, J.K., Kikkawa, K., Johns, D.C., Marban, E., and Lawrence, J.H. 1997. Ultrarapid, highly efficient viral gene transfer to the heart. *Proc. Natl. Acad. Sci. USA*. **94**:4664–4668.
- Kypson, A.P., et al. 1999. Adenoviral-mediated gene transfer of the β_2 -adrenergic receptor to donor hearts enhances cardiac function. *Gene Ther.* In press.
- Dorn, G.W., et al. 1999. Low- and high-transgenic expression of β_2 -adrenergic receptors differentially affects cardiac hypertrophy and function in $G\alpha_q$ overexpressing mice. *Proc. Natl. Acad. Sci. USA*. **96**:6400–6405.
- Liggett, S.B., et al. 1998. The ile164 β_2 -adrenergic receptor polymorphism adversely affects the outcome of congestive heart failure. *J. Clin. Invest.* **102**:1534–1539.
- Dipla, K., Mattiello, J.A., Jeevanandam, V., Houser, S.R., and Margulies, K.B. 1998. Myocyte recovery after mechanical circulatory support in humans with end-stage heart failure. *Circulation*. **97**:2316–2322.
- Zafeiridis, A., Jeevanandam, V., Houser, S.R., and Margulies, K.B. 1998. Regression of cellular hypertrophy after left ventricular assist device support. *Circulation*. **98**:656–662.