

Entomotoxic and nematotoxic lectins and protease inhibitors from fungal fruiting bodies

Jerica Sabotič¹ · Robin A. Ohm² · Markus Künzler³

Received: 24 July 2015 / Revised: 4 October 2015 / Accepted: 11 October 2015 / Published online: 31 October 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract Fruiting bodies or sporocarps of dikaryotic (ascomycetous and basidiomycetous) fungi, commonly referred to as mushrooms, are often rich in entomotoxic and nematotoxic proteins that include lectins and protease inhibitors. These protein toxins are thought to act as effectors of an innate defense system of mushrooms against animal predators including fungivorous insects and nematodes. In this review, we summarize current knowledge about the structures, target molecules, and regulation of the biosynthesis of the best characterized representatives of these fungal defense proteins, including galectins, beta-trefoil-type lectins, actinoporin-type lectins, beta-propeller-type lectins and beta-trefoil-type chimerolectins, as well as mycospin and mycocypin families of protease inhibitors. We also present an overview of the phylogenetic distribution of these proteins among a selection of fungal genomes and draw some conclusions about their evolution and physiological function. Finally, we present an outlook for future research directions in this field and their potential applications in medicine and crop protection.

Keywords Basidiomycete · Ascomycete · Sporocarp · Glycan · Insect · Nematode

Introduction

Fruiting bodies, also referred to as mushrooms or sporocarps, are important microscopic or macroscopic structures formed during the sexual reproduction cycle of the fungal subkingdom Dikarya (Hibbett et al. 2007; Taylor and Ellison 2010). These fungi are characterized by the formation of dikaryotic hyphae and include the phyla *Ascomycota* and *Basidiomycota*. Fruiting body formation is independent of the lifestyle (saprotrophic or biotrophic) of these fungi and sometimes coupled to the preceding formation of sclerotia—compact masses of hardened mycelium containing food reserves (Martin et al. 2008; Stajich et al. 2010; Teichert et al. 2014; Yin et al. 2012). Since the fruiting bodies produce and disperse the sexual spores, defense of these structures against fungivores, including predators, grazers, and parasites, is essential for fungal reproduction. Thus, dikaryotic fungi employ, in addition to a large repertoire of secondary metabolites, a plethora of proteins acting as deterrents or toxins in defense of their fruiting bodies (Spiteller 2015; Wang et al. 2002). These proteins include lectins that bind to the glycans of glycoproteins or glycolipids in the digestive tract of fungivores (Bleuler-Martinez et al. 2011), protease inhibitors that inhibit digestive proteases of fungivores (Renko et al. 2010), biotin-binding proteins that sequester this essential cofactor (Bleuler-Martinez et al. 2012), pore-forming proteins that cause cell lysis (Mancheno et al. 2010; Ota et al. 2014), RNA toxins (ribotoxins) that cleave or depurinate RNA molecules (Lacadena et al. 2007), and other enzymes including proteases, oxidases, and phospholipases (Erjavec et al. 2012). The first two groups of proteins: lectins and protease inhibitors,

Electronic supplementary material The online version of this article (doi:10.1007/s00253-015-7075-2) contains supplementary material, which is available to authorized users.

✉ Jerica Sabotič
jerica.sabotic@ijs.si

¹ Department of Biotechnology, Jožef Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia

² Department of Microbiology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

³ Institute of Microbiology, Department of Biology, ETH Zürich, Vladimir-Prelog Weg 4, 8093 Zürich, Switzerland

have been studied most thoroughly, and an ever increasing body of evidence confirms that they are part of a fungal innate defense system against predators and parasites (Künzler 2015; Sabotič et al. 2012). Many reviews on mushroom lectins and protease inhibitors have been published recently but mainly viewing their potential application in human medicine (Dunaevsky et al. 2014; Erjavec et al. 2012; Hassan et al. 2015; Kobayashi and Kawagishi 2014; Sabotič and Kos 2012; Wong et al. 2010; Xu et al. 2011). The present review focuses on fruiting body lectins and protease inhibitors that have been characterized at the genetic, molecular, structural, and functional levels and that exhibit toxicity to nematodes and/or insects (Table 1). Their common characteristics include (i) small size, solubility, and resistance toward extreme pH and temperature; (ii) the lack of a signal sequence for classical secretion; (iii) the lack or low number of cysteine residues and disulphide bridges; and (iv) the lack of glycosylation, of which (ii) to (iv) are indicators of their cytoplasmic localization. Their exceptional characteristics make these fungal defense proteins attractive reagents for protecting crops against plant pests as well as in veterinary and human medicine against parasites.

Lectins

Lectins, also known as (hem)agglutinins, are defined as proteins containing at least one domain that binds to a specific carbohydrate (glycan) structure without modifying it (Sharon and Lis 2004). Binding of the carbohydrate is usually achieved by multiple weak interactions that result in high specificity, avidity, and/or affinity of the lectin for the carbohydrate ligand (Andre et al. 2015). Lectins were first classified according to their carbohydrate-binding specificity and, later, according to their sequence homology and evolutionary relatedness. The latter classification is based on the conservation of amino acid sequence motifs and the three-dimensional structures of the carbohydrate recognition domains (CRDs) (Varki 2009).

Based on the overall organization, three types of lectins are distinguished, namely merolectins, hololectins, and chimerolectins (Peumans and Van Damme 1995). Merolectins or monovalent lectins are small proteins with a single CRD. Hololectins are composed of two or more, usually homologous, CRDs on the same or on different polypeptide chains. The multivalency of carbohydrate-binding sites of hololectins is responsible for their ability to agglutinate cells or precipitate glycoconjugates (Brewer et al. 2002). Chimerolectins are fusion proteins composed of a CRD and an unrelated domain with a well-defined catalytic or biological activity that acts independently of the carbohydrate-binding domain.

The physiological functions of lectins are almost infinite, with the common denominator that they act as recognition

molecules in cell-molecule and cell-cell interactions (Sharon and Lis 2004; Varki 2009). In fungi, lectins have been implicated in defense against fungivores (Bleuler-Martinez et al. 2011), in fungal developmental processes including fruiting body formation (Luan et al. 2010; Swamy et al. 2004; Wang et al. 1998), in molecular recognition during mycorrhization or parasitism (Guillot et al. 1994; Guillot and Konska 1997; Wang et al. 1998), and in storage of nutrients (Kellens and Peumans 1990). Most experimental evidence is in support of the defensive function of fungal lectins, which is mediated by binding of the lectin to non-self glycans on the cells of the target organism (Künzler 2015). However, despite the recent reports of the successful identification of target glycoconjugates in insects and nematodes, the mechanism of action of many of these lectins is still unclear and the mechanisms may differ for the various lectins. All the characterized fungal lectins are multivalent and thus likely to crosslink glycoconjugates on the cell surface. The lattices between lectins and glycoconjugates may affect the turnover of the involved glycoconjugates by either preventing or triggering their internalization. Alternatively, the formation of these lattices may lead to activation of intracellular signaling pathways, e.g., by lectin-mediated oligomerization of (glycosylated) signaling receptors (see Künzler 2015 and references therein).

In the last 15 years, numerous mushroom lectins have been isolated and characterized, revealing mushrooms as a rich source of lectins with unique carbohydrate-binding specificities (Goldstein and Winter 2007; Hassan et al. 2015). Six different structural families of mushroom lectins: galectin, β -trefoil-type, β -propeller-type, actinoporin-type, cyanovirin-N-type, and immunoglobulin-type, have been identified to date (Varrot et al. 2013). Here, we summarize the current knowledge about the families containing members with nematotoxic and/or entomotoxic activities.

Galectins

Galectins constitute a family of β -galactoside-binding hololectins with a characteristic fold and a signature of carbohydrate-binding residues that occurs in animals and fungi but is apparently absent from bacteria and plants (Di Lella et al. 2011). The genome of the model mushroom *Coprinopsis cinerea* contains a tandem gene array encoding two highly homologous galectins: CGL1 and CGL2 (Cooper et al. 1997) (Table 1). In addition to these two proteins, the *C. cinerea* genome codes for a homologous, galectin-like protein, CGL3, that binds LacdiNAc and chitobiose but not β -galactosides (Walti et al. 2008). Besides these proteins from *C. cinerea*, galectins from two different *Agrocybe* species: ACG from *Agrocybe cylindracea* and AAG from *Agrocybe aegerita*, have been characterized (Yagi et al. 2001; Yang et al. 2009) (Table 1). The latter two share 88 % sequence identity

Table 1 Overview of entomotoxic and/or nematotoxic lectins and protease inhibitors from mushrooms

Name	Origin	Size of monomer	Fold (PDB code)	Oligomer	No. of carbohydrate-binding sites per oligomer	In vitro specificity ^a	In vivo specificity in <i>C. elegans</i>	Toxicity	Reference
CG12	<i>Coprinopsis chinera</i>	150 aa	Galectin (1UL9)	Homotetramer	4	Gal- β 1,4-Glc Gal- β 1,4-GlcNAc Gal- β 1,3-GalNAc Gal- β 1,4-Glc	Gal- β 1,4-Fuc	Entomotoxic Nematotoxic Amoebicidal nd	(Bleuler-Martinez et al. 2011; Boulianne et al. 2000; Buschi et al. 2010; Cooper et al. 1997; Titz et al. 2009; Wälsch et al. 2004; Yan et al. 2012)
AAG	<i>Agroclype aegerita</i>	159 aa	Galectin (2ZGL)	Homodimer	2	Gal- β 1,4-GlcNAc	na	nd	(Luan et al. 2010; Yang et al. 2009; Yang et al. 2005a; Yang et al. 2005b; Yang et al. 2005c)
ACG	<i>Agroclype cylindracea</i>	160 aa	Galectin (1WW7)	Homodimer	2	Gal- β 1,4-GlcNAc Neu5Ac- α 2,3-Gal- β 1,4-GlcNAc	na	nd	(Ban et al. 2005; Inamura et al. 2011; Kuwabara et al. 2013; Liu et al. 2008; Yagi et al. 2001; Yagi et al. 1997)
RSA	<i>Rhizoctonia solani</i>	142 aa	β -trefoil (4G9M)	Homodimer	4	Gal GalNAc	na	Entomotoxic	(Hamshou et al. 2013; Hamshou et al. 2010b; Hamshou et al. 2012; Skammaki et al. 2013)
SSA	<i>Sclerotinia sclerotiorum</i>	153 aa	β -trefoil (2X2S)	Homodimer	2	Gal GalNAc	na	Entomotoxic Amoebicidal	(Bleuler-Martinez et al. 2011; Sulzenbacher et al. 2010; Van Damme et al. 2007)
CNL	<i>Clitocybe nebularis</i>	149 aa	β -trefoil (3NBC)	Homodimer	2	Gal GalNAc Gal- β 1,4-Glc GalNAc- β 1,4-GlcNAc GalNAc- α 1,3(Fuc- α 1,2)Gal- β 1,4-GlcNAc	Not clear	Entomotoxic Nematotoxic Amoebicidal	(Bleuler-Martinez et al. 2011; Pohleven et al. 2011; Pohleven et al. 2009; Pohleven et al. 2012)
CCL2	<i>Coprinopsis chinera</i>	142 aa	β -trefoil (2LIE)	Homodimer ^b	2	Gal- β 1,4(Fuc- α 1,3)-GlcNAc GlcNAc- β 1,4(Fuc- α 1,3)-GlcNAc GlcNAc	α 1,3-Fucosylated N-glycan cores	Nematotoxic	(Schubert et al. 2012; Yan et al. 2012)
MpL	<i>Macropletiota procera</i>	141 aa	β -trefoil (4ION)	Homodimer	2	Gal- β 1,4-GlcNAc Gal- β 1,3-GlcNAc	Not clear	Nematotoxic	(Žurga et al. 2014)
BEL β -trefoil	<i>Boletus edulis</i>	146 aa	β -trefoil (414R)	Homodimer	6	Gal GalNAc Gal- β 1,4-Glc Gal- β 1,3-GalNAc	na	nd	(Bovi et al. 2013)
AAL	<i>Aleuria aurantia</i>	313 aa	β -propeller (10FZ)	Homodimer	10	Fuc- α 1,2-X Fuc- α 1,3-X Fuc- α 1,6-X	Not clear	Entomotoxic Nematotoxic Amoebicidal	(Bleuler-Martinez et al. 2011; Fujihashi et al. 2003; Fukumori et al. 1990; Kochibe and Funkkawa 1980; Olausson et al. 2010; Olausson et al. 2008; Tatenno et al. 2009; Wimmerova et al. 2003)
AOL ₁	<i>Aspergillus oryzae</i>	310 aa	β -propeller	Homodimer	12	Fuc- α 1,6-GlcNAc Fuc- α 1,2-Gal- β 1,4-GlcNAc	na	nd	(Matsumura et al. 2008; Tatenno et al. 2009)
AFL	<i>Aspergillus fumigatus</i>	315 aa	β -propeller (4UOU)	Homodimer	12	Terminal Fuc	na	nd	(Houser et al. 2015; Houser et al. 2013)
PVL	<i>Psathyrella velutina</i>	401 aa	β -propeller (2BWR)	Monomer	6	GlcNAc- β 1,4-X Neu5Ac	na	nd	(Cioci et al. 2006; Ueda et al. 1999a; Ueda et al. 2002; Ueda et al. 1999b; Ueda et al. 2003)
LbTee2	<i>Laccaria bicolor</i>	227 aa	β -propeller	Homotetramer ^b	24	3-O-Me-Man 4-O-Me-Man ^b 2-O-Me-Fuc	3-O-Me-Man	Nematotoxic	(Wohlschläger et al. 2014)
XCL	<i>Xerocomus chrysenteron</i>	143 aa	Actinoporin (1X10)	Homotetramer	4+4	Gal- β 1,3-GalNAc and GlcNAc	2-O-Me-Fuc on N-glycan antenna Core 1-O-glycans and terminal GlcNAc on α 1,3-mannose of N-glycans	Entomotoxic Nematotoxic	(Birk et al. 2004; Bleuler-Martinez et al. 2011; Damian et al. 2005; Francis et al. 2003; Jaber et al. 2008; Jaber et al. 2007; Jaber et al. 2006; Marty-Detraves et al. 2004; Trigueros et al. 2003; Yan et al. 2012)

(91 % sequence similarity) and are approximately 32 % identical (43 % similar) to CGL2.


Galectins CGL1 and CGL2 are highly abundant in young fruiting bodies but hardly produced in the vegetative mycelium of *C. cinerea* (Boulianne et al. 2000; Plaza et al. 2014). Expression of the genes is, however, induced in the vegetative mycelium upon challenge of this tissue with the fungivorous nematode *Aphelenchus avenae* (Bleuler-Martinez et al. 2011). Similarly, AAG expression is high in fruiting bodies and absent from vegetative mycelium (Luan et al. 2010).

Minimal ligands of fungal galectins are Gal- β 1,4-GlcNAc (LacNAc), Gal- β 1,4-Glc (Lac), Gal- β 1,3-GalNAc, and Gal- β 1,4-Fuc (Butschi et al. 2010; Walser et al. 2004; Walti et al. 2008; Yagi et al. 1997; Yang et al. 2009). Substitutions at positions 2 and 3 of the galactose residues in these ligands can increase their affinity toward the galectins considerably. As examples, ACG shows a strong preference for NeuAc- α 2,3-Lac (Ban et al. 2005) and one of the best ligands for CGL2 is Gal- α 1,3-Lac (Walser et al. 2004).

Crystal structures of CGL2, CGL3, AAG, and ACG revealed a typical galectin fold composed of two antiparallel, six-stranded β -sheets that form a β -sandwich (Fig. 1) (Ban et al. 2005; Walser et al. 2004; Walti et al. 2008; Yang et al. 2009). Galectin CGL2 and galectin-like protein CGL3 from *C. cinerea* oligomerize into homotetramers, whereby all carbohydrate-binding sites are located on one side of the tetramer (Walser et al. 2004; Walti et al. 2008). This spatial arrangement of the carbohydrate-binding sites increases the avidity of the lectins to multivalent ligands and allows the clustering of different glycoconjugates displayed on cell surfaces (Boscher et al. 2011; Rabinovich et al. 2007). In contrast, ACG and AAG form homodimers (Ban et al. 2005; Yang et al. 2009).

Both CGL1 and CGL2 are toxic toward the bacterivorous nematode *Caenorhabditis elegans*, the mosquito *Aedes aegypti*, and the amoebozoan *Acanthamoeba castellanii* (Bleuler-Martinez et al. 2011). The toxicity toward *C. elegans* has been shown to be mediated by the binding of

Fig. 1 Structures of fungal defense lectins and protease inhibitors. *Ribbon diagrams* are shown of homotetramer of galectin CGL2 (PDB code 2WKK), homodimer of β -trefoil-type lectin CNL (PDB code 3NBD), homodimer of β -propeller-type lectin AAL (PDB code 1OFZ) and its side view, homotetramer of actinoporin-type lectin ABL (PDB code 1Y2X), homodimer of chimerolectin MOA (PDB code 3EF2), serine protease inhibitor cospin (PIC) (PDB code 3N0K), and cysteine protease inhibitor clitocypin (CLT) (PDB code 3H6R). Monomers in dimers and tetramers are shown in yellow, green, blue, and red. Bound carbohydrate ligands are shown in orange. Arrows point to loops critical for protease inhibition






Fig. 2 Distribution of families of defense lectins and protease inhibitors in the fungal kingdom. A phylogenetic tree of the 145 fungi included in this study is depicted in the center. The full names of the organisms can be found in Table S1. The lifestyles of the individual fungi are indicated with a colored circle on the leaf node: red indicates a pathogen/parasite, yellow indicates a saprotroph, green indicates a mycorrhiza/endophyte, blue indicates a nematophagous fungus, and white indicates other/unknown. Basidiomycota are shaded in green, Ascomycota are shaded in blue, and “early diverging” fungi (non-dikarya) are shaded in orange. Selected subphyla (in bold) and orders are indicated. Each ring outside the phylogenetic tree is a heat map, representing the gene counts for each type of

lectin and protease inhibitor. White indicates that this organism has no genes of that type, and the color is increasingly opaque when the organism has more genes of that type. The gene count that corresponds to fully opaque is 10, except in the case of the galectins for which it is 5. β-Propeller-type lectins are separated into AAL-like (A) and LbTec2-like (T) family members. β-Trefoil-type cysteine protease inhibitors include only clitocyprin type as there were no macrocyprin homologs found in any of the organisms in the study set. The genes were identified by the presence of a conserved domain (PFAM domain or a custom hidden Markov model) or by BLASTP, full details on the methodology can be found in Text S1. The exact gene counts can be found in Table S1 and Table S2

CGL2 to a Gal-β1,4-Fuc-α1,6 epitope on the proximal GlcNAc residue of N-glycan cores of glycoproteins of the

nematode intestinal epithelium (Butschi et al. 2010). This epitope has also been detected on N-glycans of animal-parasitic

nematodes (Paschinger and Wilson 2015) and of platyhelminths (Paschinger et al. 2011).

Interestingly, phylogenetic analysis of the galectin family in fungi (Supplementary Table S2) indicates that some representatives may not be hololectins but chimeric lectins similar to the galectins from some invertebrates (Shi et al. 2014).

β -Trefoil-type lectins

One of the most prevalent hololectin families in mushrooms is constituted by proteins with sequence and structural similarity to the B-subunit of ricin, a protein toxin from the castor bean *Ricinus communis*, and is hence referred to as β -trefoil-type lectins (Cummings and Etzler 2009; Hazes 1996) (Fig. 2). These proteins adopt the so-called β -trefoil fold with pseudo-3-fold symmetry that usually harbors three potential, so-called canonical carbohydrate-binding sites. In addition to these canonical sites, the β -trefoil fold can also harbor non-canonical carbohydrate-binding sites (Schubert et al. 2012) and, as in case of protease inhibitors, binding sites for proteases (Žurga et al. 2015) (see below). The best characterized representatives of this family of mushroom lectins are *Rhizoctonia solani* agglutinin (RSA) and *Sclerotinia sclerotiorum* agglutinin (SSA) of the plant pathogens *R. solani* (basidiomycete) (Hamshou et al. 2013) and *S. sclerotiorum* (ascomycete) (Sulzenbacher et al. 2010), as well as CNL, CCL2, MpL, and BEL β -trefoil of the homobasidio(agarico)mycetes *Clitocybe nebularis* (Pohleven et al. 2009; Pohleven et al. 2012), *C. cinerea* (Schubert et al. 2012), *Macrolepiota procera* (Žurga et al. 2014), and *Boletus edulis* (Bovi et al. 2013) (Table 1). These proteins show high sequence variability, as they share only 7 to 16 % sequence identity (25 to 35 % similarity), the exception being CNL, MpL, and RSA that are 23 to 26 % identical (30 to 40 % similar).

All these proteins were isolated from sclerotia (RSA, SSA) or fruiting bodies (CNL, CCL2, MpL, BEL β -trefoil) of the originating fungi. Expression of RSA was found to be developmentally regulated, lectin expression being low in vegetative mycelium and the protein accumulating in adult sclerotia (Hamshou et al. 2007; Kellens and Peumans 1990). Based on the accumulation of protein during sclerotium formation and its depletion during mycelium germination, a storage function for RSA in *R. solani* has been proposed. CCL2 and its paralog CCL1 exhibit a pronounced fruiting body-specific expression with almost no expression in the vegetative mycelium of *C. cinerea* (Plaza et al. 2014; Schubert et al. 2012).

The carbohydrate-binding specificity of the different β -trefoil lectins varies (Table 1). Many of these lectins bind terminal Gal or GalNAc residues of oligosaccharides by their canonical carbohydrate-binding sites. In the case of CCL2, however, a single, non-canonical binding site binds with high affinity to Gal- β 1,4-(Fuc- α 1,3)GlcNAc and to GlcNAc- β 1,

4-(Fuc- α 1,3)GlcNAc (Schubert et al. 2012). The latter epitope is called the anti-HRP epitope, since it is found in the core of plant N-glycans, e.g., on the horseradish peroxidase and glycoproteins of many invertebrates, and is a known allergen (Paschinger et al. 2009).

β -Trefoil-type lectins are composed of approximately 150 amino acid residues with acidic (CNL, MpL) or very basic (RSA) isoelectric points. The typical β -trefoil fold consists of α , β , and γ repeats built from 12 β -strands. These strands are connected by loops, arranged in pseudo-3-fold symmetry, forming a six-stranded β -barrel (Fig. 1). The canonical carbohydrate-binding sites are found on the α -, β -, and/or γ -repeats, with either none (CCL2), one (CNL and SSA), two (RSA), or all three (BEL β -trefoil) being functional. The architecture of the canonical carbohydrate-binding sites is very similar. β -Trefoil-type lectins bind β -galactosides in an orientation that differs from that for galectins. While galectins bind linear glycans in a groove parallel to the protein surface, β -trefoil-type lectins bind them in a perpendicular orientation, in which only the non-reducing end of the glycan interacts with the binding pocket. All these proteins assemble to homodimers but, interestingly, each protein uses a different interface for dimer formation (Bovi et al. 2013; Pohleven et al. 2012; Schubert et al. 2012; Skamnaki et al. 2013; Sulzenbacher et al. 2010; Žurga et al. 2014).

Many of the β -trefoil-type lectins exhibit entomotoxic activity (Table 1). The mechanism of action has been analyzed at the cellular level for RSA and SSA. The toxicity of RSA toward the cotton leafworm depends on its binding to Gal/GalNAc-containing glycans on the midgut epithelium (Hamshou et al. 2013). RSA is not taken up by the epithelial cells, and intoxicated epithelial cells show symptoms of apoptosis, possibly caused by lectin-mediated activation of the respective signaling pathways (Hamshou et al. 2012). In the case of RSA-mediated toxicity toward the red flour beetle *Tribolium castaneum*, the ability of RSA to pass through the peritrophic matrix to reach intestinal epithelial cells has been shown to be a prerequisite for toxicity (Walski et al. 2014). SSA is highly toxic toward the pea aphid *Acyrtosiphon pisum*. It binds to the insect midgut cells and, like RSA, is not internalized but causes death of midgut epithelial cells probably via a signal transduction pathway triggered by a glycoreceptor (Hamshou et al. 2010a). In contrast, CCL2 is not toxic for insects but exhibits strong toxicity toward *C. elegans*. This toxicity is mediated by binding to the anti-HRP epitope in the core of nematode N-glycans (Schubert et al. 2012). The absence of entomotoxicity may be due to spatial restriction of the anti-HRP epitope to the nervous system in insects (Paschinger et al. 2009). The mechanism of toxicity of these (and other) hololectins is not clear. The toxicity of CNL against *C. elegans* has been shown to depend on both glycan binding and dimer formation (Pohleven et al. 2012). Recently, CCL2 was shown to bind to the nematode

intestinal epithelium without being endocytosed (Stutz et al. 2015). This binding led to complete disintegration of the microvillar organization—interestingly without breaching the barrier function of the epithelium. Finally, nematotoxicity of CCL2 appears to depend on active feeding, since application of the lectin without a supply of bacterial food was not toxic to the worms. Nematotoxicity against *C. elegans* has also been demonstrated for MpL, but the *in vivo* ligand remains obscure in this case (Žurga et al. 2014).

β -Propeller-type lectins

Members of the β -propeller-type family of mushroom lectins include hololectin AAL from the orange peel mushroom *Aleuria aurantia* (Fujihashi et al. 2003; Wimmerova et al. 2003) and its homologs from various ascomycetous molds such as AOL₁ from *Aspergillus oryzae*, and AFL from *Aspergillus fumigatus* (Houser et al. 2015; Matsumura et al. 2008) as well as PVL and LbTec2 from the basidiomycetes *Psathyrella velutina* and *Laccaria bicolor* (Cioci et al. 2006; Wohlschlager et al. 2014). Basidiomycetous PVL and LbTec2 show very low sequence homology to each other and to ascomycetous AAL (below 15 % sequence identity and 30 to 45 sequence similarity), while AFL and AOL₁ are highly homologous to each other (82 % sequence identity and 90 % similarity) and 30 % identical (60 % similar) to AAL.

AAL, PVL, and LbTec2 have been isolated from fruiting bodies of the respective fungi. AAL was expressed in vegetative mycelium and in fruiting bodies (Ogawa et al. 1998). Expression of LbTec2 in vegetative mycelium has been shown to be upregulated in the presence of mycorrhizal helper bacteria (Deveau et al. 2015).

These lectins are characterized by highly repetitive, wheel-like structures made from β -strands which harbor multiple carbohydrate-binding sites. Their molecular weights range from 24 kDa (LbTec2) to 42 kDa (PVL). AAL (33.5 kDa) is folded into six propeller blades, each composed of four antiparallel β -sheets with an additional small antiparallel β -sheet that is involved in the dimerization of the protein (Fig. 1) (Fujihashi et al. 2003; Wimmerova et al. 2003). Each of the AAL protomers contains six potential binding sites for terminal fucose residues, but their affinities have been shown not to be equivalent. One binding site is not functional at all, while another has a much higher affinity than the others and was occupied by free fucose when the hololectin was isolated from fruiting bodies (Olausson et al. 2008). The primary sequence and fold of AFL are very similar to those of AAL, but, in contrast to AAL, all six binding sites, although not equivalent, are functional in that they bind fucose (Houser et al. 2015). The three homologous proteins: AAL, AOL₁ and AFL, differ slightly in their specificity for the connectivity of the bound fucose residues (Matsumura et al. 2008) and are structurally similar to fucose-binding hololectins from bacteria, including

RSL from *Ralstonia solanacearum* (Sudakevitz et al. 2002). The primary sequence of hololectin LbTec2 is unrelated to that of AAL but is similar to those of proteins from filamentous bacteria (actinobacteria), slime moulds, and animals and is predicted to also adopt a six-bladed β -propeller fold with six carbohydrate-binding sites (Wohlschlager et al. 2014). LbTec2 binds to Sepharose and is specific for 2-O-Me-fucose and 3-O-Me-mannose residues. The affinity of LbTec2 for these monosaccharides is very low (millimolar range), but, according to a commonly accepted concept in the lectin field (Shinohara et al. 1997), when they are displayed on a surface, the avidity to these carbohydrates could be very high due to the oligomerization of the protein. Finally, PVL, whose primary sequence is not related to AAL or LbTec2, sharing 8 and 11 % sequence identity (33 % similarity), respectively, folds into a monomeric, seven-blade β -propeller with a total of six binding sites for terminal GlcNAc or Neu5Ac residues located at the interfaces between the blades (Audfray et al. 2015; Cioci et al. 2006; Ueda et al. 2002).

AAL is toxic toward nematodes, insects, and amoebozoia (Bleuler-Martinez et al. 2011) and also to the mucoromycete fungus *Mucor racemosus* (Amano et al. 2012). Both nematotoxicity and entomotoxicity have been shown to depend on binding of the hololectin to fucose-containing glycoconjugates in the target organisms. Based on the lack of toxicity resistance of various *C. elegans* strains with mutations in N-glycan and glycosphingolipid biosynthesis, fucosylated O-glycans have been hypothesized as most likely target glycan of AAL in this organism. LbTec2 was recently shown to be toxic to *C. elegans* and genetic evidence, in combination with glycome analysis, showed that the nematotoxicity is dependent on 2-O-Me-fucose and 3-O-Me-mannose in *C. elegans* N-glycans (Wohlschlager et al. 2014). According to a recent report on *C. elegans* N-glycan structure, LbTec2 most likely binds to 2-O-Me-Fuc- α 1,2-Gal- β 1,4 and 3-O-Me-Man- α 1,3 on the core β -mannose of *C. elegans* N-glycans (Yan et al. 2015). To date, no nematotoxicity or entomotoxicity has been reported for PVL, but the lectin has been shown to bind truncated N-glycans on cancer cells (Audfray et al. 2015).

Actinoporin-type lectins

One of the best known and characterized mushroom lectin families is that of actinoporin-type hololectins, also referred to in the PFAM database (<http://pfam.xfam.org/>) as the fungal fruiting body lectin family (FB_lectin, PF07367). Members of this protein family show structural homology to actinoporins, a family of pore-forming proteins (cytolysins) originally isolated from sea anemones (Birck et al. 2004; Kristan et al. 2009). It was suggested that the archetypal actinoporin fold is used for specific binding to various molecules at the plasma membrane surface (Kristan et al. 2009). Characterized representatives of the actinoporin-type mushroom lectin family

include XCL, ABL (ABA), SRL, and BEL from the basidiomycetes *Xerocomus (Boletus) chrysenteron*, *Agaricus bisporus*, *Sclerotium (Athelia) rolfsii*, and *B. edulis* (Birck et al. 2004; Bovi et al. 2011; Carrizo et al. 2005; Leonidas et al. 2007) and AOL₂ and TAP1 from the ascomycetes *Arthrobotrys oligospora* and *Sordaria macrospora* (Nowrousian and Cebula 2005; Rosen et al. 1996b). These representatives of basidiomycetes share 53 to 82 % sequence identity (67 to 89 % similarity), while AOL₂ is approximately 45 % identical (62 % similar) to them and TAP1 only 34 % identical (50 % similar). Surprisingly, AOL₂ and TAP1 share only 26 % sequence identity (45 % similarity).

With the exception of AOL₂ which was isolated from nematode traps, all actinoporin-type lectins have been isolated from fruiting bodies or sclerotia, suggesting a developmental control of their synthesis. In accordance with this, the expression of TAP1 was demonstrated to be strongly upregulated during fruiting body formation (Nowrousian and Cebula 2005). Similarly, expression of SRL is much higher in sclerotia than in vegetative mycelium (Swamy et al. 2004). Using respective knockout mutants, TAP1 and AOL₂ were shown not to be essential for the formation of fruiting bodies (Nowrousian and Cebula 2005) or the function of nematode traps, respectively (Balogh et al. 2003).

The actinoporin-type lectins are small proteins of approximately 16 kDa with neutral to basic isoelectric points. They all bind specifically to N-acetyl-galactosamine (GalNAc). ABL is a dual specificity lectin harboring separate binding sites for Gal- β 1,3-GalNAc (T- or Tn-antigen) and N-acetylglucosamine (GlcNAc) (Nakamura-Tsuruta et al. 2006). Similarly, structural studies on SRL revealed two carbohydrate-binding sites: a primary one for GalNAc and secondary one for GlcNAc (Leonidas et al. 2007). Based on sequence homology and functional studies, the GlcNAc-binding site is conserved in XCL, AOL₂, and BEL but not in TAP1 (Bleuler-Martinez et al. 2011). The protein structure is, similar to those of pore-forming actinoporins, composed of two β -sheets that consist of six and four β -strands connected by a helix-loop-helix motif. XCL, ABL, and BEL form tetramers in solution that have been described as dimers of dimers (Fig. 1). SRL was shown to be a dimer but can, from a structural point of view, form similar tetramers as other representatives (Birck et al. 2004; Bovi et al. 2011; Carrizo et al. 2005; Leonidas et al. 2007).

Several representatives of this lectin family have been shown to exhibit entomotoxic and nematotoxic activity (Table 1). XCL is toxic toward the hemipterans *A. pisum* and *Myzus persicae*, the dipterans *Drosophila melanogaster* and *A. aegypti*, as well as the nematode *C. elegans* (Bleuler-Martinez et al. 2011; Jaber et al. 2008; Trigueros et al. 2003). Both the entomotoxic and nematotoxic activity of XCL are dependent on carbohydrate binding, and genetic data

suggested that XCL binds to terminal GlcNAc residues in nematode N-glycans (Jaber et al. 2008; Yan et al. 2012). TAP1 is toxic to *C. elegans*, *A. aegypti*, and the amoebozoan *A. castellanii* (Bleuler-Martinez et al. 2011). SRL is toxic toward the cotton leaf worm *Spodoptera litura* by binding to membrane proteins of the midgut epithelium, thus triggering caspase-dependent cell death (Vishwanathreddy et al. 2014). Thus, in vitro and in vivo experiments suggest that these proteins mediate their toxicity by binding, at the same time, to the Gal- β 1,3-GalNAc epitope on mucin-type O-glycans and terminal GlcNAc on N-glycans. This binding mode would lead to clustering of glycoproteins on the intestinal epithelia of insects or nematodes via two types of protein-bound glycans (Chachadi et al. 2011; Yan et al. 2012). Interestingly, actinoporin-type lectins are also found in primitive plants, where they may also have a role in defense (Peumans et al. 2007).

β -Trefoil-type chimerolectins

Besides the β -trefoil-type hololectins described above, mushrooms also produce chimerolectins in which one or several β -trefoil-type lectin domains are fused to a domain with a different function. These proteins are analogous to bacterial and plant AB toxins including the previously mentioned plant toxin ricin. The best characterized examples of this lectin family are MOA and LSL from the saprophytic mushrooms *Marasmius oreades* (Cordara et al. 2011; Grahn et al. 2007; Grahn et al. 2009; Wohlschlager et al. 2011) and *Laetiporus sulphureus* (Mancheno et al. 2005; Mancheno et al. 2010; Tateno and Goldstein 2003). These proteins consist of a single N-terminal β -trefoil-type lectin domain fused to a cysteine protease (MOA) and to an aerolysin-type pore-forming domain (LSL). The numbers and specificities of the carbohydrate-binding sites of the β -trefoil domains differ in that all three canonical binding sites of MOA are functional and bind to glycans carrying terminal Gal- α 1,3-Gal/GalNAc- β epitopes (Grahn et al. 2007; Grahn et al. 2009; Wohlschlager et al. 2011) while only two of the three canonical binding sites of LSL appear to be functional and specific for β -galactosides including lactose and LacNAc (Angulo et al. 2011; Mancheno et al. 2005). Homologs of both proteins from other dikaryotic fungi have been partially characterized (Chumkhunthod et al. 2006; Kadirvelraj et al. 2011; Plaza et al. 2014; Wohlschlager et al. 2011). The MOA homologs PSL and SCA (*Schizophyllum commune* agglutinin), from the saprophytic mushrooms *Polyporus squamosus* and *S. commune*, differ from MOA in that PSL harbors a lower number of functional carbohydrate-binding sites with a different specificity toward terminal Neu5Ac- α 2,6-Gal- β epitopes but with the catalytic triad of the cysteine protease domain conserved (Kadirvelraj et al. 2011; Wohlschlager et al. 2011). SCA exhibits the carbohydrate-binding specificity of

MOA but not its catalytic activity (Wohlschlager et al. 2011). On the other hand, the homology of LSL to fruiting body-specific proteins of *C. cinerea* is restricted to the aerolysin domain (Plaza et al. 2014). SCA and the *C. cinerea* homolog of LSL are differentially expressed in fruiting bodies compared to the vegetative mycelium in *S. commune* and in *C. cinerea* (Ohm et al. 2010; Plaza et al. 2014).

MOA and the *C. cinerea* homolog of LSL have nematotoxic and the latter also entomotoxic activity (Plaza et al. 2014; Wohlschlager et al. 2011). LSL was shown to associate into hexamers in solution and crystals (Mancheno et al. 2005). By analogy to other aerolysin-like pore-forming toxins, it has been hypothesized that LSL undergoes a conformational change in its aerolysin domain on insertion into the target membrane, thus forming a pore (Mancheno et al. 2010). In contrast, MOA forms a dumbbell-shaped dimer (Fig. 1) in which the two β -trefoil domains represent the balls of the dumbbell and the dimerized cysteine protease domain the connecting bar (Grahn et al. 2007; Grahn et al. 2009). The latter domain undergoes a conformational change on binding divalent cations, which has been shown to be necessary for activation of the protease in vitro (Cordara et al. 2011; Wohlschlager et al. 2011). Using truncations and alterations of single residues responsible for carbohydrate-binding and protease activity of MOA, it was demonstrated that both functions are necessary for full nematotoxicity of this protein (Wohlschlager et al. 2011). Exploitation of *C. elegans* mutants defective in the biosynthesis of glycosphingolipids coupled with in vitro binding assays with glycosphingolipids isolated from *C. elegans* enabled Gal- α 1,3-GalNAc- β -containing glycosphingolipid species to be pinpointed as target glycans of MOA in *C. elegans* (Wohlschlager et al. 2011). The pH and Ca^{2+} requirements of the cysteine protease activity suggest that the protein has to be internalized to be toxic, in a manner analogous to that of bacterial AB toxins (Wohlschlager et al. 2011). In the cases of both LSL and MOA, the β -trefoil domains of these chimerolectins are thought to mediate initial binding of the protein toxins to the plasma membrane.

Additional families of potential fungal defense lectins

In addition to the above-described families of fungal defense lectins, several families of potential fungal defense lectins have recently been described. From fruiting bodies of the basidiomycetes *Hygrophorus russula* and *Grifola frondosa*, two mannose-specific lectins, termed *H. russula* lectin (HRL) and *G. frondosa* lectin (GFL), were isolated and characterized (Nagata et al. 2005; Suzuki et al. 2012). The amino acid sequence of these proteins is related to that of Jacalin from plants, and some plant members of this lectin family have recently been shown to protect the producing plants from herbivorous insects (Al Atalah et al. 2014). Similarly, recent reports suggest that both ascomycetes and basidiomycetes

contain cytoplasmic homologs of GNA (*Galanthus nivalis* agglutinin or snowdrop lectin), another family of mannose-binding entomotoxic plant lectins (Fouquaert et al. 2011; Shimokawa et al. 2012). Cyanovirin-N homologs (CVNH) constitute a family of fungal cytoplasmic lectins that comprises mannose-binding lectins from various ascomycetes including *Tuber borchii* and *Neurospora crassa* (Koharudin et al. 2008). Isolation and characterization of FVE (*Flammulina velutipes* fungal immunomodulatory protein) from fruiting bodies of *F. velutipes* has identified a family of immunoglobulin (Ig)-type cytoplasmic lectins present in many basidiomycetes (Paaventhian et al. 2003). From the basidiomycete *Pholiota squarrosa*, a very small cytoplasmic lectin, PhoSL (*P. squarrosa* lectin), specific for α 1,6-linked fucose on N-glycan cores, has been identified (Kobayashi et al. 2012). The lectin is homologous to *Rhizopus stolonifer* lectin (RSL), a lectin from the zygomycete *R. stolonifer* (Oda et al. 2003) and constitutes a novel family of cytoplasmic lectins. No nematotoxicity or entomotoxicity has so far been reported for any of these lectins.

Protease inhibitors

Proteolytic enzymes (also known as proteases, proteinases, or peptidases) are present in all organisms and play essential metabolic and regulatory roles in many biological processes. Due to the essential functions of proteases in life and death processes, anomalous proteolytic activities can be very harmful. Therefore, regulation of proteolytic activity is vital and takes place on several levels, from gene expression to post-translational modification and compartmentalization, and most importantly by their interaction with protease inhibitors (Lopez-Otin and Bond 2008; Rawlings et al. 2014).

Protease inhibitors are present in all kingdoms of life and can be broadly classified into those that inhibit peptidases of more than one catalytic type, those that inhibit families of peptidases of one catalytic class, and those that inhibit peptidases belonging to one family or a single peptidase. A detailed classification of protease inhibitors based on sequence homology is available in the MEROPS database (<http://merops.sanger.ac.uk/inhibitors/>). There are two general mechanisms by which protein inhibitors inhibit peptidases: irreversible “trapping” reaction, involving a conformational change of the inhibitor, and reversible tight-binding interactions, in which the inhibitor binds with high affinity to the peptidase active site. The detailed physiological roles of many protein protease inhibitors are still unknown. They are either involved in control of endogenous proteases or in defense mechanisms. Exogenous proteases targeted in defense are either virulence factors of pathogens and parasites or digestive proteases in grazers, predators, and parasites, that are involved in nutrient acquisition for growth and development or in evasion of host

defenses (Christeller 2005; Rawlings et al. 2014; Sabotič and Kos 2012).

The MEROPS database (release 9.12) identifies more than 650 protease inhibitors (Rawlings et al. 2014). They are classified into 78 families, 22 of which include members of fungal origin and only 7 families that include members from higher fungi. The latter are all inhibitors of serine and cysteine proteases. Protein inhibitors of metalloproteases have not been identified in fungi. Two protein aspartic protease inhibitors have been isolated from *Ganoderma lucidum* (Tian and Zhang 2005) and *Coriolus versicolor* (Zhang et al. 2012), but their sequences have not been determined. Families I51 (serine carboxypeptidase Y inhibitor), I32 (survivin-like caspase inhibitor), and I4 (serpin or α -1-peptidase inhibitor) include potential inhibitors of serine and cysteine proteases identified only as homologues in genomes. Inhibitory activity against serine proteases has been established in crude protein extracts of several species of mushrooms (Gzogyan et al. 2005; Vetter 2000; Zuchowski and Grzywnowicz 2006; Zuchowski et al. 2009), but only a few protease inhibitors from mushrooms have been biochemically and structurally characterized. One of them, the peptidase A inhibitor 1 from *Pleurotus ostreatus* (POIA1) has been classified into family I9, based on sequence homology to the propeptides of subtilisin-like proteases. In addition to its inhibitory activity, this inhibitor can also act as an intramolecular chaperone, assisting the folding of the cognate protease. Its biological role is probably that of controlling endogenous proteases (Kojima et al. 2005; Sasakawa et al. 2002). Most thoroughly characterized are the serine protease inhibitors of family I66 (mycospins) and cysteine protease inhibitors of families I48 (clitocypins) and I85 (macroscypins), which all exhibit entomotoxic activity.

Mycospins, fungal inhibitors of serine proteases

Mycospins are serine protease inhibitors from mushrooms that constitute family I66 in the MEROPS classification. This family includes three biochemically characterized members: cospin from *C. cinerea* (Sabotič et al. 2012), cnispin from *C. nebularis* (Avanzo Caglič et al. 2014; Avanzo et al. 2009), and LeSPI (*Lentinula edodes* serine protease inhibitor) from *L. edodes* (Odani et al. 1999).

Cospin is expressed abundantly in fruiting bodies of *C. cinerea* in contrast to vegetative mycelium where its expression is approximately 700-fold lower. In addition to cospin (PIC1), there are three more isoproteins in the *C. cinerea* genome sharing 38 to 95 % sequence identity with cospin (Sabotič et al. 2012). Cospin (CC1G_09480) is, together with its paralog (CC1G_09479), one of the most highly upregulated genes in *C. cinerea* young fruiting bodies (Plaza et al. 2014). Sequence variability at the protein level has also been shown for the natural isolates of cnispin from

C. nebularis from which three proteins with similar biochemical properties and N-terminal amino acid sequences were isolated, indicating that cnispin could be encoded by a multigene family. There is 31 % sequence identity and 46 % sequence similarity between cnispin and cospin amino acid sequences. Comparison of expression levels of cnispin in *C. nebularis* fruiting body and in vegetative mycelium suggests that it is constitutively expressed in mycelium and fruiting body, although expression was higher in the cap of the fruiting body. It was also shown that cnispin is not secreted into the medium (Avanzo et al. 2009).

Mycospins are small proteins (16 to 18 kDa) with acidic isoelectric points that are resistant to exposure to extreme pH conditions. They have very similar inhibitory profiles, all exhibiting strong trypsin-specific inhibition. They inhibit chymotrypsin only weakly and other serine proteases not at all or very weakly. Cospin inhibits trypsin with an equilibrium constant for inhibition (K_i) in the picomolar range and cnispin in the nanomolar range, while both inhibit chymotrypsin with K_i in the micromolar range (Avanzo et al. 2009; Sabotič et al. 2012). The crystal structure of cospin reveals that these proteins are members of the β -trefoil fold protein family. This fold is composed of 12 β -strands connected by 11 loops of various lengths and composition. Surprisingly, the reactive site residues for trypsin inhibition in the two proteins differ, being Arg27 of the β 2- β 3 loop in cospin and Lys127 of the β 11- β 12 loop in cnispin. They are both classic inhibitors that bind to the protease active site in a substrate-like manner and form a tight and stable complex with trypsin. The difference between cospin and cnispin is that in vitro the former persists in complex with trypsin for over a month at 37 °C, while cnispin is completely degraded by trypsin in 24 h. It is suggested that the β 2- β 3 loop involved in the inhibition of trypsin by cospin is better optimized for trypsin inhibition, since only small conformational changes are needed for binding to trypsin and the loop is more stable than the β 11- β 12 loop of cnispin that presumably undergoes more substantial changes on binding to trypsin (Avanzo Caglič et al. 2014; Sabotič et al. 2012).

Cospin exhibits a strong entomotoxic activity against *D. melanogaster* that is mediated by specific inhibition of the fly's serine proteases based on the lack of toxicity and protease inhibition of the cospin R27N mutant (Sabotič et al. 2012). Natural isolates of cnispin also show entomotoxic activity against *D. melanogaster*, albeit lower than that for cospin (Avanzo et al. 2009), validating the entomotoxic potential of the I66 family. Furthermore, cospin showed no toxicity against the nematode *C. elegans* and amoebozoan *A. castellanii*. A biological role for cospin in the defense of fruiting bodies against Drosophilidae is further corroborated by the absence of trypsin-like protease genes in the *C. cinerea* genome and by the fact that serine proteases are the predominant digestive proteolytic enzymes in dipterans (Sabotič et al. 2012; Terra and Ferreira 2005). In addition to the defensive

role directed against exogenous serine proteases, there is some evidence that cnispin and cospin could have a dual role, also participating in the regulation of endogenous serine proteases, in fruiting body development and/or resource recycling (Avanzo et al. 2009; Sabotič et al. 2012).

Mycocypins, inhibitors of cysteine proteases

Two families of cysteine protease inhibitors—clitocypins from *C. nebularis* (family I48 in Merops) and macrocypins from *M. procera* (family I85 in Merops)—have been identified in mushrooms and are collectively called mycocypins.

Clitocypins are encoded by a small gene family in the *C. nebularis* genome with significant sequence variability. This variability however has no influence on the inhibitory activity of clitocypins. Macrocypins in *M. procera* show even higher amino acid sequence variability, their sequences being grouped into five groups (macrocypins 1–5) with 75–86 % sequence identity between groups and more than 90 % sequence identity within groups. The sequence variability is reflected in their inhibitory profiles, since different macrocypins exhibit different strengths of inhibition for proteases. Even though they have many biochemical properties in common, sequence identity between clitocypin and macrocypin amino acid sequences is low. They are approximately 23 % identical and 33 % similar (Sabotič et al. 2007a; Sabotič et al. 2006; Sabotič et al. 2009).

The expression pattern of clitocypin in *C. nebularis* fruiting bodies has revealed similar amounts of clitocypin protein throughout the fruiting body while, at the messenger RNA (mRNA) level, expression varies in different parts of fruiting bodies. Clitocypin is also expressed in the vegetative mycelium and shown not to be secreted into the medium (Sabotič et al. 2006). Macrocypins exhibit tissue-specific expression (Sabotič et al. 2011). In *M. procera*, the protein is present throughout the fruiting body, but the amount of protein is significantly higher in the veil fragments on the cap and in the ring (annulus). Again, little congruence was observed with mRNA and protein expression profiles. Regulation of mycocypin expression has been further analyzed in the model mushroom *C. cinerea*, using mycocypin promoters and a reporter gene. Clitocypin and macrocypin promoters were transcriptionally active in vegetative mycelium and in fruiting bodies of *C. cinerea*. The clitocypin promoter displayed an expression pattern similar to that of a constitutive promoter with uniform expression throughout the tissues. In contrast, the macrocypin promoter displayed tissue-specific expression during fruiting body development which was similar to the macrocypin expression pattern in *M. procera*. This difference in temporal and spatial expression indicates specific developmental or protective roles for individual mycocypins (Sabotič et al. 2011).

Mycocypins are small proteins (17–19 kDa) with similar isoelectric points (pH 4.8) and apparent stability against high temperature and extreme pH mediated by the ability to unfold reversibly. They are strong inhibitors of papain-like proteases with equilibrium constants for inhibition (K_i) of papain in the low nanomolar range. Mycocypins also strongly inhibit cysteine cathepsins: Cathepsins L, V, and S are inhibited with K_i values in the nanomolar range by both clitocypin and macrocypins, and cathepsin K is strongly inhibited by clitocypin and more weakly by macrocypins. Cathepsins B and H, that exhibit both endopeptidase and exopeptidase activity, are not or only very weakly inhibited by mycocypins. In addition to papain-like cysteine proteases (family C1), mycocypins also inhibit asparaginyl endopeptidase (AEP) also called legumain (family C13) while trypsin, but not AEP, is inhibited by macrocypin 4 (Renko et al. 2010; Sabotič et al. 2007a; Sabotič et al. 2009).

Mycocypins have a β -trefoil fold. A distinct motif for their binding to papain-like cysteine proteases was revealed by the three-dimensional structure of clitocypin in complex with cathepsin V. Two broad loops of mycocypins (β 1- β 2 and β 3- β 4) fill the active site cleft of the protease along its whole length, occluding the catalytic cysteine residue and preventing the approach of substrate molecules. The two loops are stabilized by numerous hydrogen bonds. A different site is involved in the inhibition of AEP or trypsin by mycocypins. Asparagine in the β 5- β 6 loops of clitocypin and macrocypins 1 and 3 mediates the inhibition of AEP. In macrocypin 4, the asparagine is replaced by a lysine, enabling inhibition of trypsin. Consideration of the crystal structures leads to the conclusion that the binding loops of clitocypin and macrocypins exhibit substantial conformational flexibility during binding into the active site of their target enzymes (Renko et al. 2010; Renko et al. 2012).

The sequence diversity, with sites showing positive evolutionary selection, the variations in spatial and temporal expression, the variations in inhibitory profiles, and scarcity of cysteine proteases in basidiomycetes, provides evidence that mycocypins' biological role is defense against pathogen infection and/or predation by insects or other pests. They would target exogenous cysteine proteases found in mycoviruses, nematodes, insects, mites, and slugs, all known antagonists of higher fungi (Brzin et al. 2000; Sabotič et al. 2007a; Sabotič et al. 2006; Sabotič and Kos 2012; Sabotič et al. 2007b). Entomotoxic activity of mycocypins was shown directly for the model coleopteran insect pest Colorado potato beetle. It was mediated by inhibition of specific digestive proteases, and evidence suggests that the negative effect of mycocypins on larval growth and development is mediated through multiple levels (Šmid et al. 2013; Šmid et al. 2015).

Distribution within the fungal kingdom

In the past decade, over 400 fungal genome sequences have become available, due in large part to the efforts of the US Department of Energy Joint Genome Institute (DOE JGI) in compliance with its missions in alternative energy, global carbon cycling, and biogeochemistry (Grigoriev et al. 2011). On the basis of their predicted proteomes, we selected a set of 145 genomes from the latest JGI database (<http://genome.jgi-psf.org/programs/fungi/index.jsf>) in order to examine the distribution of the above-described families of fungal defense proteins within the fungal kingdom and to draw conclusions about their evolution and physiological function. For these purposes, we calculated a phylogenetic tree and included available metadata on the lifestyle of the selected fungi. The presence or absence of fungal defense families was based either on BLASTP searches with single family representatives or on available consensus sequences using the PFAM database (see Supplementary Text for details). In addition, we performed a gene clustering analysis of the identified hits of a specific defense gene family (see Supplementary Text for details). The analysis relies inherently on the quality of the published gene predictions. The results of this analysis are summarized in Fig. 2 and Supplementary Tables S1, S2, and S3.

One of the main outcomes of the analysis is that both types of fungal defense proteins, lectins and protease inhibitors, occur more frequently in the phylum *Basidiomycota* than in the phylum *Ascomycota*. This may be partly due to the fact that most of these lectins and protease inhibitors were initially identified in basidiomycetes and they exhibit quite high sequence diversity. Thus, orthologs encoded in genomes of ascomycetes might not be detected using BLASTP searches because the sequence similarity is too low. Within *Basidiomycota*, these proteins appear to be restricted to the subphylum *Agaricomycotina* and to be almost absent from the other subphyla, *Pucciniomycotina* and *Ustilaginomycotina*. These results are in agreement with a previous, preliminary study (Bleuler-Martinez et al. 2011) and may be explained by the parasitic lifestyle of these subphyla, since the fungus may be sheltered from fungivores by the defense system of the host. Accordingly, within the phylum *Ascomycota*, the subphylum *Taphrinomycotina* (Pneji, Schpo, Tapde), which also harbors mainly parasitic fungi, also completely lacks fungal defense proteins. An alternative explanation for the lack of fungal defense proteins in these subphyla is, however, suggested by examination of the closely related subphylum *Saccharomycotina* (Yarli, Sacce, Klula, Picpa, Debha, Canal). Members of this subphylum propagate mainly as yeasts, i.e., as unicellular organisms. Since many members of the subphyla *Pucciniomycotina*, *Ustilaginomycotina*, and *Taphrinomycotina* are dimorphic, i.e., propagate as yeasts for a significant part of their lifestyle, the lack of fungal defense proteins may correlate with their unicellular morphology with the reasoning that a defense system against fungivores only

makes sense for a multicellular organism. In accordance with both of the above reasonings, the pathogenic basidiomycetous yeast *Cryptococcus neoformans*, which belongs to *Agaricomycotina*, also lacks the described defense proteins. Along these lines, the apparent difference in occurrence of fungal defense proteins between the phyla *Ascomycota* and *Basidiomycota* might also be due to the fact that most analyzed basidiomycetous species are saprotrophs, whereas most ascomycetous species are (plant) pathogens (biotrophs). On the other hand, the genomes of some of the pathogenic members of the *Agaricomycotina* (Rhiso, Armme) code for a significant number of fungal defense proteins.

Among *Agaricomycotina*, mycorrhizal species appear to be particularly rich in fungal defense proteins. In fact, there was a good correlation of the presence of lectins homologous to LbTec2 (β -propeller-type lectin T) with this lifestyle which included also the early diverging, endomycorrhizal fungus *Rhizophagus irregularis* (*Glomus intraradices*, Gloin) (Supplementary Table S3). Eventually, these fungi take over part of the defense of the root system of the host plant against root-targeting herbivores during symbiosis. In accordance with this hypothesis, genes coding for defense proteins like LbTec2 are induced in the mycorrhizal state (Martin et al. 2008) and some fungivorous nematodes are believed to be able to feed also on plant roots (Yeates et al. 1993). Similarly, the occurrence of defense proteins in the two nematophagous ascomycetes, *Arthrobotrys oligospora* (Artol) and *Monacrosporium haptotylum* (Monha) compared to that of other ascomycetes, may be explained by an increased demand of these fungi for protection from fungivorous species among the prey nematodes.

Early diverging fungi generally appear to lack defense proteins but the subphylum *Mucoromycotina* and the related, sole representative of the phylum *Glomeromycota* among the selected fungi are surprisingly rich in β -trefoil-type defense lectins. This finding suggests that this family of defense proteins is ancient.

Analysis of individual fungal genomes has confirmed previous results that the genomes often encode several paralogs of a given defense protein and that the paralogous genes are often clustered, most probably as the result of gene duplications (Supplementary Table S2). Examples are the tandem repeats of galectin- and cospin-encoding genes in *C. cinerea*. The regulation of these genes and the specificity of the encoded proteins are similar (Butschi et al. 2010; Plaza et al. 2014; Sabotič et al. 2012), but there might be cases where either the regulation of the duplicated genes or the specificity of the encoded proteins is different. Defense gene duplications and diversifications may enable the composition of species-specific armories against predators and parasites. The scattered distribution within the fungal kingdom, the conservation in bacteria, and the lack of introns in the coding regions (Supplementary Table S2) may indicate that some of these

genes were acquired from bacteria by horizontal gene transfer (HGT). As an example, Moran et al. recently presented evidence suggesting acquisition of LSL by HGT from bacteria (Moran et al. 2012).

Potential applications

The described fungal defense lectins have been considered as potential pesticidal agents in crop protection. Strong entomotoxicity affecting development and survival of the economically important plant pests cotton leafworms *Spodoptera littoralis* and *S. litura* was shown for β -trefoil-type lectin RSA and actinoporin-type lectin SRL, respectively, both binding Gal/GalNAc-containing glycans (Hamshou et al. 2010b; Vishwanathreddy et al. 2014). Additionally, RSA exhibited toxicity toward the red flour beetle *T. castaneum*, an important pest of stored products (Walski et al. 2014). Furthermore, the effect of different fungal lectins against the important sap sucking crop pests, using pea aphid *A. pisum* as a model organism, has been analyzed. The β -trefoil lectin SSA exhibited strong toxicity in feeding assays (Hamshou et al. 2010a). Actinoporin-type lectin XCL also exhibited toxicity against aphids including *A. pisum* and green peach aphid (*M. persicae*) (Jaber et al. 2007; Jaber et al. 2006; Trigueros et al. 2003).

In addition to lectins, fungal protease inhibitors have been considered as potential insecticidal agents against herbivorous insects. Both families of mycrocypins, cliticypin and macrocypins, affected larval growth and development of the major potato pest Colorado potato beetle (*Leptinotarsa decemlineata*). They acted by inhibiting specific digestive proteases without triggering adaptive responses in larval guts at transcriptional level (Šmid et al. 2013; Šmid et al. 2015).

Analogously to the application of entomotoxic fungal defense proteins against herbivorous insects, nematotoxic fungal defense proteins could be used for protecting crops from plant-parasitic nematodes, e.g., root knot nematode (*Meloidogyne incognita*), root lesion nematode (*Pratylenchus* spp.), or the pine wilt nematode (*Bursaphelenchus xylophilus*) (Jones et al. 2013). However, no toxicity of fungal defense proteins for plant-parasitic nematodes has, to our knowledge, been reported so far.

One of the issues of the use of fungal defense proteins in crop protection is the possible toxicity of the proteins for beneficial insects, humans, and livestock. As an example, XCL was endocytosed by various animal and human cell lines and caused inhibition of proliferation of some of these cell lines (Francis et al. 2003; Marty-Detraves et al. 2004). A second issue is the way the protein is applied to the crop plant. The most effective way would be to modify the crop plant genetically so that it expresses the fungal proteins in the tissue that is under attack by the herbivore. Currently, however, the use of genetically modified crops is limited due to strict

regulations and public opposition. Thus, external application of proteins as deterrents, preventing contact between the herbivore and the plant, might be a more promising option.

Another potential application of nematotoxic fungal defense proteins, in particular lectins, is in veterinary and human medicine for fighting animal and human parasitic nematodes. A recent study reported a dose-dependent toxicity of CGL2, CCL2, AAL, and MOA toward larval and adult stages of the barbers pole worm, *Haemonchus contortus* in vitro, in which toxicity of the lectins correlated with their binding to the intestinal epithelium of this animal-parasite nematode (Heim et al. 2015). Successful in vivo applications of nematotoxic, carbohydrate-binding protein toxins of microbial origin against parasitic nematodes were demonstrated for the *Bacillus thuringiensis* toxin Cry5B against the hookworm *Ancylostoma ceylanicum* in hamsters and against the large roundworm *Ascaris suum* in pigs (Cappello et al. 2006; Hu et al. 2012; Urban et al. 2013). The nematotoxicity of all those carbohydrate-binding proteins proven to be effective against parasitic nematodes had previously been demonstrated in *C. elegans*, showing the power of this model organism for detection and characterization of nematotoxic proteins. As an alternative to the direct use of nematotoxic lectins as therapeutics of infestations with parasitic nematodes, these proteins could be used as leads for a vaccination strategy against these parasites. Nematotoxicity of many of the above-described fungal defense lectins, and also of bacterial Cry5B, has been shown to rely on the binding of specific, lipid- or protein-bound glycoepitopes on the intestinal epithelium of *C. elegans*. A combination of genetic, biochemical, and toxicity assays allowed the structure of the target glycoepitope of some of these lectins in *C. elegans* to be identified (Table 1). Since some of these epitopes are conserved in parasitic nematodes with respect to both structure and localization (Heim et al. 2015; Paschinger and Wilson 2015), they represent candidate hidden antigens for vaccination. Vaccination with hidden antigens is based on the idea that epitopes that are hidden from the animal immune system during a normal infection with a pathogen or infestation with a parasite have a higher likelihood of inducing a strong and protective antibody response (Munn 1997). In support of this idea, vaccinations with partially purified, intestinal hemoglobins isolated from adult *H. contortus* significantly lowered the burden of sheep and avoided reinfections with this parasite (LeJambre et al. 2008). It is not clear at the moment whether this reported protection was mediated by carbohydrate or protein epitopes or a combination of both.

Fungal lectins are also being considered for treatment of cancers and microbial infections, based on their antitumor and immunomodulatory, antiproliferative and mitogenic, antiviral, and antimicrobial activities. Finally, fungal lectins have been considered widely as diagnostic tools based on their highly specific binding of glycoconjugates and on the altered

glycosylation profiles observed in various diseases. These potential medical applications of fungal defense lectins and protease inhibitors have been described in more detail in recent reviews (Erjavec et al. 2012; Hassan et al. 2015; Sabotič and Kos 2012; Xu et al. 2011).

Conclusions and perspectives

The abundance and diversity of toxic proteins in dikaryotic fungi is astonishing. On one hand, structurally similar proteins, for example β -trefoil-type lectins and β -trefoil-type protease inhibitors, perform different functions, and, on the other hand, structurally distinct lectins exert toxicity by binding to the same glycoepitopes. Some of these toxic proteins are directed toward very specific target organisms. Cospin, for example, appears to be toxic to *D. melanogaster* but not to other diptera, nematodes, amoebozoia, fungi, or bacteria (Sabotič et al. 2012; Sabotič, unpublished observations), whereas other toxic proteins provide a general protection against different antagonists (Bleuler-Martinez et al. 2011). LbTec2, for example, binds a glycan modification that has been observed in different phyla ranging from bacteria to plants and animals (Wohlschläger et al. 2014). Another level of diversity is found in the regulation of the biosynthesis of these fungal toxins. Most of these proteins were found to be expressed abundantly in fruiting bodies, as a form of constitutive defense of this reproductive structure. Biosynthesis of some of the proteins was demonstrated to be induced in the vegetative mycelium upon challenge with a fungivorous nematode (Bleuler-Martinez et al. 2011).

Searches of available fungal genomes for homologs of the already characterized lectins and protease inhibitors reveal that these proteins are widely present among ascomycetes and basidiomycetes. Based on the current pace of identification, of, e.g., novel fungal lectins, the proteins described in this review most probably represent only the tip of an iceberg and there is no doubt that more protein toxins from fungi still await to be identified. In order to identify novel families of fungal protein toxins, such identification should be performed not only on the basis of sequence similarity but rather on the basis of toxicity or biochemical activity (inhibition of proteases or binding to carbohydrates).

Fungal protein toxins have advantages over bacterial, animal, and most plant toxins for biotechnological applications since they are produced in the cytoplasm, which eases their expression in bacterial expression systems. They are, therefore, more readily available in recombinant form for applications in crop protection or veterinary and human medicine. Finally, fungal defense proteins are invaluable research tools.

Understanding the mechanism of action of these protein toxins will provide further insight into the general mechanism of these effectors. In the case of holotoxins, the exact toxicity

mechanism, beyond the dependence on carbohydrate binding, is still unclear. Differential expression of these effectors of a fungal innate defense system can be exploited as a readout to identify the signals and receptors that convey their induction upon challenge by the antagonist and upon developmental cues. Knowledge about these players would certainly be a big step forward in the understanding of the fungal innate defense system. At the moment, the limited availability of genetic tools, for example, for *Agaricomycotina*, reduces the pace of these investigations. Results obtained by such studies are relevant as they can be readily translated into the areas of plant and animal innate immunity. It has been shown that plants and, more recently, animals employ lectins as effector molecules in innate immunity against pathogens and parasites. Thus, learning tricks from fungi could provide an edge in combating plant pests as well as animal and human pathogens and parasites.

Acknowledgments We would like to apologize to all the people whose work could not be cited due to space limitations of this review. We are grateful to Dr. Roger H. Pain for critical reading of the manuscript.

Compliance with ethical standards

Funding The work of MK is supported by grants of the Swiss National Science Foundation (Grant No. 31003A_149512) and ETH Zürich (Grant No. ETH-34 11–2). The work of JS is supported by Slovenian Research Agency (Grant No. P4-0127).

Conflict of interest The authors declare that they have no conflict of interest.

References

- Al Atalah B, Smagghe G, Van Damme EJ (2014) Oryzata, a jacalin-related lectin from rice, could protect plants against biting-chewing and piercing-sucking insects. *Plant Sci* 221–222:21–8
- Amano K, Katayama H, Saito A, Ando A, Nagata Y (2012) *Aleuria aurantia* lectin exhibits antifungal activity against *Mucor racemosus*. *Biosci Biotechnol Biochem* 76(5):967–70. doi:10.1271/bbb.110982
- Andre S, Kaltner H, Manning JC, Murphy PV, Gabius HJ (2015) Lectins: getting familiar with translators of the sugar code. *Molecules* 20(2): 1788–1823. doi:10.3390/molecules20021788
- Angulo I, Acebron I, de las Rivas B, Munoz R, Rodriguez-Crespo I, Menendez M, Garcia P, Tateno H, Goldstein IJ, Perez-Agote B, Mancheno JM (2011) High-resolution structural insights on the sugar-recognition and fusion tag properties of a versatile beta-trefoil lectin domain from the mushroom *Laetiporus sulphureus*. *Glycobiology* 21(10):1349–61
- Audfray A, Beldjoudi M, Breiman A, Hurbin A, Boos I, Unverzagt C, Bouras M, Lantuejoul S, Coll JL, Varrot A, Le Pendu J, Busser B, Imbert A (2015) A recombinant fungal lectin for labeling truncated glycans on human cancer cells. *PLoS One* 10(6):e0128190. doi:10.1371/journal.pone.0128190
- Avanzo Caglič P, Renko M, Turk D, Kos J, Sabotič J (2014) Fungal beta-trefoil trypsin inhibitors cnispin and cospin demonstrate the

- plasticity of the beta-trefoil fold. *Biochim Biophys Acta* 1844(10): 1749–56. doi:10.1016/j.bbapap.2014.07.004
- Avanzo P, Sabotić J, Anžlovar S, Popović T, Leonardi A, Pain RH, Kos J, Brzin J (2009) Trypsin-specific inhibitors from the basidiomycete *Clitocybe nebularis* with regulatory and defensive functions. *Microbiology* 155(12):3971–3981. doi:10.1099/mic.0.032805-0
- Balogh J, Tunlid A, Rosen S (2003) Deletion of a lectin gene does not affect the phenotype of the nematode-trapping fungus *Arthrobotrys oligospora*. *Fungal Genet Biol* 39(2):128–35
- Ban M, Yoon HJ, Demirkan E, Utsumi S, Mikami B, Yagi F (2005) Structural basis of a fungal galectin from *Agrocybe cylindracea* for recognizing sialoconjugate. *J Mol Biol* 351(4):695–706
- Birck C, Damian L, Marty-Detraves C, Lougarre A, Schulze-Briese C, Koehl P, Fournier D, Paquereau L, Samama JP (2004) A new lectin family with structure similarity to actinoporins revealed by the crystal structure of *Xerocomus chrysenteron* lectin XCL. *J Mol Biol* 344(5):1409–20
- Bleuler-Martinez S, Butschi A, Garbani M, Walti MA, Wohlschlagel T, Potthoff E, Sabotić J, Pohleven J, Luthy P, Hengartner MO, Aebi M, Künzler M (2011) A lectin-mediated resistance of higher fungi against predators and parasites. *Mol Ecol* 20(14):3056–70. doi:10.1111/j.1365-294X.2011.05093.x
- Bleuler-Martinez S, Schmieder S, Aebi M, Künzler M (2012) Biotin-binding proteins in the defense of mushrooms against predators and parasites. *Appl Environ Microbiol* 78(23):8485–7
- Boscher C, Dennis JW, Nabi IR (2011) Glycosylation, galectins and cellular signaling. *Curr Opin Cell Biol* 23(4):383–92
- Boulianne RP, Liu Y, Aebi M, Lu BC, Kues U (2000) Fruiting body development in *Coprinus cinereus*: regulated expression of two galectins secreted by a non-classical pathway. *Microbiology* 146(Pt 8):1841–53
- Bovi M, Carrizo ME, Capaldi S, Perduca M, Chiarelli LR, Galliano M, Monaco HL (2011) Structure of a lectin with antitumoral properties in king bolete (*Boletus edulis*) mushrooms. *Glycobiology* 21(8):1000–9
- Bovi M, Cenci L, Perduca M, Capaldi S, Carrizo ME, Civiero L, Chiarelli LR, Galliano M, Monaco HL (2013) BEL beta-trefoil: a novel lectin with antineoplastic properties in king bolete (*Boletus edulis*) mushrooms. *Glycobiology* 23(5):578–92
- Brewer CF, Miceli MC, Baum LG (2002) Clusters, bundles, arrays and lattices: novel mechanisms for lectin-saccharide-mediated cellular interactions. *Curr Opin Struct Biol* 12(5):616–23
- Brzin J, Rogelj B, Popović T, Štrukelj B, Ritonja A (2000) Clitocyprin, a new type of cysteine proteinase inhibitor from fruit bodies of mushroom *Clitocybe nebularis*. *J Biol Chem* 275(26):20104–9
- Butschi A, Titz A, Walti MA, Olieric V, Paschinger K, Nobauer K, Guo X, Seiberger PH, Wilson IB, Aebi M, Hengartner MO, Künzler M (2010) *Caenorhabditis elegans* N-glycan core beta-galactoside confers sensitivity towards nematotoxic fungal galectin CGL2. *PLoS Pathog* 6(1):e1000717. doi:10.1371/journal.ppat.1000717
- Cappello M, Bungi RD, Harrison LM, Bischof LJ, Griffiths JS, Barrows BD, Aroian RV (2006) A purified *Bacillus thuringiensis* crystal protein with therapeutic activity against the hookworm parasite *Ancylostoma ceylanicum*. *Proc Natl Acad Sci U S A* 103(41): 15154–9. doi:10.1073/pnas.0607002103
- Carrizo ME, Capaldi S, Perduca M, Irazoqui FJ, Nores GA, Monaco HL (2005) The antineoplastic lectin of the common edible mushroom (*Agaricus bisporus*) has two binding sites, each specific for a different configuration at a single epimeric hydroxyl. *J Biol Chem* 280(11):10614–23. doi:10.1074/jbc.M411989200
- Chachadi VB, Inamdar SR, Yu LG, Rhodes JM, Swamy BM (2011) Exquisite binding specificity of *Sclerotium rolfii* lectin toward TF-related O-linked mucin-type glycans. *Glycoconj J* 28(1):49–56. doi:10.1007/s10719-011-9323-8
- Christeller JT (2005) Evolutionary mechanisms acting on proteinase inhibitor variability. *FEBS J* 272(22):5710–22. doi:10.1111/j.1742-4658.2005.04975.x
- Chumkhunthod P, Rodtong S, Lambert SJ, Fordham-Skelton AP, Rizkallah PJ, Wilkinson MC, Reynolds CD (2006) Purification and characterization of an N-acetyl-d-galactosamine-specific lectin from the edible mushroom *Schizophyllum commune*. *Biochim Biophys Acta Gen Subj* 1760(3):326–332
- Cioci G, Mitchell EP, Chazalet V, Debray H, Oscarson S, Lahmann M, Gautier C, Breton C, Perez S, Imbert A (2006) [beta]-Propeller crystal structure of *Psathyrella velutina* lectin: an integrin-like fungal protein interacting with monosaccharides and calcium. *J Mol Biol* 357(5):1575–1591
- Cooper DN, Boulianne RP, Charlton S, Farrell EM, Sucher A, Lu BC (1997) Fungal galectins, sequence and specificity of two isolectins from *Coprinus cinereus*. *J Biol Chem* 272(3):1514–21
- Cordara G, Egge-Jacobsen W, Johansen HT, Winter HC, Goldstein IJ, Sandvig K, Krenzel U (2011) *Marasmius oreades* agglutinin (MOA) is a chimerolectin with proteolytic activity. *Biochem Biophys Res Commun* 408(3):405–10. doi:10.1016/j.bbrc.2011.04.031
- Crenshaw RW, Harper SN, Moyer M, Privalle LS (1995) Isolation and characterization of a cDNA clone encoding a lectin gene from *Agaricus bisporus*. *Plant Physiol* 107(4):1465–6
- Cummings RD, Etzler ME (2009) R-type Lectins. In: Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, Hart GW, Etzler ME (eds) *Essentials of Glycobiology*. 2010/03/20 edn. Cold Spring Harbor Laboratory Press
- Damian L, Fournier D, Winterhalter M, Paquereau L (2005) Determination of thermodynamic parameters of *Xerocomus chrysenteron* lectin interactions with N-acetyl-galactosamine and Thomsen-Friedenreich antigen by isothermal titration calorimetry. *BMC Biochem* 6:11. doi:10.1186/1471-2091-6-11
- Deveau A, Barret M, Diedhiou AG, Leveau J, de Boer W, Martin F, Samiguet A, Frey-Klett P (2015) Pairwise transcriptomic analysis of the interactions between the ectomycorrhizal fungus *Laccaria bicolor* S238N and three beneficial, neutral and antagonistic soil bacteria. *Microb Ecol* 69(1):146–59. doi:10.1007/s00248-014-0445-y
- Di Lella S, Sundblad V, Cerliani JP, Guardia CM, Estrin DA, Vasta GR, Rabinovich GA (2011) When galectins recognize glycans: from biochemistry to physiology and back again. *Biochemistry* 50(37): 7842–57. doi:10.1021/bi201121m
- Dunaevsky YE, Popova VV, Semenova TA, Beliakova GA, Belozersky MA (2014) Fungal inhibitors of proteolytic enzymes: classification, properties, possible biological roles, and perspectives for practical use. *Biochimie* 101:10–20. doi:10.1016/j.biochi.2013.12.007
- Erjavec J, Kos J, Ravnikar M, Dreot T, Sabotić J (2012) Proteins of higher fungi—from forest to application. *Trends Biotechnol* 30(5):259–273. doi:10.1016/j.tibtech.2012.01.004
- Fouquaert E, Peumans WJ, Gheysen G, Van Damme EJ (2011) Identical homologs of the *Galanthus nivalis* agglutinin in *Zea mays* and *Fusarium verticillioides*. *Plant Physiol Biochem* 49(1):46–54. doi:10.1016/j.plaphy.2010.09.018
- Francis F, Marty-Detraves C, Poincloux R, Baricault L, Fournier D, Paquereau L (2003) Fungal lectin, XCL, is internalized via clathrin-dependent endocytosis and facilitates uptake of other molecules. *Eur J Cell Biol* 82(10):515–22
- Fujihashi M, Peapus DH, Kamiya N, Nagata Y, Miki K (2003) Crystal structure of fucose-specific lectin from *Aleuria aurantia* binding ligands at three of its five sugar recognition sites. *Biochemistry* 42(38):11093–9. doi:10.1021/bi034983z
- Fukumori F, Takeuchi N, Hagiwara T, Ohbayashi H, Endo T, Kochibe N, Nagata Y, Kobata A (1990) Primary structure of a fucose-specific lectin obtained from a mushroom, *Aleuria aurantia*. *J Biochem* 107(2):190–6
- Goldstein IJ, Winter HC (2007) Mushroom lectins. In: Kamerling JP (ed) *Comprehensive glycoscience: from chemistry to systems biology*, vol 3. Elsevier Ltd., Amsterdam

- Grahn E, Askarieh G, Holmner A, Tateno H, Winter HC, Goldstein IJ, Krengel U (2007) Crystal structure of the *Marasmius oreades* mushroom lectin in complex with a xenotransplantation epitope. *J Mol Biol* 369(3):710–21. doi:10.1016/j.jmb.2007.03.016
- Grahn EM, Winter HC, Tateno H, Goldstein IJ, Krengel U (2009) Structural characterization of a lectin from the mushroom *Marasmius oreades* in complex with the blood group B trisaccharide and calcium. *J Mol Biol* 390(3):457–66. doi:10.1016/j.jmb.2009.04.074
- Grigoriev IV, Cullen D, Goodwin SB, Hibbett D, Jeffries TW, Kubicek CP, Kuske C, Magnuson JK, Martin F, Spatafora JW, Tsang A, Baker SE (2011) Fueling the future with fungal genomics. *Mycologia* 2(3):192–209
- Guillot J, Konska G (1997) Lectins in higher fungi. *Biochem Syst Ecol* 25(3):203–230
- Guillot J, Giollant M, Damez M, Dusser M (1994) Involvement of fungal lectins in recognition between mushroom and tree during the early stages of mycorrhizae formation. *Acta Botanica Gallica* 141(4):443–447
- Gzogyán LA, Proskuryakov MT, Ievleva EV, Valueva TA (2005) Trypsin-like proteinases and trypsin inhibitors in fruiting bodies of higher fungi. *Appl Biochem Microbiol* 41(6):538–541
- Hamshou M, Smaghe G, Van Damme EJ (2007) Analysis of lectin concentrations in different *Rhizoctonia solani* strains. *Commun Agric Appl Biol Sci* 72(3):639–44
- Hamshou M, Smaghe G, Shahidi-Noghabi S, De Geyter E, Lannoo N, Van Damme EJ (2010a) Insecticidal properties of *Sclerotinia sclerotiorum* agglutinin and its interaction with insect tissues and cells. *Insect Biochem Mol Biol* 40(12):883–90. doi:10.1016/j.ibmb.2010.08.008
- Hamshou M, Van Damme EJ, Smaghe G (2010b) Entomotoxic effects of fungal lectin from *Rhizoctonia solani* towards *Spodoptera littoralis*. *Fungal Biology* 114(1):34–40
- Hamshou M, Van Damme EJ, Vandenborre G, Ghesquiere B, Trooskens G, Gevaert K, Smaghe G (2012) GalNAc/Gal-binding *Rhizoctonia solani* agglutinin has antiproliferative activity in *Drosophila melanogaster* S2 cells via MAPK and JAK/STAT signaling. *PLoS One* 7(4):e33680
- Hamshou M, Van Damme EJ, Caccia S, Cappelle K, Vandenborre G, Ghesquiere B, Gevaert K, Smaghe G (2013) High entomotoxicity and mechanism of the fungal GalNAc/Gal-specific *Rhizoctonia solani* lectin in pest insects. *J Insect Physiol* 59(3):295–305. doi:10.1016/j.jinsphys.2012.12.003
- Hassan MA, Rouf R, Tiralongo E, May TW, Tiralongo J (2015) Mushroom lectins: specificity, structure and bioactivity relevant to human disease. *Int J Mol Sci* 16(4):7802–38. doi:10.3390/ijms16047802
- Hazes B (1996) The (QxW)₃ domain: a flexible lectin scaffold. *Protein Sci* 5(8):1490–501. doi:10.1002/pro.5560050805
- Heim C, Hertzberg H, Butschi A, Bleuler-Martinez S, Aebi M, Deplazes P, Künzler M, Štefanić S (2015) Inhibition of *Haemonchus contortus* larval development by fungal lectins. *Parasites & Vectors* 8:425. doi:10.1186/s13071-015-1032-x
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, Huhndorf S, James T, Kirk PM, Lucking R, Thorsten Lumsch H, Lutzoni F, Matheny PB, McLaughlin DJ, Powell MJ, Redhead S, Schoch CL, Spatafora JW, Stalpers JA, Vilgalys R, Aime MC, Aptroot A, Bauer R, Begerow D, Benny GL, Castlebury LA, Crous PW, Dai YC, Gams W, Geiser DM, Griffith GW, Gueidan C, Hawksworth DL, Hestmark G, Hosaka K, Humber RA, Hyde KD, Ironside JE, Koljalg U, Kurtzman CP, Larsson KH, Lichtwardt R, Longcore J, Miadlikowska J, Miller A, Moncalvo JM, Mozley-Standridge S, Oberwinkler F, Parmasto E, Reeb V, Rogers JD, Roux C, Ryvarden L, Sampaio JP, Schussler A, Sugiyama J, Thorn RG, Tibell L, Untereiner WA, Walker C, Wang Z, Weir A, Weiss M, White MM, Winka K, Yao YJ, Zhang N (2007) A higher-level phylogenetic classification of the fungi. *Mycol Res* 111(Pt 5):509–47. doi:10.1016/j.mycres.2007.03.004
- Houser J, Komarek J, Kostlanova N, Cioci G, Varrot A, Kerr SC, Lahmann M, Balloy V, Fahy JV, Chignard M, Imbert A, Wimmerova M (2013) A soluble fucose-specific lectin from *Aspergillus fumigatus* conidia—structure, specificity and possible role in fungal pathogenicity. *PLoS One* 8(12):e83077. doi:10.1371/journal.pone.0083077
- Houser J, Komarek J, Cioci G, Varrot A, Imbert A, Wimmerova M (2015) Structural insights into *Aspergillus fumigatus* lectin specificity: AFL binding sites are functionally non-equivalent. *Acta Crystallogr D Biol Crystallogr* 71(Pt 3):442–53. doi:10.1107/S1399004714026595
- Hu Y, Zhan B, Keegan B, Yiu YY, Miller MM, Jones K, Aroian RV (2012) Mechanistic and single-dose in vivo therapeutic studies of Cry5B anthelmintic action against hookworms. *PLoS Negl Trop Dis* 6(11):e1900. doi:10.1371/journal.pntd.0001900
- Imamura K, Takeuchi H, Yabe R, Tateno H, Hirabayashi J (2011) Engineering of the Glycan-binding Specificity of *Agrocybe cylindracea* Galectin toward {α}(2,3)-linked Sialic Acid by Saturation Mutagenesis. *J Biochem* 150(5):545–52
- Inbar J, Chet I (1994) A newly isolated lectin from the plant pathogenic fungus *Sclerotium rolfsii*: purification, characterization and role in mycoparasitism. *Microbiology* 140(Pt 3):651–7
- Jaber K, Paquereau L, Fournier D, Haubruge E, Francis F (2006) Use of artificial diet system to assess the potential bio-insecticide effect of a fungal lectin from *Xerocomus chrysenteron* (XCL) on *Myzus persicae*. *Commun Agric Appl Biol Sci* 71(2 Pt B):497–505
- Jaber K, Francis F, Paquereau L, Fournier D, Haubruge E (2007) Effect of a fungal lectin from *Xerocomus chrysenteron* (XCL) on the biological parameters of aphids. *Commun Agric Appl Biol Sci* 72(3):629–38
- Jaber K, Cuartero Diaz G, Haubruge E, Francis F (2008) Investigation of carbohydrate binding property of a fungal lectin from *Xerocomus chrysenteron* and potential use on *Myzus persicae* aphid. *Commun Agric Appl Biol Sci* 73(3):629–38
- Jones JT, Haegeman A, Danchin EG, Gaur HS, Helder J, Jones MG, Kikuchi T, Manzanilla-Lopez R, Palomares-Rius JE, Wesemael WM, Perry RN (2013) Top 10 plant-parasitic nematodes in molecular plant pathology. *Mol Plant Pathol* 14(9):946–61. doi:10.1111/mpp.12057
- Kadirvelraj R, Grant OC, Goldstein IJ, Winter HC, Tateno H, Fadda E, Woods RJ (2011) Structure and binding analysis of *Polyporus squamosus* lectin in complex with the Neu5Ac{α}2-6Gal{β}1-4GlcNAc human-type influenza receptor. *Glycobiology* 21(7):973–84. doi:10.1093/glycob/cwr030
- Kellens JTC, Peumans W (1990) Developmental accumulation of lectin in *Rhizoctonia solani*: a potential role as a storage protein. *J Gen Microbiol* 136:2489–2495
- Kobayashi Y, Kawagishi H (2014) Fungal lectins: a growing family. *Methods Mol Biol* 1200:15–38. doi:10.1007/978-1-4939-1292-6_2
- Kobayashi Y, Tateno H, Dohra H, Moriwaki K, Miyoshi E, Hirabayashi J, Kawagishi H (2012) A novel core fucose-specific lectin from the mushroom *Pholiota squarrosa*. *J Biol Chem* 287(41):33973–82. doi:10.1074/jbc.M111.327692
- Kochibe N, Furukawa K (1980) Purification and properties of a novel fucose-specific hemagglutinin of *Aleuria aurantia*. *Biochemistry* 19(13):2841–6
- Koharudin LM, Viscomi AR, Jee JG, Ottonello S, Gronenborn AM (2008) The evolutionarily conserved family of cyanovirin-N homologs: structures and carbohydrate specificity. *Structure* 16(4):570–84. doi:10.1016/j.str.2008.01.015
- Kojima S, Iwahara A, Yanai H (2005) Inhibitor-assisted refolding of protease: a protease inhibitor as an intramolecular chaperone. *FEBS Lett* 579(20):4430–6. doi:10.1016/j.febslet.2005.06.083
- Kristan KC, Viero G, Dalla Serra M, Macek P, Anderluh G (2009) Molecular mechanism of pore formation by actinoporins. *Toxicon* 54(8):1125–34. doi:10.1016/j.toxicon.2009.02.026

- Künzler M (2015) Hitting the sweet spot: glycans as targets of fungal defense effector proteins. *Molecules* 20(5):8144–8167. doi:10.3390/molecules20058144
- Kuwabara N, Hu D, Tateno H, Makyio H, Hirabayashi J, Kato R (2013) Conformational change of a unique sequence in a fungal galectin from *Agrocybe cylindracea* controls glycan ligand-binding specificity. *FEBS Lett* doi:10.1016/j.febslet.2013.08.046
- Lacadena J, Alvarez-Garcia E, Carreras-Sangra N, Herrero-Galan E, Alegre-Cebollada J, Garcia-Ortega L, Onaderra M, Gavilanes JG, Martinez del Pozo A (2007) Fungal ribotoxins: molecular dissection of a family of natural killers. *FEMS Microbiol Rev* 31(2):212–37. doi:10.1111/j.1574-6976.2006.00063.x
- LeJambre LF, Windon RG, Smith WD (2008) Vaccination against *Haemonchus contortus*: performance of native parasite gut membrane glycoproteins in Merino lambs grazing contaminated pasture. *Vet Parasitol* 153(3–4):302–12. doi:10.1016/j.vetpar.2008.01.032
- Leonidas DD, Swamy BM, Hatzopoulos GN, Gonchigar SJ, Chachadi VB, Inamdar SR, Zographos SE, Oikonomakos NG (2007) Structural basis for the carbohydrate recognition of the *Sclerotium rolfsii* lectin. *J Mol Biol* 368(4):1145–61. doi:10.1016/j.jmb.2007.02.092
- Liu C, Zhao X, Xu XC, Li LR, Liu YH, Zhong SD, Bao JK (2008) Hemagglutinating activity and conformation of a lactose-binding lectin from mushroom *Agrocybe cylindracea*. *Int J Biol Macromol* 42(2):138–44. doi:10.1016/j.ijbiomac.2007.10.017
- Lopez-Otin C, Bond JS (2008) Proteases: Multifunctional Enzymes in Life and Disease. *J Biol Chem* 283(45):30433–30437
- Luan R, Liang Y, Chen Y, Liu H, Jiang S, Che T, Wong B, Sun H (2010) Opposing developmental functions of *Agrocybe aegerita* galectin (AAL) during mycelia differentiation. *Fungal Biol* 114(8):599–608. doi:10.1016/j.funbio.2010.05.001
- Mancheno JM, Tateno H, Goldstein IJ, Martinez-Ripoll M, Hermoso JA (2005) Structural analysis of the *Laetiporus sulphureus* hemolytic pore-forming lectin in complex with sugars. *J Biol Chem* 280(17):17251–9. doi:10.1074/jbc.M413933200
- Mancheno JM, Tateno H, Sher D, Goldstein IJ (2010) *Laetiporus sulphureus* lectin and aerolysin protein family. *Adv Exp Med Biol* 677:67–80
- Martin F, Aerts A, Ahren D, Brun A, Danchin EG, Duchaussoy F, Gibon J, Kohler A, Lindquist E, Pereda V, Salamov A, Shapiro HJ, Wuyts J, Blaudez D, Buee M, Brokstein P, Canback B, Cohen D, Courty PE, Coutinho PM, Delaruelle C, Detter JC, Deveau A, DiFazio S, Duplessis S, Fraissinet-Tachet L, Lucic E, Frey-Klett P, Fourrey C, Feussner I, Gay G, Grimwood J, Hoegger PJ, Jain P, Kilaru S, Labbe J, Lin YC, Legue V, Le Tacon F, Marmeisse R, Melayah D, Montanini B, Muratet M, Nehls U, Niculita-Hirzel H, Oudot-Le Secq MP, Peter M, Quesneville H, Rajashekar B, Reich M, Rouhier N, Schmutz J, Yin T, Chalot M, Henrissat B, Kues U, Lucas S, Van de Peer Y, Podila GK, Polle A, Pukkila PJ, Richardson PM, Rouze P, Sanders IR, Stajich JE, Tunlid A, Tuskan G, Grigoriev IV (2008) The genome of *Laccaria bicolor* provides insights into mycorrhizal symbiosis. *Nature* 452(7183):88–92. doi:10.1038/nature06556
- Marty-Detraves C, Francis F, Baricault L, Fournier D, Paquereau L (2004) Inhibitory action of a new lectin from *Xerocomus chrysenteron* on cell-substrate adhesion. *Mol Cell Biochem* 258(1–2):49–55
- Matsumura K, Higashida K, Hata Y, Kominami J, Nakamura-Tsuruta S, Hirabayashi J (2008) Comparative analysis of oligosaccharide specificities of fucose-specific lectins from *Aspergillus oryzae* and *Aleuria aurantia* using frontal affinity chromatography. *Anal Biochem* doi:10.1016/j.ab.2008.11.044
- Mo H, Winter HC, Goldstein IJ (2000) Purification and characterization of a Neu5Acalpha2-6Galbeta1-4Glc/GlcNAc-specific lectin from the fruiting body of the polypore mushroom *Polyporus squamosus*. *J Biol Chem* 275(14):10623–9
- Moran Y, Fredman D, Szczesny P, Grynberg M, Technau U (2012) Recurrent horizontal transfer of bacterial toxin genes to eukaryotes. *Mol Biol Evol* 29(9):2223–30. doi:10.1093/molbev/mss089
- Munn EA (1997) Rational design of nematode vaccines: hidden antigens. *Int J Parasitol* 27(4):359–66
- Nagata Y, Yamashita M, Honda H, Akabane J, Uehara K, Saito A, Sumisa F, Nishibori K, Oodaira Y (2005) Characterization, occurrence, and molecular cloning of a lectin from *Grifola frondosa*: jacalin-related lectin of fungal origin. *Biosci Biotechnol Biochem* 69(12):2374–80. doi:10.1271/bbb.69.2374
- Nakamura-Tsuruta S, Kominami J, Kuno A, Hirabayashi J (2006) Evidence that *Agaricus bisporus* agglutinin (ABA) has dual sugar-binding specificity. *Biochem Biophys Res Commun* 347(1):215–220
- Nowrousian M, Cebula P (2005) The gene for a lectin-like protein is transcriptionally activated during sexual development, but is not essential for fruiting body formation in the filamentous fungus *Sordaria macrospora*. *BMC Microbiol* 5:64. doi:10.1186/1471-2180-5-64
- Oda Y, Senaha T, Matsuno Y, Nakajima K, Naka R, Kinoshita M, Honda E, Furuta I, Kakehi K (2003) A new fungal lectin recognizing alpha(1-6)-linked fucose in the N-glycan. *J Biol Chem* 278(34):32439–47. doi:10.1074/jbc.M305181200
- Odani S, Tominaga K, Kondou S, Hori H, Koide T, Hara S, Isemura M, Tsunasawa S (1999) The inhibitory properties and primary structure of a novel serine proteinase inhibitor from the fruiting body of the basidiomycete, *Lentinus edodes*. *Eur J Biochem* 262(3):915–23
- Ogawa S, Nakajima E, Nagao H, Ohtoshi M, Ando A, Nagata Y (1998) Synthesis of a lectin in both mycelia and fruit bodies of the ascomycete mushroom *Aleuria aurantia*. *Biosci Biotechnol Biochem* 62(5):915–918. doi:10.1271/Bbb.62.915
- Ohm RA, de Jong JF, Lugones LG, Aerts A, Kothe E, Stajich JE, de Vries RP, Record E, Levasseur A, Baker SE, Bartholomew KA, Coutinho PM, Erdmann S, Fowler TJ, Gathman AC, Lombard V, Henrissat B, Knabe N, Kues U, Lilly WW, Lindquist E, Lucas S, Magnuson JK, Piumi F, Raudaskoski M, Salamov A, Schmutz J, Schwarze FW, vanKuyk PA, Horton JS, Grigoriev IV, Wosten HA (2010) Genome sequence of the model mushroom *Schizophyllum commune*. *Nat Biotechnol* 28(9):957–63. doi:10.1038/nbt.1643
- Olausson J, Tibell L, Jonsson BH, Pahlsson P (2008) Detection of a high affinity binding site in recombinant *Aleuria aurantia* lectin. *Glycoconj J* 25(8):753–62. doi:10.1007/s10719-008-9135-7
- Olausson J, Astrom E, Jonsson BH, Tibell LA, Pahlsson P (2010) Production and characterization of a monomeric and a single site form of *Aleuria aurantia* lectin. *Glycobiology* 21(1):34–44. doi:10.1093/glycob/cwq129
- Ota K, Butala M, Viero G, Dalla Serra M, Sepcic K, Macek P (2014) Fungal MACPF-like proteins and aegerolysins: bi-component pore-forming proteins? *Subcell Biochem* 80:271–91. doi:10.1007/978-94-017-8881-6_14
- Paaventhnan P, Joseph JS, Seow SV, Vaday S, Robinson H, Chua KY, Kolatkar PR (2003) A 1.7 Å structure of Fve, a member of the new fungal immunomodulatory protein family. *J Mol Biol* 332(2):461–70
- Paschinger K, Wilson IB (2015) Two types of galactosylated fucose motifs are present on N-glycans of *Haemonchus contortus*. *Glycobiology* 25(6):585–90. doi:10.1093/glycob/cwv015
- Paschinger K, Rendic D, Wilson IB (2009) Revealing the anti-HRP epitope in *Drosophila* and *Caenorhabditis*. *Glycoconj J* 26(3):385–95. doi:10.1007/s10719-008-9155-3
- Paschinger K, Razzazi-Fazeli E, Furukawa K, Wilson IB (2011) Presence of galactosylated core fucose on N-glycans in the planaria *Dugesia japonica*. *J Mass Spectrom* 46(6):561–7. doi:10.1002/jms.1925
- Peumans WJ, Van Damme EJ (1995) Lectins as plant defense proteins. *Plant Physiol* 109(2):347–52
- Peumans WJ, Fouquaert E, Jauneau A, Rouge P, Lannoo N, Hamada H, Alvarez R, Devreese B, Van Damme EJ (2007) The liverwort

- Marchantia polymorpha* expresses orthologs of the fungal *Agaricus bisporus* agglutinin family. *Plant Physiol* 144(2):637–47. doi:10.1104/pp.106.087437
- Plaza DF, Lin CW, van der Velden NS, Aebi M, Künzler M (2014) Comparative transcriptomics of the model mushroom *Coprinopsis cinerea* reveals tissue-specific armories and a conserved circuitry for sexual development. *BMC Genomics* 15:492. doi:10.1186/1471-2164-15-492
- Pohleven J, Obermajer N, Sabotić J, Anžlovar S, Sepčić K, Kos J, Kralj B, Štrukelj B, Brzin J (2009) Purification, characterization and cloning of a ricin B-like lectin from mushroom *Clitocybe nebularis* with antiproliferative activity against human leukemic T cells. *Biochim Biophys Acta Gen Subj* 1790(3):173–181. doi:10.1016/j.bbagen.2008.11.006
- Pohleven J, Brzin J, Vrabec L, Leonardi A, Cokl A, Štrukelj B, Kos J, Sabotić J (2011) Basidiomycete *Clitocybe nebularis* is rich in lectins with insecticidal activities. *Appl Microbiol Biotechnol* 91(4):1141–8. doi:10.1007/s00253-011-3236-0
- Pohleven J, Renko M, Magister S, Smith DF, Künzler M, Štrukelj B, Turk D, Kos J, Sabotić J (2012) Bivalent carbohydrate binding is required for biological activity of *Clitocybe nebularis* lectin (CNL), the N, N'-diacetyllactosediamine (GalNAcbeta1-4GlcNAc, LacdiNAc)-specific lectin from basidiomycete *C. nebularis*. *J Biol Chem* 287(13):10602–12. doi:10.1074/jbc.M111.317263
- Presant CA, Kornfeld S (1972) Characterization of the cell surface receptor for the *Agaricus bisporus* hemagglutinin. *J Biol Chem* 247(21):6937–45
- Rabinovich GA, Toscano MA, Jackson SS, Vasta GR (2007) Functions of cell surface galectin-glycoprotein lattices. *Curr Opin Struct Biol* 17(5):513–20. doi:10.1016/j.sbi.2007.09.002
- Rawlings ND, Waller M, Barrett AJ, Bateman A (2014) MEROPS: the database of proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Res* 42(Database issue):D503–9. doi:10.1093/nar/gkt953
- Rempel BP, Winter HC, Goldstein IJ, Hindsgaul O (2002) Characterization of the recognition of blood group B trisaccharide derivatives by the lectin from *Marasmius oreades* using frontal affinity chromatography-mass spectrometry. *Glycoconj J* 19(3):175–80. doi:10.1023/A:1024297623445
- Renko M, Sabotić J, Mihelić M, Brzin J, Kos J, Turk D (2010) Versatile loops in mycocybins inhibit three protease families. *J Biol Chem* 285(1):308–16. doi:10.1074/jbc.M109.043331
- Renko M, Sabotić J, Turk D (2012) β -Trefol inhibitors—from the work of Kunitz onward. *Biol Chem* 393(10):1043. doi:10.1515/hsz-2012-0159
- Rosen S, Ek B, Rask L, Tunlid A (1992) Purification and characterization of a surface lectin from the nematode-trapping fungus *Arthrobotrys oligospora*. *J Gen Microbiol* 138(12):2663–72
- Rosen S, Bergstrom J, Karlsson KA, Tunlid A (1996a) A multispecific saline-soluble lectin from the parasitic fungus *Arthrobotrys oligospora*—similarities in the binding specificities compared with a lectin from the mushroom *Agaricus bisporus*. *Eur J Biochem* 238(3):830–837. doi:10.1111/j.1432-1033.1996.0830w.x
- Rosen S, Kata M, Persson Y, Lipniunas PH, Wikstrom M, Van Den Hondel CAMJJ, Van Den Brink JM, Rask L, Heden LO, Tunlid A (1996b) Molecular characterization of a saline-soluble lectin from a parasitic fungus—extensive sequence similarities between fungal lectins. *Eur J Biochem* 238(3):822–829. doi:10.1111/j.1432-1033.1996.0822w.x
- Rosen S, Sjollem K, Veenhuis M, Tunlid A (1997) A cytoplasmic lectin produced by the fungus *Arthrobotrys oligospora* functions as a storage protein during saprophytic and parasitic growth. *Microbiol-Uk* 143:2593–2604
- Sabotić J, Kos J (2012) Microbial and fungal protease inhibitors—current and potential applications. *Appl Microbiol Biotechnol* 93(4):1351–75. doi:10.1007/s00253-011-3834-x
- Sabotić J, Gaser D, Rogelj B, Gruden K, Štrukelj B, Brzin J (2006) Heterogeneity in the cysteine protease inhibitor cliticypin gene family. *Biol Chem* 387(12):1559–1566. doi:10.1515/bc.2006.194
- Sabotić J, Galeša K, Popović T, Leonardi A, Brzin J (2007a) Comparison of natural and recombinant cliticypins, the fungal cysteine protease inhibitors. *Protein Expr Purif* 53(1):104–111. doi:10.1016/j.pep.2006.11.015
- Sabotić J, Trček T, Popović T, Brzin J (2007b) Basidiomycetes harbour a hidden treasure of proteolytic diversity. *J Biotechnol* 128(2):297–307. doi:10.1016/j.jbiotec.2006.10.006
- Sabotić J, Popović T, Puizdar V, Brzin J (2009) Macrocybins, a family of cysteine protease inhibitors from the basidiomycete *Macrolepiota procera*. *FEBS J* 276(16):4334–45. doi:10.1111/j.1742-4658.2009.07138.x
- Sabotić J, Kilaru S, Budić M, Gašparić Buh M, Gruden K, Bailey AM, Foster GD, Kos J (2011) Protease inhibitors cliticypin and macrocybin are differentially expressed within basidiomycete fruiting bodies. *Biochimie* 93(10):1685–93. doi:10.1016/j.biochi.2011.05.034
- Sabotić J, Bleuler-Martinez S, Renko M, Avanzo Caglić P, Kallert S, Štrukelj B, Turk D, Aebi M, Kos J, Künzler M (2012) Structural basis of trypsin inhibition and entomotoxicity of cospin, serine protease inhibitor involved in defense of *Coprinopsis cinerea* fruiting bodies. *J Biol Chem* 287(6):3898–907. doi:10.1074/jbc.M111.285304
- Sasakawa H, Yoshinaga S, Kojima S, Tamura A (2002) Structure of POIA1, a homologous protein to the propeptide of subtilisin: implication for protein foldability and the function as an intramolecular chaperone. *J Mol Biol* 317(1):159–67. doi:10.1006/jmbi.2002.5412
- Schubert M, Bleuler-Martinez S, Buttschi A, Wälti MA, Egloff P, Stutz K, Yan S, Wilson IBH, Hengartner MO, Aebi M, Allain FHT, Künzler M (2012) Plasticity of the β -trefol protein fold in the recognition and control of invertebrate predators and parasites by a fungal defence system. *PLoS Pathog* 8(5):e1002706. doi:10.1371/journal.ppat.1002706
- Sharon N, Lis H (2004) History of lectins: from hemagglutinins to biological recognition molecules. *Glycobiology* 14(11):53R–62R. doi:10.1093/glycob/cwh122
- Shi XZ, Wang L, Xu S, Zhang XW, Zhao XF, Vasta GR, Wang JX (2014) A galectin from the kuruma shrimp (*Marsupenaeus japonicus*) functions as an opsonin and promotes bacterial clearance from hemolymph. *PLoS One* 9(3):e91794. doi:10.1371/journal.pone.0091794
- Shimokawa M, Fukudome A, Yamashita R, Minami Y, Yagi F, Tateno H, Hirabayashi J (2012) Characterization and cloning of GNA-like lectin from the mushroom *Marasmius oreades*. *Glycoconj J* 29(7):457–65. doi:10.1007/s10719-012-9401-6
- Shinohara Y, Hasegawa Y, Kaku H, Shibuya N (1997) Elucidation of the mechanism enhancing the avidity of lectin with oligosaccharides on the solid phase surface. *Glycobiology* 7(8):1201–8
- Skamnaki VT, Peumans WJ, Kantsadi AL, Cubeta MA, Plas K, Pakala S, Zographos SE, Smaghe G, Nierman WC, Van Damme EJ, Leonidas DD (2013) Structural analysis of the *Rhizoctonia solani* agglutinin reveals a domain-swapping dimeric assembly. *FEBS J* 280(8):1750–63. doi:10.1111/febs.12190
- Šmid I, Gruden K, Buh Gašparić M, Koruza K, Petek M, Pohleven J, Brzin J, Kos J, Žel J, Sabotić J (2013) Inhibition of the growth of Colorado potato beetle larvae by macrocybins, protease inhibitors from the parasol mushroom. *J Agric Food Chem* 61(51):12499–509. doi:10.1021/jf403615f
- Šmid I, Rotter A, Gruden K, Brzin J, Buh Gašparić M, Kos J, Žel J, Sabotić J (2015) Cliticypin, a fungal cysteine protease inhibitor, exerts its insecticidal effect on Colorado potato beetle larvae by inhibiting their digestive cysteine proteases. *Pestic Biochem Physiol* 122:59–66. doi:10.1016/j.pestbp.2014.12.022
- Spiteller P (2015) Chemical ecology of fungi. *Nat Prod Rep* 32(7):971–993. doi:10.1039/c4np00166d

- Stajich JE, Wilke SK, Ahren D, Au CH, Birren BW, Borodovsky M, Burns C, Canback B, Casselton LA, Cheng CK, Deng J, Dietrich FS, Fargo DC, Farman ML, Gathman AC, Goldberg J, Guigo R, Hoegger PJ, Hooker JB, Huggins A, James TY, Kamada T, Kilaru S, Kodira C, Kues U, Kupfer D, Kwan HS, Lomsadze A, Li W, Lilly WW, Ma LJ, Mackey AJ, Manning G, Martin F, Muraguchi H, Natvig DO, Palmerini H, Ramesh MA, Rehmeier CJ, Roe BA, Shenoy N, Stanke M, Ter-Hovhannisyan V, Tunlid A, Velagapudi R, Vision TJ, Zeng Q, Zolan ME, Pukkila PJ (2010) Insights into evolution of multicellular fungi from the assembled chromosomes of the mushroom *Coprinopsis cinerea* (*Coprinus cinereus*). *Proc Natl Acad Sci U S A* 107(26):11889–94. doi:10.1073/pnas.1003391107
- Stutz K, Kaech A, Aebi M, Künzler M, Hengartner MO (2015) Disruption of the *C. elegans* intestinal brush border by the fungal lectin CCL2 phenocopies dietary lectin toxicity in mammals. *PLoS One* 10(6):e0129381. doi:10.1371/journal.pone.0129381
- Sudakevitz D, Imberty A, Gilboa-Garber N (2002) Production, properties and specificity of a new bacterial L-fucose- and D-arabinose-binding lectin of the plant aggressive pathogen *Ralstonia solanacearum*, and its comparison to related plant and microbial lectins. *J Biochem* 132(2):353–8
- Sulzenbacher G, Roig-Zamboni V, Peumans WJ, Rouge P, Van Damme EJ, Bourne Y (2010) Crystal structure of the GalNAc/Gal-specific agglutinin from the phytopathogenic ascomycete *Sclerotinia sclerotiorum* reveals novel adaptation of a beta-trefoil domain. *J Mol Biol* 400(4):715–23. doi:10.1016/j.jmb.2010.05.038
- Suzuki T, Sugiyama K, Hirai H, Ito H, Morita T, Dohra H, Murata T, Usui T, Tateno H, Hirabayashi J, Kobayashi Y, Kawagishi H (2012) Mannose-specific lectin from the mushroom *Hygrophorus russula*. *Glycobiology* 22(5):616–29. doi:10.1093/glycob/cwr187
- Swamy BM, Bhat AG, Hegde GV, Naik RS, Kulkarni S, Inamdar SR (2004) Immunolocalization and functional role of *Sclerotium rolfii* lectin in development of fungus by interaction with its endogenous receptor. *Glycobiology* 14(11):951–7. doi:10.1093/glycob/cwh130
- Tateno H, Goldstein IJ (2003) Molecular cloning, expression, and characterization of novel hemolytic lectins from the mushroom *Laetiporus sulphureus*, which show homology to bacterial toxins. *J Biol Chem* 278(42):40455–63. doi:10.1074/jbc.M306836200
- Tateno H, Goldstein IJ (2004) Partial identification of carbohydrate-binding sites of a Gal α 1,3Gal β 1,4GlcNAc-specific lectin from the mushroom *Marasmius oreades* by site-directed mutagenesis. *Arch Biochem Biophys* 427(1):101–9. doi:10.1016/j.abb.2004.04.013
- Tateno H, Winter HC, Goldstein IJ (2004) Cloning, expression in *Escherichia coli* and characterization of the recombinant Neu5A α 2,6Gal β 1,4GlcNAc-specific high-affinity lectin and its mutants from the mushroom *Polyporus squamosus*. *Biochem J* 382(Pt 2):667–75. doi:10.1042/BJ20040391
- Tateno H, Nakamura-Tsuruta S, Hirabayashi J (2009) Comparative analysis of core-fucose-binding lectins from *Lens culinaris* and *Pisum sativum* using frontal affinity chromatography. *Glycobiology* 19(5):527–36. doi:10.1093/glycob/cwp016
- Taylor JW, Ellison CE (2010) Mushrooms: morphological complexity in the fungi. *Proc Natl Acad Sci U S A* 107(26):11655–6. doi:10.1073/pnas.1006430107
- Teichert I, Nowrousian M, Poggeler S, Kuck U (2014) The filamentous fungus *Sordaria macrospora* as a genetic model to study fruiting body development. *Adv Genet* 87:199–244. doi:10.1016/B978-0-12-800149-3.00004-4
- Terra WR, Ferreira C (2005) 4.5—Biochemistry of digestion. In: Lawrence IG, Kostas I, Sarjeet SG (eds) *Comprehensive Molecular Insect Science*. Elsevier, Amsterdam, pp 171–224
- Tian Y, Zhang K (2005) Purification and characteristic of proteinase inhibitor GLPIA2 from *Ganoderma lucidum* by submerged fermentation. *Se Pu* 23(3):267–9
- Titz A, Butschi A, Henrissat B, Fan YY, Hennet T, Razzazi-Fazeli E, Hengartner MO, Wilson IB, Künzler M, Aebi M (2009) Molecular basis for galactosylation of core fucose residues in invertebrates: identification of *Caenorhabditis elegans* N-glycan core { α }1,6-fucoside { β }1,4-galactosyltransferase GALT-1 as a member of a novel glycosyltransferase family. *J Biol Chem* 284(52):36223–33. doi:10.1074/jbc.M109.058354
- Trigueros V, Lougarre A, Ali-Ahmed D, Rahbe Y, Guillot J, Chavant L, Fournier D, Paquereau L (2003) *Xerocomus chrysenteron* lectin: identification of a new pesticidal protein. *Biochim Biophys Acta* 1621(3):292–8
- Ueda H, Kojima K, Saitoh T, Ogawa H (1999a) Interaction of a lectin from *Psathyrella velutina* mushroom with N-acetylneuraminic acid. *FEBS Lett* 448(1):75–80
- Ueda H, Saitoh T, Kojima K, Ogawa H (1999b) Multi-specificity of a *Psathyrella velutina* mushroom lectin: heparin/pectin binding occurs at a site different from the N-acetylglucosamine/N-acetylneuraminic acid-specific site. *J Biochem* 126(3):530–7
- Ueda H, Matsumoto H, Takahashi N, Ogawa H (2002) *Psathyrella velutina* mushroom lectin exhibits high affinity toward sialoglycoproteins possessing terminal N-acetylneuraminic acid α -2,3-linked to penultimate galactose residues of trisialyl N-glycans. comparison with other sialic acid-specific lectins. *J Biol Chem* 277(28):24916–25. doi:10.1074/jbc.M110727200
- Ueda H, Takahashi N, Ogawa H (2003) *Psathyrella velutina* lectin as a specific probe for N-acetylneuraminic acid in glycoconjugates. *Methods Enzymol* 363:77–90. doi:10.1016/S0076-6879(03)01044-9
- Urban JF Jr, Hu Y, Miller MM, Scheib U, Yiu YY, Aroian RV (2013) *Bacillus thuringiensis*-derived Cry5B has potent anthelmintic activity against *Ascaris suum*. *PLoS Negl Trop Dis* 7(6):e2263. doi:10.1371/journal.pntd.0002263
- Van Damme EJ, Nakamura-Tsuruta S, Hirabayashi J, Rouge P, Peumans WJ (2007) The *Sclerotinia sclerotiorum* agglutinin represents a novel family of fungal lectins remotely related to the *Clostridium botulinum* non-toxin haemagglutinin HA33/A. *Glycoconj J* 24(2–3):143–56. doi:10.1007/s10719-006-9022-z
- Varki A (2009) *Essentials of glycobiology*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y
- Varrot A, Basheer SM, Imberty A (2013) Fungal lectins: structure, function and potential applications. *Curr Opin Struct Biol* 23(5):678–85. doi:10.1016/j.sbi.2013.07.007
- Vetter J (2000) Trypsin inhibitory activity of basidiomycetous mushrooms. *Eur Food Res Technol* 211:346–348
- Vishwanathreddy H, Bhat GG, Inamdar SR, Gudihal RK, Swamy BM (2014) *Sclerotium rolfii* lectin exerts insecticidal activity on *Spodoptera litura* larvae by binding to membrane proteins of midgut epithelial cells and triggering caspase-3-dependent apoptosis. *Toxicon* 78:47–57. doi:10.1016/j.toxicon.2013.11.012
- Walser PJ, Haebel PW, Künzler M, Sargent D, Kues U, Aebi M, Ban N (2004) Structure and functional analysis of the fungal galectin CGL2. *Structure* 12(4):689–702. doi:10.1016/j.str.2004.03.002
- Walski T, Van Damme EJ, Smagghe G (2014) Penetration through the peritrophic matrix is a key to lectin toxicity against *Tribolium castaneum*. *J Insect Physiol* 70:94–101. doi:10.1016/j.jinsphys.2014.09.004
- Walt MA, Walser PJ, Thore S, Grunler A, Bednar M, Künzler M, Aebi M (2008) Structural basis for chitotetraose coordination by CGL3, a novel galectin-related protein from *Coprinopsis cinerea*. *J Mol Biol* 379(1):146–59. doi:10.1016/j.jmb.2008.03.062
- Wang HX, Ng TB, Ooi VEC (1998) Lectins from mushrooms. *Mycol Res* 102:897–906
- Wang M, Trigueros V, Paquereau L, Chavant L, Fournier D (2002) Proteins as active compounds involved in insecticidal activity of mushroom fruitbodies. *J Econ Entomol* 95(3):603–7

- Wimmerova M, Mitchell E, Sanchez JF, Gautier C, Imberty A (2003) Crystal structure of fungal lectin: six-bladed beta-propeller fold and novel fucose recognition mode for *Aleuria aurantia* lectin. *J Biol Chem* 278(29):27059–67. doi:10.1074/jbc.M302642200
- Winter HC, Mostafapour K, Goldstein IJ (2002) The mushroom *Marasmius oreades* lectin is a blood group type B agglutinin that recognizes the Gal α 1,3Gal and Gal α 1,3Gal β 1,4GlcNAc porcine xenotransplantation epitopes with high affinity. *J Biol Chem* 277(17):14996–5001. doi:10.1074/jbc.M200161200
- Wohlschlager T, Butschi A, Zurfluh K, Vonesch SC, Auf dem Keller U, Gehrig P, Bleuler-Martinez S, Hengartner MO, Aebi M, Künzler M (2011) Nematotoxicity of *Marasmius Oreades* agglutinin (MOA) depends on glycolipid-binding and cysteine protease activity. *J Biol Chem* 286:30337–43. doi:10.1074/jbc.M111.258202
- Wohlschlager T, Butschi A, Grassi P, Sutov G, Gauss R, Hauck D, Schmieder SS, Knobel M, Titz A, Dell A, Haslam SM, Hengartner MO, Aebi M, Künzler M (2014) Methylated glycans as conserved targets of animal and fungal innate defense. *Proc Natl Acad Sci U S A* 111(27):E2787–96. doi:10.1073/pnas.1401176111
- Wong JH, Ng TB, Cheung RC, Ye XJ, Wang HX, Lam SK, Lin P, Chan YS, Fang EF, Ngai PH, Xia LX, Ye XY, Jiang Y, Liu F (2010) Proteins with antifungal properties and other medicinal applications from plants and mushrooms. *Appl Microbiol Biotechnol* 87(4):1221–35. doi:10.1007/s00253-010-2690-4
- Xu X, Yan H, Chen J, Zhang X (2011) Bioactive proteins from mushrooms. *Biotechnol Adv* 29(6):667–74. doi:10.1016/j.biotechadv.2011.05.003
- Yagi F, Miyamoto M, Abe T, Minami Y, Tadera K, Goldstein IJ (1997) Purification and carbohydrate-binding specificity of *Agrocybe cylindracea* lectin. *Glycoconj J* 14(2):281–8
- Yagi F, Hiroyama H, Kodama S (2001) *Agrocybe cylindracea* lectin is a member of the galectin family. *Glycoconj J* 18(10):745–9
- Yan S, Bleuler-Martinez S, Plaza Gutierrez DF, Künzler M, Aebi M, Joachim A, Razzazi-Fazeli E, Jantsch V, Geyer R, Wilson IB, Paschinger K (2012) Galactosylated fucose epitopes in nematodes: increased expression in a *Caenorhabditis* mutant associated with altered lectin sensitivity and occurrence in parasitic species. *J Biol Chem* doi:10.1074/jbc.M112.353128
- Yan S, Brecker L, Jin C, Titz A, Dragosits M, Karlsson N, Jantsch V, Wilson IB, Paschinger K (2015) Bisecting galactose as a feature of N-glycans of wild-type and mutant *Caenorhabditis elegans*. *Mol Cell Proteomics* 14(8):2111–25. doi:10.1074/mcp.M115.049817
- Yang N, Liang Y, Xiang Y, Zhang Y, Sun H, Wang DC (2005a) Crystallization and preliminary crystallographic studies of an antitumor lectin from the edible mushroom *Agrocybe aegerita*. *Protein Pept Lett* 12(7):705–7
- Yang N, Tong X, Xiang Y, Zhang Y, Liang Y, Sun H, Wang DC (2005b) Molecular character of the recombinant antitumor lectin from the edible mushroom *Agrocybe aegerita*. *J Biochem* 138(2):145–50. doi:10.1093/jb/mvi109
- Yang N, Tong X, Xiang Y, Zhang Y, Sun H, Wang DC (2005c) Crystallization and preliminary crystallographic studies of the recombinant antitumor lectin from the edible mushroom *Agrocybe aegerita*. *Biochim Biophys Acta* 1751(2):209–12. doi:10.1016/j.bbapap.2005.06.003
- Yang N, Li DF, Feng L, Xiang Y, Liu W, Sun H, Wang DC (2009) Structural basis for the tumor cell apoptosis-inducing activity of an antitumor lectin from the edible mushroom *Agrocybe aegerita*. *J Mol Biol* 387(3):694–705. doi:10.1016/j.jmb.2009.02.002
- Yeates GW, Bongers T, De Goede RG, Freckman DW, Georgieva SS (1993) Feeding habits in soil nematode families and genera—an outline for soil ecologists. *J Nematol* 25(3):315–31
- Yin Y, Yu G, Chen Y, Jiang S, Wang M, Jin Y, Lan X, Liang Y, Sun H (2012) Genome-wide transcriptome and proteome analysis on different developmental stages of *Cordyceps militaris*. *PLoS One* 7(12):e51853. doi:10.1371/journal.pone.0051853
- Yu L, Fernig DG, Smith JA, Milton JD, Rhodes JM (1993) Reversible inhibition of proliferation of epithelial cell lines by *Agaricus bisporus* (edible mushroom) lectin. *Cancer Res* 53(19):4627–32
- Yu LG, Fernig DG, White MR, Spiller DG, Appleton P, Evans RC, Grierson I, Smith JA, Davies H, Gerasimenko OV, Petersen OH, Milton JD, Rhodes JM (1999) Edible mushroom (*Agaricus bisporus*) lectin, which reversibly inhibits epithelial cell proliferation, blocks nuclear localization sequence-dependent nuclear protein import. *J Biol Chem* 274(8):4890–9
- Zhang B, Palcic MM, Mo H, Goldstein IJ, Hindsgaul O (2001) Rapid determination of the binding affinity and specificity of the mushroom *Polyporus squamosus* lectin using frontal affinity chromatography coupled to electrospray mass spectrometry. *Glycobiology* 11(2):141–7
- Zhang GQ, Zhang QP, Sun Y, Tian YP, Zhou ND (2012) Purification of a novel pepsin inhibitor from *Coriolus versicolor* and its biochemical properties. *J Food Sci* 77(3):C293–7. doi:10.1111/j.1750-3841.2011.02581.x
- Zuchowski J, Grzywnowicz K (2006) Partial purification of proteinase K inhibitors from liquid-cultured mycelia of the white rot basidiomycete *Trametes versicolor*. *Curr Microbiol* 53(4):259–64
- Zuchowski J, Jaszek M, Grzywnowicz K (2009) Novel trypsin inhibitors from the white rot fungus *Abortiporus biennis*. Partial purification and characterization. *Biochemistry (Mosc)* 74(2):226–30
- Žurga S, Pohleven J, Renko M, Bleuler-Martinez S, Sosnowski P, Turk D, Künzler M, Kos J, Sabotič J (2014) A novel beta-trefoil lectin from the parasol mushroom (*Macrolepiota procera*) is nematotoxic. *FEBS J* 281:3489–3506. doi:10.1111/febs.12875
- Žurga S, Pohleven J, Kos J, Sabotič J (2015) beta-Trefoil structure enables interactions between lectins and protease inhibitors that regulate their biological functions. *J Biochem* 158(1):83–90. doi:10.1093/jb/mvv025