



Environmental pollution and COVID-19 outbreak: insights from Germany

Bilal¹ · Muhammad Farhan Bashir² · Maroua Benghouli³ · Umar Numan² · Awais Shakoor⁴ · Bushra Komal⁵ · Muhammad Adnan Bashir⁶ · Madiha Bashir⁷ · Duoqiao Tan¹

Received: 13 June 2020 / Accepted: 27 July 2020 / Published online: 3 August 2020
© Springer Nature B.V. 2020

Abstract

The impact of environmental pollutants and climate indicators on the outbreak of COVID-19 has gained considerable attention in the recent literature. However, specific investigation of industrial economies like Germany is not available. This provides us motivation to examine the association between environmental pollutants, climate indicators and the COVID-19 cases, recoveries, and deaths in Germany using daily data from February 24, 2020, to July 02, 2020. The correlation analysis and wavelet transform coherence (WTC) approach are the analytical tools, which are used to explore the association between variables included in the study. Our findings indicate that PM_{2.5}, O₃, and NO₂ have a significant relationship with the outbreak of COVID-19. In addition, temperature is the only significant climate indicator which has significant correlation with the spread of COVID-19. Finally, PM₁₀, humidity, and environmental quality index have a significant relationship only with the active cases from COVID-19 pandemic. Our findings conclude that Germany's successful response to COVID-19 is attributed to environmental legislation and the medical care system, which oversaw significant overhaul after the SARS and MERS outbreaks. The current study implicates that other industrial economies, especially European economies, that are still facing COVID-19 outbreak can follow the German model for pandemic response.

Keywords COVID-19 · Germany · Environmental pollution

Introduction

The World Health Organization (WHO) declared the outbreak of the novel coronavirus COVID-19 as a global pandemic on

March 11, 2020, as it became a global health threat (Organization 2020). The prevailing COVID-19 pandemic is spreading at a continuous pace with 12,507,849 confirmed cases as of July 12, 2020, worldwide. The European region

✉ Muhammad Farhan Bashir
farhan.paks@csu.edu.cn; farhan.paks89@gmail.com

Bilal
bilal@hbue.edu.cn

Maroua Benghouli
benghouliarwa@gmail.com

Umar Numan
umamuman@csu.edu.cn

Awais Shakoor
awais.shakoor@udl.cat

Bushra Komal
bushrakomal@hotmail.com

Muhammad Adnan Bashir
adnanbashir2034@gmail.com

Madiha Bashir
madihacheema16@gmail.com

Duoqiao Tan
tanduoqiao@hbue.edu.cn

¹ Hubei University of Economics, Wuhan, Hubei, People's Republic of China

² School of Business, Central South University, Changsha, Hunan, People's Republic of China

³ Anadolu University, Eskişehir, Turkey

⁴ Department of Environment and Soil Sciences, University of Lleida, Avinguda Alcalde Rovira Roure 191, 25198 Lleida, Spain

⁵ Business School, University of International Business and Economics, Beijing, People's Republic of China

⁶ School of Economics, Nankai University, Tianjin, People's Republic of China

⁷ Education Department, Government of The Punjab, Sialkot, Pakistan

was one of the worst hits by COVID-19 pandemic as several European countries, including Germany and France, reported imported cases of COVID-19 from travelers returning from Italy (Fattorini and Regoli 2020) and Spain (Ceylan 2020). The high infectious rate of COVID-19 in Europe has posed significant challenges for the health system across the globe as epidemiological surveillance capabilities played a key role in detecting COVID-19 cases in European countries. Early research suggests that person-to-person infectious transmission from COVID-19 requires up to two weeks of incubation time (Dutheil et al. 2020). However, the current situation requires drastic measures to protect the further spread of novel coronavirus. Currently, only experimental treatments are available; therefore, effective planning and upgrading health facilities are the only options to control the spread of COVID-19.

Germany is a leading industrial economy in western Europe with one of the highest GDP per capita income in the entire world (Bank 2020). Germany is also the front-runner in adopting clean energy initiatives, e.g., renewable energy significantly increased from 6% in 2000 to 34% in 2016 (Mez 2020). Germany plans to expand this share by up to 95% until the year 2050. Such initiatives have drastically increased environmental quality and have reduced air pollution in Germany. Germany also has one of the advanced healthcare systems among European countries with its proactive healthcare strategies, such as early-stage diagnosis of COVID-19 patients, extensive testing, and availability of medical facilities ensured limiting COVID-19 outbreak. Germany, until July 12, 2020, reported 199,802 confirmed cases but, unlike other epicenters of COVID-19, had the lowest death ratio despite having 21% of the total population over the age of 65 (Bank 2020). The successful response of Germany to the COVID-19 pandemic motivates us to study Germany to provide policy implications for other countries.

The current study investigated the impact of air pollutants and climate indicators' association with the COVID-19 pandemic in Germany. Our study is based on the fact that environmental and pollution indicators are crucial in the outbreak of COVID-19 pandemic (Bashir et al. 2020a). Since the late 1970s, Germany began implementing structural reforms to combat environmental pollution (Mez 2020) i.e., to help the East German government to reduce sulfur dioxide, nitrogen dioxide, and mercury by providing innovative coal-burning technology to filter mercury and hydrocarbons in contaminated water (O'Riordan 1989). Because of these policy changes, Germany has become a global front-runner in green technology with a \$400 billion of domestic clean energy industry. Also, domestic policy initiatives such as "Energiewende" have been implemented to battle chemical and air pollution in Germany.

Our research contributes in the economic literature in two ways. First, we are the first study to examine the effect of environmental pollutants in COVID-19 pandemic in an

industrial economy like Germany. Current literature is mainly focused on the impact of air pollutants (Anderson et al. 2013) in industrial economies (Bashir et al. 2020b; Cheung et al. 2020). The premise of high-pollution in industrial economies is vital because the air and environmental pollution significantly affect the mortality rate from COVID-19 (Fofana et al. 2020; Lalive et al. 2018; Popp 2006). Analytical structure of prior environmental pollutant studies is mainly based on correlation analysis and the results suffer from data endogeneity issues (Kessler et al. 2001; Schulz et al. 2007; Zijlema et al. 2016; Zimmermann et al. 2003). Gehrsitz (2017), one of the few exceptions to investigate the importance of environmental pollutants in Germany, suggested that low emissions zones have overseen 4–8% improvement in air quality in Germany. However, he recommended that these environmental changes are too small to carry a significant effect over human health. The current study adopts a different approach from existing literature by utilizing wavelet transform coherence (WTC) approach to identify the exact channels through which the COVID-19 outbreak is influenced by pollution indicators.

The current study suggests that an advanced healthcare system and institutional reforms can significantly reduce the impact of environmental and air pollutants with COVID-19 pandemic. In recent past, German authorities have introduced significant organizational reforms and invested in medical care facilities to improve medical facilities for unforeseen medical emergencies such as COVID-19. Our findings indicate that well-established state institutions can moderate the impact of air pollutants, and industrial and emerging economies should focus on introducing policies to reduce environmental pollution and the medical care system.

The rest of the sections as follows: the "Air and environmental pollution in Germany" section provides literature review about environmental pollutants in Germany; the "Research methodology" section presents research and data analysis methodology; the "Results and discussion" section narrates empirical results and lastly, the "Conclusion" section provides conclusion and policy recommendations.

Air and environmental pollution in Germany

Air pollution is a pollutant that enters the ambient air, instigated by an individual or blend of agents, i.e., chemical, radioactive, or biological matter. Existing research indicates that 82% of European Union residents are exposed to air pollution (Gehrsitz 2017). According to EEA (2017), lower air quality is a major environmental issue, which affects humans through environmental and air pollutants i.e., nitrogen dioxide, sulfur dioxide, carbon dioxide, and particulate matter (Collivignarelli et al. 2020). In Germany, urban areas are more affected by air pollution as traffic circulation and demographic

density contribute in lower environmental quality (Li et al. 2018; Qiu et al. 2019; Silva et al. 2018).

According to the EEA (2017), PM10 and PM2.5 contribute about 13% of total air pollutants in Germany, mainly due to a higher proportion of diesel trucks emit NO₂. Which is why, German federal court's recent rulings have validated recent changes in environmental policies, i.e., ban on diesel vehicles in urban areas to prevent further GHG emissions (Giesberts 2018; Schmitz et al. 2018). Air quality in Germany is one of the key environmental challenges for German government as it encourages sustainable environment, hence signifying the importance of environmental issues to address environmental issues (Zambrano et al. 2010). Currently, there is a lack of scientific studies that empirically analyze whether or not the concentration level of air pollutants can contribute to the outbreak of COVID-19.

Research methodology

In the current study, the COVID-19 outbreak is represented by total confirmed cases, total deaths, current active cases, and the number of recovered patients with data taken from the German Federal Ministry of Health (Fig. 1). Air and environmental pollution are measured through PM2.5, PM10, Ozone (O₃), and Nitrogen dioxide (NO₂) and data for these pollutants has been collected from the Federal Ministry for the Environment, Nature Conservation and Nuclear Safety in Germany. Humidity and average daily temperature represent climate indicators with daily data taken from Deutscher Wetterdienst. Lastly, Environmental Quality Index (EQI) is included as a proxy for environmental pollution, with data taken from the Federal Ministry for the Environment, Nature Conservation and Nuclear Safety. Dataset for all the variables has been taken from February 24, 2020, to July 2, 2020, to provide a comprehensive analysis for COVID-19 pandemic in Germany.

We have employed the correlation analysis and the wavelet transform coherence (WTC) approach as analytical tools to examine the association between COVID-19 outbreak, and air pollution and climate indicators in Germany. As a prerequisite of correlation analysis, we use Kolmogorov-Smirnov *Z* test to check data normality. As indicated by values in Table 1, our data is non-normal; hence, we use Spearman correlation analysis to examine the correlation; furthermore, we apply WTC to provide graphical association among variables of the study. Wavelet analysis is one of the most widely used analytical approaches in geosciences, environment, weather, and economics (Afshan et al. 2018; Wu et al. 2019). The wavelet function is preferred as an analytical tool as it reveals weak transients and peculiarities within time series datasets (Arneodo et al. 1995; Muzy et al. 1994). The current study has selected wavelet methodology because of its ability to analyze non-normal and non-stationary present in time series data distributions (Masset 2008; Struzik 2001). Moreover, it enables us to study associations between data variables at different frequencies, which helps in identifying the positive and negative associations at multiple frequencies.

Results and discussion

Table 1 presents the results of the Kolmogorov-Smirnov *Z* test, which indicates that the variables are not normally distributed. Thus, non-parametric tests are more appropriate for data analysis. The median values of total cases, recoveries, active patients, and deaths of COVID-19 are 160,726, 118,900, 12,306, and 6391, respectively, in our sample data.

Table 2 presents the Spearman's correlation analysis, which indicates that PM2.5, O₃, and NO₂ are significantly associated with the total cases, recovered, active cases, and deaths from COVID-19. Temperature has a significant association with total cases, recoveries, and deaths. Lastly, PM10, humidity, and EQI only have significant association with active cases from COVID-19 pandemic in Germany. Thus, our

Fig. 1 COVID-19 outbreak in Germany




Table 1 Descriptive statistics

Variables	N	Mean	Median	Kolmogorov-Smirnov Z	P value
Cases	130	123,248	160,726	3.15	0.00
Recoveries	130	95,863	118,900	2.51	0.00
Active cases	130	22,427	12,306	2.35	0.00
Deaths	130	4957	6391	3.43	0.00
PM2.5	130	36	33	3.45	0.00
PM10	130	15	13	3.42	0.00
O ₃	130	36	35	4.91	0.00
NO ₂	130	08	07	3.01	0.00
Temperature	130	54	53	5.34	0.00
Humidity	130	61	58	5.10	0.00
EQI	130	123,248	160,726	7.11	0.00

findings indicate that among air pollutants, PM_{2.5}, NO₂, and O₃ are the key determinants of COVID-19 in Germany. On the other hand, temperature is the only climate indicator, which has a significant influence over COVID-19 pandemic.

Wu et al. (2020) investigated the association of COVID-19 outbreak with fine particulate matter and recommended that a mere increase of 1 µg/m³ results in 15% increase in the mortality cases from COVID-19. Likewise, Fann and Risley 2013, Wang et al. (2020), and Piazzalunga-Expert (2020) arrived at the same conclusions for the USA, China, and Italy, respectively. Recent trends in urbanization have dramatically changed the demographic characteristics of urban areas, which is why a growing number of scientific studies are investigating the association of chronic illnesses with air pollutants such as NO₂ and O₃. Pandey et al. (2005) studied the health risk associated with NO₂ concentration level in New Delhi, India, and suggested that NO₂ concentration in industrial and residential areas poses severe health challenges for the elderly and children. They further articulated that HR values for NO₂ are 22 times worse than other air pollutants in New Delhi. Similarly, Xue et al. (2020) studied spatio-temporal variations for Shanghai and suggested that increased focus on urban residential planning and

introduction of emission management policies have significantly reduced NO₂ concentrations in Shanghai. Bernardini et al. (2020) explored the impact of air pollutants, i.e., O₃, and suggested that a number of patients into the emergency section have direct correlation with air pollutants. Furthermore, Adach et al. (2020) and Ha (2020) registered similar findings.

Shi et al. (2020), based on Chinese provincial data, evaluated the effect of climate indicators and determined that temperature is a key element, which affects the viral outbreak of COVID-19. Previous studies, to examine the association of temperature and spread of infectious diseases, also suggested that temperature influences infectious diseases with low temperature generally attributed with higher cases from the pandemic (Tan et al. 2005; Vandini et al. 2013). Another key indicator is the humidity. Current literature concludes that high temperature and humidity reduce the spread of coronavirus; furthermore, the arrival of summer significantly reduces the COVID-19 (Luo et al. 2020; Ma et al. 2020; Oliveiros et al. 2020). Environmental quality index serves as a comprehensive indicator to analyze the impact of environmental quality on human health. Based on Chinese provincial dataset from December 01, 2019, to March 20, 2020, Han et al. (2020) concluded that lower environmental quality is negatively correlated with human immune system and as a consequence, it is contributing towards COVID-19 pandemic in countries with poor environmental quality. Similarly, Venter et al. (2020) reinforced comparable discoveries and recommended that better environmental quality lowers the risk of getting infected from COVID-19.

Wavelet transform coherence

The current study has used the wavelet transform coherence to portray a graphical correlation between variables, which is one of the most extensively used as an econometric tool. In WTC approach, small arrows indicate the direction of correlation

Table 2 Spearman correlation analysis

Variables	Cases	Recoveries	Active cases	Deaths
PM2.5	−0.272***	−0.273***	0.269***	−0.281***
PM10	0.013	0.012	0.457***	0.005
O ₃	0.214**	0.216**	0.467***	0.215**
NO ₂	0.615***	0.614***	−0.228***	0.614***
Temperature	0.876***	0.878***	−0.132	0.877***
Humidity	−0.119	−0.118	−0.555***	−0.119
EQI	−0.016	−0.016	0.493***	−0.017

***, **, and * represent 1%, 5%, and 10% level of significance

running from one variable to the other. The colors inside the circles indicate the strength of such a relationship; furthermore, the strength of the association goes from high to low as we move from yellow to blue color. Cone of influence separates the significant region from the rest with a thick black lining drawn from top to bottom on both sides.

Figure 2a portrays a mild coherence between PM_{2.5} and total confirmed cases from COVID-19. None of these variables is leading/lagging in this scenario shown by a 0° angle of the arrows. Other yellow areas can be seen scattered

throughout the figure but do not represent a significant relationship. Similar coherence can be found in Fig. 2b between PM_{2.5} and deaths from COVID-19, with only significant coherence between 50 and 60 days of time frame, hence signifying that PM_{2.5} has no significant association with deaths from COVID-19 pandemic.

Figure 2c represents coherence between active cases and PM_{2.5}, though it represents mild coherence but a significant impact can be observed between 50 and 70 days and 90–110 days time scale, signalling that PM_{2.5} played an active




Fig. 2 a Coherence between PM_{2.5} and total confirmed cases. b Coherence between PM_{2.5} and deaths. c Coherence between active cases and PM_{2.5}. d Coherence of recoveries and PM_{2.5}

role in COVID-19 pandemic in Germany for such time periods. This implies that decreased PM_{2.5} contributes to COVID-19 cases but only during a short period of time frequency contribution. Figure 2d shows the coherence of recoveries from COVID-19 and PM_{2.5} at the start of the 60–80 frequency range displays weak coherence, which indicates that PM_{2.5} has a weak impact in reducing the recoveries from COVID-19.

Figure 3a reports a mild coherence with negative correlation in some limited areas, between total confirmed cases from COVID-19 and O₃. The notable yellow areas with black borders can be seen at the start of the 20–40 and 70–80 frequency range. Next, we analyze coherence between deaths from COVID-19 and O₃ with negative correlation areas in Fig. 3b. The frequency range of 40–50 and 70–80 demonstrates that these variables have mild coherence between them.




Fig. 3 a Coherence between total confirmed cases and O₃. b Coherence between deaths and O₃. c Coherence between active cases and O₃. d Coherence of recoveries and O₃

Additionally, Fig. 3c showcases mild coherence with negative correlation. Lastly, Fig. 3d showcases mild coherence with indecisive correlation between recoveries from COVID-19 and O₃ in Germany.

Figure 4 reports the coherence between confirmed cases, deaths, active cases, and recoveries from COVID-19, and temperature. Figure 4a indicates mild positive coherence between total confirmed cases and temperature throughout the 40–60

and 90–100 frequency range. Figure 4b shows coherence with indecisive correlation between deaths and temperature, indicating that deaths from COVID-19 are not significantly affected by the change in temperatures in Germany. Next, Fig. 4c shows coherence with negative correlation between active cases and temperature. Lastly, Fig. 4d highlights weak positive coherence between recovered cases and temperature, showing temperature variation plays a marginal role in recoveries from COVID-19.




Fig. 4 a Coherence between total confirmed cases and temperature. b Coherence between deaths and temperature. c Coherence between active cases and temperature. d Coherence of recoveries and temperature

Finally, we analyze coherence between NO_2 and total confirmed cases, deaths, active cases, and recoveries from COVID-19. Figure 5a shows moderate coherence between total confirmed cases and NO_2 with a notable yellow area between 110 and 120 frequency range. Next, Fig. 5b shows that deaths from COVID-19 have a positive but mild coherence with NO_2 , visible between

40 and 90 frequency range. Next, Fig. 5c shows negative correlation between active cases and NO_2 , which shows that NO_2 has not played a significant role in increased active cases during the COVID-19 pandemic. Finally, the coherence between recoveries and NO_2 is visible from Fig. 5d, which highlights mild coherence with indecisive correlation.




Fig. 5 a Coherence between total confirmed cases and NO_2 . b Coherence between deaths and NO_2 . c Coherence between active cases and NO_2 . d Coherence of recoveries and NO_2

Conclusion

Industrial European economies such as Germany are one of the epicenters of the COVID-19 pandemic, as the number of infections from COVID-19 has an ever-increasing trend; thus, the risk of contracting the virus in these regions is high. Germany offers a unique scenario, as it faced a high infectious rate but still a very marginal mortality rate from COVID-19. In the current study, we have examined the association of COVID-19 outbreak (e.g., total cases, recoveries, and deaths) with environmental and climate indicators in Germany. Our analysis reveals that PM₁₀, O₃, temperature, and humidity are the main determinants of the COVID-19 outbreak in Germany. Data analysis supports the fact that lower environmental pollution level has contributed significantly to combat coronavirus pandemic as in the recent past, Germany has introduced significant policy initiatives in its pandemic response strategies after the outbreak of SARS so that environmental policies complement medical initiatives. Statistically, lower exposure to environmental and air pollutants and access to medical facilities significantly reduce the pandemic threat as vulnerable age groups, i.e., elderly and people with chronic diseases are more susceptible to the pandemic. Our study also provides policy implications for health and policy regulators around the globe. We suggest that other countries can take insights from Germany's response by focusing simultaneously on the development of the medical care system and implementing state and federal policies to reduce environmental pollution.

Although our study has interesting findings, the following limitations exist within our work. First, our study is limited to Germany. Furthermore, the current study is based on daily data with limited frequency range; we recommend the researchers to explore the long-term association of environmental and climate indicators with global pandemics by analyzing empirical association within panel dataset of advanced industrial economies. Finally, the wavelet transform coherence serves as a unique analytical tool in comparison against time series and correlation, but research findings should be cautiously interpreted. Also, we encourage the researchers to study inter-regional response towards COVID-19 pandemic to provide a better outlook for understanding the spread of COVID-19 as a global pandemic.

References

- Adach W, Błaszczak M, Olas B (2020) Carbon monoxide and its donors-chemical and biological properties. *Chem Biol Interact* 108973:108973. <https://doi.org/10.1016/j.cbi.2020.108973>
- Afshan S, Sharif A, Loganathan N, Jammazi R (2018) Time–frequency causality between stock prices and exchange rates: further evidences from cointegration and wavelet analysis. *Phys A Stat Mech Appl* 495:225–244. <https://doi.org/10.1016/j.physa.2017.12.033>
- Anderson HR, Favarato G, Atkinson RW (2013) Long-term exposure to air pollution and the incidence of asthma: meta-analysis of cohort studies. *Air Qual Atmos Health* 6(1):47–56. <https://doi.org/10.1007/s11869-012-0184-5>
- Arneodo A, Bacry E, Muzy J (1995) The thermodynamics of fractals revisited with wavelets. *Phys A Stat Mech Appl* 213(1–2):232–275. [https://doi.org/10.1016/0378-4371\(94\)00163-N](https://doi.org/10.1016/0378-4371(94)00163-N)
- Bank W (2020) World Development Indicators Retrieved from <https://datacatalog.worldbank.org/dataset/world-development-indicators>
- Bashir MF, Ma B, Bilal, Komal B, Bashir MA, Tan D, Bashir M (2020a) Correlation between climate indicators and COVID-19 pandemic in New York, USA. *Sci Total Environ* 728:138835. <https://doi.org/10.1016/j.scitotenv.2020.138835>
- Bashir MF, Ma BJ, Bilal, Komal B, Bashir MA, Farooq TH, Iqbal N, Bashir M (2020b) Correlation between environmental pollution indicators and COVID-19 pandemic: a brief study in Californian context. *Environ Res* 187:109652. <https://doi.org/10.1016/j.envres.2020.109652>
- Bernardini F, Attademo L, Trezzi R, Gobbi C, Balducci P, Del Bello V, Menculini G, Pauselli L, Piselli M, Sciarra T (2020) Air pollutants and daily number of admissions to psychiatric emergency services: evidence for detrimental mental health effects of ozone. *Epidemiol Psychiatr Sci* 29. <https://doi.org/10.1017/S2045796019000623>
- Ceylan Z (2020) Estimation of COVID-19 prevalence in Italy, Spain, and France. *Sci Total Environ* 729:138817. <https://doi.org/10.1016/j.scitotenv.2020.138817>
- Cheung CW, He G, Pan Y (2020) Mitigating the air pollution effect? The remarkable decline in the pollution-mortality relationship in Hong Kong. *J Environ Ecol Manag* 102316:102316. <https://doi.org/10.1016/j.jeem.2020.102316>
- Collivignarelli MC, Abbà A, Bertanza G, Pedrazzani R, Ricciardi P, Miino MC (2020) Lockdown for CoViD-2019 in Milan: what are the effects on air quality? *Sci Total Environ* 732:139280. <https://doi.org/10.1016/j.scitotenv.2020.139280>
- Dutheil F, Baker JS, Navel V (2020) COVID-19 as a factor influencing air pollution? *Environ Poll (Barking, Essex: 1987)* 263:114466. <https://doi.org/10.1016/j.envpol.2020.114466>
- EEA (2017) Air quality in Europe — 2017 report, EEA technical report. <https://doi.org/10.2800/358908>
- Fann N, Risley D (2013) The public health context for PM_{2.5} and ozone air quality trends. *Air Qual Atmos Health* 6(1):1–11. <https://doi.org/10.1007/s11869-10-0125-0>
- Fattorini D, Regoli F (2020) Role of the atmospheric pollution in the Covid-19 outbreak risk in Italy. medRxiv. <https://doi.org/10.1101/2020.04.23.20076455>
- Fofana NK, Latif F, Sarfraz S, Bilal, Bashir MF, Komal B (2020) Fear and agony of the pandemic leading to stress and mental illness: an emerging crisis in the novel coronavirus (COVID-19) outbreak. *Psychiatry Res* 113230:113230. <https://doi.org/10.1016/j.psychres.2020.113230>
- Gehrsitz M (2017) The effect of low emission zones on air pollution and infant health. *J Environ Ecol Manag* 83:121–144. <https://doi.org/10.1016/j.jeem.2017.02.003>
- Giesberts L (2018) Diesel-Verkehrsverbote" ausnahmsweise moeglich! NEUE ZEITSCHRIFT FUER VERWALTUNGSRECHT, 37(17)
- Ha E (2020) Impact of air pollution hazards on human development health impacts of developmental exposure to environmental chemicals. Springer, pp 223–245. https://doi.org/10.1007/978-981-15-0520-1_10
- Han Y, Lam J, Li V, Guo P, Zhang Q, Wang A, Crowcroft J, Wang S, Fu J, Gilani Z (2020). The effects of outdoor air pollution concentrations and lockdowns on Covid-19 infections in Wuhan and other provincial capitals in China. <https://doi.org/10.20944/preprints202003.0364.v1>

- Kessler C, Brücher W, Memmesheimer M, Kerschgens M, Ebel A (2001) Simulation of air pollution with nested models in North Rhine-Westphalia. *Atmos Environ* 35:S3–S12. [https://doi.org/10.1016/S1352-2310\(01\)00137-6](https://doi.org/10.1016/S1352-2310(01)00137-6)
- Lalive R, Luechinger S, Schmutzler A (2018) Does expanding regional train service reduce air pollution? *J Environ Ecol Manag* 92:744–764. <https://doi.org/10.1016/j.jeem.2017.09.003>
- Li Y, Chiu YH, Lu LC (2018) Energy and AQI performance of 31 cities in China. *Energy Policy* 122:194–202. <https://doi.org/10.1016/j.enpol.2018.07.037>
- Luo W, Majumde, M, Liu D, Poirier C, Mandl K, Lipsitch M, Santillana M (2020) The role of absolute humidity on transmission rates of the COVID-19 outbreak. <https://doi.org/10.1101/2020.02.12.20022467>
- Ma Y, Zhao Y, Liu J, He X, Wang B, Fu S, Yan J, Niu J, Zhou J, Luo B (2020) Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Sci Total Environ* 138226:138226. <https://doi.org/10.1016/j.scitotenv.2020.138226>
- Masset P (2008) Analysis of financial time-series using Fourier and wavelet methods. Available at SSRN 1289420. <https://doi.org/10.2139/ssrn.1289420>
- Mez L (2020) 40 years promoting renewable energy in Germany the ecological modernization capacity of Japan and Germany. Springer, pp 119–136. https://doi.org/10.1007/978-3-658-27405-4_9
- Muzy JF, Bacry E, Arneodo A (1994) The multifractal formalism revisited with wavelets. *Int J Bifurc Chaos* 4(02):245–302. <https://doi.org/10.1142/S0218127494000204>
- O’Riordan T (1989) Air pollution legislation and regulation in the European Community: a review essay. *Atmos Environ* (1967) 23(2):293–306. [https://doi.org/10.1016/0004-6981\(89\)90578-7](https://doi.org/10.1016/0004-6981(89)90578-7)
- Oliveiros B, Caramelo L, Ferreira NC, Caramelo F (2020). Role of temperature and humidity in the modulation of the doubling time of COVID-19 cases. medRxiv. <https://doi.org/10.1101/2020.03.05.20031872>
- Organization WH (2020) Rational use of personal protective equipment (PPE) for coronavirus disease (COVID-19): interim guidance, 19 March 2020
- Pandey JS, Kumar R, Devotta S (2005) Health risks of NO₂, SPM and SO₂ in Delhi (India). *Atmos Environ* 39(36):6868–6874. <https://doi.org/10.1016/j.atmosenv.2005.08.004>
- Piazzalunga-Expert A (2020) Evaluation of the potential relationship between particulate matter (PM) pollution and COVID-19 infection spread in Italy
- Popp D (2006) International innovation and diffusion of air pollution control technologies: the effects of NO_x and SO₂ regulation in the US, Japan, and Germany. *J Environ Ecol Manag* 51(1):46–71. <https://doi.org/10.1016/j.jeem.2005.04.006>
- Qiu G, Song R, He S (2019) The aggravation of urban air quality deterioration due to urbanization, transportation and economic development—panel models with marginal effect analyses across China. *Sci Total Environ* 651:1114–1125. <https://doi.org/10.1016/j.scitotenv.2018.09.219>
- Schmitz S, Weiand L, Becker S, Niehoff N, Schwartzbach F, von Schneidmesser E (2018) An assessment of perceptions of air quality surrounding the implementation of a traffic-reduction measure in a local urban environment. *Sustain Cities Soc* 41:525–537. <https://doi.org/10.1016/j.scs.2018.06.011>
- Schulz C, Conrad A, Becker K, Kolossa-Gehring M, Seiwert M, Seifert B (2007) Twenty years of the German Environmental Survey (GerES): human biomonitoring – temporal and spatial (West Germany/East Germany) differences in population exposure. *Int J Hyg Environ Health* 210(3):271–297. <https://doi.org/10.1016/j.ijheh.2007.01.034>
- Shi P, Dong Y, Yan H, Li X, Zhao C, Liu W, He M, Tang S, Xi S (2020) The impact of temperature and absolute humidity on the coronavirus disease 2019 (COVID-19) outbreak—evidence from China. medRxiv. <https://doi.org/10.1101/2020.03.22.20038919>
- Silva BN, Khan M, Han K (2018) Towards sustainable smart cities: a review of trends, architectures, components, and open challenges in smart cities. *Sustain Cities Soc* 38:697–713. <https://doi.org/10.1016/j.scs.2018.01.053>
- Struzik ZR (2001) Wavelet methods in (financial) time-series processing. *Phy A Stat Mech Appl* 296(1–2):307–319. [https://doi.org/10.1016/S0378-4371\(01\)00101-7](https://doi.org/10.1016/S0378-4371(01)00101-7)
- Tan J, Mu L, Huang J, Yu S, Chen B, Yin J (2005) An initial investigation of the association between the SARS outbreak and weather: with the view of the environmental temperature and its variation. *J Epidemiol Community Health* 59(3):186–192. <https://doi.org/10.1136/jech.2004.020180>
- Vandini S, Corvaglia L, Alessandrini R, Aquilano G, Marsico C, Spinelli M, Lanari M, Faldella G (2013) Respiratory syncytial virus infection in infants and correlation with meteorological factors and air pollutants. *Ital J Pediatr* 39(1):1. <https://doi.org/10.1186/1824-7288-39-1>
- Venter ZS, Aunan K, Chowdhury S, and Lelieveld J (2020). COVID-19 lockdowns cause global air pollution declines with implications for public health risk. medRxiv. DOI: <https://doi.org/10.1101/2020.04.10.20060673>
- Wang P, Chen K, Zhu S, Wang P, Zhang H (2020) Severe air pollution events not avoided by reduced anthropogenic activities during COVID-19 outbreak. *Res Conserv Recycl* 158:104814. <https://doi.org/10.1016/j.resconrec.2020.104814>
- Wu Q, Tan J, Guo F, Li H, Chen S (2019) Multi-scale relationship between land surface temperature and landscape pattern based on wavelet coherence: the case of metropolitan Beijing, China. *Remote Sens* 11(24):3021. <https://doi.org/10.3390/rs11243021>
- Wu X, Nethery RC, Sabath BM, Braun D, Dominici F (2020) Exposure to air pollution and COVID-19 mortality in the United States. medRxiv. <https://doi.org/10.1101/2020.04.05.20054502>
- Xue R, Wang S, Li D, Zou Z, Chan KL, Valks P, Saiz-Lopez A, Zhou B (2020) Spatio-temporal variations in NO₂ and SO₂ over Shanghai and Chongming Eco-Island measured by Ozone Monitoring Instrument (OMI) during 2008–2017. *J Clean Prod* 258:120563. <https://doi.org/10.1016/j.jclepro.2020.120563>
- Zambrano AMA, Broadbent EN, Durham WH (2010) Social and environmental effects of ecotourism in the Osa Peninsula of Costa Rica: the Lapa Rios case. *J Ecotour* 9(1):62–83. <https://doi.org/10.1080/14724040902953076>
- Zijlema WL, Wolf K, Emeny R, Ladwig KH, Peters A, Kongsgård H, Hveem K, Kvaløy K, Yli-Tuomi T, Rosmalen JGM (2016) The association of air pollution and depressed mood in 70,928 individuals from four European cohorts. *Int J Hyg Environ Health* 219(2): 212–219. <https://doi.org/10.1016/j.ijheh.2015.11.006>
- Zimmermann F, Lux H, Maenhaut W, Matschullat J, Plessow K, Reuter F, Wienhaus O (2003) A review of air pollution and atmospheric deposition dynamics in southern Saxony, Germany, Central Europe. *Atmos Environ* 37(5):671–691. [https://doi.org/10.1016/S1352-2310\(02\)00829-4](https://doi.org/10.1016/S1352-2310(02)00829-4)

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.