

Review Article

Epidemiology of metabolic syndrome in Asia

Wen-Harn Pan PhD^{1,2,3}, Wen-Ting Yeh MS¹ and Lu-Chen Weng MS¹¹*Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan*²*Institute of Epidemiology, School of Public Health, National Taiwan University, Taipei, Taiwan*³*Institute of Microbiology and Biochemistry, National Taiwan University, Taipei, Taiwan*

Metabolic syndrome (MS) is a rising disease entity characterized by a clustering of metabolic conditions. Although prevalence of obesity as defined by the World Health Organization (WHO) is relatively low in Asia compared to western countries, metabolic syndrome is growing into a significant public health problem. Comparative studies indicate that metabolic responses to obesity may be greater in South and East Asians than their western counterparts at given Body Mass Indexes (BMIs). Higher percentage body fat in Asians at given BMIs and over-responsiveness to obesity may in part explain the phenomenon for which the underlying causes are not clear. Furthermore, aborigines may be at an even greater MS risk. The metabolic syndrome definition itself as well as whether it should be defined are controversial. The National Cholesterol Education Program-Adult Treatment Panel III (NCEP-ATP III) gives equal weight to each component disorder, while the International Diabetes Federation (IDF) takes central obesity as a pre-requisite. Both criteria adopt ethnic-specific cut-off points for waist circumference. Asian data favour the new NCEP-ATP III definition, as individuals that were selected through the NCEP criteria but disregarded by the IDF criteria had similar Framingham cardiovascular disease risk scores to those picked by both definitions. Prospective data show that the metabolic syndrome not only increases the risk of coronary artery disease but also cerebrovascular disease in Asians. Macronutrient composition and the quality of the diet are associated with the risk of metabolic syndrome. More research is needed to relate diet and metabolic syndrome in Asians.

Key Words: metabolic syndrome, Asian, definition, diet, epidemiology

INTRODUCTION

The significance of metabolic syndrome has unravelled gradually in recent years. The metabolic syndrome is a condition characterized by a constellation of metabolic disorders including: abdominal obesity, insulin resistance/glucose intolerance, atherogenic dyslipidemia (elevated TG, and lower HDL-C), raised blood pressure, proinflammatory and prothrombotic state. It was referred to as the "X syndrome" by Kylin in the 1920's and described as a phenomenon of the clustering of obesity, hypertension, and gout.¹ The concept of syndrome X was reintroduced by Reaven² in the late 1980's for the clustering of cardiovascular risks.

Operational definitions of metabolic syndrome

A unified operational definition of the syndrome was first proposed in "Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation" in 1998³. According to this definition, a person with type 2 DM, impaired fasting glucose, impaired glucose tolerance, or insulin resistance has the metabolic syndrome if two or more of the following component criteria are satisfied: dyslipidemia (elevated plasma triglyceride or low HDL-C), obesity (high BMI or waist-to-hip ratio), hypertension, and microalbuminuria.

On the other hand, the criteria suggested by the National Cholesterol Education Program-Adult Treatment Panel III (NCEP-ATP III) in 2001 utilized primarily readily available biochemical parameters: fasting plasma triglyceride, HDL-C, glucose, and blood pressure, whose cut-off points

were lowered to the higher end of the normal range.⁴ In addition, waist circumference was used to define obesity, focusing on abdominal adiposity. A person is diagnosed as having the metabolic syndrome if three or more of the criteria are satisfied. Furthermore, the International Diabetes Federation (IDF) in 2005⁵ proposed yet another criteria which mandates central obesity as the essential condition in addition to two other disorders out of the four stated. Cut-off points similar to the NCEP-ATP III criteria were used except for fasting glucose which was lowered to 100mg/dl and ethnic-specific cut-off points for waist circumference were adopted. In the fall of 2005, the revised NCEP-ATP III criteria also adopted the same glucose and waist cut-offs while maintaining the equality principle for all five component disorders.⁶

Although most experts recognize that obesity-related insulin resistance may be the core cause of metabolic syndrome, each medical society has its emphasis in defining the syndrome. The definition used in the WHO-diabetes report centres on diabetes and insulin resistance, whereas IDF focuses on central obesity. While the NCEP-ATP III

Corresponding Author: Dr W.H Pan, Institute of Biomedical Sciences, Academia Sinica, 128 Sec. 2, Academia Rd., Nankang, Taipei 11529, Taiwan.

Tel: 011-886-2-2789-9121; Fax: 011-886-2-2782-3047

Email: pan@ibms.sinica.edu.tw

Manuscript received 9 September 2007. Accepted 3 December 2007.




Figure 1. Prevalence rates of metabolic syndrome in Asians

guideline gives equal weights to abdominal adiposity, high blood pressure, hyperglycemia, elevated triglyceridemia and low-HDL cholesterol. Prevalence rate estimated from different definitions would vary inevitably and long-term cardiovascular risks of these definitions may also differ.

Prevalence of metabolic syndrome in Asian countries

Most prevalence data in literature used either the 2001 or the 2005 version of the NCEP-ATP III definition (Figure 1). According to the 2005 version; China,⁷ Taiwan,^{8, 9} Hong Kong¹⁰, and Thailand¹¹ had similar prevalence rates ranging between 10-15%. However, it was much lower (around 5%) in southern rural China.¹² On the other hand, rates for Koreans,¹³ approximately one quarter, were higher than the Chinese and Thais, even though their average body mass index (BMI) was similar to other East Asian countries. India¹⁴ had strikingly high prevalence rates compared to the rest of Asia, approaching that of US population. Metabolic syndrome prevalence in Turkey¹⁵ and Iran¹⁶ was comparable to that of the US with a large excess in females. Considering the relatively low mean BMI and the low prevalence of obesity (BMI > 30) in east and south Asians, the magnitude of the metabolic syndrome prevalence is unusually high.

Excess metabolic risk in Asians

A series of comparative data pointed out that under fixed BMI or fatness, metabolic risk was much greater in South Asians compared to Caucasians in terms of diabetes mellitus, insulin resistance and hypertriglyceridemia.¹⁷ Our study comparing data from NHANES II and from Nutritional and Health Survey in Taiwan also showed that Taiwanese, a group of East Asians, experienced higher absolute and relative risk of hypertension, hyperglycemia, dyslipidemia, and hyperuricemia than non-hispanic US white and US blacks.¹⁸ An integrated graph of this phenomenon relating BMI and metabolic risks is provided in figure 2. This phenomenon is most likely due to a higher




Figure 2. Age-standardized prevalence (1a) and relative prevalence ratio (1b) in the presence of two or more metabolic risk factors† by BMI levels in three populations: Taiwanese, non-hispanic US whites, and US blacks. †High blood pressure: SBP>130, DBP>85 or on antihypertensives, hyperglycemia: fasting glucose> 100mg/dl or on lipid-lowering drug Low HDL-C: HDL-C< † 40/ ‡ 50 mg/dl.

percentage of body fat accumulation in Asians than in Caucasians at fixed BMI level as pointed out by a series of studies from Deurenberg et al,^{19,20} comparing data among Indonesians and Singaporeans with Caucasians.




Figure 3. Odds ratio of aboriginal ethnicity on the presence of metabolic syndrome and hyperuricemia, Nutrition Health Survey in Taiwan 1993-1996. Potential confounding factors controlled for both health conditions: age, physical activity, presence of physical disability, education level, and amount of alcohol consumption; additional confounding factors controlled for hyperuricemia: BMI, and waist circumference.

Recent study from Taiwan²¹, Korea²², and Australian aborigines²³ further supported this hypothesis.

Our study²⁴ (figure 3) found that aborigines residing in Taiwan had a much greater risk of ascertaining metabolic syndrome than Han Chinese, despite controlling for potential intermediate factors such as age, physical activity, education level and alcohol consumption. This indicated the potential effect of genetics or the interaction between genetics and the environment. Studies showed that insulin sensitivity was poorer in Asian Indians than Caucasians at fixed body fat percentages.²⁵ Increased body fat percentage may affect individuals differently due to differences in genetic make-up, intra-uterine (developmental) environment, and dietary and physical activity patterns.

Applicability of available definitions of metabolic syndrome to Asians

ATP III and IDF definitions are two of the most popular definitions for metabolic syndrome. However, it is controversial as to which one is better in defining metabolic syndrome. We addressed the differences between these

two definitions with regard to prevalence estimation and its relations with cardiovascular risks.

Comparing data from the Nutrition and Health Survey in Taiwan²⁶ and NHANES III (figure 2), prevalence estimates from the two definitions did not differ much in US populations,²⁷ but with Taiwanese data the IDF estimate for women was 25% lower (12.6% vs 16.5%) and for men it was 50% lower (6.2% vs 11.6%) than that of the ATP III. The discrepancy is likely due to the proportion of men and women with metabolic syndrome defined by the revised ATP III who do not have elevated waist circumferences (figure 4). It is speculated by this author that this phenomenon may be due to either lack of proper indicators for abdominal obesity or a inappropriate waist circumference cut-off point for Asians. Furthermore, we found that there is a greater proportion of Taiwanese that have four or more metabolic disorders clustered or high blood pressure, but only a smaller proportion of people that have high fasting glucose values. Further research is needed to understand the rational and consequences of these different combination patterns.

With regard to cardiovascular risk, when comparing people picked up by both guidelines (with central obesity) with those identified by NCEP guideline only (without central obesity),⁸ no significant differences were observed for levels of blood pressure, HDL-C, fasting triglyceride and glucose, and Framingham coronary artery disease risk score. And these cardiovascular risks were much stronger in both groups of metabolic syndrome patients than people with no such syndrome. This data indicate that a good proportion of people with metabolic syndrome may be missed by the IDF definition even though it may be sound for IDF to consider central obesity appearing early in the chain of events in the pathogenesis of metabolic syndrome. In addition, our recent work²⁸ comparing the 2001 and 2005 versions of NECP-ATP III definitions found that the new version corresponded to a sharper dose response curve relating the number of metabolic disorders to the risk of ischemic stroke in the Two-township studies in Taiwan. Taken together, among the




Figure 4. Clustering patterns of metabolic disorders in US non-Hispanic whites (WT), African Americans (AA), and Taiwanese (TW)

available definitions, the new NCEP-ATP III guideline may be more advantageous than others to predict future cardiovascular risks and may be used for screening in Asians.

Metabolic syndrome and cardiovascular events

It is well documented in literature that metabolic syndrome is associated with increased risk of all cause mortality and cardiovascular events. Ford et al²⁹ conducted a meta-analysis on 12 prospective studies primarily for Caucasian populations and demonstrated a 74% increase in cardiovascular risk. We have obtained estimates from five prospective studies from Japan,^{30,31} Korean,¹³ and Taiwan^{28,32} that examined the relations between baseline metabolic syndrome status and incidental events on coronary artery disease or cerebrovascular disease, using either the 2001 NCEP-ATP III guideline, 2005 NCEP-ATP III guideline, or the modified criteria using BMI cut-off point to define obesity.

As shown in Table 1, their hazard ratios are in the vicinity of 2 which is very close to the estimate obtained from Asia Pacific Cohort Study Collaboration³³ which gathered individual data from 35 cohorts from mainland China, Hong Kong, Japan, south Korean, Taiwan, Thailand, and Singapore. The APCSC data further showed that the more component disorders one had, the higher the

risk in acquiring cardiovascular death.

Diet and metabolic syndrome

A great body of information have accumulated with regard to the relations between diet and each of the component disorder of metabolic syndrome. In general, it is accepted that the Mediterranean diet with fish, vegetable, fruit, red wine, and olive oil may protect against the development of metabolic syndrome.³⁴⁻³⁶ But up to this point, only a dozen of cross-sectional studies and a relatively small number of prospective studies have tried to determine the association between diet and the metabolic syndrome and comprised of primarily Caucasian studies (Framingham,³⁷⁻³⁹ WHS,^{40,41} NHANES,⁴² Boston,⁴³ ATTICA-greece⁴⁴), one Japanese-Brazilian,⁴⁵ and a few Iranian study.⁴⁶⁻⁴⁹ Overall speaking, increased consumption of whole grains, dairy products, vegetables, fruits, calcium, magnesium, fiber from cereals, linoleic acid, and the maintenance of a healthy dietary pattern (high in fruits, tomatoes, poultry, legumes, cruciferous and green leafy vegetables, other vegetables, tea, fruit juices, and whole grains) have shown to be protective. The consumption of refined grains, soft or sweetened drinks, alcoholic beverage and fat as well as keeping to a empty calorie diet, potato and meat diet have been associated with elevated

Table 1. Hazard ratios relating metabolic syndrome to cardiovascular events in Asian prospective studies

Study	Subjects	Age	Outcomes	MS definition†	Hazard ratio
KNHANES ¹³	Korean (N=4452)	20+	CHD/Stroke	ATPIII.8090.110	♂ 0.8 (0.4-1.7)
					♀ 2.4 (1.1-5.1) RR
Hisayama study ³⁰	Japanese (N=2452)	40+	CHD/Stroke	ATPIII.8090.110	♂ 1.86(1.32-2.62)
					♀ 1.70(1.22-2.36)
					♂ 1.94 (1.19-3.17)
					♀ 2.86 (1.56-5.24)
Cardiovascular Risk Survey ³¹	Japanese (N=9087)	40-69	CHD/ IS	ATPIII.BMI25.110	♂ 2.4 (1.4-4.0)
					♀ 2.3 (1.2-4.3)
			IS Stroke		♂ 2.0 (1.3-3.1)
					♀ 1.5 (1.0-2.3)
CVDFACTS ²⁸	Taiwanese (N=3453)	20+	IS stroke	ATPIII.8090.110	♂ 2.12 (1.32-3.4)
					♀ 1.61 (0.95-2.75)
Kin-Shan study ³²	Taiwanese (N=3602)	35+	CHD	ATPIII.8090.100	♂ 2.34 (1.46-3.73)
					♀ 1.41 (0.83-2.39)
APCSC ³³	Asians (N=329,166)	30-75	CHD(fatal)	ATPIII.BMI27.100	1.83 (1.16-2.9)
					Stroke
APCSC ³³	Asians (N=329,166)	30-75	CHD(fatal)	ATPIII.BMI28.110	2.29 (1.28-4.10)
					2.05 (1.13-3.72)
					-more adjustment

† ATPIII.8090.110: obesity: waist circumference \geq 80 cm for women, \geq 90 cm for men; blood glucose \geq 110 mg/dl; ATPIII.BMI25.110: obesity: BMI \geq 25; blood glucose \geq 110 mg/dl; ATPIII.8090.100: obesity: waist circumference \geq 80 cm for women, \geq 90 cm for men; blood glucose \geq 100 mg/dl; ATPIII.BMI27.110: obesity: BMI \geq 27; blood glucose \geq 110 mg/dl; ATPIII.BMI28.110: obesity: BMI \geq 28; blood glucose \geq 110 mg/dl

risk. In addition, our yet unpublished meta-analysis consisting of 21 dietary intervention studies showed that a low carbohydrate diet may be associated with a reduced risk of hypertriglyceridemia, low-HDL-C, but an elevated risk of hypercholesterolemia compared to the American Heart Diet.⁵⁰ To the best of our knowledge, there has yet to be any dietary studies of this kind from South and East Asians in western literature.

CONCLUSIONS

When obesity becomes a pandemic in the world, the significance of the metabolic syndrome is unravelled, in particular in South and East Asians who seem to react to a greater degree to metabolic risk at a given level of obesity. This may be due to higher body fat composition and an over-responsiveness to obesity in Asians. More research is needed to understand the gene-environment interaction in this observed discrepancy between Asians and Caucasians.

The magnitude of the cardiovascular risk associated with the metabolic syndrome in Asians is similar to, if not larger than, that of Caucasians. In addition, among the several available definitions, the 2005 NCEP-ATP III criteria is most suitable for Asians in terms of the dose-response relationship with ischemic stroke. The IDF definition will miss a sizable number of Asians if used in screening probably due to the inadequacy of current methods to identify central obesity.

Very little has been done to relate dietary characteristics and metabolic syndrome in Asians. More prospective investigations and dietary intervention studies are required to understand the relations between diet and the metabolic syndrome.

Overall speaking, the metabolic syndrome has caught the attention of health professionals in Asian. One of the challenges is to understand its pathogenesis and to identify effective measures to prevent it either at the population level or at the individual level.

AUTHOR DISCLOSURES

Wen-Harn Pan, Wen-Ting Yeh and Lu-Chen Weng, no conflicts of interest.

REFERENCES

- Kylin E. Studien ueber das hypertonie-hyperglykemie-hyperurikemiesyndrome. *Zentralblatt Fuer Innere Med.* 1923;44:105-27.
- Reaven GM. Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes.* 1988;37:1595-607.
- Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med.* 1998;15:539-53.
- Executive Summary of the Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA.* 2001;285:2486-97.
- Alberti KG, Zimmet P, Shaw J. The metabolic syndrome--a new worldwide definition. *Lancet.* 2005;366:1059-62.
- Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC, Jr., Spertus JA, Costa F. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation.* 2005;112:2735-52.
- Gu D, Reynolds K, Wu X, Chen J, Duan X, Reynolds RF, Whelton PK, He J. Prevalence of the metabolic syndrome and overweight among adults in China. *Lancet.* 2005;365:1398-405.
- Chen HJ, Pan WH. Probable blind spot in the International Diabetes Federation definition of metabolic syndrome. *Obesity.* (Silver Spring, Md 2007;15:1096-100.
- Chuang SY, Chen CH, Chou P. Prevalence of metabolic syndrome in a large health check-up population in Taiwan. *J Chin Med Assoc.* 2004;67:611-20.
- Ko GT, Cockram CS, Chow CC, Yeung V, Chan WB, So WY, Chan NN, Chan JC. High prevalence of metabolic syndrome in Hong Kong Chinese--comparison of three diagnostic criteria. *Diabetes Res Clin Pract.* 2005;69:160-8.
- Lohsoonthorn V, Dhanamun B, Williams MA. Prevalence of metabolic syndrome and its relationship to white blood cell count in a population of Thai men and women receiving routine health examinations. *Am J Hypertens.* 2006;19:339-45.
- Feng Y, Hong X, Li Z, Zhang W, Jin D, Liu X, Zhang Y, Hu FB, Wei LJ, Zang T, Xu X, Xu X. Prevalence of metabolic syndrome and its relation to body composition in a Chinese rural population. *Obesity.* (Silver Spring, Md 2006;14:2089-98.
- Kim HM, Kim DJ, Jung IH, Park C, Park J. Prevalence of the metabolic syndrome among Korean adults using the new International Diabetes Federation definition and the new abdominal obesity criteria for the Korean people. *Diabetes Res Clin Pract.* 2007;77:99-106.
- Gupta R, Deedwania PC, Gupta A, Rastogi S, Panwar RB, Kothari K. Prevalence of metabolic syndrome in an Indian urban population. *Int J Cardiol.* 2004;97:257-61.
- Kozan O, Oguz A, Abaci A, Erol C, Ongen Z, Temizhan A, Celik S. Prevalence of the metabolic syndrome among Turkish adults. *Eur J Clin Nutr.* 2007;61:548-53.
- Zabetian A, Hadaegh F, Azizi F. Prevalence of metabolic syndrome in Iranian adult population, concordance between the IDF with the ATP III and the WHO definitions. *Diabetes Res Clin Pract.* 2007;77:251-7.
- Simmons D, Williams DR, Powell MJ. The Coventry Diabetes Study: prevalence of diabetes and impaired glucose tolerance in Europeans and Asians. *Q J Med.* 1991;81:1021-30.
- Pan WH, Flegal KM, Chang HY, Yeh WT, Yeh CJ, Lee WC. Body mass index and obesity-related metabolic disorders in Taiwanese and US whites and blacks: implications for definitions of overweight and obesity for Asians. *Am J Clin Nutr.* 2004;79:31-9.
- Deurenberg P, Deurenberg Yap M, Wang J, Lin FP, Schmidt G. The impact of body build on the relationship between body mass index and percent body fat. *Int J Obes Relat Metab Disord.* 1999;23:537-42.
- Gurrici S, Hartriyanti Y, Hautvast JG, Deurenberg P. Differences in the relationship between body fat and body mass index between two different Indonesian ethnic groups: the effect of body build. *Eur J Clin Nutr.* 1999;53:468-72.
- Chang CJ, Wu CH, Chang CS, Yao WJ, Yang YC, Wu JS, Lu FH. Low body mass index but high percent body fat in Taiwanese subjects: implications of obesity cutoffs. *Int J Obes Relat Metab Disord.* 2003;27:253-9.

22. Chung S, Song MY, Shin HD, Kim DY, He Q, Heshka S, Wang J, Thornton J, LaFerrere B, Pi-Sunyer FX, Gallagher D. Korean and Caucasian overweight premenopausal women have different relationship of body mass index to percent body fat with age. *J Appl Physiol*. 2005;99:103-7.
23. Piers LS, Rowley KG, Soares MJ, O'Dea K. Relation of adiposity and body fat distribution to body mass index in Australians of Aboriginal and European ancestry. *Eur J Clin Nutr*. 2003;57:956-63.
24. Chang YP, Yeh WT, Cheng YY, Pan WH. Excess metabolic syndrome and hyperuricemia in Taiwanese aborigines: environmental or genetic? *J Genet Mol Biol*. 2007;18:29-33.
25. Chandalia M, Abate N, Garg A, Stray-Gundersen J, Grundy SM. Relationship between generalized and upper body obesity to insulin resistance in Asian Indian men. *J Clin Endocrinol Metab*. 1999;84:2329-35.
26. Pan WH, Chang HY, Yeh WT, Hsiao SY, Hung YT. Prevalence, awareness, treatment and control of hypertension in Taiwan: results of Nutrition and Health Survey in Taiwan (NAHSIT) 1993-1996. *J Hum Hypertens*. 2001;15:793-8.
27. Ford ES. Prevalence of the metabolic syndrome defined by the International Diabetes Federation among adults in the U.S. *Diabetes care*. 2005;28:2745-9.
28. Chen HJ, Bai CH, Yeh WT, Chiu HC, Pan WH. Influence of metabolic syndrome and general obesity on the risk of ischemic stroke. *Stroke*. 2006;37:1060-4.
29. Ford ES. Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: a summary of the evidence. *Diabetes care*. 2005;28:1769-78.
30. Ninomiya T, Kubo M, Doi Y, Yonemoto K, Tanizaki Y, Rahman M, Arima H, Tsuruyama K, Iida M, Kiyohara Y. Impact of metabolic syndrome on the development of cardiovascular disease in a general Japanese population: the Hisayama study. *Stroke*. 2007;38:2063-9.
31. Iso H, Sato S, Kitamura A, Imano H, Kiyama M, Yamagishi K, Cui R, Tanigawa T, Shimamoto T. Metabolic syndrome and the risk of ischemic heart disease and stroke among Japanese men and women. *Stroke*. 2007;38:1744-51.
32. Chien KL, Hsu HC, Sung FC, Su TC, Chen MF, Lee YT. Metabolic syndrome as a risk factor for coronary heart disease and stroke: An 11-year prospective cohort in Taiwan community. *Atherosclerosis*. 2007;194:214-221.
33. Patel A, Barzi F, Woodard M, Ni Mhurchu C, Ohkubo T, Lam TH, Welborn T. An evaluation of metabolic risks for coronary death in the Asia Pacific region. *Diabetes Res Clin Pract*. 2006;74:274-81.
34. Tortosa A, Bes-Rastrollo M, Sanchez-Villegas A, Basterra-Gortari FJ, Nunez-Cordoba JM, Martinez-Gonzalez MA. Mediterranean Diet Inversely Associated with the Incidence of Metabolic Syndrome: the Sun Prospective Cohort. *Diabetes care*. 2007;30(11):2957-2959.
35. Meydani M. A Mediterranean-style diet and metabolic syndrome. *Nutrition reviews* 2005;63:312-4.
36. Alvarez Leon EE, Henriquez P, Serra-Majem L. Mediterranean diet and metabolic syndrome: a cross-sectional study in the Canary Islands. *Public Health Nutr*. 2006;9:1089-98.
37. McKeown NM, Meigs JB, Liu S, Saltzman E, Wilson PW, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. *Diabetes care*. 2004;27:538-46.
38. Dhingra R, Sullivan L, Jacques PF, Wang TJ, Fox CS, Meigs JB, D'Agostino RB, Gaziano JM, Vasan RS. Soft drink consumption and risk of developing cardio-metabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation*. 2007;116:480-8.
39. Sonnenberg L, Pencina M, Kimokoti R, Quatromoni P, Nam BH, D'Agostino R, Meigs JB, Ordovas J, Cobain M, Millen B. Dietary patterns and the metabolic syndrome in obese and non-obese Framingham women. *Obes Res*. 2005;13:153-62.
40. Liu S, Song Y, Ford ES, Manson JE, Buring JE, Ridker PM. Dietary calcium, vitamin D, and the prevalence of metabolic syndrome in middle-aged and older U.S. women. *Diabetes care*. 2005;28:2926-32.
41. Song Y, Ridker PM, Manson JE, Cook NR, Buring JE, Liu S. Magnesium intake, C-reactive protein, and the prevalence of metabolic syndrome in middle-aged and older U.S. women. *Diabetes care*. 2005;28:1438-44.
42. Ford ES, Li C, McGuire LC, Mokdad AH, Liu S. Intake of dietary magnesium and the prevalence of the metabolic syndrome among U.S. adults. *Obesity*. 2007;15:1139-46.
43. Sahyoun NR, Jacques PF, Zhang XL, Juan W, McKeown NM. Whole-grain intake is inversely associated with the metabolic syndrome and mortality in older adults. *Am J Clin Nutr*. 2006;83:124-31.
44. Panagiotakos DB, Pitsavos C, Skoumas Y, Stefanadis C. The association between food patterns and the metabolic syndrome using principal components analysis: The ATTICA Study. *American Dietetic Association*. 2007;107:979-87; quiz 97.
45. Freire RD, Cardoso MA, Gimeno SG, Ferreira SR. Dietary fat is associated with metabolic syndrome in Japanese Brazilians. *Diabetes care*. 2005;28:1779-85.
46. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, Willett WC. Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *Am J Clin Nutr*. 2006;84:1489-97.
47. Esmailzadeh A, Mirmiran P, Azizi F. Whole-grain consumption and the metabolic syndrome: a favorable association in Tehranian adults. *Eur J Clin Nutr*. 2005;59:353-62.
48. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, Willett WC. Dietary patterns, insulin resistance, and prevalence of the metabolic syndrome in women. *Am J Clin Nutr*. 2007;85:910-8.
49. Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F. Dairy consumption is inversely associated with the prevalence of the metabolic syndrome in Tehranian adults. *Am J Clin Nutr*. 2005;82:523-30.
50. Hsueh YP. The effect of Low-carbohydrate Diets on cardiovascular risk Factors: Meta-Analysis. Master Thesis. Graduate Institute of Epidemiology, College of Public Health, Taiwan. 2007.