

Epidithiodiketopiperazines : strain-promoted thiol-mediated cellular uptake at the highest tension

ZONG, Lili, *et al.*

Abstract

The disulfide dihedral angle in epidithiodiketopiperazines (ETPs) is near 0°. Application of this highest possible ring tension to strain-promoted thiol-mediated uptake results in efficient delivery to the cytosol and nucleus. Compared to the previous best asparagusic acid (AspA), ring-opening disulfide exchange with ETPs occurs more efficiently even with nonactivated thiols, and the resulting thiols exchange rapidly with nonactivated disulfides. ETP-mediated cellular uptake is more than 20 times more efficient compared to AspA, occurs without endosomal capture, depends on temperature, and is “unstoppable” by inhibitors of endocytosis and conventional thiol-mediated uptake, including siRNA against the transferrin receptor. These results suggest that ETP-mediated uptake not only maximizes delivery to the cytosol and nucleus but also opens the door to a new multitarget hopping mode of action.

Reference

ZONG, Lili, *et al.* Epidithiodiketopiperazines : strain-promoted thiol-mediated cellular uptake at the highest tension. *ACS central science*, 2017, vol. 3, no. 5, p. 449-453

DOI : 10.1021/acscentsci.7b00080

Available at:

<http://archive-ouverte.unige.ch/unige:95404>

Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ
DE GENÈVE

Epidithiodiketopiperazines: Strain-Promoted Thiol-Mediated Cellular Uptake at the Highest Tension


Lili Zong,[†] Eline Bartolami,[†] Daniel Abegg, Alexander Adibekian,[‡] Naomi Sakai, and Stefan Matile*[‡]

Department of Organic Chemistry, University of Geneva, CH-1211 Geneva, Switzerland

National Centre of Competence in Research (NCCR) Chemical Biology, Geneva, Switzerland

S Supporting Information

ABSTRACT: The disulfide dihedral angle in epidithiodiketopiperazines (ETPs) is near 0°. Application of this highest possible ring tension to strain-promoted thiol-mediated uptake results in efficient delivery to the cytosol and nucleus. Compared to the previous best asparagusic acid (AspA), ring-opening disulfide exchange with ETPs occurs more efficiently even with nonactivated thiols, and the resulting thiols exchange rapidly with nonactivated disulfides. ETP-mediated cellular uptake is more than 20 times more efficient compared to AspA, occurs without endosomal capture, depends on temperature, and is “unstoppable” by inhibitors of endocytosis and conventional thiol-mediated uptake, including siRNA against the transferrin receptor. These results suggest that ETP-mediated uptake not only maximizes delivery to the cytosol and nucleus but also opens the door to a new multitarget hopping mode of action.



INTRODUCTION

Epidithiodiketopiperazines (ETPs) such as verticillin **1** are an intriguing family of natural products with a broad variety of biological activities (Figure 1A).^{1–8} Their complex structures have attracted considerable interest in synthetic organic chemistry. The distinguishing feature of ETPs is the bicyclic disulfide with the CSSC dihedral angle $\theta \approx 0^\circ$ (5.7° and 8.6° have been observed in crystals, Figure 1B).^{5,6} This is remarkable because relaxed disulfides have $\theta \approx 90^\circ$.⁹ Despite having the highest possible strain energy, ETPs are stable,





Figure 1. (A) Structure of verticillin **1**, a representative ETP natural product, and 1,2-dithietane **2**. (B) Structure of ETP transporter **6** with AspA control **5** and examples for decreasing disulfide ring tension.

unlike 1,2-dithietanes, which occur only as reactive intermediates except for rare and remarkable exceptions such as dithiatopazine **2**.¹⁰

We became interested in disulfide ring tension with regard to cellular uptake.^{11–14} Disulfides in general are increasingly recognized to enter cells by thiol-mediated uptake, i.e., covalent attachment by disulfide exchange with exofacial thiols followed by efficient uptake via diverse, to a good part unknown mechanisms.^{11–23} The emergence of thiol-mediated uptake called for the application of ring tension.¹¹ Uptake efficiencies were found to increase with ring tension from relaxed disulfides **3** with $\theta \approx 90^\circ$ to lipoic acid derivatives **4** with $\theta = 35^\circ$ and asparagusic acid derivatives **5** with $\theta = 27^\circ$.^{12,13} The most efficient “AspA tag” as in **5** allowed the delivery of functional peptides,¹⁴ liposomes and polymersomes¹³ into cells, and the transferrin receptor (TFRC) has been identified as one of the targets.¹⁴ The power and promise of strain-promoted thiol-mediated uptake at $\theta = 27^\circ$ provided a compelling incentive to drive disulfide ring tension to the extreme. To tackle this challenge, ETPs appeared just perfect. Their high reactivity in disulfide exchange reactions was predicted computationally and demonstrated experimentally to be crucial for the function of some natural ETPs.^{1–9,23} Here, we introduce “ETP tags” for the “unstoppable” strain-promoted delivery of model probes **6** to the cytosol and nucleus, and reveal a new mechanism with distinct characteristics.

Received: February 17, 2017

Published: April 6, 2017

Scheme 1^a


^a(a) K_2CO_3 , CH_3CN , rt, 12 h, 54%; (b) DCC, DMAP, Et_3N , CH_2Cl_2 , rt, 24 h, 74%; (c) TFA, CH_2Cl_2 , 0 °C to rt, 30 min; (d) toluene, reflux, 6 h, 79% (from 14); (e) NaH, THF, 0 °C to rt, 12 h, 87%; (f) 1. NBS, AIBN, cyclohexane, reflux, 2 h, 2. 12, CH_2Cl_2 , rt, 12 h; 34% (7.5% *trans*); (g) 1. NH_3 , MeOH, rt, 30 min, 2. 1, CH_2Cl_2 , rt, 30 min, 63%; (h) TFA, CH_2Cl_2 , rt, 2 h; (i) NHS, DCC, THF, rt, 24 h; (j) R- NH_2 (see Figure 1), DMF, rt, 2 h, 68% (three steps from 19).

RESULTS AND DISCUSSION

The ETP tag 7 was synthesized, as in biosynthesis, using exclusively C_2 building blocks derived from acetate, i.e., 8–12 (Scheme 1). At the beginning, chloroacetate 8 was reacted with ethylamine 9. The resulting secondary amine 13 was coupled with Boc-protected glycine 10. Liberation of the amine in the obtained dipeptide 14 prepared for the cyclization of 15. The resulting diketopiperazine heterocycle 16 was alkylated with bromoacetate 11. With dilactam 17, a key intermediate was reached. The sulfur atoms were introduced via radical bromination followed by substitution with thioacetate 12.²⁴ The *cis* isomer 18 was obtained as the major product (4.5:1), easily separated from the *trans* isomer, and assigned by a strong NOE between the two remaining endocyclic hydrogens. Hydrolysis of the thioesters 18 with ammonia afforded the free thiols, which were immediately oxidized with molecular iodine to afford the high-tension ETP disulfide 19 in excellent 63% yield as a pale yellow solid. The bicyclic ETP scaffold remained intact during the acid-catalyzed removal of the *t*Bu protecting group in 19, the activation of the resulting acid 20 with *N*-hydroxysuccinimide (NHS), and reaction of the resulting ETP tag 7 with amines of free choice, here a fluorescent model substrate, under mildest conditions, to give the CF-ETP conjugate 6 in 68% yield.

In D_2O at pD 8.0, equimolar DTT reduced 5 mM ETP 20 instantaneously and completely to dithiol 21 (Table 1, entry a, Figure S10). This was also true at pD 5.5 and with 2 equiv of glutathione (GSH) at pD 8.0 (Table 1, entries b and c, Figures S9 and S14). At pD 5.5 with GSH, the consumption of the hyperstrained disulfide 20 reached 50% within the time needed to set up and record an 1H NMR spectrum (Table 1, entry d, Figure S13). In sharp contrast, AspA 23 reacted slowly with DTT and failed to react with GSH under these conditions (Table 1, entries a–d, Figures S11, S12, S15, and S16).

To explore the formation of strained disulfides by disulfide exchange, dithiols 21 and 24 were prepared in situ by 1 equiv of TCEP. Subsequent addition of 1 equiv of DTNB 25 in neutral water gave rise to the strained ETP 20 and AspA 23 instantaneously (Table 1, entry e, Figures S17 and S18). With 2 equivalents of oxidized glutathione GSSG, a much less reactive disulfide, the reduced ETP 21 exchanged rapidly into the tension-free mixed disulfide 22 (Table 1, entry f, Figure S19), and, with time, ring closure into hyperstrained ETP 20

Table 1. Disulfide Exchange Cycles^a

E ^b	S ^c	ETP ^a			AspA ^a	
		pD ^d	t ^e	η (%) ^f	t ^e	η (%) ^f
20 → 21						
a)	DTT	8.0	<5 min	100	30 min	98
b)	DTT	5.5	<5 min	100	60 min	14
c)	GSH	8.0	<5 min	100	18 h	0
d)	GSH	5.5	<5 min	50 ^g	18 h	0
21 → 22 → 20 ^h						
e)	DTNB	7.2	<5 min	100 (20)	<5 min	100
f)	GSSG	7.2	<5 min	100 (22)	30 min	0
g)	GSSG	7.2	16 h	100 ⁱ	16 h	70
24 → 23 ^h						

^aFor ETP 20 and AspA 23 (5 mM), determined by 1H NMR kinetics; for original spectra, see Figures S9–S22. ^bEntry, letters refer to reaction scheme. ^cSubstrates, DTT: 1,4-dithiothreitol, 5 mM (1 equiv); GSH: glutathione, 10 mM (2 equiv), DTNB: 5,5-dithio-bis(2-nitrobenzoic acid) (25, Figure 3), 5 mM (1 equiv); GSSG: oxidized GSH, 5 mM (1 equiv). ^dpD in 0.1 M aqueous (D_2O) sodium phosphate buffer. ^eReaction time at rt, in D_2O . ^fConversion, determined from integration of NMR signals. Unless specified, only designated starting products are formed. ^gUnidentified product formed. ^hFully reduced starting materials 21 and 24 were prepared in situ from 20 and 23 with 1 equiv TCEP (tris(2-carboxyethyl)phosphine, entries e–g). ⁱETP 20 (20%) could be identified from the mixture of products at least partially arisen from the decomposition of 20.

could be observed (Table 1 entry g, Figure S20). The high reactivity of reduced ETP 21 could be ascribed to the lower than usual pK_a of thiols due to the presence of lactam nitrogen and carbonyl groups on the α position. Besides high tension,

this increased acidity also explained the ease of ring-opening disulfide exchange ($20 \rightarrow 21/22$) and the reluctance of ring closure ($22 \rightarrow 20$). In comparison, the thiols of the reduced AspA control **24** were much less reactive toward nonactivated disulfides (Table 1, entry f, Figure S21), whereas formation of the less strained dithiolane ring was faster (Table 1, entry g, Figure S22). Control experiments without GSSG resulted in very little auto-oxidative ring closure to **20** or **23**, thus demonstrating that the rings form through the mixed disulfides, such as **22**, by disulfide exchange reactions. In summary, compared to AspAs, ETPs are (1) more reactive in ring-opening disulfide exchange with nonactivated thiols, also under acidic conditions, (2) more reactive in their reduced form with nonactivated disulfides, and (3) less efficient in ring-closing disulfide exchange to go full cycle and reproduce the hyperstrained ETPs in neutral water (Table 1).

The uptake of the green-fluorescent CF-ETP conjugate **6** into HeLa Kyoto cells was monitored by confocal laser scanning microscopy (CLSM). Incubation with $10 \mu\text{M}$ **6** in Leibovitz medium for 1 h at 37°C resulted in intense homogeneous emission from the cytosol and particularly from the nuclei, including nuclei that were poorly stained by Hoechst 33342 (Figure 2C). This result contrasted sharply from the




Figure 2. CLSM images of HeLa Kyoto cells after 1 h of incubation with $10 \mu\text{M}$ CF-NH₂ (A), CF-AspA **5** (B), and CF-ETP **6** (C) in Leibovitz medium at 37°C (top), together with Hoechst 33342 to stain the nuclei (bottom).

uptake of the AspA control **5**, which failed to reach the nucleus and produced mostly punctate emission at much lower intensity (Figure 2B). The same distinct differences between ETP **6** and AspA **5** were observed in several other cell lines (Figure 3A,B).

The punctate emission obtained with AspA **5** can be assigned with confidence to receptor-mediated delivery into endosomes.¹⁴ The absence of punctate emission suggested that contrary to AspA **5**, ETP **6** does not suffer from endosomal capture and is delivered exclusively to the cytosol and nucleus. Different from the polycationic CPDs,¹¹ accumulation of the overall anionic ETPs in the nuclei is not driven by ion pairing and thus not limited to the DNA-rich nucleoli. Possibly, the presence of target proteins with reactive thiols, such as histone methyl transferase,^{7,8} dictates the intracellular distribution of ETPs. ETPs continued to deliver efficiently at concentrations as low as 500 nM , whereas detectable uptake of AspAs stopped below $5 \mu\text{M}$ (Figure 3C,D). Still higher intensities obtained




Figure 3. Microscopic images of (A, C) ETP **6** and (B, D) AspA **5** after (A, B) incubation (Leibovitz medium, 37°C) at $10 \mu\text{M}$ with A431, Huh7, MCF7, and PC-3 cells (left to right, automated microscope images) and (C, D) incubation with HeLa Kyoto cells at $10 \mu\text{M}$, $5 \mu\text{M}$, $1 \mu\text{M}$, and 500 nM (left to right; CLSM images merged with differential interference contrast (DIC)).

with ETPs at 500 nM than with AspAs at $10 \mu\text{M}$ suggested that ETPs are at least 20 times more active (Figure 3C,D).

As many natural ETPs are toxins, the MTT assay was employed to assess the toxicity of ETP tags in HeLa Kyoto cells. This assay reports on the enzymatic conversion of the tetrazolium dye MTT into formazan, that is, the metabolic activity of the cells.²⁵ The positive control, polyarginine (pR), was confirmed to be cytotoxic at $10 \mu\text{M}$ (Figure 4A).¹¹ Under the same conditions, ETP **6** and AspA **5** were not toxic (Figure 4A).

Flow cytometry analysis confirmed the impression from CLSM images that the hyperstrained ETP **6** is much more active than the AspA control **5** (Figure 4B). The loss of essentially all activity at 4°C is commonly interpreted as indication of uptake by endocytosis (Figure 4B). However, other possible explanations such as changes in disulfide exchange kinetics, membrane fluidity, etc., should not be forgotten, particularly since all common endocytosis inhibitors were inactive. Namely, insensitivity toward chlorpromazine (CPZ) excluded clathrin-mediated endocytosis, methyl- β -cyclodextrin (m β CD) caveolae-mediated endocytosis, and wortmannin and cytochalasin B (cytoB) ruled out macropinocytosis (Figure 4B).^{11,22,26–28}

Contrary to AspA controls,^{12,13} the removal of thiols on cell surfaces with maleimide **26**,²² iodoacetamide **27**,¹² and the most powerful hypervalent iodine reagent **28**²⁹ failed to inactivate ETP **6** (Figure 4C). Similarly, the presence of 10% serum²² caused only a minor $\sim 25\%$ reduction of ETP uptake (Figures 4C and S5). DTNB **25** was special because this reagent converts thiols on cell surfaces into activated disulfides. After preincubation with 1.2 mM DTNB, uptake activity of ETP **6** indeed dropped to $\sim 65\%$ (Figure 4C). However, unlike AspA tags,¹² ETP activity increased rather than decreased with further increasing DTNB concentration to reach saturation




Figure 4. (A) Cell viability from MTT assay for 10 μM transporters in HeLa Kyoto cells; pR: polyarginine. (B) Flow cytometry data for HeLa Kyoto cells and **6** with endocytosis inhibitors (CPZ, m β CD, wort, cytoB), temperature dependence, and comparison to **5**, normalized to 1 for **6**. (C) Flow cytometry data for **6** and HeLa Kyoto cells that were preincubated with inhibitors (**25**–**28**, 0.02–2 mM) and activators (DTT, TCEP, 2 mM) of thiol-mediated uptake, with 10% serum, and with TFRC siRNA (quantified using automated microscope). Shown are average values \pm error.

near 80%. Also unlike AspA controls,¹² preincubation of the cells with DTT or TCEP did not strongly increase the activity of ETPs (Figure 4C). Most importantly, the knockdown of the transferrin receptor (TFRC) with siRNA inhibited the uptake of AspA controls¹⁴ but failed to inhibit ETP-mediated uptake. The observed partial inactivation by TFRC knockdown down to \sim 65% was most revealing (Figures 4C and S7). It supported that (1) ETPs operate by thiol-mediated uptake, that is, dynamic covalent disulfide exchange on the cell surface, (2) ETPs do not depend on single targets such as the transferrin receptor, and (3) ETPs have access to targets that are inaccessible to AspA controls.

CONCLUSIONS

In this report, we introduce ETP-mediated cellular uptake. Epidithiodiketopiperazines attracted our attention to drive ring tension in cyclic disulfides to the maximum, i.e., a CSSC dihedral angle of $\sim 0^\circ$. However, rather than simply maximizing the efficiency of strain-promoted thiol mediated uptake,^{12–14} completely new, exceptionally promising properties emerged. ETP-mediated uptake excels with the efficient, nontoxic delivery to cytosol and particularly nucleus, without any endosomal capture, sensitive to temperature but “unstoppable” by all conventional inhibitors of endocytosis and thiol-mediated uptake. This poor responsiveness to inhibitors and activators such as cytochalasin B, DTT, Ellman’s reagent, TFRC siRNA, or serum indicated that the unique reactivity of ETPs is decisive for function. High reactivity of ETPs in both oxidized and reduced form allows for covalent capture by nonactivated thiols and disulfides in cellular targets²⁹ that are otherwise beyond reach (Table 1, entries a–d, f, h–i). Moreover, the possibility of repeated disulfide-exchange cycles in neutral water suggested that ETPs can change targets during uptake (Table 1, entries a–d, e–g). Such a multitarget hopping mechanism could explain the characteristics found for ETP-mediated uptake: namely, efficient delivery to cytosol and nucleus, without endosomal capture, without toxicity. These stunning characteristics invite the highest expectations with regard to the general,

covalent, charge-free delivery of substrates of biological and medicinal relevance.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscentsci.7b00080.

Detailed experimental procedures (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: stefan.matile@unige.ch.

ORCID

Alexander Adibekian: 0000-0001-6453-0244

Stefan Matile: 0000-0002-8537-8349

Author Contributions

[†]L.Z. and E.B. contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the group of A. Roux for access to and assistance with cell culture and imaging, the group of J. Waser for probes, the NMR, the MS, and the Bioimaging platforms for services, and the University of Geneva, the Swiss National Centre of Competence in Research (NCCR) Chemical Biology, the NCCR Molecular Systems Engineering and the Swiss NSF for financial support.

REFERENCES

- (1) Kim, J.; Movassaghi, M. Biogenetically-inspired total synthesis of epidithiodiketopiperazines and related alkaloids. *Acc. Chem. Res.* **2015**, *48*, 1159–1171.
- (2) Borthwick, A. D. 2,5-Diketopiperazines: synthesis, reactions, medicinal chemistry, and bioactive natural products. *Chem. Rev.* **2012**, *112*, 3641–3716.
- (3) Jiang, C.-S.; Müller, W. E. G.; Schröder, H. C.; Guo, Y.-W. Disulfide- and polysulfide-containing metabolites from marine organisms. *Chem. Rev.* **2012**, *112*, 2179–2207.
- (4) Waring, P.; Chai, C. L. L. The multiple properties of gliotoxin and other epipolythiodioxopiperazine metabolites. *Aust. J. Chem.* **2015**, *68*, 178–183.
- (5) Park, H. B.; Kwon, H. C.; Lee, C.-H.; Yang, H. O. Glionitrin A, an antibiotic-antitumor metabolite derived from competitive interaction between abandoned mine microbes. *J. Nat. Prod.* **2009**, *72*, 248–252.
- (6) Overman, L. E.; Sato, T. Construction of epidithiodioxopiperazines by directed oxidation of hydroxyproline-derived dioxopiperazines. *Org. Lett.* **2007**, *9*, 5267–5270.
- (7) Iwasa, E.; Hamashima, Y.; Sodeoka, M. Epipolythiodiketopiperazine alkaloids: total synthesis and biological activities. *Isr. J. Chem.* **2011**, *51*, 420–433.
- (8) Cherblanc, F. L.; Chapman, K. L.; Reid, J.; Borg, A. J.; Sundriyal, S.; Alcazar-Fuoli, L.; Bignell, E.; Demetriades, M.; Schofield, C. J.; DiMaggio, P. A., Jr; Brown, R.; Fuchter, M. J. On the histone lysine methyltransferase activity of fungal metabolite chaetocin. *J. Med. Chem.* **2013**, *56*, 8616–8625.
- (9) Singh, R.; Whitesides, G. M. Comparisons of rate constants for thiolate-disulfide interchange in water and in polar aprotic solvents using dynamic protons NMR line shape analysis. *J. Am. Chem. Soc.* **1990**, *112*, 1190–1197.
- (10) Nicolau, K. C.; Hwang, C.-K.; DeFrees, S.; Stylianides, N. A. Novel chemistry of dithiatopazine. *J. Am. Chem. Soc.* **1988**, *110*, 4868–4869.

(11) Gasparini, G.; Bang, E.-K.; Molinard, G.; Tulumello, D. V.; Ward, S.; Kelley, S. O.; Roux, A.; Sakai, N.; Matile, S. Cellular uptake of substrate-initiated cell-penetrating poly(disulfide)s. *J. Am. Chem. Soc.* **2014**, *136*, 6069–6074.

(12) Gasparini, G.; Sargsyan, G.; Bang, E.-K.; Sakai, N.; Matile, S. Ring tension applied to thiol-mediated cellular uptake. *Angew. Chem., Int. Ed.* **2015**, *54*, 7328–7331.

(13) Chuard, N.; Gasparini, G.; Moreau, D.; Lörcher, S.; Palivan, C.; Meier, W.; Sakai, N.; Matile, S. Strain-promoted thiol-mediated cellular uptake of giant substrates: liposomes and polymersomes. *Angew. Chem., Int. Ed.* **2017**, *56*, 2947–2950.

(14) Abegg, D.; Gasparini, G.; Hoch, D. G.; Shuster, A.; Bartolami, E.; Matile, S.; Adibekian, A. Strained cyclic disulfides enable cellular uptake by reacting with the transferrin receptor. *J. Am. Chem. Soc.* **2017**, *139*, 231–238.

(15) Aubry, S.; Burlina, F.; Dupont, E.; Delaroche, D.; Joliot, A.; Lavielle, S.; Chassaing, G.; Sagan, S. Cell-surface thiols affect cell entry of disulfide-conjugated peptides. *FASEB J.* **2009**, *23*, 2956–2967.

(16) Oupický, D.; Li, J. Bioreducible polycations in nucleic acid delivery: past, present, and future trends. *Macromol. Biosci.* **2014**, *14*, 908–922.

(17) Ling, Y. Y.; Ren, J.; Li, T.; Zhao, Y. B.; Wu, C. L. POEGMA-based disulfide-containing fluorescent probes for imitating and tracing noninternalization-based intracellular drug delivery. *Chem. Commun.* **2016**, *52*, 4533–4536.

(18) Fu, J.; Yu, C.; Li, L.; Yao, S. Q. Intracellular delivery of functional proteins and native drugs by cell-penetrating poly(disulfide)s. *J. Am. Chem. Soc.* **2015**, *137*, 12153–12160.

(19) Brülisauer, L.; Kathriner, N.; Prenrecaj, M.; Gauthier, M. A.; Leroux, J.-C. Tracking the bioreduction of disulfide-containing cationic dendrimers. *Angew. Chem., Int. Ed.* **2012**, *51*, 12454–12458.

(20) Kichler, A.; Remy, J. S.; Boussif, O.; Frisch, B.; Boeckler, C.; Behr, J.-P.; Schuber, F. Efficient gene delivery with neutral complexes of lipospermine and thiol-reactive phospholipids. *Biochem. Biophys. Res. Commun.* **1995**, *209*, 444–450.

(21) Feener, E. P.; Shen, W. C.; Ryser, H. J. P. Cleavage of disulfide bonds in endocytosed macromolecules. A processing not associated with lysosomes or endosomes. *J. Biol. Chem.* **1990**, *265*, 18780–18785.

(22) Li, T.; Takeoka, S. Enhanced cellular uptake of maleimide-modified liposomes via thiol-mediated transport. *Int. J. Nanomed.* **2014**, *9*, 2849–2861.

(23) Bernardo, P. H.; Brasch, N.; Chai, C. L. L.; Waring, P. A novel redox mechanism for the glutathione-dependent reversible uptake of a fungal toxin in cells. *J. Biol. Chem.* **2003**, *278*, 46549–46555.

(24) Kang, S. W.; Kang, D. H.; Lee, D. J. Epidithiodioxopiperazine compound or its derivatives, and the use of thereof. WO 2014/189343 A1.

(25) Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55–63.

(26) Khalil, I. A.; Kogure, K.; Akita, H.; Harashima, H. Uptake pathways and subsequent intracellular trafficking in nonviral gene delivery. *Pharmacol. Rev.* **2006**, *58*, 32–45.

(27) Tang, H.; Yin, L.; Kim, K. H.; Cheng, J. Helical poly(arginine) mimics with superior cell-penetrating and molecular transporting properties. *Chem. Sci.* **2013**, *4*, 3839–3844.

(28) McNaughton, B. R.; Cronican, J. J.; Thompson, D. B.; Liu, D. R. Mammalian cell penetration, siRNA transfection, and DNA transfection by supercharged proteins. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 6111–6116.

(29) Abegg, D.; Frei, R.; Cerato, L.; Hari, D. P.; Wang, C.; Waser, J.; Adibekian, A. Proteome-wide profiling of targets of cysteine reactive small molecules by using ethynyl benziodoxolone reagents. *Angew. Chem., Int. Ed.* **2015**, *54*, 10852–10857.