

RESEARCH ARTICLE

Epinephrine Activation of the β_2 -Adrenoceptor Is Required for IL-13-Induced Mucin Production in Human Bronchial Epithelial Cells

Nour Al-Sawalha^{1†}, Indira Pokkunuri¹, Ozozoma Omoluabi², Hosu Kim¹, Vaidehi J. Thanawala¹, Adrian Hernandez², Richard A. Bond^{1,2}, Brian J. Knoll^{1,2*}

1 Department of Pharmacological and Pharmaceutical Sciences, University of Houston, 4800 Calhoun Road, Houston, Texas, 77204, United States of America, **2** Department of Biology and Biochemistry, University of Houston, 4800 Calhoun Road, Houston, Texas, 77204, United States of America

† Current address: Department of Clinical Pharmacy, College of Pharmacy, Jordan University of Science and Technology, Irbid, Jordan

* bkno1@Central.UH.edu



CrossMark
click for updates

OPEN ACCESS

Citation: Al-Sawalha N, Pokkunuri I, Omoluabi O, Kim H, Thanawala VJ, Hernandez A, et al. (2015) Epinephrine Activation of the β_2 -Adrenoceptor Is Required for IL-13-Induced Mucin Production in Human Bronchial Epithelial Cells. PLoS ONE 10(7): e0132559. doi:10.1371/journal.pone.0132559

Editor: Shama Ahmad, University of Alabama at Birmingham, UNITED STATES

Received: July 29, 2014

Accepted: June 17, 2015

Published: July 10, 2015

Copyright: © 2015 Al-Sawalha et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: BJK and RAB were supported by National Institutes of Health RO1A179236. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Brian J. Knoll is an Academic Editor for PLOS ONE. RA Bond is a co-inventor on a provisional patent application 2011, joint between MD Anderson and the University of Houston - "The steroid-sparing effects of betaadrenergic inverse

Abstract

Mucus hypersecretion by airway epithelium is a hallmark of inflammation in allergic asthma and results in airway narrowing and obstruction. Others have shown that administration a T_H2 cytokine, IL-13 is sufficient to cause mucus hypersecretion *in vivo* and *in vitro*. Asthma therapy often utilizes β_2 -adrenoceptor (β_2AR) agonists, which are effective acutely as bronchodilators, however chronic use may lead to a worsening of asthma symptoms. In this study, we asked whether β_2AR signaling in normal human airway epithelial (NHBE) cells affected mucin production in response to IL-13. This cytokine markedly increased mucin production, but only in the presence of epinephrine. Mucin production was blocked by ICI-118,551, a preferential β_2AR antagonist, but not by CGP-20712A, a preferential β_1AR antagonist. Constitutive β_2AR activity was not sufficient for IL-13 induced mucin production and β -agonist-induced signaling is required. A clinically important long-acting β -agonist, formoterol, was as effective as epinephrine in potentiating IL-13 induced MUC5AC transcription. IL-13 induced mucin production in the presence of epinephrine was significantly reduced by treatment with selective inhibitors of ERK1/2 (FR180204), p38 (SB203580) and JNK (SP600125). Replacement of epinephrine with forskolin + IBMX resulted in a marked increase in mucin production in NHBE cells in response to IL-13, and treatment with the inhibitory cAMP analogue Rp-cAMPS decreased mucin levels induced by epinephrine + IL-13. Our findings suggest that β_2AR signaling is required for mucin production in response to IL-13, and that mitogen activated protein kinases and cAMP are necessary for this effect. These data lend support to the notion that β_2AR -agonists may contribute to asthma exacerbations by increasing mucin production via activation of β_2AR s on epithelial cells.

agonists." At this point this patent has not been assigned and has no value or license agreement in place or pending issuance. RA Bond is also a minor shareholder in Invion, a company in which he was scientific co-founder. RA Bond sold off almost his entire stock ownership and now owns ~\$10,000 worth of stock (as of 12 June 2015). RA Bond has no other role than shareholder with Invion, and is not on any board and has no consulting agreement. This does not alter the authors' adherence to PLOS ONE Editorial policies and criteria.

Introduction

Asthma is a chronic inflammatory disease characterized by airway hyperreactivity, subepithelial fibrosis, airway smooth muscle hyperplasia and mucous metaplasia [1]. Mucous metaplasia is an increase in the number of mucus-secreting goblet cells in the epithelium [2] that results in increased mucus synthesis and secretion. Excessive accumulation of airway mucus leads to the formation of mucous plugs that reduce the effective airway diameter and increase airway resistance. Patients who die of severe asthma attacks often exhibit goblet cell hyperplasia, mucus accumulation and large mucus plugs of unusual solidity due to high mucin content in their peripheral airways compared to asthmatic patients who did not die of acute attacks [3].

Allergic asthma has properties of a type I hypersensitivity, in which type 2 T-helper lymphocytes and type 2 innate lymphoid cells contribute to produce a distinctive set of cytokines in the airways, including IL-5, IL-9 and IL-13 [4]. Although the allergic airway also contains diverse hematopoietic and parenchymal cells, and factors secreted by them, airway epithelial overexpression of IL-13 or airway instillation of IL-13 is sufficient to induce mucous metaplasia in mice [5, 6]. Airway epithelium is essential and sufficient for mucous metaplasia induced by IL-13, and this is dependent on the expression of STAT6 (which mediates the action of IL-13) in the epithelium [7]. The epithelium has been suggested by many studies to play an important role in the pathogenesis of asthma [7–10] as well as being a key contributor to the mucus plugs responsible for asthma mortality [11]. Sputum of asthmatic patients show elevated levels of IL-13 and its presence is negatively associated with therapeutic responsiveness [12].

The MUC5AC gene encodes the major component of mucin in human airways, and induction of MUC5AC transcription by IL-13 is observed in cultured human airway epithelium [10, 13]. However other factors also are required for MUC5AC transcription in these cells. Signaling through the EGF [10] and TGF- β 2 [14] receptors is required for IL-13 to induce MUC5AC transcription, and in the promoter region of the gene there are binding sites for numerous diverse transcription factors [15], although none for STAT-6, suggesting the action of multiple intersecting and possibly indirect pathways. Recently, we found evidence for the involvement of β_2 -adrenoceptor (β_2 AR) signaling in the pathogenesis of asthma. Pharmacologic or genetic ablation of β_2 AR signaling causes reductions in mucous metaplasia, airway cellularity and airway hyperreactivity (AHR) in a murine asthma model [16, 17]. Thus, among the other pathways already mentioned, some that are initiated or influenced by β_2 ARs may also be involved in the regulation of MUC5AC transcription.

Due to the complexity of the signaling pathways that are involved in mediating mucus production, and the involvement of diverse cell types in whole animal models, we undertook to study human airway epithelial cells cultured in low concentrations of retinoic acid, conditions where mucin expression is normally minimal. We investigated the requirement for β_2 AR signaling in the transcription of the MUC5AC gene, the expression of MUC5AC protein and intracellular mucin accumulation in response to IL-13. In addition, we examined signaling components downstream of β_2 AR that may be required for this response.

Materials and Methods

Cell Culture

Normal human bronchial epithelial (NHBE) cells were obtained from Lonza (Walkersville, MD). The cells were seeded on Transwell-culture inserts (0.45 μ m pore size) at 2×10^4 / cm^2 and grown in 5% CO_2 /95% air at 37°C in differentiation medium: 50% bronchial epithelial basal media, 50% DMEM high glucose and supplemented with 30 μ g/ml bovine pituitary extract, 0.5 μ g/ml BSA, 0.5 μ g/ml epinephrine, 50 μ g/ml gentamycin, 50 ng/ml amphotericin B,

0.5 ng/ml human EGF, 0.5 μ g/ml hydrocortisone, 5 μ g/ml insulin, 7 ng/ml triiodothyronine, 10 μ g/ml transferrin and 0.1 ng/ml retinoic acid. The cells were cultured with epinephrine for ~8 days until they reached confluence, then the apical medium was removed and air-liquid interface (ALI) was established. The cells were then treated with 20 ng/ml of IL-13 combined with different antagonists or inhibitors. In some experiments, the cells were grown in the absence of epinephrine 72 hours before they reached ALI. Compound-related toxicity was assessed through the dryness of the apical surface of the cultured NHBE cells [18]. Cells grown in the absence or presence of epinephrine formed tight junctions that produced equal increases in transepithelial electrical resistance (TEER) suggesting similar epithelial barrier function, consistent with previously published findings [19]. The formation of tight junctions was also confirmed by the presence of ZO-1 on the cell peripheries by immunofluorescence (data not shown).

Real-Time PCR Analysis of MUC5AC Expression

Total RNA was extracted from cells using Trizol according to manufacturer's protocol. cDNA was generated from 5 μ g of total RNA and MUC5AC and 18S mRNA were quantified using the Taqman Gene Expression Assay (Applied Biosystems, Grand Island, NY) and analyzed by real time quantitative PCR (ABI PRISM 7000 Sequence Detection System, Applied Biosystems). The threshold cycles (C_t) for MUC5AC of treated groups was compared to the control groups and normalized to 18S. Relative MUC5AC expression was calculated using Delta-Delta CT method.

Immunoblotting

NHBE cells were lysed in a buffer containing 20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM Na_2EDTA , 1 mM EGTA, 1% Triton, 2.5 mM sodium pyrophosphate, 1 mM β -glycerophosphate, 1 mM Na_3VO_4 and 1 μ g/ml leupeptin (Cell Signaling, Danvers, MA) combined with protease inhibitor cocktail tablet (Roche Applied Sciences, Indianapolis, IN). Total protein concentration was determined by BCA Protein Assay Kit (Thermo Fisher Scientific Inc, Rockford, IL), according to the manufacturer's protocols. Thirty micrograms of protein per lane were subjected to SDS-PAGE using 10% Tris-HCl gels (Bio-Rad Laboratories, Hercules, CA) and transblotted to PVDF membrane (EMD Millipore, Billerica, MA). Membranes were blocked in 3% BSA for 1 hour at room temperature and then incubated with antibodies against phospho-ERK1/2 (Cell Signaling, #4370S), phospho-p38 (Santa Cruz Biotechnology, #sc-17852-R) or phospho-c-Jun (EMD Millipore, #06-659) overnight at 4°C followed by treatment for 1 hour with HRP conjugated secondary antibody. The protein bands were developed using SuperSignal West Pico chemiluminescent substrate (Thermo Fisher Scientific) according to manufacturer's recommendations. A CCD camera (Fluorochem 8800) was used to collect the digital images and AlphaEase software (ProteinSimple, Santa Clara, CA) to quantify band density. The membranes were then stripped and probed with GAPDH antibody (EMD Millipore, #MAB374). The signal density of the phosphorylated proteins was normalized with that of GAPDH.

Periodic Acid Fluorescent Schiff's (PAFS) Staining

The apical surfaces of NHBE cells were washed with PBS, fixed in 4% paraformaldehyde (PFA) and permeabilized with Triton X-100. The inserts were stained with PAFS as described previously [20]. Red fluorescence of mucin was detected when the slides were excited at 380–580 nm and observed at 600–650 nm while the nucleic acid and cytoplasm of the cells fluorescence green when observed at a lower wavelength (380–500 nm and 450–475 nm excitation and

emission wavelengths respectively) [20]. Images were captured using an Olympus DUS spinning disc confocal microscope maintained in the College of Pharmacy Imaging Core. To correct for insert background, empty inserts were stained and used as a negative control. To maintain consistency in subsequent image analysis, we used the same channel-specific threshold when capturing all images. The ratio of integrated mucin density to integrated nucleic acid and cytoplasm density was calculated using Image J (NIH).

Immunofluorescence Labeling for Mucin 5AC

The apical surfaces of NHBE cells were washed with PBS, fixed in 4% paraformaldehyde (PFA) and permeabilized with Triton X-100. The inserts were incubated for 15 minutes with 10% normal goat serum at room temperature followed by Mucin 5AC Antibody (H-160, Santa Cruz Biotechnology) [21] overnight at 4°C. After washing the inserts with PBS, they were incubated with Alexa 594 goat anti-rabbit secondary antibody for 1 hour at room temperature. DAPI, at final concentration of 1 $\mu\text{g/ml}$, was used to counterstain the nuclei. Slides incubated with primary antibody diluents were used as negative controls. Images were captured by confocal microscopy and the same channel-specific threshold was maintained when capturing the images. The ratio of integrated mucin 5AC density of each group to the integrated mucin 5AC density of the corresponding control group was calculated using Image J software (NIH).

Statistical Analysis

Data are presented as means \pm SEM. All experiments were done with NHBEs from 3 donors ($N = 3$). One-way ANOVA followed by Tukey's multicomparison test for multiple group statistical analysis was performed using GraphPad Prism 4 software. $p < 0.05$ was considered statistically significant.

Results

To determine whether epinephrine increases mucus production in NHBE cells, we cultured them for 72 hours before reaching ALI in a medium that lacks epinephrine and then stimulated with 20 ng/ml of IL-13 for 14 days, also in the absence of epinephrine. Under these conditions, IL-13 did not increase the expression of MUC5AC mRNA, and in the presence of epinephrine alone, MUC5AC mRNA was scarcely detectable above baseline. But with IL-13 treatment in the presence of epinephrine, the MUC5AC expression level was increased by ~ 15 fold (Fig 1A). Similar results were obtained with cells grown in the presence of 50 nM retinoic acid (S1 Fig). To correlate mRNA expression levels with intracellular mucin 5AC accumulation and mucin glycoprotein, immunofluorescence with anti-mucin 5AC antibody and PAFS staining were used respectively. Epinephrine was required for IL-13-induced increases in the intracellular mucin 5AC and total mucin glycoproteins (Fig 1B and 1C).

To determine if a therapeutic β -agonist showed an effect similar to epinephrine, we substituted formoterol. NHBE cells were cultivated with physiological levels of RA to more closely mimic the *in vivo* state. Under these conditions, IL-13 again did not significantly increase MUC5AC transcription (though levels were slightly higher). However the addition of formoterol increased MUC5Ac transcripts to a similar degree as epinephrine (S2 Fig).

To determine the β AR subtype involved in MUC5AC expression in response to IL-13 in the presence of epinephrine, NHBE cells were incubated with either a preferential β_2 AR antagonist (1 μM ICI-118,551) or a preferential β_1 AR antagonist (3 μM CGP-20712A). ICI-118,551 completely abolished ($>99\%$) IL-13 induced MUC5AC expression (0.039 ± 0.038 fold vs 15.99 ± 1.48 fold increase by IL-13. $p < 0.05$). On the other hand, CGP-20712A did not affect the MUC5AC expression level (14.75 ± 0.96 fold vs 15.99 ± 1.48 fold increase by IL-13,




Fig 1. Epinephrine is required for mucin production in response to IL-13 in NHBE cells. **A:** NHBE cells were grown in the presence or absence of 3 μ M epinephrine. At ALI, the cells were treated with 20 ng/ml IL-13 for 14 days, total RNA was harvested and then MUC5AC transcripts were measured by qRT-PCR. Data are presented as fold change compared to the corresponding treatment control (in the absence of IL13). **B:** Representative images of immunofluorescence with a rabbit antibody against human mucin 5AC (red) (scale bar = 100 μ m). The Transwell membranes were incubated with DAPI to counterstain the nuclei (blue). Incubation with antibody diluent showed no red fluorescence (data not shown). The ratio of integrated fluorescence density of each group to the integrated mucin 5AC density of the corresponding control group

was calculated and expressed as fold change. **C:** PAFS staining of NHBE cells to quantify total intracellular mucin glycoproteins. Representative images are shown. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to the corresponding control cells (in the absence of IL-13 treatment). Data are presented as means \pm SEM from three donors. *, † and ¥ indicate $p < 0.05$ significance as compared to + epinephrine, -epinephrine and -epinephrine + IL-13 treated cells respectively.

doi:10.1371/journal.pone.0132559.g001

$p > 0.05$) (Fig 2A). CGP-20712A did not affect the intracellular mucin levels induced by IL-13 while ICI-118,551 brought the levels back to baseline (Fig 2B and 2C; for representative images see S3A and S3B Fig).

We next asked if the increased MUC5AC expression in response to IL-13 is due to agonist induced or constitutive β_2 AR signaling. NHBE cells were treated with 10 μ M nadolol, a non-selective β AR ligand with inverse agonist activity at β_2 ARs that blocks both constitutive and agonist-induced receptor activity, or with 10 μ M alprenolol, a non-selective β AR antagonist with no inverse agonist activity, for 14 days in combination with IL-13 and in the presence of epinephrine. Treatment with nadolol reduced IL-13 induced MUC5AC expression (3.36 ± 4.10 fold vs 25.37 ± 16.30 fold increase by IL-13, $p < 0.05$), intracellular mucin 5AC protein and mucin content (Fig 3A, 3B and 3C; for representative images see S4A and S4B Fig). Treatment with alprenolol reduced IL-13-induced MUC5AC expression to a similar extent (3.19 ± 3.73 fold vs 25.37 ± 16.30 fold increase by IL-13, $p < 0.05$) and also reduced intracellular mucin 5AC and mucin content (Fig 3A, 3B and 3C, and S4A and S4B Fig for representative images).

To investigate the role of mitogen activated protein kinases (MAPKs), we examined their activation using antibodies specific for phosphorylated (activated) MAPKs. In the absence of epinephrine, IL-13 did not affect the phosphorylation of ERK1/2 (Fig 4A), c-Jun (Fig 4B) or p38 (Fig 4C) as compared to their corresponding controls. When epinephrine was included in the medium, IL-13 induced an approximately 3-fold increase in the phosphorylation of ERK1/2 and c-Jun when compared to their corresponding controls (Fig 4A and 4B). However, phosphorylation of p38 was unaffected by IL-13 even in the presence of epinephrine (Fig 4C). Next, we treated NHBE cells with 3 μ M FR180204, SP600125 or SB203580 (inhibitors of ERK1/2, JNK and p38 respectively) in combination with IL-13 and epinephrine for 14 days. All three MAPKs inhibitors significantly reduced MUC5AC gene expression (15.18 ± 3.76 fold increase by IL-13 vs 1.82 ± 0.68 , 0.77 ± 0.39 and 0.80 ± 0.65 fold by FR180204, SP600125 and SB203580 respectively) (Fig 4D). While all MAPK inhibitors reduced the intracellular mucin 5AC protein (see Fig 4E and S5A Fig for representative images), only FR180204 and SP600125 reduced intracellular mucin content when compared to IL-13 treated cells (see Fig 4F and S5B Fig for representative images).

To explore a possible role for PKA in the induction of MUC5AC, we treated NHBE cells with a competitive cAMP analogue, Rp-cAMPS, for 14 days in combination with IL-13 and epinephrine. Rp-cAMPS did not significantly reduce the levels of MUC5AC expression at 50 μ M (5.97 ± 4.29 fold vs 12.50 ± 5.38 fold increase by IL-13, $p > 0.05$), while at 100 μ M, there was a significant reduction (2.35 ± 1.63 fold vs 12.50 ± 5.38 fold increase by IL-13, $p < 0.05$) (Fig 5A). The intracellular mucin 5AC protein level was significantly reduced when the cells were treated with 100 μ M Rp-cAMPS but not at 50 μ M, while mucin glycoproteins levels were reduced at both concentrations (see Fig 5B and 5C, and S6A and S6B Fig for representative images).

To provide more evidence for a role for cAMP in mucin production in response to IL-13, we treated cells with 10 μ M forskolin combined with 100 μ M 3-isobutyl-1-methylxanthine (IBMX), in the absence of epinephrine. This treatment caused a dramatic increase in MUC5AC expression (75.73 ± 66.59 fold vs 0.56 ± 0.40 fold increase by IL-13, $p < 0.05$) (Fig 6A) when the cells were treated with IL-13. The same trend was also observed at the level of intracellular mucin




Fig 2. β_2 ARs are required for mucin production in response to IL-13 in NHBE cells. NHBE cells were grown in the presence of 3 μ M epinephrine, then at ALI, they were treated with 20 ng/ml IL-13 in combination with 3 μ M CGP-20712A (a preferential β_1 AR antagonist) or 1 μ M ICI-118,551 (a preferential β_2 AR antagonist) for 14 days. **A:** MUC5AC transcripts were quantified from extracted total RNA by qRT-PCR. Data are presented as fold change compared to cells grown in the presence of epinephrine only. **B:** Quantification of intracellular mucin 5AC content. The ratio of mucin 5AC integrated density of each group to the integrated density of the cells grown in the presence of epinephrine alone (control cells) was calculated and expressed as fold change. See the supplement for the representative images (S3A Fig). **C:** Quantification of intracellular mucin glycoproteins in response to different ligands. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to control

cells. See the supplement for the representative images (S3B Fig). Data are presented as means \pm SEM from three donors. *, # and @ indicate $p < 0.05$ significance as compared to + epinephrine, + epinephrine + IL-13 and + epinephrine + IL-13 + CGP-20721A treated cells respectively.

doi:10.1371/journal.pone.0132559.g002

5AC protein accumulation and mucin content of NHBE cells (Fig 6B and 6C; for representative images see S7A and S7B Fig).

Discussion

IL-13 plays an important role in the mucus over-production characteristic of bronchial asthma, and MUC5AC is the major mucin gene that is overexpressed by airway epithelium in asthmatic patients [22, 23]. We present evidence that signaling through β_2 AR increases IL-13 induced mucin production in human bronchial epithelium cultured in low RA concentrations. Most previous studies using these cells included epinephrine in the medium (e.g., [10, 18, 24, 25]) and to our knowledge, this is the first study showing an important role for β_2 AR signaling in the IL-13 mediated induction of MUC5AC expression and mucin content in NHBE cells.

These results are consistent with our earlier findings that β_2 AR signaling is required for mucous metaplasia in the airways of mice [16, 17, 26]. The clinical relevance of these findings is shown by the ability of formoterol, a commonly used long-acting β -agonist bronchodilator, to mimic epinephrine in the stimulation of MUC5AC transcripts by IL-13 (S2 Fig).

β_2 AR is the principal subtype present in human airway epithelium [27]. To verify that this subtype mediates the effects of IL-13 in cultured NHBE cells, we used preferential β AR antagonists. The selectivity of ICI-118,551 toward β_2 AR, and of CGP 20712A toward β_1 AR is least 500 fold or greater [28]. Treating NHBE cells with the preferential β_2 AR antagonist completely abolished MUC5AC expression in response to IL-13 and epinephrine. In contrast, the preferential β_1 AR antagonist did not alter MUC5AC expression levels under the same conditions. Intracellular mucin 5AC protein levels and mucin glycoprotein were not induced by IL-13 + epinephrine treatment in the presence of ICI-118,551 while CGP 20712A had no effect, paralleling the MUC5AC mRNA expression results (Fig 2A–2C). Our *in vitro* results are also consistent with our recent animal data where genetic ablation of β_2 ARs in mice resulted in attenuation of mucous metaplasia in an allergic murine model of asthma [16, 26].

β_2 AR signaling can proceed from agonist-activated receptors and from constitutively active receptors [29]. Cells that were grown in the absence of epinephrine showed no response to IL-13 treatment in terms of MUC5AC expression, mucin 5AC protein levels and mucin content while cells grown in the presence of epinephrine showed an increase in all three parameters (Fig 1A–1C), suggesting that agonist-induced β_2 AR activation is necessary. To further distinguish between constitutive versus agonist induced β_2 AR signaling, we treated NHBE cells with nadolol or alprenolol in combination with IL-13 in the presence of epinephrine. Nadolol is a non-selective β AR ligand that has full inverse agonist activity at β_2 ARs, thus it blocks both induced and constitutive β_2 AR signaling. Alprenolol lacks inverse agonist activity [29] but has weak β_2 AR agonist activity [30] and behaves as an antagonist in the presence of epinephrine, thus preserving constitutive β_2 AR signaling. Both nadolol and alprenolol blocked the effect of IL-13 on MUC5AC mRNA expression, protein levels and intracellular mucin content to similar extents (Fig 3A–3C). Thus, we provide evidence that constitutive β_2 AR receptor activity is not sufficient to drive the increase in mucin in response to IL-13 in human bronchial epithelial cells. Again, our *in vitro* results are also consistent with our recent mouse data where genetic or pharmacological depletion of epinephrine in mice resulted in attenuation of mucous metaplasia in an antigen-driven murine model of asthma [16, 17, 26]




Fig 3. Agonist induced β_2 AR signaling is required for mucin production in response to IL-13 in NHBE cells. NHBE cells were grown in the presence of 3 μ M epinephrine, then at ALI, they were treated with 20 ng/ml IL-13 in combination with 10 μ M nadolol (a non-selective inverse agonist of β ARs) or 10 μ M alprenolol (a non-selective β_2 AR blocker with no inverse agonist activity) for 14 days. **A:** MUC5AC transcripts were measured by qRT-PCR. Data are presented as fold change compared to cells grown in the presence of epinephrine only. **B:** Quantification of intracellular mucin 5AC content. The ratio of mucin 5AC integrated

density of each group to the integrated density of the cells grown in the presence of epinephrine alone (control cells) was calculated and expressed as fold change. See the supplement for the representative images (S4A Fig). **C:** Quantification of intracellular mucin glycoproteins in response to different ligands. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to control cells. See the supplement for the representative images (S4B Fig). Data are presented as means \pm SEM from three donors. * and # indicate $p < 0.05$ significance as compared to + epinephrine and + epinephrine + IL-13 treated cells respectively.

doi:10.1371/journal.pone.0132559.g003

We have begun determining the basis for the interaction between the β_2 AR and IL-13 signaling pathways. Since β_2 AR can activate MAPKs through cAMP-dependent and independent pathways, we tested the role of these kinases in the expression of MUC5AC. Since cells that were grown in the absence of epinephrine did not show induction of MUC5AC expression by IL-13, we examined only cells grown in the presence of epinephrine. All three families of MAPKs, JNKs, ERKs and p38, are known to be involved in the production of cytokine and chemokines by airway epithelium [31]. Phosphorylated p38 and ERK1/2 are detectable in the epithelium of asthmatic patients and the level of phosphorylation correlates with asthma severity [32]. In NHBE cells, IL-13 induced the phosphorylation of ERK1/2 and the ERK1/2 selective inhibitor, FR180204, attenuated MUC5AC expression, mucin 5AC protein levels and mucin content in response to IL-13 (Fig 4A, 4D–4F). A selective MEK 1/2 inhibitor, U0126, also




Fig 4. MAPK signaling is required for mucin production in response to IL-13 in NHBE cells. **A-C:** NHBE cells were grown in the presence or absence of epinephrine or IL-13 as indicated for 14 days after ALI before harvesting for total proteins. Immunoblots were performed using antibodies to the indicated phosphorylated MAP-kinases. The signal densities of the phosphorylated proteins were normalized to GAPDH protein density and the relative intensities were reported as the degree of activation of the protein. The data presented as fold change compared to the corresponding control cells. **D-F:** NHBE cells were grown in the presence of 3 μ M epinephrine, then at ALI, they were treated with 20 ng/ml IL-13 in combination with 3 μ M FR180204 (ERK1/2 inhibitor), 3 μ M SP600125 (JNK inhibitor) or 3 μ M SB203580 (p38 inhibitor) for 14 days. **D:** MUC5AC transcripts were measured by qRT-PCR, and the data presented as fold change compared to cells grown in the presence of epinephrine only. **E:** Quantification of intracellular mucin 5AC content. The ratio of mucin 5AC integrated density of each group to the integrated density of the cells grown in the presence of epinephrine alone (control cells) was calculated and expressed as fold change. See the supplement for the representative images (S5A Fig). **F:** Quantification of total intracellular mucin glycoproteins. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to control cells. See the supplement for the representative images (S5B Fig) Data are presented as means \pm SEM from three donors. * and # indicate $p < 0.05$ significance as compared to +epinephrine and + epinephrine + IL-13 treated cells respectively.

doi:10.1371/journal.pone.0132559.g004




Fig 5. Inhibiting PKA signaling reduced mucin production in response to IL-13 in NHBE cells. Cells were grown in the presence of 3 μ M epinephrine, then after ALI, they were treated with 20 ng/ml IL-13 in combination with 50 μ M or 100 μ M Rp-cAMP (cAMP-dependent protein kinase inhibitor) for 14 days. **A:** MUC5AC transcripts were measured by qRT-PCR, and the data presented as fold change compared to cells grown in the absence of inhibitor. **B:** Quantification of intracellular mucin 5AC content. The ratio of mucin 5AC integrated density of each group to the integrated density of the cells grown in the presence of epinephrine alone (control cells) was calculated and expressed as fold change. See the supplement for the representative images (S6A Fig). **C:** Quantification of intracellular mucin glycoproteins. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to control cells. See the supplement for the representative images (S6B Fig). Data are presented as means \pm SEM from three donors. * and # indicate $p < 0.05$ significance as compared to + epinephrine and + epinephrine + IL-13 treated cells respectively.

doi:10.1371/journal.pone.0132559.g005




Fig 6. cAMP potentiates mucin production in response to IL-13 in NHBE cells. Cells were grown in the absence of epinephrine, then at ALI, they were incubated with or without 20 ng/ml IL-13 with or without 10 μ M forskolin and 100 μ M IBMX for 14 days. **A:** MUC5AC transcripts were measured by qRT-PCR, and the data presented as fold change compared to cells grown in the absence of IL-13, IBMX and forskolin (control cells) **B:** Quantification of intracellular mucin 5AC content. The ratio of mucin 5AC integrated density of each group to the integrated density of the cells grown in the presence of epinephrine alone (control cells) was calculated and expressed as fold change. See the supplement for the representative images ([S7A Fig](#)). **C:** Quantification of intracellular mucin glycoproteins. The ratio of mucin integrated density and nucleic acid/cytoplasm integrated density was calculated and the data presented as fold change compared to control cells. See the supplement for the representative images ([S7B Fig](#)). Data are presented as means \pm SEM from three donors. \dagger and \yen indicate $p < 0.05$ significance as compared to $-$ epinephrine and $-$ epinephrine + IL-13 treated cells respectively.

doi:10.1371/journal.pone.0132559.g006

reduced MUC5AC expression in NHBE cells in response to IL-13 (Nguyen et al, in preparation). These results are consistent with other similar studies in NHBE cells [18].

JNK is involved in regulating pro-inflammatory gene expression and remodeling in airway diseases [33]. In NHBE cells, IL-13 induced the phosphorylation of c-Jun, a distinct JNK downstream target, and a JNK inhibitor, SP600125, inhibited MUC5AC expression and mucin content response to IL-13 (Fig 4B, 4D–4F). Consistent with this finding, attenuation of the asthma phenotype, including mucous metaplasia, by SP600125 has been shown in allergen driven murine model of asthma [34].

The other MAPK that we examined, p38, has four known isoforms: α , β , γ and δ [31]. The transcript levels of the α and β isoforms in the human lung is higher than that of other p38 isoforms of [35]. Inhaled p38 α antisense oligonucleotide attenuates mucus production in IL-13 transgenic mice [36]. We did not observe an increase in p38 phosphorylation in response to IL-13 (Fig 4C), however a selective inhibitor of both α and β isoforms [37], SB203580, reduced levels of MUC5AC expression and mucin 5AC content induced by IL-13 and epinephrine (Fig 4D–4F). These data provide evidence for the involvement of p38 in IL-13 induced mucous metaplasia, consistent with previous studies [18, 38]. Even though p38 phosphorylation was not affected by IL-13, basal levels of p38 activation may still be required for MUC5AC expression and mucin 5AC production, perhaps by integrating signals from the β_2 - and IL-13 receptors. There was no significant decrease in total mucin content after inhibiting p38, perhaps because mucin 5AC production decreases while the production of other mucin glycoproteins increases. All of these MAPKs (ERK1/2, JNK and p38) can be activated by β_2 AR via the G_s -adenylyl cyclase pathway, and by G protein-independent, β -arrestin2-dependent pathways [39]. MAPKs then act through downstream components that lead to the activation or translocation to the nucleus of transcriptional factors that act upon the MUC5AC and other genes [40].

To assess the possible involvement of cAMP mediated mechanisms, we used the Rp-isomer of adenosine-3', 5'-cyclic monophosphorothioate (Rp-cAMPS). Rp-cAMPS binds to PKA preventing its activation by cAMP [41]. MUC5AC expression, intracellular mucin 5AC protein levels and mucin content induced by IL-13 and epinephrine were not significantly inhibited by 50 μ M Rp-cAMPS ($IC_{50} \sim 10 \mu$ M for inhibiting PKA *in vitro*), but were inhibited by 100 μ M Rp-cAMPS (Fig 5A–5C). We then examined the effects of raising cAMP levels by treating NHBE cells with forskolin, a direct activator of adenylyl cyclase, combined with the non-specific phosphodiesterase inhibitor (PDE), IBMX. This treatment resulted in significant increases in MUC5AC transcripts in response to IL-13 (Fig 6). Taken together the data support a role of cAMP in regulating MUC5AC transcription as induced by IL-13. However, this data must be taken in the context of the high concentration of Rp-cAMPS required for the inhibition of MUC5AC expression, and that we used a compound (forskolin) that causes a 'global' intracellular increase in the activity of all adenylyl cyclases, and an inhibitor of all PDE subtypes (IBMX) that regulate the breakdown of cAMP. Given the emergence of data demonstrating distinct subcellular compartmentalization of cAMP associated with specific PDE subtypes [42] we cannot conclusively assert that epinephrine-induced cAMP increases cause the induction of MUC5AC. There are also studies suggesting that, at least with prolonged agonist exposures, other G_s -coupled receptors such as the adenosine A2B receptor can enhance lung injury and damage epithelial cell integrity [43, 44]. While our experiments could have eliminated a role for cAMP mediating mucin production, the data instead only suggest a role, and additional experiments using more refined pharmacologic tools that are PDE subtype specific, and genetic approaches will be needed to further investigate the possible role of β_2 AR-mediated cAMP increases in regulating mucin production. Given that MAP-kinases can be activated by cAMP mediated mechanisms, it would be interesting to test whether the effect of forskolin + IBMX is negated by inhibition of a MAP kinase, such as ERK or JNK, each of which show increased

phosphorylated upon treatment with IL-13 + epinephrine. It may also be suggested that cAMP is needed for p38 to integrate signals from the β_2 - and IL-13 receptors.

How do β_2 AR and IL-13 signaling interact? Due to the multiplicity of pathways proceeding from each receptor, it is not possible at present to firmly identify nodes of cross-talk. However, one potential factor could be an isoform of phosphoinositide 3-kinase (PI3K). Mice deficient in PI3K γ show reductions in most indices of airway inflammation in OVA-sensitized and challenged mice, though there was disagreement over effects on mucin production [45, 46]. Similar results were obtained, including decreases in mucin production, after treatment of mouse airways with inhibitors of PI3K γ or δ [47–50]. Studies in NHBE cells showed that pharmacologic inhibition of PI3K γ reduces IL-13 induced increases in goblet cell density [18]. Possibly, β_2 AR could amplify the IL-13 induction of PI3K isoforms by way of G $_i$ [51] or G β/γ [52]. An argument also could be made for p38, which by being constitutive in activity in this system, might therefore be a key transducer in more than one signaling pathway. These possibilities are currently under examination.

In the experiments reported here, only IL-13 and β_2 AR ligands were manipulated, and this reductionist approach does not take into account numerous other cytokines, chemokines and hormones present in the allergic airways. Also, leucocytes and other lung parenchymal cell types are absent from NHBE cultures. Nevertheless, the requirement for β_2 AR signaling in the induction of MUC5AC by IL-13 shown here is consistent with results we have obtained previously with β_2 AR knockout mice, mice chronically treated with β blockers, and mice deficient in epinephrine [16, 17, 26]. A further limitation regards possible off target effects from the use of chemical inhibitors and activators. Future studies utilizing more specific genetic approaches will be needed to delineate the precise signaling pathways involved. In particular, a role for β -arrestin2 cannot be excluded from the present data, and information from *in vivo* studies suggests that there is an involvement of β -arrestin2 in the pathogenesis of asthma [53, 54].

In conclusion, our results reveal an important role for β_2 AR signaling in mediating mucus production in response to IL-13 in NHBE cells. Constitutive β_2 AR activity alone is not sufficient to mediate this effect and it requires agonist activation of the receptor. Moreover, the three major MAPK signaling molecules (ERK1/2, JNK and p38) play a role in mediating the effects of IL-13 in NHBE cells in the presence of epinephrine. Our data also suggests that a cAMP activated signaling cascade may be involved in mediating the inflammatory effect of IL-13. The present report supports the notion that use of selective β_2 AR antagonists could be of value in the treatment of mucus overproduction in asthma and other similar disorders such as COPD [55].

Supporting Information

S1 Fig. Epinephrine is required for mucin production in response to IL-13 in NHBE cell grown with 50 nM retinoic acid. NHBE cells were grown in the presence or absence of 3 μ M epinephrine. At ALI, the cells were treated with 20 ng/ml IL-13 for 14 days, total RNA was harvested and then MUC5AC transcripts were measured by qRT-PCR. Data are presented as fold change compared to the corresponding treatment control (in the absence of IL13). *, indicates $p < 0.05$ significance as compared to + epinephrine, —epinephrine and –epinephrine + IL-13 treated cells respectively. N = 3.

(PNG)

S2 Fig. Formoterol also potentiates mucin production in response to IL-13. NHBE cells were cultured as described in S1 Fig, except that 10 nM formoterol was used in place of epinephrine.

(PNG)

S3 Fig. β_2 ARs are required for mucin production in response to IL-13 in NHBE cells. Representative images of data quantified in Fig 2. **A:** Intracellular MUC5AC content. **B:** Intracellular mucin glycoproteins.

(PNG)

S4 Fig. Agonist induced β_2 AR signaling is required for mucin production in response to IL-13 in NHBE cells. Representative images of data quantified in Fig 3. **A:** Intracellular MUC5AC content. **B:** Intracellular mucin glycoproteins.

(PNG)

S5 Fig. MAPK signaling is required for mucin production in response to IL-13 in NHBE cells. Representative images of data quantified in Fig 4. **A:** Intracellular MUC5AC content. **B:** Intracellular mucin glycoproteins.

(PNG)

S6 Fig. Inhibiting PKA signaling reduced mucin production in response to IL-13 in NHBE cells. Representative images of data quantified in Fig 5. **A:** Intracellular MUC5AC content. **B:** Intracellular mucin glycoproteins.

(PNG)

S7 Fig. cAMP potentiates mucin production in response to IL-13 in NHBE cells. Representative images of data quantified in Fig 6. **A:** Intracellular MUC5AC content. **B:** Intracellular mucin glycoproteins.

(PNG)

Acknowledgments

The authors thank Gloria Forkuo for reviewing the manuscript.

Author Contributions

Conceived and designed the experiments: BJK RAB NS. Performed the experiments: NS IP AH OO HK. Analyzed the data: NS IP RAB BJK VT. Wrote the paper: NS VT RAB BJK.

References

1. Bai TR, Knight DA. Structural changes in the airways in asthma: Observations and consequences. *Clin Sci (Lond)*. 2005; 108(6):463–77. Epub 2005/05/18. doi: [10.1042/CS20040342](https://doi.org/10.1042/CS20040342) PMID: [15896192](https://pubmed.ncbi.nlm.nih.gov/15896192/).
2. Evans CM, Kim K, Tuvim MJ, Dickey BF. Mucus hypersecretion in asthma: causes and effects. *Curr Opin Pulm Med*. 2009; 15(1):4–11. Epub 2008/12/17. doi: [10.1097/MCP.0b013e32831da8d3](https://doi.org/10.1097/MCP.0b013e32831da8d3) PMID: [19077699](https://pubmed.ncbi.nlm.nih.gov/19077699/); PubMed Central PMCID: PMC2709596.
3. Aikawa T, Shimura S, Sasaki H, Ebina M, Takishima T. Marked goblet cell hyperplasia with mucus accumulation in the airways of patients who died of severe acute asthma attack. *Chest*. 1992; 101(4):916–21. Epub 1992/04/01. PMID: [1555462](https://pubmed.ncbi.nlm.nih.gov/1555462/).
4. Licona-Limón P, Kim LK, Palm NW, Flavell RA. TH2, allergy and group 2 innate lymphoid cells. *Nature immunology*. 2013; 14(6):536–42. Epub 2013/05/21. doi: [10.1038/ni.2617](https://doi.org/10.1038/ni.2617) PMID: [23685824](https://pubmed.ncbi.nlm.nih.gov/23685824/).
5. Wills-Karp M, Luyimbazi J, Xu X, Schofield B, Neben TY, Karp CL, et al. Interleukin-13: central mediator of allergic asthma. *Science*. 1998; 282(5397):2258–61. PMID: [9856949](https://pubmed.ncbi.nlm.nih.gov/9856949/).
6. Grünig G, Warnock M, Wakil AE, Venkayya R, Brombacher F, Rennick DM, et al. Requirement for IL-13 independently of IL-4 in experimental asthma. *Science*. 1998; 282(5397):2261–3. PMID: [9856950](https://pubmed.ncbi.nlm.nih.gov/9856950/).
7. Kuperman DA, Huang X, Koth LL, Chang GH, Dolganov GM, Zhu Z, et al. Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat Med*. 2002; 8(8):885–9. PMID: [12091879](https://pubmed.ncbi.nlm.nih.gov/12091879/).
8. Williams OW, Sharafkhaneh A, Kim V, Dickey BF, Evans CM. Airway mucus: From production to secretion. *Am J Respir Cell Mol Biol*. 2006; 34(5):527–36. PMID: [16415249](https://pubmed.ncbi.nlm.nih.gov/16415249/); PubMed Central PMCID: PMC <http://ajrcmb.atsjournals.org/cgi/content/full/34/5/527>.

9. Holgate ST, Davies DE, Puddicombe S, Richter A, Lackie P, Lordan J, et al. Mechanisms of airway epithelial damage: epithelial-mesenchymal interactions in the pathogenesis of asthma. *Eur Respir J Suppl.* 2003; 44:24s–9s. PMID: [14582897](#).
10. Zhen G, Park SW, Nguyenvu LT, Rodriguez MW, Barbeau R, Paquet AC, et al. IL-13 and epidermal growth factor receptor have critical but distinct roles in epithelial cell mucin production. *Am J Respir Cell Mol Biol.* 2007; 36(2):244–53. PMID: [16980555](#); PubMed Central PMCID: PMC1899314.
11. Lambrecht BN, Hammad H. The airway epithelium in asthma. *Nature medicine.* 2012; 18(5):684–92. Epub 2012/05/09. doi: [10.1038/nm.2737](#) PMID: [22561832](#).
12. Park SW, Jangm HK, An MH, Min JW, Jang AS, Lee JH, et al. Interleukin-13 and interleukin-5 in induced sputum of eosinophilic bronchitis: comparison with asthma. *Chest.* 2005; 128(4):1921–7. Epub 2005/10/21. doi: [10.1378/chest.128.4.1921](#) PMID: [16236836](#).
13. Yasuo M, Fujimoto K, Tanabe T, Yaegashi H, Tsushima K, Takasuna K, et al. Relationship between calcium-activated chloride channel 1 and MUC5AC in goblet cell hyperplasia induced by interleukin-13 in human bronchial epithelial cells. *Respiration; international review of thoracic diseases.* 2006; 73(3):347–59. Epub 2006/02/09. doi: [10.1159/000091391](#) PMID: [16465045](#).
14. Chu HW, Balzar S, Seedorf GJ, Westcott JY, Trudeau JB, Silkoff P, et al. Transforming growth factor- β_2 induces bronchial epithelial mucin expression in asthma. *Am J Pathol.* 2004; 165(4):1097–106. Epub 2004/10/07. 165/4/1097 [pii]. PMID: [15466377](#); PubMed Central PMCID: PMC1618635.
15. Young HW, Williams OW, Chandra D, Bellinghausen LK, Perez G, Suarez A, et al. Central role of Muc5ac expression in mucous metaplasia and its regulation by conserved 5' elements. *Am J Respir Cell Mol Biol.* 2007; 37(3):273–90. PMID: [17463395](#); PubMed Central PMCID: PMC1994232.
16. Nguyen LP, Lin R, Parra S, Omoluabi O, Hanania NA, Tuvim MJ, et al. β_2 -adrenoceptor signaling is required for the development of an asthma phenotype in a murine model. *Proc Natl Acad Sci U S A.* 2009; 106:2435–40. Epub 2009/01/28. 0810902106 [pii] doi: [10.1073/pnas.0810902106](#) PMID: [19171883](#).
17. Nguyen LP, Omoluabi O, Parra S, Frieske JM, Clement C, Ammar-Aouchiche Z, et al. Chronic exposure to beta-blockers attenuates inflammation and mucin content in a murine asthma model. *Am J Respir Cell Mol Biol.* 2008; 38(3):256–62. Epub 2007/12/22. doi: [10.1165/rcmb.2007-0279RC](#) PMID: [18096872](#); PubMed Central PMCID: PMC2258446.
18. Atherton HC, Jones G, Danahay H. IL-13-induced changes in the goblet cell density of human bronchial epithelial cell cultures: MAP kinase and phosphatidylinositol 3-kinase regulation. *Am J Physiol Lung Cell Mol Physiol.* 2003; 285(3):L730–9. Epub 2003/06/10. [pii]. PMID: [12794003](#).
19. Lin H, Li H, Cho HJ, Bian S, Roh HJ, Lee MK, et al. Air-liquid interface (ALI) culture of human bronchial epithelial cell monolayers as an in vitro model for airway drug transport studies. *J Pharm Sci.* 2007; 96(2):341–50. Epub 2006/11/03. doi: [10.1002/jps.20803](#) PMID: [17080426](#).
20. Piccotti L, Dickey BF, Evans CM. Assessment of intracellular mucin content in vivo. *Methods Mol Biol.* 2012; 842:279–95. Epub 2012/01/20. doi: [10.1007/978-1-61779-513-8_17](#) PMID: [22259143](#).
21. Fujisawa T, Velichko S, Thai P, Hung LY, Huang F, Wu R. Regulation of airway MUC5AC expression by IL-1 β and IL-17A; the NF- κ B paradigm. *J Immunol.* 2009; 183(10):6236–43. Epub 2009/10/21. doi: [10.4049/jimmunol.0900614](#) PMID: [19841186](#).
22. Rose MC, Voynow JA. Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol Rev.* 2006; 86(1):245–78. Epub 2005/12/24. doi: [10.1152/physrev.00010.2005](#) PMID: [16371599](#).
23. Ordoñez CL, Khashayar R, Wong HH, Ferrando R, Wu R, Hyde DM, et al. Mild and moderate asthma is associated with airway goblet cell hyperplasia and abnormalities in mucin gene expression. *Am J Respir Crit Care Med.* 2001; 163(2):517–23. PMID: [11179133](#).
24. Yoshisue H, Hasegawa K. Effect of MMP/ADAM inhibitors on goblet cell hyperplasia in cultured human bronchial epithelial cells. *Biosci Biotechnol Biochem.* 2004; 68(10):2024–31. Epub 2004/10/27. PMID: [15502346](#).
25. Kono Y, Nishiuma T, Okada T, Kobayashi K, Funada Y, Kotani Y, et al. Sphingosine kinase 1 regulates mucin production via ERK phosphorylation. *Pulm Pharmacol Ther.* 2010; 23(1):36–42. Epub 2009/10/20. doi: [10.1016/j.pupt.2009.10.005](#) PMID: [19835973](#).
26. Thanawala VJ, Forkuo GS, Al-Sawalha N, Azzegagh Z, Nguyen LP, Eriksen JL, et al. β_2 -adrenoceptor agonists are required for development of the asthma phenotype in a murine model. *Am J Respir Cell Mol Biol.* 2013; 48:220–9. Epub 2012/12/04. doi: [10.1165/rcmb.2012-0364OC](#) PMID: [23204390](#).
27. Davis PB, Silski CL, Kerckmar CM, Infeld M. β -adrenergic receptors on human tracheal epithelial cells in primary culture. *Am J Physiol.* 1990; 258(1 Pt 1):C71–6. Epub 1990/01/01. PMID: [1689114](#).
28. Baker JG. The selectivity of β -adrenoceptor antagonists at the human β_1 , β_2 and β_3 adrenoceptors. *British journal of pharmacology.* 2005; 144:317–22. PMID: [15655528](#)

29. Bond RA, Leff P, Johnson TD, Milano CA, Rockman HA, McMinn TR, et al. Physiological effects of inverse agonists in transgenic mice with myocardial overexpression of the beta 2-adrenoceptor. *Nature*. 1995; 374(6519):272–6. PMID: [7885448](#).
30. Wisler JW, DeWire SM, Whalen EJ, Violin JD, Drake MT, Ahn S, et al. A unique mechanism of β -blocker action: carvedilol stimulates β -arrestin signaling. *Proc Natl Acad Sci U S A*. 2007; 104(42):16657–62. Epub 2007/10/11. 0707936104 [pii]doi: [10.1073/pnas.0707936104](#) PMID: [17925438](#); PubMed Central PMCID: PMC2034221.
31. Pelaia G, Cuda G, Vatrella A, Gallelli L, Caraglia M, Marra M, et al. Mitogen-activated protein kinases and asthma. *J Cell Physiol*. 2005; 202(3):642–53. Epub 2004/08/19. doi: [10.1002/jcp.20169](#) PMID: [15316926](#).
32. Liu W, Liang Q, Balzar S, Wenzel S, Gorska M, Alam R. Cell-specific activation profile of extracellular signal-regulated kinase 1/2, Jun N-terminal kinase, and p38 mitogen-activated protein kinases in asthmatic airways. *J Allergy Clin Immunol*. 2008; 121(4):893–902 e2. Epub 2008/04/09. doi: [10.1016/j.jaci.2008.02.004](#) PMID: [18395552](#).
33. Bennett BL. c-Jun N-terminal kinase-dependent mechanisms in respiratory disease. *The European respiratory journal: official journal of the European Society for Clinical Respiratory Physiology*. 2006; 28(3):651–61. Epub 2006/09/02. doi: [10.1183/09031936.06.00012106](#) PMID: [16946096](#).
34. Nath P, Eynott P, Leung SY, Adcock IM, Bennett BL, Chung KF. Potential role of c-Jun NH2-terminal kinase in allergic airway inflammation and remodelling: effects of SP600125. *Eur J Pharmacol*. 2005; 506(3):273–83. Epub 2005/01/04. doi: [10.1016/j.ejphar.2004.11.040](#) PMID: [15627438](#).
35. Jiang Y, Gram H, Zhao M, New L, Gu J, Feng L, et al. Characterization of the structure and function of the fourth member of p38 group mitogen-activated protein kinases, p38delta. *J Biol Chem*. 1997; 272(48):30122–8. Epub 1997/12/31. PMID: [9374491](#).
36. Ma JY, Medicherla S, Kerr I, Mangadu R, Protter AA, Higgins LS. Selective p38 α mitogen-activated protein kinase inhibitor attenuates lung inflammation and fibrosis in IL-13 transgenic mouse model of asthma. *J Asthma Allergy*. 2008; 1:31–44. Epub 2008/01/01. PMID: [21436983](#); PubMed Central PMCID: PMC3121334.
37. Chung KF. p38 mitogen-activated protein kinase pathways in asthma and COPD. *Chest*. 2011; 139(6):1470–9. Epub 2011/06/10. doi: [10.1378/chest.10-1914](#) PMID: [21652557](#).
38. Fujisawa T, Ide K, Holtzman MJ, Suda T, Suzuki K, Kuroishi S, et al. Involvement of the p38 MAPK pathway in IL-13-induced mucous cell metaplasia in mouse tracheal epithelial cells. *Respirology*. 2008; 13(2):191–202. PMID: [18339016](#). doi: [10.1111/j.1440-1843.2008.01237.x](#)
39. DeWire SM, Ahn S, Lefkowitz RJ, Shenoy SK. β -arrestins and cell signaling. *Annu Rev Physiol*. 2007; 69:483–510. Epub 2007/02/20. doi: [10.1146/annurev.ph.69.013107.100021](#) PMID: [17305471](#).
40. Thai P, Loukoianov A, Wachi S, Wu R. Regulation of airway mucin gene expression. *Annu Rev Physiol*. 2008; 70:405–29. Epub 2007/10/27. doi: [10.1146/annurev.physiol.70.113006.100441](#) PMID: [17961085](#).
41. Dostmann WR. (RP)-cAMPS inhibits the cAMP-dependent protein kinase by blocking the cAMP-induced conformational transition. *FEBS Lett*. 1995; 375(3):231–4. Epub 1995/11/20. PMID: [7498506](#).
42. Stangherlin A, Zaccolo M. Phosphodiesterases and subcellular compartmentalized cAMP signaling in the cardiovascular system. *Am J Physiol Heart Circ Physiol*. 2012; 302(2):H379–90. Epub 2011/11/01. doi: [10.1152/ajpheart.00766.2011](#) PMID: [22037184](#).
43. Karmouty-Quintana H, Xia Y, Blackburn MR. Adenosine signaling during acute and chronic disease states. *J Mol Med (Berl)*. 2013; 91(2):173–81. Epub 2013/01/24. doi: [10.1007/s00109-013-0997-1](#) PMID: [23340998](#); PubMed Central PMCID: PMC3606047.
44. Lu Q, Sakhatskyy P, Newton J, Shamirian P, Hsiao V, Curren S, et al. Sustained adenosine exposure causes lung endothelial apoptosis: a possible contributor to cigarette smoke-induced endothelial apoptosis and lung injury. *Am J Physiol Lung Cell Mol Physiol*. 2013; 304(5):L361–70. Epub 2013/01/15. doi: [10.1152/ajplung.00161.2012](#) PMID: [23316066](#); PubMed Central PMCID: PMC3602741.
45. Lim DH, Cho JY, Song DJ, Lee SY, Miller M, Broide DH. PI3K γ -deficient mice have reduced levels of allergen-induced eosinophilic inflammation and airway remodeling. *American journal of physiology Lung cellular and molecular physiology*. 2009; 296(2):L210–9. Epub 2008/11/26. doi: [10.1152/ajplung.90275.2008](#) PMID: [19028980](#); PubMed Central PMCID: PMC2643991.
46. Takeda M, Ito W, Tanabe M, Ueki S, Kato H, Kihara J, et al. Allergic airway hyperresponsiveness, inflammation, and remodeling do not develop in phosphoinositide 3-kinase γ -deficient mice. *J Allergy Clin Immunol*. 2009; 123(4):805–12. Epub 2009/02/24. doi: [10.1016/j.jaci.2008.11.047](#) PMID: [19232703](#).
47. Myou S, Leff AR, Myo S, Boetticher E, Tong J, Meliton AY, et al. Blockade of inflammation and airway hyperresponsiveness in immune-sensitized mice by dominant-negative phosphoinositide 3-kinase-TAT. *J Exp Med*. 2003; 198(10):1573–82. doi: [10.1084/jem.20030298](#) PMID: [14623911](#); PubMed Central PMCID: PMC2194122.

48. Doukas J, Eide L, Stebbins K, Racanelli-Layton A, Dellamary L, Martin M, et al. Aerosolized phosphoinositide 3-kinase γ/δ inhibitor TG100-115 [3-[2,4-diamino-6-(3-hydroxyphenyl)pteridin-7-yl]phenol] as a therapeutic candidate for asthma and chronic obstructive pulmonary disease. *J Pharmacol Exp Ther*. 2009; 328(3):758–65. Epub 2008/12/06. doi: [10.1124/jpet.108.144311](https://doi.org/10.1124/jpet.108.144311) PMID: [19056934](https://pubmed.ncbi.nlm.nih.gov/19056934/).
49. Duan W, Aguinaldo Datiles AM, Leung BP, Vlahos CJ, Wong WS. An anti-inflammatory role for a phosphoinositide 3-kinase inhibitor LY294002 in a mouse asthma model. *Int Immunopharmacol*. 2005; 5(3):495–502. PMID: [15683846](https://pubmed.ncbi.nlm.nih.gov/15683846/).
50. Kwak YG, Song CH, Yi HK, Hwang PH, Kim JS, Lee KS, et al. Involvement of PTEN in airway hyperresponsiveness and inflammation in bronchial asthma. *J Clin Invest*. 2003; 111(7):1083–92. doi: [10.1172/JCI16440](https://doi.org/10.1172/JCI16440) PMID: [12671058](https://pubmed.ncbi.nlm.nih.gov/12671058/); PubMed Central PMCID: PMC152583.
51. Chesley A, Lundberg MS, Asai T, Xiao RP, Ohtani S, Lakatta EG, et al. The β_2 -adrenergic receptor delivers an antiapoptotic signal to cardiac myocytes through Gi-dependent coupling to phosphatidylinositol 3'-kinase. *Circ Res*. 2000; 87(12):1172–9. PMID: [11110775](https://pubmed.ncbi.nlm.nih.gov/11110775/).
52. Leopoldt D, Hanck T, Exner T, Maier U, Wetzker R, Nurnberg B. G $\beta\gamma$ stimulates phosphoinositide 3-kinase- γ by direct interaction with two domains of the catalytic p110 subunit. *J Biol Chem*. 1998; 273(12):7024–9. Epub 1998/04/18. PMID: [9507010](https://pubmed.ncbi.nlm.nih.gov/9507010/).
53. Nichols HL, Saffeddine M, Theriot BS, Hegde A, Polley D, El-Mays T, et al. β -Arrestin-2 mediates the proinflammatory effects of proteinase-activated receptor-2 in the airway. *Proc Natl Acad Sci U S A*. 2012; 109(41):16660–5. Epub 2012/09/27. doi: [10.1073/pnas.1208881109](https://doi.org/10.1073/pnas.1208881109) PMID: [23012429](https://pubmed.ncbi.nlm.nih.gov/23012429/); PubMed Central PMCID: PMC3478622.
54. Hollingsworth JW, Theriot BS, Li Z, Lawson BL, Sunday M, Schwartz DA, et al. Both hematopoietic-derived and non-hematopoietic-derived β -arrestin-2 regulates murine allergic airway disease. *Am J Respir Cell Mol Biol*. 2010; 43(3):269–75. Epub 2009/10/07. doi: [10.1165/rcmb.2009-0198OC](https://doi.org/10.1165/rcmb.2009-0198OC) PMID: [19805483](https://pubmed.ncbi.nlm.nih.gov/19805483/); PubMed Central PMCID: PMC2933545.
55. Hanania NA, Singh S, El-Wali R, Flashner M, Franklin AE, Garner WJ, et al. The safety and effects of the β -blocker, nadolol, in mild asthma: an open-label pilot study. *Pulm Pharmacol Ther*. 2008; 21(1):134–41. PMID: [17703976](https://pubmed.ncbi.nlm.nih.gov/17703976/); PubMed Central PMCID: PMC2254137.