


FIG. 2 Three successive rounds of DNA shuffling were done and the cells were grown on increasing cefotaxime levels. The MIC of cefotaxime (Sigma) for *E. coli* XL1-blue (Stratagene, San Diego) carrying wild-type p182Sfi is $0.02 \mu\text{g ml}^{-1}$. A mutant with a 16,000-fold increased resistance to cefotaxime was obtained ($\text{MIC} = 320 \mu\text{g ml}^{-1}$). This mutant was backcrossed twice, by shuffling with a 40-fold excess of wild-type DNA. The backcrossed mutant was 32,000-fold more resistant than the wild type ($\text{MIC} = 640 \mu\text{g ml}^{-1}$). After selection, the plasmid of selected clones was transferred back into wild-type XL-1 blue cells to ensure that none of the measured drug resistance



was due to chromosomal mutations. DNA sequencing showed that both mutants had 9 single-base-pair mutations.

7. Arkin, A. & Youvan, D. C. *Proc. natn. Acad. Sci. U.S.A.* **89**, 7811–7815 (1992).
8. Delagrave, S. & Youvan, D. C. *Biotechnology* **11**, 1548–1552 (1993).
9. Palzkill, T. & Botstein, D. J. *Bact.* **174**, 5237–5243 (1992).
10. Oiphant, A. R., Nussbaum, A. L. & Struhl, K. *Gene* **44**, 177–183 (1986).
11. Hermes, J. D., Blacklow, S. C. & Knowles, J. R. *Proc. natn. Acad. Sci. U.S.A.* **87**, 696–700 (1990).
12. Leung, D. W., Chen, E. & Goeddel, D. V. *Technique* **1**, 11–15 (1989).
13. Caldwell, R. C. & Joyce, G. F. *PCR Meth. Applic.* **2**, 28–33 (1992).
14. Kauffman, S. A. *The Origins of Order* (Oxford Univ. Press, New York, 1993).
15. Kauffman, S. A. *J. theor. Biol.* **157**, 1–7 (1992).
16. Bartel, D. P. & Szostak, J. W. *Science* **261**, 1411–1418 (1993).

17. Tuerk, C. & Gold, L. *Science* **249**, 505–510 (1990).
18. Joyce, G. F. *Scient. Am.* **267**, 90–97 (1992).
19. Jacoby, G. A. & Medeiros, A. A. *Antimicrob. Ag. Chemother.* **35**, 1697–1704 (1991).
20. Collatz, E., Labia, R. & Gutmann, L. *Molec. Microbiol.* **4**, 1615–1620 (1990).
21. Philippon, A., Labia, R. & Jacoby, G. *Antimicrob. Ag. Chemother.* **33**, 1131–1136 (1989).
22. McCafferty, J., Griffiths, A. D., Winter, G. & Chiswell, D. J. *Nature* **348**, 552–554 (1990).
23. Huse, W. D., Sastry, L., Iverson, S. A. & Kang, A. S. *Science* **246**, 1275–1278 (1989).
24. Watson, N. *Gene* **70**, 399–403 (1988).
25. Ambler, R. P. et al. *Biochem. J.* **278**, 269–272 (1991).

ACKNOWLEDGEMENTS. We thank A. Crameri for assistance with DNA sequencing, P. Schatz for suggesting the model system and W. Dower for valuable discussions.


ERRATA

Parthenogenetic activation of oocytes in c-mos-deficient mice

Naohiro Hashimoto, Nobumoto Watanabe, Yasuhide Furuta, Hiroyuki Tamemoto, Noriyuki Sagata, Minesuke Yokoyama, Kenji Okazaki, Mariko Nagayoshi, Naoki Takeda, Yoji Ikawa & Shinichi Alzawa

Nature **370**, 68–71 (1994)

FIGURE 3b and c of this Letter was an early version that should not have been published. The correct version of this figure is shown here. □



Degradation of trifluoroacetate in oxic and anoxic sediments

Pieter T. Visscher, Charles W. Culbertson & Ronald S. Oremland

Nature **369**, 729–731 (1994)

In the last sentence of the opening paragraph of this Letter, an error was introduced during editing in which trifluoroform was referred to as a “potential ozone-depleting compound.” In fact, trifluoroform as well as other HFCs were recently shown by Ravishankara *et al.*¹ to have “negligibly small” ozone depletion potentials. □

1. Ravishankara, A. R. et al. *Science* **263**, 71–75 (1994).

Miller-Dieker lissencephaly gene encodes a subunit of brain platelet-activating factor acetylhydrolase

Mitsuharu Hattori, Hideki Adachi, Masafumi Tsujimoto, Hiroyuki Arai & Keizo Inoue

Nature **370**, 216–218 (1994)

The word ‘acetylhydrolase’ was accidentally omitted from the end of the title of this paper. The correct title should read “Miller-Dieker lissencephaly gene encodes a subunit of brain platelet-activating factor acetylhydrolase”. □