

ERRATUM

Open Access



Erratum to: Cardiac ischemia in patients with septic shock randomized to vasopressin or norepinephrine

Sangeeta Mehta^{1*}, John Granton², Anthony C. Gordon³, Deborah J. Cook⁴, Stephen Lapinsky¹, Gary Newton⁵, Kris Bandayrel¹, Anjuli Little¹, Chuin Siau⁶, Dieter Ayers⁷, Joel Singer⁷, Terry C. K. Lee⁷, Keith R. Walley⁸, Michelle Storms⁸, D. James Cooper⁹, Cheryl L. Holmes¹⁰, Paul Hebert¹¹, Jeffrey Presneill¹², James A. Russell⁸ and for the Vasopressin and Septic Shock Trial (VASST) Investigators

Erratum

Unfortunately this article [1] published without its additional files (Additional files 1, 2, 3, 4, 5, 6 and 7). Please see these listed below and attached as additional files to this erratum.

Additional files

Additional file 1: Participating sites. (DOCX 13 kb)

Additional file 2: Table S7. Baseline characteristics and outcomes in vasopressin and norepinephrine-treated patients, grouped by serum troponin levels. (DOCX 17 kb)

Additional file 3: Table S8. Specific ECG diagnoses and patient outcomes. (DOCX 14 kb)

Additional file 4: Table S9. Outcomes related to ECG ischemia for all patients and for patients with troponin elevation. (DOCX 15 kb)

Additional file 5: Figure S1. Comparison of mean arterial pressure and heart rate in the norepinephrine group and vasopressin group. (DOCX 122 kb)

Additional file 6: Figure S2. Rates of total norepinephrine infusion (open-label and study drug) in the vasopressin treated group and the norepinephrine treated group. (DOCX 60 kb)

Additional file 7: Figure S3. Serum creatinine in the vasopressin treated group and the norepinephrine treated group. (DOCX 64 kb)

Author details

¹Department of Medicine and Interdepartmental Division of Critical Care Medicine, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada. ²University Health Network-General Division, University of Toronto, Toronto, Ontario, Canada. ³Intensive Care, Charing Cross Hospital, Imperial, College, London, UK. ⁴Departments of Medicine, Clinical Epidemiology & Biostatistics, McMaster University, Hamilton, Ontario, Canada. ⁵Department of Medicine and Division of Cardiology, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada. ⁶Department of Respiratory Medicine, Changi General Hospital, Singapore, Singapore. ⁷School of Population and Public Health, Centre for Health Evaluation and Outcome Sciences, and University of British Columbia, Vancouver, British Columbia, Canada. ⁸Critical Care Research, Laboratories, Heart

and Lung Institute, St. Paul's Hospital, and University of British Columbia, Vancouver, British Columbia, Canada. ⁹Alfred Hospital, Monash University, Melbourne, Australia. ¹⁰Kelowna General Hospital, Kelowna, and University of British Columbia, Kelowna, British Columbia, Canada. ¹¹Ottawa Hospital-General Campus, and University of Ottawa, Ottawa, Ontario, Canada. ¹²Mater Health Services, University of Queensland and Monash University, South Brisbane, Queensland, Australia.

Received: 28 March 2017 Accepted: 28 March 2017

Published online: 04 May 2017

Reference

1. Mehta S, et al. Cardiac ischemia in patients with septic shock randomized to vasopressin or norepinephrine. *Crit Care*. 2013;17:R117.

* Correspondence: smehta@mtsinai.on.ca

¹Department of Medicine and Interdepartmental Division of Critical Care Medicine, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada