

## Erratum to: Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment

Tjeerd Blacquière · Guy Smagghe ·  
Cornelis A. M. van Gestel · Veerle Mommaerts

Published online: 3 April 2012  
© Springer Science+Business Media, LLC 2012

### Erratum to: Ecotoxicology DOI 10.1007/s10646-012-0863-x

There is an incorrect text in the original publication of the article. The authors would like to correct the error, and the corrected sentence should read as given below.

Within the different insecticide classes, the neonicotinoid insecticides, which include imidacloprid, acetamiprid,

clothianidin, thiamethoxam, thiacloprid, dinotefuran and nitenpyram, are an important group of neurotoxins specifically acting as agonists of the insect nicotinic acetylcholine receptors (nAChR) (Matsuda et al. Trends Pharmacol Sci 22:573–580, 2001; Elbert et al. Pest Manag Sci 64:1099–1105, 2008).

---

The online version of the original article can be found under doi:  
[10.1007/s10646-012-0863-x](https://doi.org/10.1007/s10646-012-0863-x).

---

T. Blacquière (✉)  
Plant Research International, Wageningen University &  
Research, PO Box 69, 6700 AB Wageningen, The Netherlands  
e-mail: tjeerd.blacquiere@wur.nl

G. Smagghe · V. Mommaerts (✉)  
Department of Crop Protection, Faculty of Bioscience  
Engineering, Ghent University, Coupure Links 653, 9000 Ghent,  
Belgium  
e-mail: veerle.mommaerts@ugent.be

C. A. M. van Gestel  
Department of Animal Ecology, Faculty of Earth and Life  
Sciences, VU University, De Boelelaan 1085, 1081 HV  
Amsterdam, The Netherlands