

Erratum to: Update with level 1 studies of the European Hernia Society guidelines on the treatment of inguinal hernia in adult patients

M. Miserez · E. Peeters · T. Aufenacker · J. L. Bouillot · G. Campanelli · J. Conze · R. Fortelny · T. Heikkinen · L. N. Jorgensen · J. Kukleta · S. Morales-Conde · P. Nordin · V. Schumpelick · S. Smedberg · M. Smietanski · G. Weber · M. P. Simons

Published online: 30 April 2014
© Springer-Verlag France 2014

Erratum to: Hernia (2014) 18:151–163
DOI 10.1007/s10029-014-1236-6

The original publication of this article unfortunately contained the wrong figure 2 (page 154). The correct figure is shown below.

The online version of the original article can be found under doi:[10.1007/s10029-014-1236-6](https://doi.org/10.1007/s10029-014-1236-6).

M. Miserez (✉) · E. Peeters
Department of Abdominal Surgery, University Hospitals
Leuven, Herestraat 49, 3000 Leuven, Belgium
e-mail: marc.miserez@uzleuven.be

T. Aufenacker
Department of Surgery, Rijnstate Hospital, PO Box 9555,
6800 TA Arnhem, The Netherlands

J. L. Bouillot
Department of General, Digestive and Metabolic Surgery,
Ambroise Paré Teaching Hospital, 9, Avenue Charles de Gaulle,
92100 Boulogne, France

G. Campanelli
University of Insubria, Via Quadronno, 29, 20122 Milan, Italy

J. Conze
Hernienzentrum, Arabellastrabe 7, 81925 Munich, Germany

R. Fortelny
Department of General, Visceral and Oncological Surgery,
Wilhelminenspital, 1171 Vienna, Austria

T. Heikkinen
Oulu University Hospital, Otavansakara 7, 90630 Oulu, Finland

L. N. Jorgensen
Digestive Disease Center, Bispebjerg Hospital, University of
Copenhagen, 2400 Copenhagen, NV, Denmark

J. Kukleta
Department of Surgery, Klinik Im Park, Grossmuensterplatz 9,
8001 Zurich, Switzerland

S. Morales-Conde
University Hospital Sevilla, Betis 65, 41010 Seville, Spain

P. Nordin
Department of Surgical and Perioperative Science, Umeå
University, 901 87 Umeå, Sweden

V. Schumpelick
Falkensteiner Ufer 34, 22587 Hamburg, Germany




Fig. 2 Pooled data of seven studies on severe chronic pain with follow-up >4 years after endoscopic vs. Lichtenstein inguinal hernia repair

S. Smedberg
Department of Surgery, Helsingborg/Ängelholm Hospital,
Ängelholm, Sweden

M. Smietanski
Department of General and Vascular Surgery, Ceynowa
Hospital, Jagalskiego 10, 84-200 Wejherowo, Poland

G. Weber
Department of Surgical Research and Techniques, Semmelweis
University, 1089 Budapest, Hungary

M. P. Simons
Onze Lieve Vrouwe Hospital, Postbus 95500,
1040 HM Amsterdam, The Netherlands