

ERROR BOUNDS FOR MONOTONE APPROXIMATION SCHEMES FOR PARABOLIC HAMILTON-JACOBI-BELLMAN EQUATIONS

GUY BARLES AND ESPEN R. JAKOBSEN

ABSTRACT. We obtain non-symmetric upper and lower bounds on the rate of convergence of general monotone approximation/numerical schemes for parabolic Hamilton Jacobi Bellman Equations by introducing a new notion of consistency. We apply our general results to various schemes including finite difference schemes, splitting methods and the classical approximation by piecewise constant controls.

1. INTRODUCTION

In this article, we are interested in the rate of convergence of general monotone approximation/numerical schemes for time-dependent Hamilton Jacobi Bellman (HJB) Equations.

In order to be more specific, the HJB Equations we consider are written in the following form

$$(1.1) \quad u_t + F(t, x, u, Du, D^2u) = 0 \quad \text{in } Q_T := (0, T] \times \mathbb{R}^N,$$

$$(1.2) \quad u(0, x) = u_0(x) \quad \text{in } \mathbb{R}^N,$$

where

$$F(t, x, r, p, X) = \sup_{\alpha \in \mathcal{A}} \{ \mathcal{L}^\alpha(t, x, r, p, X) \},$$

with

$$\mathcal{L}^\alpha(t, x, r, p, X) := -\text{tr}[a^\alpha(t, x)X] - b^\alpha(t, x)p - c^\alpha(t, x)r - f^\alpha(t, x).$$

The coefficients a^α , b^α , c^α , f^α and the initial data u_0 take values respectively in \mathbb{S}^N , the space of $N \times N$ symmetric matrices, \mathbb{R}^N , \mathbb{R} , \mathbb{R} , and \mathbb{R} . Under suitable assumptions (see **(A1)** in Section 2), the initial value problem (1.1)-(1.2) has a unique, bounded, Hölder continuous, viscosity solution u which is the value function of a finite horizon, optimal stochastic control problem.

We consider approximation/numerical schemes for (1.1)-(1.2) written in the following abstract way

$$(1.3) \quad \begin{aligned} S(h, t, x, u_h(t, x), [u_h]_{t,x}) &= 0 & \text{in } \mathcal{G}_h^+ &:= \mathcal{G}_h \setminus \{t = 0\}, \\ u_h(0, x) &= u_{h,0}(x) & \text{in } \mathcal{G}_h^0 &:= \mathcal{G}_h \cap \{t = 0\}, \end{aligned}$$

Date: June 24, 2005.

Key words and phrases. Hamilton-Jacobi-Bellman Equations, switching system, viscosity solution, approximation schemes, finite difference methods, splitting methods, convergence rate, error bound.

Jakobsen was supported by the Research Council of Norway, grant no. 151608/432.

where S is, loosely speaking, a consistent, monotone and uniformly continuous approximation of the equation (1.1) defined on a grid/mesh $\mathcal{G}_h \subset \overline{Q_T}$. The approximation parameter h can be multi-dimensional, e.g. h could be $(\Delta t, \Delta x)$, Δt , Δx denoting time and space discretization parameters, Δx can be itself multi-dimensional. The approximate solution is $u_h : \mathcal{G}_h \rightarrow \mathbb{R}$, $[u_h]_{t,x}$ is a function defined from u_h representing, typically, the value of u_h at other points than (t, x) . We assume that the total scheme including the initial value is well-defined on some appropriate subset of the space of bounded continuous functions on \mathcal{G}_h .

The abstract notation was introduced by Barles and Souganidis [3] to display clearly the monotonicity of the scheme. One of the main assumptions is that S is non-decreasing in u_h and non-increasing in $[u_h]_{t,x}$ with the classical ordering of functions. The typical approximation schemes we have in mind are various finite differences numerical scheme (see e.g. Kushner and Dupuis [13] and Bonnans and Zidani [5]) and control schemes based on the dynamic programming principle (see e.g. Camilli and Falcone [6]). However, for reasons explained below, we will not discuss control schemes in this paper.

The aim of this paper is to obtain estimates on the rate of the convergence of u_h to u . To obtain such results, one faces the double difficulty of having to deal with both *fully nonlinear* equations and *non-smooth* solutions. Since these equations may be also degenerate, the (viscosity) solutions are expected to be no more than Hölder continuous in general.

Despite of these difficulties, in the 80's, Crandall & Lions [10] provided the first optimal rates of convergence for first-order equations. We refer to Souganidis [27] for more general results in this direction. For technical reasons, the problem turns out to be more difficult for second-order equations, and the question remained open for a long time.

The breakthrough came in 1997 and 2000 with Krylov's papers [20, 21], and by now there exists several papers based on and extending his ideas, e.g. [1, 2, 11, 18, 22, 23]. The main idea of Krylov is a method named by himself "shaking the coefficients". Combined with a standard mollification argument, it allows one to get smooth *subsolutions* of the equation which approximate the solution. Then classical arguments involving consistency and monotonicity of the scheme yield a *one-sided bound* on the error. This method uses in a crucial way the convexity of the equation in u , Du , and D^2u .

It is much more difficult to obtain the other bound and essentially there are two main approaches. The first one consists of interchanging the role of the scheme and the equation. By applying the above explained ideas, one gets a sequence of appropriate smooth subsolutions of the scheme and concludes by consistency and the comparison principle for the equation. This idea was used in different articles, see [1, 11, 18, 20, 23]. Here, the key difficulty is to obtain a "continuous dependence" result for the scheme. Even though it is now standard to prove that the solutions of the HJB Equation with "shaken coefficients" remain close to the solution of the original equation, such type of results are not known for numerical schemes in general. We mention here the nice paper of Krylov [23] where such kind of results are obtained by a tricky Bernstein type of argument. However, these results along with the corresponding error bounds, only hold for equations and schemes with special structures.

The second approach consists of considering some approximation of the equation or the associated control problem and to obtain the other bound either by probabilistic arguments (as Krylov first did using piecewise constant controls, [22, 21]) or by building a sequence of appropriate “smooth supersolution” of the equation (see [2] where, as in the present paper, approximations by switching are considered).

The first approach leads to better error bounds than the second one but it seems to work only for very specific schemes and with restrictions on the equations. The second approach yields error bounds in “the general case” but at the expense of lower rates.

In this paper we use the second approach by extending the methods introduced in [2]. Compared with the various results of Krylov, we obtain better rates in most cases, our results apply to more general schemes, and we use a simpler, purely analytical approach. In fact our method is robust in the sense that it applies to “general” schemes without any particular form and under rather natural assumptions. However, we mention again that in certain situations the first approach can be used to get better rates, see in particular [23].

The results in [2] apply to stationary HJB equations set in whole space \mathbb{R}^N . In this paper we extend these results to initial value problems for time-dependent HJB equations. The latter case is much more interesting in view of applications, and from a mathematical point of view, slightly more difficult. However, in our opinion the most important difference between the two papers lays in the formulation of the consistency requirements and the main (abstract) results. Here we introduce a new (and more general) formulation that emphasizes more the non-symmetrical feature of the upper and lower bounds and their proofs. It is a kind of a recipe on how to obtain error bounds in different situations, one which we feel is easier to apply to new problems and gives better insight into how the error bounds are produced. We also present several technical improvements and simplifications in the proofs and, finally, several new applications, some for which error bounds have not appeared before: Finite difference methods (FDMs) using the θ -method for time discretization, semidiscrete splitting methods, and approximation by piecewise constant controls.

The results for finite difference approximations can be compared with the ones obtained by Krylov in [21, 22]. As in [2], we get the rate $1/5$ for monotone FDMs while the corresponding result in [22] is $1/21$. Of course, in special situations the rate can be improved to $1/2$ which is the optimal rate under our assumptions. We refer to [23] for the most general results in that direction, and to [12] for the optimality of the rate $1/2$. The results for semidiscrete splitting methods are new, while the ones for the control approximation we get $1/10$ which is worse than $1/6$ obtained by Krylov in [22]. It would be interesting to understand why Krylov is doing better than us here but not in the other cases.

We conclude this introduction by explaining the notations we will use throughout this paper. By $|\cdot|$ we mean the standard Euclidean norm in any \mathbb{R}^p type space (including the space of $N \times P$ matrices). In particular, if $X \in \mathbb{S}^N$, then $|X|^2 = \text{tr}(XX^T)$ where X^T denotes the transpose of X .

If w is a bounded function from some set $Q' \subset \overline{Q}_\infty$ into either \mathbb{R} , \mathbb{R}^M , or the space of $N \times P$ matrices, we set

$$|w|_0 = \sup_{(t,y) \in Q'} |w(t,y)|.$$

Furthermore, for $\delta \in (0, 1]$, we set

$$[w]_\delta = \sup_{(t,x) \neq (s,y)} \frac{|w(t,x) - w(s,y)|}{(|x-y| + |t-s|^{1/2})^\delta} \quad \text{and} \quad |w|_\delta = |w|_0 + [w]_\delta.$$

Let $C_b(Q')$ and $\mathcal{C}^{0,\delta}(Q')$, $\delta \in (0, 1]$, denote respectively the space of bounded continuous functions on Q' and the subset of $C_b(Q')$ in which the norm $|\cdot|_\delta$ is finite. Note in particular the choices $Q' = Q_T$ and $Q' = \mathbb{R}^N$. In the following we always suppress the domain Q' when writing norms.

We denote by \leq the component by component ordering in \mathbb{R}^M and the ordering in the sense of positive semi-definite matrices in \mathbb{S}^N . For the rest of this paper we let ρ denotes the same, fixed, positive smooth function with support in $\{0 < t < 1\} \times \{|x| < 1\}$ and mass 1. From this function ρ , we define the sequence of mollifiers $\{\rho_\varepsilon\}_{\varepsilon>0}$ as follows,

$$\rho_\varepsilon(t, x) = \frac{1}{\varepsilon^{N+2}} \rho\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right) \quad \text{in} \quad \overline{Q}_\infty.$$

The rest of this paper is organized as follows: In the next section we present results on the so-called switching approximation for the problem (1.1)-(1.2). As in [2], these results are crucial to obtain the general results on the rate of convergence of approximation/numerical schemes and are of an independent interest. Section 3 is devoted to state and prove the main result on the rate of convergence. Finally we present various applications to classical finite difference schemes, splitting method and on the classical approximation by piecewise constant controls.

2. CONVERGENCE RATE FOR A SWITCHING SYSTEM

In this section, we obtain the rate of convergence for a certain switching system approximations to the HJB equation (1.1). Such approximations have been studied in [14, 7], and a viscosity solutions theory of switching systems can be found in [28, 17, 16]. We consider the following type of switching systems,

$$(2.1) \quad \begin{aligned} F_i(t, x, v, \partial_t v_i, Dv_i, D^2 v_i) &= 0 && \text{in } Q_T, \quad i \in \mathcal{I} := \{1, \dots, M\}, \\ v(0, x) &= v_0(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where the solution $v = (v_1, \dots, v_M)$ is in \mathbb{R}^M , and for $i \in \mathcal{I}$, $(t, x) \in Q_T$, $r = (r_1, \dots, r_M) \in \mathbb{R}^M$, $p_t \in \mathbb{R}$, $p_x \in \mathbb{R}^N$, and $X \in \mathcal{S}^N$, F_i is given by

$$F_i(t, x, r, p_t, p_x, X) = \max \left\{ p_t + \sup_{\alpha \in \mathcal{A}_i} \mathcal{L}^\alpha(t, x, r_i, p_x, X); r_i - \mathcal{M}_i r \right\},$$

where the \mathcal{A}_i 's are subsets of \mathcal{A} , \mathcal{L}^α is defined below (1.1), and for $k > 0$,

$$\mathcal{M}_i r = \min_{j \neq i} \{r_j + k\}.$$

Finally for the initial data, we are interested here in the case when $v_0 = (u_0, \dots, u_0)$.

Under suitable assumptions on the data (See **(A1)** below), we have existence and uniqueness of a solution v of this system. Moreover, it is not so difficult to see that, as $k \rightarrow 0$, every component of v converge locally uniformly to the solution of the following HJB equation

$$(2.2) \quad \begin{aligned} u_t + \sup_{\alpha \in \mathcal{A}} \mathcal{L}^\alpha(x, u, Du, D^2 u) &= 0 && \text{in } Q_T, \\ u(0, x) &= u_0(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where $\tilde{\mathcal{A}} = \cup_i \mathcal{A}_i$.

The objective of this section is to obtain an error bound for this convergence. For the sake of simplicity, we restrict ourselves to the situation where the solutions are in $\mathcal{C}^{0,1}(Q_T)$, i.e. when they are bounded, Lipschitz continuous in x , and Hölder 1/2 in t . Such type of regularity is natural in this context. However, it is not difficult to adapt our approach to more general situations, and we give results in this direction in Section 6.

We will use the following assumption

(A1) For any $\alpha \in \mathcal{A}$, $a^\alpha = \frac{1}{2}\sigma^\alpha \sigma^{\alpha T}$ for some $N \times P$ matrix σ^α . Moreover, there is a constant K independent of α such that

$$|u_0|_1 + |\sigma^\alpha|_1 + |b^\alpha|_1 + |c^\alpha|_1 + |f^\alpha|_1 \leq K.$$

Assumption (A1) ensures the well-posedness of all the equations and systems of equations we consider in this paper; we refer the reader to the Appendix for a (partial) proof of this claim. In the present situation, we have the following well-posedness and regularity result.

Proposition 2.1. *Assume (A1). Then there exist unique solutions v and u of (2.1) and (2.2) respectively, satisfying*

$$|v|_1 + |u|_1 \leq C,$$

where the constant C only depends on T and K appearing in (A1).

Furthermore, if w_1 and w_2 are sub- and supersolutions of (2.1) or (2.2) satisfying $w_1(0, \cdot) \leq w_2(0, \cdot)$, then $w_1 \leq w_2$.

Remark 2.1. The functions $\sigma^\alpha, b^\alpha, c^\alpha, f^\alpha$ are a priori only defined for times $t \in [0, T]$. But they can easily be extended to times $[-r, T+r]$ for any $r \in \mathbb{R}^+$ in such a way that (A1) still holds. In view of Proposition 2.1 we can then solve our initial value problems (2.1) and (2.2) either up to time $T+r$ and even, by using a translation in time, on time intervals of the form $[-r, T+r]$. We will use this fact several times below.

In order to obtain the rate of convergence for the switching approximation, we use a regularization procedure introduced by Krylov [21, 1]. This procedure requires the following auxiliary system

$$(2.3) \quad \begin{aligned} F_i^\varepsilon(t, x, v^\varepsilon, \partial_t v_i^\varepsilon, Dv_i^\varepsilon, D^2 v_i^\varepsilon) &= 0 && \text{in } Q_{T+\varepsilon^2}, \quad i \in \mathcal{I}, \\ v^\varepsilon(0, x) &= v_0(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where $v^\varepsilon = (v_1^\varepsilon, \dots, v_M^\varepsilon)$,

$$F_i^\varepsilon(t, x, r, p_t, p_x, M) = \max \left\{ p_t + \sup_{\substack{\alpha \in \mathcal{A}_i \\ 0 \leq s \leq \varepsilon^2, |e| \leq \varepsilon}} \mathcal{L}^\alpha(t+s, x+e, r_i, p_x, X); r_i - \mathcal{M}_i r \right\},$$

and \mathcal{L} and \mathcal{M} are defined below (1.1) and (2.1) respectively. Note that we use here the extension mentioned in Remark 2.1.

By Theorems A.1 and A.3 in the Appendix, we have the following result:

Proposition 2.2. *Assume (A1). Then there exist a unique solution $v^\varepsilon : \overline{Q}_{T+\varepsilon^2} \rightarrow \mathbb{R}$ of (2.3) satisfying*

$$|v^\varepsilon|_1 + \frac{1}{\varepsilon} |v^\varepsilon - v|_0 \leq C,$$

where v solves (2.1) and the constant C only depends on T and K from (A1).

Furthermore, if w_1 and w_2 are sub- and supersolutions of (2.3) satisfying $w_1(0, \cdot) \leq w_2(0, \cdot)$, then $w_1 \leq w_2$.

We are now in a position to state and prove the main result of this section.

Theorem 2.3. *Assume (A1) and $v_0 = (u_0, \dots, u_0)$. If u and v are the solutions of (2.2) and (2.1) respectively, then for k small enough,*

$$0 \leq v_i - u \leq Ck^{1/3} \quad \text{in } Q_T, \quad i \in \mathcal{I},$$

where C only depends on T and K from (A1).

Proof. Since $w = (u, \dots, u)$ is a subsolution of (2.1), comparison for (2.1) (Proposition 2.1) yields $u \leq v_i$ for $i \in \mathcal{I}$.

To get the other bound, we use an argument suggested by P.-L. Lions [24] together with the regularization procedure of Krylov [21]. Consider first system (2.3). It follows that, for every $0 \leq s \leq \varepsilon^2$, $|e| \leq \varepsilon$,

$$\partial_t v_i^\varepsilon + \sup_{\alpha \in \mathcal{A}_i} \mathcal{L}^\alpha(t+s, x+e, v_i^\varepsilon(t, x), Dv_i^\varepsilon, D^2v_i^\varepsilon) \leq 0 \quad \text{in } Q_{T+\varepsilon^2}, \quad i \in \mathcal{I}.$$

After a change of variables, we see that for every $0 \leq s \leq \varepsilon^2$, $|e| \leq \varepsilon$, $v^\varepsilon(t-s, x-e)$ is a subsolution of the following system of uncoupled equations

$$(2.4) \quad \partial_t w_i + \sup_{\alpha \in \mathcal{A}_i} \mathcal{L}^\alpha(t, x, w_i, Dw_i, D^2w_i) = 0 \quad \text{in } Q_T^\varepsilon, \quad i \in \mathcal{I},$$

where $Q_T^\varepsilon := (\varepsilon^2, T) \times \mathbb{R}^N$. Define $v_\varepsilon := v^\varepsilon * \rho_\varepsilon$ where $\{\rho_\varepsilon\}_\varepsilon$ is the sequence of mollifiers defined at the end of the introduction. A Riemann-sum approximation shows that $v_\varepsilon(t, x)$ can be viewed as the limit of convex combinations of $v^\varepsilon(t-s, x-e)$'s for $0 < s < \varepsilon^2$ and $|e| < \varepsilon$. Since the $v^\varepsilon(t-s, x-e)$'s are subsolutions of the *convex* equation (2.4), so are the convex combinations. By the stability result for viscosity subsolutions we can now conclude that v_ε is itself a subsolution of (2.4). We refer to the Appendix in [1] for more details.

On the other hand, since v^ε is a continuous subsolution of (2.3), we have

$$v_i^\varepsilon \leq \min_{j \neq i} v_j^\varepsilon + k \quad \text{in } Q_{T+\varepsilon^2}, \quad i \in \mathcal{I}.$$

It follows that $\max_i v_i^\varepsilon(t, x) - \min_i v_i^\varepsilon(t, x) \leq k$ in $Q_{T+\varepsilon^2}$, and hence

$$|v_i^\varepsilon - v_j^\varepsilon|_0 \leq k, \quad i, j \in \mathcal{I}.$$

Then, by the definition and properties of v_ε , we have

$$|\partial_t v_{\varepsilon i} - \partial_t v_{\varepsilon j}|_0 \leq C \frac{k}{\varepsilon^2}, \quad |D^n v_{\varepsilon i} - D^n v_{\varepsilon j}|_0 \leq C \frac{k}{\varepsilon^n}, \quad n \in \mathbb{N}, \quad i, j \in \mathcal{I},$$

where C depends only on ρ and the uniform bounds on $v_{\varepsilon i}$ and $Dv_{\varepsilon i}$, i.e. on T and K given in (A1). Furthermore, from these bounds, we see that for $\varepsilon < 1$,

$$\left| \partial_t v_{\varepsilon j} + \sup_{\alpha \in \mathcal{A}_i} \mathcal{L}^\alpha[v_{\varepsilon j}] - \partial_t v_{\varepsilon i} - \sup_{\alpha \in \mathcal{A}_i} \mathcal{L}^\alpha[v_{\varepsilon i}] \right| \leq C \frac{k}{\varepsilon^2} \quad \text{in } Q_T^\varepsilon, \quad i, j \in \mathcal{I}.$$

Here, as above, C only depends on ρ , T and K . Since v_ε is a subsolution of (2.4), this means that,

$$\partial_t v_{\varepsilon i} + \sup_{\alpha \in \overline{\mathcal{A}}} \mathcal{L}^\alpha(x, v_{\varepsilon i}, Dv_{\varepsilon i}, D^2v_{\varepsilon i}) \leq C \frac{k}{\varepsilon^2} \quad \text{in } Q_T^\varepsilon, \quad i \in \mathcal{I}.$$

From assumption (A1) and the structure of the equation, we see that $v_{\varepsilon i} - te^{Kt}C\frac{k}{\varepsilon^2}$ is a subsolution of equation (2.2) restricted to Q_T^ε .

Comparison for (2.2) restricted to Q_T^ε (Proposition 2.1) yields

$$v_{\varepsilon i} - u \leq e^{Kt} \left(|v_{\varepsilon i}(\varepsilon^2, \cdot) - u(\varepsilon^2, \cdot)|_0 + Ct\frac{k}{\varepsilon^2} \right) \quad \text{in } Q_T^\varepsilon, \quad i \in \mathcal{I}.$$

Regularity of u and v_i (Proposition 2.1) implies that

$$|u(t, \cdot) - v_i(t, \cdot)|_0 \leq ([u]_1 + [v_i]_1)\varepsilon \quad \text{in } [0, \varepsilon^2].$$

Hence by Proposition 2.2, regularity of u and v_i^ε , and properties of mollifiers, we have

$$v_i - u \leq v_i - v_{\varepsilon i} + v_{\varepsilon i} - u \leq C\left(\varepsilon + \frac{k}{\varepsilon^2}\right) \quad \text{in } Q_T^\varepsilon, \quad i \in \mathcal{I}.$$

Minimizing w.r.t. ε now yields the result. \square

3. CONVERGENCE RATE FOR THE HJB EQUATION

In this section we derive our main result, an error bound for the convergence of the solution of the scheme (1.3) to the solution of the HJB Equation (1.1)-(1.2). As in [2], this result is general and derived using only PDE methods, and it extends and improves earlier results by Krylov [20, 21], Barles and Jakobsen [1, 18]. Compared to [2], we consider here the time-dependent case and introduce a new, improved, formulation of the consistency requirement.

Throughout this section, we assume that (A1) holds and we recall that, by Proposition 2.1, there exists a unique $C^{0,1}$ -solution u of (1.1) satisfying $|u|_1 \leq C$, where the constant C only depends on T and K from (A1). In Section 6, we will weaken assumption (A1) and give results for $C^{0,\beta}$ solutions, $\beta \in (0, 1)$.

In order to get a lower bound on the error, we have to require a technical assumption: If $\{\alpha_i\}_{i \in \mathcal{I}} \subset \mathcal{A}$ is a sufficiently refined grid for \mathcal{A} , the solution associated to the control set $\{\alpha_i\}_{i \in \mathcal{I}}$ is close to u . In fact for this to be true we need to assume that the coefficients $\sigma^\alpha, b^\alpha, c^\alpha, f^\alpha$ can be approximated uniformly in (t, x) by $\sigma^{\alpha_i}, b^{\alpha_i}, c^{\alpha_i}, f^{\alpha_i}$. The precise assumption is:

(A2) For every $\delta > 0$, there are $M \in \mathbb{N}$ and $\{\alpha_i\}_{i=1}^M \subset \mathcal{A}$, such that for any $\alpha \in \mathcal{A}$,

$$\inf_{1 \leq i \leq M} (|\sigma^\alpha - \sigma^{\alpha_i}|_0 + |b^\alpha - b^{\alpha_i}|_0 + |c^\alpha - c^{\alpha_i}|_0 + |f^\alpha - f^{\alpha_i}|_0) < \delta.$$

We point out that this assumptions is automatically satisfied if either \mathcal{A} is a finite set or if \mathcal{A} is compact and $\sigma^\alpha, b^\alpha, c^\alpha, f^\alpha$ are uniformly continuous functions of t, x , and α .

Next we introduce the following assumptions for the scheme (1.3).

(S1) (Monotonicity) There exists $\lambda, \mu \geq 0, h_0 > 0$ such that if $|h| \leq h_0, u \leq v$ are functions in $C_b(\mathcal{G}_h)$, and $\phi(t) = e^{\mu t}(a + bt) + c$ for $a, b, c \geq 0$, then

$$S(h, t, x, r + \phi(t), [u + \phi]_{t,x}) \geq S(h, t, x, r, [v]_{t,x}) + b/2 - \lambda c \quad \text{in } \mathcal{G}_h^+.$$

(S2) (Regularity) For every h and $\phi \in C_b(\mathcal{G}_h)$, the function $(t, x) \mapsto S(h, t, x, \phi(t, x), [\phi]_{t,x})$ is bounded and continuous in \mathcal{G}_h^+ and the function $r \mapsto S(h, t, x, r, [\phi]_{t,x})$ is uniformly continuous for bounded r , uniformly in $(t, x) \in \mathcal{G}_h^+$.

Remark 3.1. In (S1) and (S2) we may replace $C_b(\mathcal{G}_h)$ by any relevant subset of this space. The point is that (1.3) has to make sense for the class of functions used. In Section 4, $C_b(\mathbb{R}^N)$ is itself the relevant class of functions, while, in Section 5, it is $C(\{0, 1, \dots, n_T\}; \mathcal{C}^{0,1}(\mathbb{R}^N))$ (since $\mathcal{G}_h = \{0, 1, \dots, n_T\} \times \mathbb{R}^N$).

Assumptions (S1) and (S2) imply a comparison result for the scheme (1.3), see Lemma 3.2 below.

Let us now state the key consistency conditions.

(S3)(i) (Sub-consistency) There exists a function $E_1(\tilde{K}, h, \varepsilon)$ such that for any sequence $\{\phi_\varepsilon\}_{\varepsilon>0}$ of smooth functions satisfying

$$|\partial_t^{\beta_0} D^{\beta'} \phi_\varepsilon(x, t)| \leq \tilde{K} \varepsilon^{1-2\beta_0-|\beta'|} \quad \text{in } \bar{Q}_T, \quad \text{for any } \beta_0 \in \mathbb{N}, \beta' = (\beta'_i)_i \in \mathbb{N}^N,$$

where $|\beta'| = \sum_{i=1}^N \beta'_i$, the following inequality holds:

$$S(h, t, x, \phi_\varepsilon(t, x), [\phi_\varepsilon]_{t,x}) \leq \phi_{\varepsilon t} + F(t, x, \phi, D\phi_\varepsilon, D^2\phi_\varepsilon) + E_1(\tilde{K}, h, \varepsilon) \quad \text{in } \mathcal{G}_h^+.$$

(S3)(ii) (Super-consistency) There exists a function $E_2(\tilde{K}, h, \varepsilon)$ such that for any sequence $\{\phi_\varepsilon\}_\varepsilon$ of smooth functions satisfying

$$|\partial_t^{\beta_0} D^{\beta'} \phi_\varepsilon(x, t)| \leq \tilde{K} \varepsilon^{1-2\beta_0-|\beta'|} \quad \text{in } \bar{Q}_T, \quad \text{for any } \beta_0 \in \mathbb{N}, \beta' \in \mathbb{N}^N,$$

the following inequality holds:

$$S(h, t, x, \phi_\varepsilon(t, x), [\phi_\varepsilon]_{t,x}) \geq \phi_{\varepsilon t} + F(t, x, \phi, D\phi_\varepsilon, D^2\phi_\varepsilon) - E_2(\tilde{K}, h, \varepsilon) \quad \text{in } \mathcal{G}_h^+.$$

Typically the ϕ_ε we have in mind in (S3) are of the form $\chi_\varepsilon * \rho_\varepsilon$ where $(\chi_\varepsilon)_\varepsilon$ is a sequence of uniformly bounded functions in $\mathcal{C}^{0,1}$ and ρ_ε is the mollifier defined at the end of the introduction.

The main result in this paper is the following:

Theorem 3.1. *Assume (A1), (S1), (S2) and that (1.3) has a unique solution u_h in $C_b(\mathcal{G}_h)$. Let u denote the solution of (1.1)-(1.2), and let h be sufficiently small.*

(a) **(Upper bound)** *If (S3)(i) holds, then there exists a constant C depending only μ, K in (S1), (A1) such that*

$$u - u_h \leq e^{\mu t} |(u_0 - u_{0,h})^+|_0 + C \min_{\varepsilon>0} \left(\varepsilon + E_1(\tilde{K}, h, \varepsilon) \right) \quad \text{in } \mathcal{G}_h,$$

where $\tilde{K} = |u|_1$.

(b) **(Lower bound)** *If (S3)(ii) and (A3) holds, then there exists a constant C depending only μ, K in (S1), (A1) such that*

$$u - u_h \geq -e^{\mu t} |(u_0 - u_{0,h})^-|_0 - C \min_{\varepsilon>0} \left(\varepsilon^{1/3} + E_2(\tilde{K}, h, \varepsilon) \right) \quad \text{in } \mathcal{G}_h,$$

where $\tilde{K} = |u|_1$.

The motivation for this new formulation of the upper and lower bounds is three-fold: (i) in some applications, $E_1 \neq E_2$ and therefore it is natural to have such disymmetry in the consistency requirement (see Section 5), (ii) from the proof it can be seen that the upper bound (a) is proven independently of the lower bound (b), and most importantly, (iii) the new formulation describes completely how the bounds are obtained from the consistency requirements. The good h -dependence and the bad ε dependence of E_1 and E_2 are combined in the minimization process to give the final bounds, see Remark 3.2 below.

Since the minimum is achieved for $\varepsilon \ll 1$, the upper bound is in general much better than the lower bound (in particular in cases where $E_1 = E_2$).

Finally note that the existence of a u_h in $C_b(\mathcal{G}_h)$ must be proved for each particular scheme S . We refer to [20, 21, 1, 18] for examples of such arguments.

Remark 3.2. In the case of a finite difference method with a time step Δt and maximal mesh size in space Δx , a standard formulation of the consistency requirement would be

(S3') There exist finite sets $I \subset \mathbb{N} \times \mathbb{N}_0^N, \bar{I} \subset \mathbb{N}_0 \times \mathbb{N}^N$ and constants $K_c \geq 0, k_\beta, \bar{k}_{\bar{\beta}}$ for $\beta = (\beta_0, \beta') \in I, \bar{\beta} = (\bar{\beta}_0, \bar{\beta}') \in \bar{I}$ such that for every $h = (\Delta t, \Delta x) > 0, (t, x) \in \mathcal{G}_h^+$, and smooth functions ϕ :

$$\begin{aligned} & |\phi_t + F(t, x, \phi, D\phi, D^2\phi) - S(h, t, x, \phi(t, x), [\phi]_{t,x})| \\ & \leq K_c \sum_{\beta \in I} |\partial_t^{\beta_0} D^{\beta'} \phi|_0 \Delta t^{k_\beta} + K_c \sum_{\bar{\beta} \in \bar{I}} |\partial_t^{\bar{\beta}_0} D^{\bar{\beta}'} \phi|_0 \Delta x^{\bar{k}_{\bar{\beta}}}. \end{aligned}$$

The corresponding version of (S3) is obtained by plugging ϕ_ε into (S3') and using the estimates on its derivatives. The result is

$$\begin{aligned} E_1(\tilde{K}, h, \varepsilon) &= E_2(\tilde{K}, h, \varepsilon) \\ &= \tilde{K} K_c \sum_{\beta \in I} \varepsilon^{1-2\beta_0-|\beta'|} \Delta t^{k_\beta} + \tilde{K} K_c \sum_{\bar{\beta} \in \bar{I}} \varepsilon^{1-2\bar{\beta}_0-|\bar{\beta}'|} \Delta x^{\bar{k}_{\bar{\beta}}}. \end{aligned}$$

From this formula we see that the dependence in the small parameter ε is bad since all the exponents of ε are negative, while the dependence on $\Delta t, \Delta x$ is good since their exponents are positive.

Remark 3.3. Assumption (S1) contains two different kinds of information. First, by taking $\phi \equiv 0$ it implies that the scheme is nondecreasing with respect to the $[u]$ argument. Second, by taking $u \equiv v$ it indicates that a parabolic equation – an equation with a u_t term – is being approximated. Both these points play a crucial role in the proof of the comparison principle for (1.3) (Lemma 3.2 below).

To better understand that assumption (S1) implies parabolicity of the scheme, consider the following more restrictive assumption:

(S1') (Monotonicity) There exists $\lambda \geq 0, \bar{K} > 0$ such that if $u \leq v, u, v \in C_b(\mathcal{G}_h)$, and $\phi : [0, T] \rightarrow \mathbb{R}$ is smooth, then

$$\begin{aligned} & S(h, t, x, r + \phi(t), [u + \phi]_{t,x}) \\ & \geq S(h, t, x, r, [v]_{t,x}) + \phi'(t) - \bar{K} \Delta t |\phi''|_0 - \lambda \phi^+(t) \quad \text{in } \mathcal{G}_h^+. \end{aligned}$$

Here $h = (\Delta t, h')$ where h' representing a small parameter related to e.g. the space discretization. It is easy to see that (S1') implies (S1), e.g. with the same value for λ and the following values of μ and h_0 :

$$\mu = \lambda + 1 \quad \text{and} \quad h_0^{-1} = 2\bar{K}e^{(\lambda+1)T}(\lambda + 1)(2 + (\lambda + 1)T).$$

Assumption (S1') is satisfied for all *monotone* finite difference in time approximations of (1.1), e.g. *monotone* Runge-Kutta methods and *monotone* multi-step methods, both explicit and implicit methods. We have emphasized the word *monotone* because whereas many Runge Kutta methods actually lead to monotone schemes for (1.1) (possibly under a CFL condition), it seems that the most commonly used

multistep methods (Adams-Bashforth, BDS) do not. We refer to [26] for an example of a multistep method that yields a monotone approximation of (1.1).

Proof of Theorem 3.1. We start by proving that conditions (S1) and (S2) imply a comparison result for bounded continuous sub and supersolutions of (1.3).

Lemma 3.2. *Assume (S1), (S2), and that $u, v \in C_b(\mathcal{G}_h)$ satisfy*

$$S(h, t, x, u(t, x), [u]_{t,x}) \leq g_1 \quad \text{in } \mathcal{G}_h^+,$$

$$S(h, t, x, v(t, x), [v]_{t,x}) \geq g_2 \quad \text{in } \mathcal{G}_h^+,$$

where $g_1, g_2 \in C_b(\mathcal{G}_h)$. Then

$$u - v \leq e^{\mu t} |(u(0, \cdot) - v(0, \cdot))^+|_0 + 2te^{\mu t} |(g_1 - g_2)^+|_0,$$

where λ and μ are given by (S1).

Proof. 1. First, we notice that it suffices to prove the lemma in the case

$$(3.1) \quad u(0, x) - v(0, x) \leq 0 \quad \text{in } \mathcal{G}_h^0,$$

$$(3.2) \quad g_1(t, x) - g_2(t, x) \leq 0 \quad \text{in } \mathcal{G}_h.$$

The general case follows from this result after noting that, by (S1),

$$w = v + e^{\mu t} (|(u(0, \cdot) - v(0, \cdot))^+|_0 + 2t|(g_1 - g_2)^+|_0),$$

satisfies $S(h, t, x, w(t, x), [w]_{t,x}) \geq g_1$ in \mathcal{G}_h^+ and $u(0, x) - w(0, x) \leq 0$ in \mathcal{G}_h^0 .

2. We assume that (3.1) and (3.2) hold and, for $b \geq 0$, we set $\psi_b(t) = e^{\mu t} 2bt$ where μ is given by (S1) and

$$M(b) = \sup_{\mathcal{G}_h} \{u - v - \psi_b\}.$$

We have to prove that $M(0) \leq 0$ and we argue by contradiction assuming that $M(0) > 0$.

3. First we consider some $b \geq 0$ for which $M(b) > 0$ and take a sequence $\{(t_n, x_n)\}_n \subset \mathcal{G}_h$ such that

$$\delta_n := M(b) - (u - v - \psi_b)(t_n, x_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Since $M(b) > 0$ and (3.1) holds, $t_n > 0$ for all sufficiently large n and for such n , we have

$$\begin{aligned} g_1 &\geq S(h, t_n, x_n, u, [u]_{t_n, x_n}) && (u \text{ subsolution}) \\ &\geq S(h, t_n, x_n, v + \psi_b + M(b) - \delta_n, [v + \psi_b + M(b)]_{t_n, x_n}) && (S1), \phi \equiv 0 \\ &\geq \omega(\delta_n) \\ &\quad + S(h, t_n, x_n, v + \psi_b + M(b), [v + \psi_b + M(b)]_{t_n, x_n}) && (S2) \\ &\geq \omega(\delta_n) + b - \lambda M(b) + S(h, t_n, x_n, v, [v]_{t_n, x_n}) && (S1), \phi = \psi + M \\ &\geq \omega(\delta_n) + b - \lambda M(b) + g_2, && (v \text{ supersolution}) \end{aligned}$$

where we have dropped the dependence in t_n, x_n of u, v and ψ_b for the sake of simplicity of notation. Recalling (3.2) and sending $n \rightarrow \infty$ lead to

$$b - \lambda M(b) \leq 0.$$

4. Since $M(b) \leq M(0)$, the above inequality yields a contradiction for b large, so for such b , $M(b) \leq 0$. On the other hand, since $M(b)$ is a continuous function of b and $M(0) > 0$, there exists a minimal solution $\bar{b} > 0$ of $M(\bar{b}) = 0$. For $\delta > 0$

satisfying $\bar{b} - \delta > 0$, we have $M(\bar{b} - \delta) > 0$ and $M(\bar{b} - \delta) \rightarrow 0$ as $\delta \rightarrow 0$. But, by 3 we have

$$\bar{b} - \delta \leq \lambda M(\bar{b} - \delta),$$

which is a contradiction for δ small enough since $\bar{b} > 0$. \square

Now we turn to the **proof of the upper bound**, i.e. of (a). We just sketch it since it relies on the regularization procedure of Krylov which is used in Section 2. We also refer to Krylov [20, 21], Barles and Jakobsen [1, 18] for more details. The main steps are:

1. Introduce the solution u^ε of

$$\begin{aligned} u_t^\varepsilon + \sup_{0 \leq s \leq \varepsilon^2, |e| \leq \varepsilon} F(t+s, x+e, u^\varepsilon(t, x), Du^\varepsilon, D^2u^\varepsilon) &= 0 && \text{in } Q_{T+\varepsilon^2}, \\ u^\varepsilon(x, 0) &= u_0(x) && \text{in } \mathbb{R}^N. \end{aligned}$$

Essentially as a consequence of Proposition 2.1, it follows that u^ε belongs to $\mathcal{C}^{0,1}(Q_T)$ with a uniform $\mathcal{C}^{0,1}(Q_T)$ -bound \bar{K} .

2. By analogous arguments to the ones used in Section 2, it is easy to see that $u_\varepsilon := u^\varepsilon * \rho_\varepsilon$ is a subsolution of (1.1). By combining regularity and continuous dependence results (Theorem A.3 in the Appendix), we also have $|u_\varepsilon - u|_0 \leq C\varepsilon$ where C only depends T and K in (A1).

3. Plugging u_ε into the scheme and using (S3)(i) and the uniform estimates on u^ε we get

$$S(h, t, x, u_\varepsilon(t, x), [u_\varepsilon]_{t,x}) \leq E_1(\bar{K}, h, \varepsilon) \quad \text{in } \mathcal{G}_h^+,$$

where \bar{K} is the above mentioned $\mathcal{C}^{0,1}$ -uniform estimate on u^ε which depends only on the data and is essentially the same as for u .

4. Use Lemma 3.2 to compare u_ε and u_h and conclude by using the control we have on $u - u_\varepsilon$ and by taking the minimum in ε .

We now provide the **proof of the lower bound**, i.e. of (b). Unfortunately, contrarily to the proof of (a), we do not know how to obtain a sequence of approximate, global, smooth supersolutions. As in [2], we are going to obtain approximate “almost smooth” supersolutions which are in fact supersolutions which are smooth at the “right points”. We build them by considering the following switching system approximation of (1.1):

$$(3.3) \quad \begin{aligned} F_i^\varepsilon(t, x, v^\varepsilon, \partial_t v_i^\varepsilon, Dv_i^\varepsilon, D^2v_i^\varepsilon) &= 0 && \text{in } Q_{T+2\varepsilon^2}, \quad i \in \mathcal{I} := \{1, \dots, M\}, \\ v^\varepsilon(0, x) &= v_0(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where $v^\varepsilon = (v_1^\varepsilon, \dots, v_M^\varepsilon)$, $v_0 = (u_0, \dots, u_0)$,

$$(3.4) \quad \begin{aligned} F_i^\varepsilon(t, x, r, p_t, p_x, X) &= \\ \max \left\{ p_t + \min_{0 \leq s \leq \varepsilon^2, |e| \leq \varepsilon} \mathcal{L}^{\alpha_i}(t+s-\varepsilon^2, x+e, r_i, p_x, X); r_i - \mathcal{M}_i r \right\}, \end{aligned}$$

and \mathcal{L} and \mathcal{M} are defined below (1.1) and (2.1) respectively. The solution of this system is expected to be close to the solution of (1.1) if k and ε are small and $\{\alpha_i\}_{i \in \mathcal{I}} \subset \mathcal{A}$ is a sufficiently refined grid for \mathcal{A} . This is where the assumption (A2) plays a role.

For equation (3.3), we have the following result.

Lemma 3.3. *Assume (A1).*

(a) *There exists a unique solution v^ε of (3.3) satisfying $|v^\varepsilon|_1 \leq \bar{K}$, where \bar{K} only depends on T and K from (A1).*

(b) *Assume in addition (A2) and let u denote the solution of (1.1). For $i \in \mathcal{I}$, we introduce the functions $\bar{v}_i^\varepsilon : [-\varepsilon^2, T + \varepsilon^2] \times \mathbb{R}^N \rightarrow \mathbb{R}$ defined by*

$$\bar{v}_i^\varepsilon(t, x) := v_i^\varepsilon(t - \varepsilon^2, x).$$

Then, for any $\delta > 0$, there are $M \in \mathbb{N}$ and $\{\alpha_i\}_{i=1}^M \subset \mathcal{A}$ such that

$$\max_i |u - \bar{v}_i^\varepsilon|_0 \leq C(\varepsilon + k^{1/3} + \delta),$$

where C only depends on T and K from (A1).

In order to simplify the arguments of the proof of the lower bound (to have the simplest possible formulation of Lemma 3.5 below), we need the solutions of the equation with “shaken coefficients” to be defined in a slightly larger domain than Q_T . More precisely on

$$Q_T^\varepsilon := (-\varepsilon^2, T + \varepsilon^2] \times \mathbb{R}^N.$$

This is the role of the \bar{v}_i^ε 's. In fact they solve the same system of equations as the v_i^ε 's but on Q_T^ε and with $\mathcal{L}^{\alpha_i}(t + s - \varepsilon^2, x + e, r_i, p_x, X)$ being replaced by $\mathcal{L}^{\alpha_i}(t + s, x + e, r_i, p_x, X)$ in (3.4).

The (almost) smooth supersolutions of (1.1) we are looking for are built out of the \bar{v}_i^ε 's by mollification. Before giving the next lemma, we remind the reader that the sequence of mollifiers $\{\rho_\varepsilon\}_\varepsilon$ is defined at the end of the introduction.

Lemma 3.4. *Assume (A1) and define $v_{\varepsilon i} := \rho_\varepsilon * \bar{v}_i^\varepsilon : \bar{Q}_{T+\varepsilon^2} \rightarrow \mathbb{R}$ for $i \in \mathcal{I}$.*

(a) *There is a constant C depending only on T and K from (A1), such that*

$$|v_{\varepsilon j} - \bar{v}_j^\varepsilon| \leq C(k + \varepsilon) \quad \text{in } Q_{T+\varepsilon^2}, \quad i, j \in \mathcal{I}.$$

(b) *Assume in addition that $\varepsilon \leq (8 \sup_i [v_i^\varepsilon]_1)^{-1} k$. For every $(t, x) \in Q_T$, if $j := \operatorname{argmin}_{i \in \mathcal{I}} v_{\varepsilon i}(t, x)$, then*

$$\partial_t v_{\varepsilon j}(t, x) + \mathcal{L}^{\alpha_j}(t, x, v_{\varepsilon j}(t, x), Dv_{\varepsilon j}(t, x), D^2 v_{\varepsilon j}(t, x)) \geq 0.$$

The proofs of these two lemmas will be given at the end of this section.

The key consequence is the following result which is the corner-stone of the proof of the lower bound.

Lemma 3.5. *Assume (A1) and that $\varepsilon \leq (8 \sup_i [v_i^\varepsilon]_1)^{-1} k$. Then the function $w := \min_{i \in \mathcal{I}} v_{\varepsilon i}$ is an approximate supersolution of the scheme (1.3) in the sense that*

$$S(h, t, x, w(t, x), [w]_{t,x}) \geq -E_2(\bar{K}, h, \varepsilon) \quad \text{in } \mathcal{G}_h^+,$$

where \bar{K} comes from Lemma 3.3.

Proof. Let $(t, x) \in Q_T$ and j be as in Lemma 3.4 (b). We see that $w(t, x) = v_{\varepsilon j}(t, x)$ and $w \leq v_{\varepsilon j}$ in \mathcal{G}_h , and hence the monotonicity of the scheme (cf. (S1)) implies that

$$S(h, t, x, w(t, x), [w]_{t,x}) \geq S(h, t, x, v_{\varepsilon j}(t, x), [v_{\varepsilon j}]_{t,x}).$$

But then, by (S3)(ii),

$$\begin{aligned} & S(h, t, x, w(t, x), [w]_{t,x}) \\ & \geq \partial_t v_{\varepsilon j}(t, x) + \mathcal{L}^{\alpha_j}(t, x, v_{\varepsilon j}(t, x), Dv_{\varepsilon j}(t, x), D^2 v_{\varepsilon j}(t, x)) - E_2(\bar{K}, h, \varepsilon), \end{aligned}$$

and the proof complete by applying Lemma 3.4 (b). \square

It is now straightforward to conclude the proof of the lower bound, we simply choose $k = 8 \sup_i [v_i^\varepsilon]_1 \varepsilon$ and use Lemma 3.2 to compare u_h and w . This yields

$$u_h - w \leq e^{\mu t} |(u_{h,0} - w(0, \cdot))^+|_0 + 2te^{\mu t} E_2(\bar{K}, h, \varepsilon) \quad \text{in } \mathcal{G}_h.$$

But, by Lemmas 3.3 (b) and 3.4 (a), we have

$$|w - u|_0 \leq C(\varepsilon + k + k^{1/3} + \delta),$$

and therefore

$$u_h - u \leq e^{\mu t} |(u_{h,0} - u_0)^+|_0 + 2te^{\mu t} E_2(\bar{K}, h, \varepsilon) + C(\varepsilon + k + k^{1/3} + \delta) \quad \text{in } \mathcal{G}_h,$$

for some constant C . In view of our choice of k , we conclude the proof by minimizing w.r.t ε .

Now we give the proofs of Lemmas 3.3 and 3.4.

Proof of Lemma 3.3. **1.** We first approximate (1.1) by

$$\begin{aligned} v_t + \sup_{i \in \mathcal{I}} \mathcal{L}^{\alpha_i}(t, x, v, Dv, D^2v) &= 0 && \text{in } Q_T, \\ v(0, x) &= u_0(x) && \text{in } \mathbb{R}^N. \end{aligned}$$

From assumption (A2) and Lemmas A.1 and A.3 in the Appendix, it follows that there exists a unique solution v of the above equation satisfying

$$|v - u|_0 \leq C\delta,$$

where C only depends on T and K from (A1).

2. We continue by approximating the above equation by the following switching system

$$\begin{aligned} \max \left\{ \partial_t v_i + \mathcal{L}^{\alpha_i}(t, x, v_i, Dv_i, D^2v_i); v_i - \mathcal{M}_i v \right\} &= 0 && \text{in } Q_T, \quad i \in \mathcal{I}, \\ v(0, x) &= v_0(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where $v_0 = (u_0, \dots, u_0)$ and \mathcal{M} is defined below (2.1). From Proposition 2.1 and Theorem 2.3 in Section 2 we have existence and uniqueness of a solution $\bar{v} = (\bar{v}_1, \dots, \bar{v}_M)$ of the above system satisfying

$$|\bar{v}_i - v|_0 \leq Ck^{1/3}, \quad i \in \mathcal{I},$$

where C only depends on the mollifier ρ , T , and K from (A1).

3. The switching system defined in the previous step is nothing but (3.3) with $\varepsilon = 0$ or (2.3) with the \mathcal{A}_i 's being singletons. Theorems A.1 and A.3 in the Appendix yield the existence and uniqueness of a solution $v^\varepsilon : \bar{Q}_{T+2\varepsilon^2} \rightarrow \mathbb{R}$ of (3.3) satisfying

$$|v^\varepsilon|_1 + \frac{1}{\varepsilon} |v^\varepsilon - \bar{v}|_0 \leq C,$$

where C only depends on T and K from (A1).

4. The proof is complete by combining the estimates in steps 1 – 3, and noting that $|v^\varepsilon - \bar{v}^\varepsilon| \leq [v^\varepsilon]_1 \varepsilon$ in $Q_{T+\varepsilon^2}$ and (A2) is only needed in step 1. \square

Proof of Lemma 3.4. We start by (a). From the properties of mollifiers and the Hölder continuity of \bar{v}^ε , it is immediate that

$$(3.5) \quad |v_{\varepsilon i} - \bar{v}_i^\varepsilon| \leq C\varepsilon \quad \text{in } \bar{Q}_{T+\varepsilon^2}, \quad i \in \mathcal{I},$$

where $C = 2 \max_i [\bar{v}_i^\varepsilon]_1 = 2 \max_i [v_i^\varepsilon]_1$ depends only on T and K from (A1). Furthermore as we pointed out after the statement of Lemma 3.3, \bar{v}^ε solves a switching system in Q_T^ε , so arguing as in the proof of Theorem 2.3 in Section 2 leads to

$$0 \leq \max_i \bar{v}_i^\varepsilon - \min_i \bar{v}_i^\varepsilon \leq k \quad \text{in } Q_T^\varepsilon.$$

From these two estimates, (a) follows.

Now consider (b). Fix an arbitrary point $(t, x) \in Q_T$ and set

$$j = \operatorname{argmin}_{i \in \mathcal{I}} v_{\varepsilon i}(t, x).$$

Then, by definition of \mathcal{M} and j , we have

$$v_{\varepsilon j}(t, x) - \mathcal{M}_j v_\varepsilon(t, x) = \max_{i \neq j} \{v_{\varepsilon j}(t, x) - v_{\varepsilon i}(t, x) - k\} \leq -k,$$

and the bound (3.5) leads to

$$\bar{v}_j^\varepsilon(t, x) - \mathcal{M}_j \bar{v}^\varepsilon(t, x) \leq -k + 2 \max_i [v_i^\varepsilon]_1 2\varepsilon.$$

Next, by using the Hölder continuity of \bar{v}^ε (Lemma 3.3), for any $(\bar{t}, \bar{x}) \in Q_T^\varepsilon$, we have

$$\bar{v}_j^\varepsilon(\bar{t}, \bar{x}) - \mathcal{M}_j \bar{v}^\varepsilon(\bar{t}, \bar{x}) \leq -k + 2 \max_i [v_i^\varepsilon]_1 (2\varepsilon + |x - \bar{x}| + |t - \bar{t}|^{1/2}).$$

From this we conclude that if $|x - \bar{x}| < \varepsilon$, $|t - \bar{t}| < \varepsilon^2$, and $\varepsilon \leq (8 \max_i [v_i^\varepsilon]_1)^{-1} k$, then

$$\bar{v}_j^\varepsilon(\bar{t}, \bar{x}) - \mathcal{M}_j \bar{v}^\varepsilon(\bar{t}, \bar{x}) < 0,$$

and by equation (3.3) and the definition of \bar{v}^ε , $\bar{v}^\varepsilon(t, x) = v^\varepsilon(t - \varepsilon^2, x)$,

$$\partial_t \bar{v}_j^\varepsilon(\bar{t}, \bar{x}) + \inf_{0 \leq s \leq \varepsilon^2, |e| \leq \varepsilon} \mathcal{L}^{\alpha j}(\bar{t} + s, \bar{x} + e, \bar{v}_j^\varepsilon(\bar{t}, \bar{x}), D\bar{v}_j^\varepsilon(\bar{t}, \bar{x}), D^2\bar{v}_j^\varepsilon(\bar{t}, \bar{x})) = 0.$$

After a change of variables, we see that, for every $0 \leq s < \varepsilon^2$, $|e| < \varepsilon$,

$$(3.6) \quad \begin{aligned} & \partial_t \bar{v}_j^\varepsilon(t - s, x - e)(t, x) \\ & + \mathcal{L}^{\alpha j}(t, x, \bar{v}_j^\varepsilon(t - s, x - e), D\bar{v}_j^\varepsilon(t - s, x - e), D^2\bar{v}_j^\varepsilon(t - s, x - e)) \geq 0. \end{aligned}$$

In other words, for every $0 \leq s < \varepsilon^2$, $|e| < \varepsilon$, $\bar{v}_j^\varepsilon(t - s, x - e)$ is a (viscosity) supersolution at (t, x) of

$$(3.7) \quad \chi_t + \mathcal{L}^{\alpha j}(t, x, \chi, D\chi, D^2\chi) = 0.$$

By mollifying (3.6) (w.r.t. the (s, e) -argument) we see that $v_{\varepsilon j}$ is also a smooth supersolution of (3.7) at (t, x) and hence a (viscosity) supersolution of the HJB equation (1.1) at (t, x) . This is correct since $v_{\varepsilon j}$ can be viewed as the limit of convex combinations of supersolutions $\bar{v}_j^\varepsilon(t - s, x - e)$ of the linear and hence concave equation (3.7), we refer to the proof of Theorem 2.3 and to the Appendix in [1] for the details. We conclude the proof by noting that since $v_{\varepsilon j}$ is smooth, it is in fact a classical supersolution of (1.1) at x . \square

4. MONOTONE FINITE DIFFERENCE METHODS

In this section, we apply our main result to finite difference approximations of (1.1) based on the ϑ -method approximation in time and two different approximations in space: One proposed by Kushner [13] which is monotone when a is diagonal dominant and a (more) general approach based on directional second derivatives proposed by Bonnans and Zidani [5], but see also Dong and Krylov [11]. For simplicity we take $h = (\Delta t, \Delta x)$ and consider the uniform grid

$$\mathcal{G}_h = \Delta t\{0, 1, \dots, n_T\} \times \Delta x \mathbb{Z}^N.$$

4.1. Discretization in space. To explain the methods we first write equation (1.1) like

$$u_t + \sup_{\alpha \in \mathcal{A}} \left\{ -L^\alpha u - c^\alpha(t, x)u - f^\alpha(t, x) \right\} = 0 \quad \text{in } Q_T,$$

where

$$L^\alpha \phi(t, x) = \text{tr}[a^\alpha(t, x)D^2\phi(t, x)] + b^\alpha(t, x)D\phi(t, x).$$

To obtain a discretization in space we approximate L by a finite difference operator L_h , which we will take to be of the form

$$(4.1) \quad L_h^\alpha \phi(t, x) = \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta)(\phi(t, x + \beta\Delta x) - \phi(t, x)),$$

for $(t, x) \in \mathcal{G}_h$, where the *stencil* \mathcal{S} is a finite subset of $\mathbb{Z}^N \setminus \{0\}$, and where

$$(4.2) \quad C_h^\alpha(t, x, \beta) \geq 0 \quad \text{for all } \beta \in \mathcal{S}, (t, x) \in \mathcal{G}_h^+, h = (\Delta x, \Delta t) > 0, \alpha \in \mathcal{A}.$$

The last assumption says that the difference approximation is of *positive* type. This is a sufficient assumption for monotonicity in the stationary case.

(i) The approximation of Kushner.

We denote by $\{e_i\}_{i=1}^N$ the standard basis in \mathbb{R}^N and define

$$(4.3) \quad L_h^\alpha \phi = \sum_{i=1}^N \left[\frac{a_{ii}^\alpha}{2} \Delta_{ii} + \sum_{j \neq i} \left(\frac{a_{ij}^{\alpha+}}{2} \Delta_{ij}^+ - \frac{a_{ij}^{\alpha-}}{2} \Delta_{ij}^- \right) + b_i^{\alpha+} \delta_i^+ - b_i^{\alpha-} \delta_i^- \right] \phi,$$

where $b^+ = \max\{b, 0\}$, $b^- = (-b)^+$ ($b = b^+ - b^-$), and

$$\begin{aligned} \delta_i^\pm w(x) &= \pm \frac{1}{\Delta x} \{w(x \pm e_i \Delta x) - w(x)\}, \\ \Delta_{ii} w(x) &= \frac{1}{\Delta x^2} \{w(x + e_i \Delta x) - 2w(x) + w(x - e_i \Delta x)\}, \\ \Delta_{ij}^+ w(x) &= \frac{1}{2\Delta x^2} \{2w(x) + w(x + e_i \Delta x + e_j \Delta x) + w(x - e_i \Delta x - e_j \Delta x)\} \\ &\quad - \frac{1}{2\Delta x^2} \{w(x + e_i \Delta x) + w(x - e_i \Delta x) + w(x + e_j \Delta x) + w(x - e_j \Delta x)\}, \\ \Delta_{ij}^- w(x) &= -\frac{1}{2\Delta x^2} \{2w(x) + w(x + e_i \Delta x - e_j \Delta x) + w(x - e_i \Delta x + e_j \Delta x)\} \\ &\quad + \frac{1}{2\Delta x^2} \{w(x + e_i \Delta x) + w(x - e_i \Delta x) + w(x + e_j \Delta x) + w(x - e_j \Delta x)\}. \end{aligned}$$

The stencil is $\mathcal{S} = \{\pm e_i, \pm(e_i \pm e_j) : i, j = 1, \dots, N\}$, and it is easy to see that the coefficients in (4.1) are

$$\begin{aligned} C_h^\alpha(t, x, \pm e_i) &= \frac{a_{ii}^\alpha(x)}{2\Delta x^2} - \sum_{j \neq i} \frac{|a_{ij}^\alpha(x)|}{4\Delta x^2} + \frac{b_i^{\alpha\pm}(x)}{\Delta x}, \\ C_h^\alpha(t, x, e_i h \pm e_j h) &= \frac{a_{ij}^{\alpha\pm}(x)}{2\Delta x^2}, \quad i \neq j, \\ C_h^\alpha(t, x, -e_i h \pm e_j h) &= \frac{a_{ij}^{\alpha\mp}(x)}{2\Delta x^2}, \quad i \neq j. \end{aligned}$$

The approximation is of positive type (4.2) if and only if a is diagonal dominant, i.e.

$$(4.4) \quad a_{ii}^\alpha(t, x) - \sum_{j \neq i} |a_{ij}^\alpha(t, x)| \geq 0 \quad \text{in } Q_T, \quad \alpha \in \mathcal{A}, \quad i = 1, \dots, N.$$

(ii) The approximation of Bonnans and Zidani.

We assume that there is a (finite) stencil $\bar{\mathcal{S}} \subset \mathbb{Z}^N \setminus \{0\}$ and a set of positive coefficients $\{\bar{a}_\beta : \beta \in \bar{\mathcal{S}}\} \subset \mathbb{R}_+$ such that

$$(4.5) \quad a^\alpha(t, x) = \sum_{\beta \in \bar{\mathcal{S}}} \bar{a}_\beta^\alpha(t, x) \beta^T \beta \quad \text{in } Q_T, \quad \alpha \in \mathcal{A}.$$

Under assumption (4.5) we may rewrite the operator L using second order directional derivatives $D_\beta^2 = \text{tr}[\beta \beta^T D^2] = (\beta \cdot D)^2$,

$$L^\alpha \phi(t, x) = \sum_{\beta \in \bar{\mathcal{S}}} \bar{a}_\beta^\alpha(t, x) D_\beta^2 \phi(t, x) + b^\alpha(t, x) D \phi(t, x).$$

The approximation of Bonnans and Zidani is given by

$$(4.6) \quad L_h^\alpha \phi = \sum_{\beta \in \bar{\mathcal{S}}} \bar{a}_\beta^\alpha \Delta_\beta \phi + \sum_{i=1}^N \left[b_i^{\alpha+} \delta_i^+ - b_i^{\alpha-} \delta_i^- \right] \phi,$$

where Δ_β is an approximation of D_β^2 given by

$$\Delta_\beta w(x) = \frac{1}{|\beta|^2 \Delta x^2} \{w(x + \beta \Delta x) - 2w(x) + w(x - \beta \Delta x)\}.$$

In this case, the stencil is $\mathcal{S} = \pm \bar{\mathcal{S}} \cup \{\pm e_i : i = 1, \dots, N\}$ and the coefficients corresponding to (4.1) are given by

$$\begin{aligned} C_h^\alpha(t, x, \pm e_i) &= \frac{b_i^{\alpha\pm}(x)}{\Delta x}, & i = 1, \dots, N, \\ C_h^\alpha(t, x, \pm \beta) &= \frac{\bar{a}_\beta^\alpha(t, x)}{|\beta|^2 \Delta x^2}, & \beta \in \bar{\mathcal{S}}, \end{aligned}$$

and the sum of the two whenever $\beta = e_i$. Under assumption (4.5), which is more general than (4.4) (see below), this approximation is always of positive type.

For both approximations there is a constant $C > 0$, independent of Δx , such that, for every $\phi \in C^4(\mathbb{R}^N)$ and $(t, x) \in \mathcal{G}_h^+$,

$$(4.7) \quad |L^\alpha \phi(t, x) - L_h^\alpha \phi(t, x)| \leq C(|b^\alpha|_0 |D^2 \phi|_0 \Delta x + |a^\alpha|_0 |D^4 \phi|_0 \Delta x^2).$$

4.2. The fully discrete scheme. To obtain a fully discrete scheme, we apply the ϑ -method, $\vartheta \in [0, 1]$, to discretize the time derivative. The result is the following scheme,

$$(4.8) \quad \begin{aligned} u(t, x) &= u(t - \Delta t, x) \\ &\quad - (1 - \vartheta)\Delta t \sup_{\alpha \in \mathcal{A}} \{-L_h^\alpha u - c^\alpha u - f^\alpha\}(t - \Delta t, x) \\ &\quad - \vartheta\Delta t \sup_{\alpha \in \mathcal{A}} \{-L_h^\alpha u - c^\alpha u - f^\alpha\}(t, x) \quad \text{in } \mathcal{G}_h^+. \end{aligned}$$

The case $\vartheta = 0$ and $\vartheta = 1$ correspond to the forward and backward Euler time-discretizations respectively, while for $\vartheta = 1/2$ the scheme is a generalization of the second order in time Crank-Nicholson scheme. Note that the scheme is implicit except for the value $\vartheta = 0$. We may write (4.8) in the form (1.3) by setting

$$\begin{aligned} S(h, t, x, r, [u]_{t,x}) &= \sup_{\alpha \in \mathcal{A}} \left\{ \left[\frac{1}{\Delta t} - \vartheta c^\alpha + \vartheta \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta) \right] r \right. \\ &\quad - \left[\frac{1}{\Delta t} + (1 - \vartheta)c^\alpha - (1 - \vartheta) \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta) \right] [u]_{t,x}(-\Delta t, 0) \\ &\quad \left. - \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta) \left[\vartheta [u]_{t,x}(0, \beta\Delta x) + (1 - \vartheta)[u]_{t,x}(-\Delta t, \beta\Delta x) \right] \right\}, \end{aligned}$$

where $[u]_{t,x}(s, y) = u(t + s, x + y)$. Under assumption (4.2) the scheme (4.8) is monotone (i.e. satisfies (S1) or (S1')) provided the following CFL conditions hold

$$(4.9) \quad \Delta t (1 - \vartheta) \left(-c^\alpha(t, x) + \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta) \right) \leq 1,$$

$$(4.10) \quad \Delta t \vartheta \left(c^\alpha(t, x) - \sum_{\beta \in \mathcal{S}} C_h^\alpha(t, x, \beta) \right) \leq 1.$$

Furthermore, in view of (A1) and (4.7), Taylor expansion in (4.8) yields the following consistency result for smooth functions ϕ and $(t, x) \in \mathcal{G}_h^+$,

$$\begin{aligned} &|\phi_t + F(t, x, \phi, D\phi, D^2\phi) - S(h, t, x, \phi, [\phi]_{t,x})| \\ &\leq C(\Delta t |\phi_{tt}|_0 + \Delta x |D^2\phi|_0 + \Delta x^2 |D^4\phi|_0 + (1 - \vartheta)\Delta t (|D\phi_t|_0 + |D^2\phi_t|_0)). \end{aligned}$$

The $(1 - \vartheta)\Delta t$ -term is a non-standard term coming from the fact that we need the equation and the scheme to be satisfied in the *same* point, see assumption (S3). The necessity of this assumption follows from the proof of Theorem 3.1.

We have seen that if (4.2) and (4.7) hold along with the CFL conditions (4.9) and (4.10) then the scheme (4.8) satisfies assumptions (S1) – (S3) in Section 3. Theorem 3.1 therefore yields the following error bound:

Theorem 4.1. *Assume (A1), (A2), (4.2), (4.7), (4.9), (4.10) hold. If $u_h \in C_b(\mathcal{G}_h)$ is the solution of (4.8) and u is the solution of (1.1), then there is $C > 0$ such that in \mathcal{G}_h*

$$-e^{\mu t} |(u_0 - u_{0,h})^-|_0 - C|h|^{\frac{1}{5}} \leq u - u_h \leq e^{\mu t} |(u_0 - u_{0,h})^+|_0 + C|h|^{\frac{1}{2}},$$

where $|h| := \sqrt{\Delta x^2 + \Delta t}$.

Remark 4.1. Except when $\vartheta = 1$, the CFL condition (4.9) essentially implies that $\Delta t \leq C\Delta x^2$. Therefore Δt and Δx^2 play essentially the same role. Also note that the CFL condition (4.10) is satisfied if e.g. $\Delta t \leq (\sup_\alpha |(c^\alpha)^+|_0)^{-1}$.

Remark 4.2. Even though the above consistency relationship is not quite the “standard” one, it gives the correct asymptotic behavior of our scheme. First of all note that the new term, the $(1 - \vartheta)$ -term, behaves just like the Δt and Δx^2 terms. To see this, we note that according to (S3) we only need the above relation when ϕ is replaced by ϕ_ε defined in (S3). But for ϕ_ε we have $|\phi_{\varepsilon,tt}|_0 \approx |D^4\phi_\varepsilon|_0 \approx |D^2\phi_{\varepsilon,t}|_0 \approx \tilde{K}\varepsilon^{-3}$. By the CFL conditions (4.9) and (4.10) we have essentially that $\Delta x^2 \approx \Delta t$, so

$$\Delta t|\phi_{\varepsilon,tt}|_0 \approx \Delta x^2|D^4\phi_\varepsilon|_0 \approx \Delta t|D^2\phi_{\varepsilon,t}|_0 \approx \tilde{K}\Delta x^2\varepsilon^{-3}.$$

Next note that for $\vartheta = 1/2$ (the Crank-Nicholson case) the scheme is formally second order in time. However this is no longer the case for the monotone version. It is only first order in time due to the CFL condition which implies that $\Delta x^2\|D^4\phi\| = C\Delta t\|D^4\phi\|$.

Proof. In this case

$$\begin{aligned} E_1(\bar{K}, h, \varepsilon) &= E_2(\bar{K}, h, \varepsilon) \\ &= C(\Delta t\varepsilon^{-3} + \Delta x\varepsilon^{-1} + \Delta x^2\varepsilon^{-3} + (1 - \vartheta)\Delta t(\varepsilon^{-2} + \varepsilon^{-3})). \end{aligned}$$

So we have to minimize w.r.t. ε the following functions

$$\begin{aligned} \varepsilon + C(\Delta t\varepsilon^{-3} + \Delta x\varepsilon^{-1} + \Delta x^2\varepsilon^{-3}), \\ \varepsilon^{1/3} + C(\Delta t\varepsilon^{-3} + \Delta x\varepsilon^{-1} + \Delta x^2\varepsilon^{-3}). \end{aligned}$$

By minimizing separately in Δt and Δx , one finds that ε has to be like $\Delta t^{1/4}$ and $\Delta x^{1/2}$ in the first case, and that $\varepsilon^{1/3}$ has to be like $\Delta t^{1/10}$ and $\Delta x^{1/5}$ in the second case. The result now follows by taking $\varepsilon = \max(\Delta t^{1/4}, \Delta x^{1/2})$ in the first case and $\varepsilon^{1/3} = \max(\Delta t^{1/10}, \Delta x^{1/5})$ in the second case. \square

4.3. Remarks. For approximations of nonlinear equations monotonicity is a key property since it ensures (along with consistency) that the approximate solutions converge to the *correct* generalized solution of the problem (the viscosity solution in our case). This is not the case for nonmonotone methods, at least not in any generality.

However, the monotonicity requirement poses certain problems. Monotone schemes are low order schemes, and maybe more importantly, it is not always possible to find consistent monotone approximations for a given problem. To see the last point we note that in general the second derivative coefficient matrix a is only positive semidefinite, while the monotone schemes of Kushner and Bonnans/Zidani require the stronger assumptions (4.4) and (4.5) respectively. In fact, in Dong and Krylov [11] it was proved that if an operator L admits an approximation L_h of the form (4.1) which is of positive type, then a has to satisfy (4.5) (at least if a is bounded).

This is a problem in real applications, e.g. in finance, and it was this problem was the motivation behind the approximation of Bonnans and Zidani. First of all we note that their condition (4.5) is more general than (4.4) because any symmetric $N \times N$ matrix a can be decomposed as

$$a = \sum_{i=1}^N \sum_{j \neq i} (a_{ii} - |a_{ij}|) e_i e_i^T + \frac{a_{ij}^+}{2} (e_i + e_j)(e_i + e_j)^T + \frac{a_{ij}^-}{2} (e_i - e_j)(e_i - e_j)^T,$$

where the coefficients are nonnegative if and only if a is diagonal dominant. More importantly, it turns out that any symmetric positive semidefinite matrix can be approximated by a sequence of matrices satisfying (4.5). In Bonnans, Ottenwaelter,

and Zidani [4], this was proved in the case of symmetric 2×2 matrices along with an explicit error bound and an algorithm for computing the approximate matrices. Because of continuous dependence results for the equations, convergence of the coefficients immediately imply convergence of the solutions of the corresponding equations. Hence the Bonnans/Zidani approximation yields a way of approximating general problems where a is only positive semidefinite.

5. SEMIGROUP APPROXIMATIONS AND SPLITTING METHODS

In this section, we consider various approximations of semigroups obtained by a semi-discretization in time. In order to simplify the presentation we start by specializing Theorem 3.1 to the semigroup setting. To be precise we consider one-step in time approximations of (1.1) given by

$$(5.1) \quad \begin{aligned} u_h(t_n, x) &= S_h(t_n, t_{n-1})u_h(t_{n-1}, x) && \text{in } \mathbb{R}^N, \\ u_h(0, x) &= u_{h,0}(x) && \text{in } \mathbb{R}^N, \end{aligned}$$

where $t_0 = 0 < t_1 < \dots < t_n < \dots < t_{n_T} = T$, $h := \max_n(t_{n+1} - t_n)$, and the approximation semigroup S_h satisfies the following sub and superconsistency requirements: There exist a constant K_c , a subset I of $\mathbb{N} \times \mathbb{N}^N$, and constants $\gamma_\beta, \delta_\beta$ for $\beta \in I$ such that for any smooth functions ϕ ,

$$(5.2) \quad \begin{aligned} &\frac{1}{\Delta t} \left[S_h(t_n, t_{n-1}) - 1 \right] \phi(t_{n-1}, x) - F(t, x, \phi, D\phi, D^2\phi)_{t=t_n} \\ &\leq K_c \sum_{\beta \in I} |\partial_t^{\beta_0} D^{\beta'} \phi|_0^{\gamma_\beta} \Delta t^{\delta_\beta}, \end{aligned}$$

where $\beta = (\beta_0, \beta') \in I$ for $\beta_0 \in \mathbb{N}$ and $\beta' \in \mathbb{N}^N$, and in a similar way

$$(5.3) \quad \begin{aligned} &\frac{1}{\Delta t} \left[S_h(t_n, t_{n-1}) - 1 \right] \phi(t_{n-1}, x) - F(t, x, \phi, D\phi, D^2\phi)_{t=t_n} \\ &\geq -\bar{K}_c \sum_{\beta \in \bar{I}} |\partial_t^{\beta_0} D^{\beta'} \phi|_0^{\bar{\gamma}_\beta} \Delta t^{\bar{\delta}_\beta}, \end{aligned}$$

with corresponding data $\bar{K}_c, \bar{I}, \bar{\gamma}_\beta, \bar{\delta}_\beta$. We say that the semigroup is *monotone* if

$$\phi \leq \psi \quad \Rightarrow \quad S_h(t_n, t_{n-1})\phi \leq S_h(t_n, t_{n-1})\psi, \quad n = 1, \dots, n_T,$$

for all continuous bounded functions ϕ, ψ for which $S_h(t)\phi$ and $S_h(t)\psi$ are well defined.

We have the following corollary to Theorem 3.1.

Proposition 5.1. *Assume (A1), (A2), and that S_h is a monotone semigroup satisfying (5.2) and (5.3) and which is defined on a subset of $C_b(\mathbb{R}^N)$. If u is the solution of (1.1) and u_h is the solution of (5.1), then*

$$-C(|u_0 - u_{h,0}|_0 + \Delta t^{\frac{1}{10} \wedge r_1}) \leq u - u_h \leq C(|u_0 - u_{h,0}|_0 + \Delta t^{\frac{1}{4} \wedge r_2})$$

in \mathbb{R}^N , where

$$\begin{aligned} r_1 &:= \min_{\beta \in I} \left\{ \frac{\delta_\beta}{3(2\beta_0 + |\beta'| - 1)\gamma_\beta + 1} \right\}, \\ r_2 &:= \min_{\beta \in \bar{I}} \left\{ \frac{\bar{\delta}_\beta}{(2\beta_0 + |\beta'| - 1)\bar{\gamma}_\beta + 1} \right\}, \end{aligned}$$

where $|\beta'|$ denotes the sum of the components of β' .

Proof. We define

$$S(h, t_n, x, u_h, [u_h]_{t_n, x}) = \frac{1}{\Delta t} \left(u_h(t_n, x) - [u_h]_{t_n, x} \right),$$

where

$$[u_h]_{t_n, x} = S_h(t_{n-1}, t_n) u_h(t_{n-1}, x).$$

To apply Theorem 3.1, we just have to check that (S1) – (S3) hold and this is clear for (S1) and (S2) (see Remark 3.1). For (S3)(i), note that by (5.2) we have

$$\begin{aligned} & \phi_t + F(t_n, x, \phi, D\phi, D^2\phi) - S(h, t_n, x, \phi, [\phi]_{t_n, x}) \\ & \leq \frac{1}{2} |\partial_t^2 \phi|_0 \Delta t + K_c \sum_{\beta \in I} |\partial_t^{\beta_0} D^{\beta'} \phi|_0^{\gamma_\beta} \Delta t^{\delta_\beta}, \end{aligned}$$

which leads to

$$E_1(\bar{K}, h, \varepsilon) = \frac{1}{2} \bar{K} \varepsilon^{1-4} \Delta t + K_c \sum_{\beta \in I} (\bar{K} \varepsilon^{1-2\beta_0-|\beta'|})^{\gamma_\beta} \Delta t^{\delta_\beta}.$$

The upper bound now follows by optimizing with respect to ε as in the proof of Theorem 4.1. In a similar way we may use (5.3) to define E_2 and then conclude the lower bound. \square

Remark 5.1. In view of the consistency requirements (5.2) and (5.3), for schemes like (5.1) it is natural to think that only the x -variable is really playing a role and that one can get results on the rate of convergence by using this special “semi-group type” structure. More specifically, one might think that a different proof using a mollification of the solution with respect to the space variable only, can produce the estimates in an easier and maybe better way. We tried this strategy but we could not avoid using the short time expansion of the solution of the HJB Equation associated with smooth initial data (the short time expansion of the semi-group), and this leads to worse rates, even in cases where F is smooth. One way of understanding this – without justifying it completely – consists of looking at our estimates for the ϕ_{tt} -term (cf. (S3)(i) and (ii)). The present approach leads to an estimate of order ε^{-3} , while if we use the short time expansion, we are lead to a worse estimate of order ε^{-4} . We refer the reader to Subsection 5.1 and in particular to Lemma 5.6 below, where short time expansions for semi-groups are obtained and used to study the rate of convergence for splitting problems.

5.1. Semidiscrete splitting. We consider an equation of the form

$$(5.4) \quad u_t + F_1(D^2u) + F_2(D^2u) = 0 \quad \text{in } Q_T,$$

where

$$F_j(X) = \sup_{\alpha \in \mathcal{A}} \{-\text{tr}[a_j^\alpha X] - f_j^\alpha\}, \quad j = 1, 2,$$

and $a_j^\alpha \geq 0$ are matrices and f_j^α real numbers. We assume that they are both uniformly bounded in α and are independent of (t, x) . It follows that F_1 and F_2 are Lipschitz continuous and that (A1) is satisfied.

Let S denote the semigroup of (5.4), i.e. $S(\Delta t)\phi$ is the solution at time $t = \Delta t$ of (5.4) with initial value ϕ . Similarly, let S_1 and S_2 denote the semigroups associated with the equations $u_t + F_1(D^2u) = 0$ and $u_t + F_2(D^2u) = 0$.

We can define a semidiscrete splitting method by taking (5.1) with $t_n := n\Delta t$ and

$$(5.5) \quad S_h(t_{n-1}, t_n) = S_1(\Delta t)S_2(\Delta t).$$

Under the current assumptions all these semigroups map $W^{1,\infty}(\mathbb{R}^N)$ into itself, they are monotone, and they are nonexpansive,

$$|\bar{S}(t)\phi|_0 \leq |\phi|_0,$$

for $\phi \in W^{1,\infty}(\mathbb{R}^N)$ and where \bar{S} denotes one of the semigroups above.

As soon as we know the consistency relation for this scheme, we can find an error bound using Theorem 5.1. However, contrarily to the case of finite different schemes in the previous section, here the precise form of the consistency requirement is not well known. We are going to provide such results under different assumptions on F_1, F_2 . Our first result is the following:

Lemma 5.2. *Under the above assumptions, if in addition $|DF_1| \in W^{1,\infty}(\mathbb{S}^N)$ and $|DF_2| \in W^{3,\infty}(\mathbb{S}^N)$, then*

$$\begin{aligned} & -C(\Delta t|D^2\phi_t|_0 + \Delta t^2|D^3\phi|_0^4) - h.o.t. \\ & \leq \frac{1}{\Delta t}[S_h(t) - 1]\phi(t_{n-1}, x) + F_1(D^2\phi(t_n, x)) + F_2(D^2\phi(t_n, x)) \\ & \leq C(\Delta t|D^2\phi_t|_0 + \Delta t|D^3\phi|_0^2) + h.o.t. \end{aligned}$$

for all smooth functions ϕ , where “h.o.t.” stands for “higher order terms”.

Remark 5.2. Due to the convexity of the equation, in this example the upper and lower bounds are different.

Remark 5.3. We have only stated the principal error terms, the terms deciding the rate. The other terms are put in the “h.o.t.” category. Since the principal error terms need not be the lowest order terms (see the first inequality in Lemma 5.2), maybe a better name than “h.o.t.” would be the “less important terms”.

A direct consequence of Proposition 5.1 is the following result:

Corollary 5.3. *Let u_h denote the solution of (5.1) where S_h is defined in (5.5) and $u_{h,0} = u_0$, and let u be the solution of (5.4) with initial value u_0 . Under the assumptions of Lemma 5.2 we have*

$$-C\Delta t^{\frac{1}{13}} \leq u - u_h \leq C\Delta t^{\frac{2}{9}} \quad \text{in } \Delta t\{0, 1, 2, \dots, n_T\} \times \mathbb{R}^N.$$

Next, we give the result when F_1 and F_2 are assumed to be only Lipschitz continuous (which is the natural regularity assumption here). In this case the consistency relation is:

Lemma 5.4. *Under the above assumptions, if F_1 and F_2 are only Lipschitz continuous, we have*

$$\begin{aligned} & \left| \frac{1}{\Delta t}[S_h(t) - 1]\phi(t_{n-1}, x) + F_1(D^2\phi(t_n, x)) + F_2(D^2\phi(t_n, x)) \right| \\ & \leq C\Delta t|D^2\phi_t|_0 + C\Delta t^{\frac{1}{2}}|D^3\phi|_0 + h.o.t. \end{aligned}$$

for all smooth functions ϕ .

Again as a direct consequence of Proposition 5.1 we have the following error bound:

Corollary 5.5. *Under the assumptions of Corollary 5.3 but where F_1 and F_2 are only assumed to be Lipschitz continuous, we have*

$$-C\Delta t^{\frac{1}{14}} \leq u - u_h \leq C\Delta t^{\frac{1}{6}} \quad \text{in } \Delta t\{0, 1, 2, \dots, n_T\} \times \mathbb{R}^N.$$

Remark 5.4. We see a slight reduction of the rates in the Lipschitz case but not as important as one might have guessed. For first order equations these methods lead to the same rates in the smooth and Lipschitz cases.

Remark 5.5. If we change operators S_1, S_2 so that $S_1(t)\phi$ and $S_2(t)\phi$ denote the viscosity solutions of

$$\begin{aligned} u(x) &= \phi(x) - tF_1(D^2u(x)) \quad \text{in } \mathbb{R}^N, \\ u(x) &= \phi(x) - tF_2(D^2u(x)) \quad \text{in } \mathbb{R}^N, \end{aligned}$$

respectively, then the statements of Corollary 5.3 and 5.5 still hold.

In the proofs of Lemmas 5.2 and 5.4 we will use the following lemma:

Lemma 5.6. *Let \bar{S} be the semigroup associated to the equation*

$$u_t + \bar{F}(D^2u) = 0,$$

where \bar{F} is Lipschitz, convex, and non-increasing. Define \bar{F}_δ by

$$\bar{F}_\delta = \bar{F} * \bar{\rho}_\delta,$$

where $\bar{\rho}_\delta(X) = \delta^{-N} \bar{\rho}(X/\delta)$ and $\bar{\rho}$ is a smooth function on $\mathbb{S}(N)$ with mass one and support in $B(0, 1)$. Then for any smooth function ϕ ,

$$\bar{S}(t)\phi - \phi + t\bar{F}_\delta(D^2\phi) \leq t\delta|D\bar{F}|_0 + \frac{1}{2}t^2|D\bar{F}|_0|D\bar{F}_\delta|_0|D^4\phi|_0,$$

and

$$\bar{S}(t)\phi - \phi + t\bar{F}_\delta(D^2\phi) \geq -\frac{1}{2}t^2|D\bar{F}|_0(|D^2\bar{F}_\delta|_0|D^3\phi|_0^2 + |D\bar{F}_\delta|_0|D^4\phi|_0).$$

The proof of this result will be given after the proofs of Lemmas 5.2 and 5.4.

Proofs of Lemmas 5.2 and 5.4. In order to treat the two results at the same time, we mollify F_1 and F_2 and consider $F_{1,\delta}$ and $F_{2,\delta}$ (see Lemma 5.6 for the definitions). By Lemma 5.6 we have the following (small time) expansions:

$$(5.6) \quad S_j(t)\phi - \phi + tF_{j,\delta}(D^2\phi) \leq t\delta|DF_j|_0 + \frac{1}{2}t^2|DF_j|_0|DF_{j,\delta}|_0|D^4\phi|_0,$$

$$(5.7) \quad \begin{aligned} S_j(t)\phi - \phi + tF_{j,\delta}(D^2\phi) \\ \geq -\frac{1}{2}t^2|DF_j|_0(|D^2F_{j,\delta}|_0|D^3\phi|_0^2 + |DF_{j,\delta}|_0|D^4\phi|_0), \end{aligned}$$

for smooth functions ϕ and $j = 1, 2$.

Now we want to find an (small time) expansion for S_h . We write

$$\begin{aligned} S_h(t)\phi - \phi + t(F_1 + F_2)(D^2\phi) \\ = [S_1(t)S_2(t)\phi - S_1(t)(\phi - tF_{2,\delta}(D^2\phi))] \\ + [S_1(t)(\phi - tF_{2,\delta}(D^2\phi)) - \phi + tF_{1,\delta}(D^2\phi) + tF_{2,\delta}(D^2\phi)] \\ + t[(F_1 + F_2)(D^2\phi) - (F_{1,\delta} + F_{2,\delta})(D^2\phi)]. \end{aligned}$$

In view of the Lipschitz regularity and convexity of F_1 and F_2 , the last term on right hand side is between $-Ct\delta$ (Lipschitz regularity) and 0 (convexity), while the

first 2 terms can be estimated using non-expansiveness and small time expansions for S_1 and S_2 . The principal error term comes from the small time expansion for the term $S_1(t)(\phi - tF_{2,\delta}(D^2\phi))$. In view of (5.6) and (5.7),

$$t\delta|DF_1|_0 + \frac{1}{2}t^2|DF_1|_0|DF_{1,\delta}|_0|D^4\{\phi - tF_{2,\delta}(D^2\phi)\}|_0$$

is an upper bound on the principal error term, while

$$\frac{1}{2}t^2|DF_1|_0\left[|D^2F_{1,\delta}|_0|D^3\{\phi - tF_{2,\delta}(D^2\phi)\}|_0^2 + |DF_{1,\delta}|_0|D^4\{\phi - tF_{2,\delta}(D^2\phi)\}|_0\right]$$

is a lower bound. Expanding out these expressions keeping only the ‘‘worst terms’’ and bearing in mind the Lipschitz regularity of F_1 and F_2 , lead to the following upper and lower bounds respectively,

$$C(t\delta + t^2|D^4\phi|_0 + t^3|D^4F_{2,\delta}|_0|D^3\phi|_0^4) \quad \text{and} \\ C|D^2F_{1,\delta}|_0\left(t^2|D^3\phi|^2 + t^3|D^3F_{2,\delta}|_0|D^3\phi|_0^4 + t^4|D^3F_{2,\delta}|^2|D^3\phi|_0^6\right).$$

To conclude the proofs of the upper bounds in Lemmas 5.2 and 5.4, we note that

$$\frac{1}{\Delta t}[S_h(t) - 1]\phi(t_{n-1}, x) + F_1(D^2\phi(t_n, x)) + F_2(D^2\phi(t_n, x)) \\ \leq \Delta t(|DF_1|_0 + |DF_2|_0)|D^2\phi_t|_0 \\ + \left[\frac{1}{\Delta t}[S_h(t) - 1]\phi + F_1(D^2\phi) + F_2(D^2\phi)\right]_{(t_{n-1}, x)}.$$

In view of the above estimates the right hand side can be upper bounded by

$$(5.8) \quad C\left[\Delta t|D^2\phi_t|_0 + \delta + \Delta t|D^4\phi|_0 + \Delta t^2|D^4F_{2,\delta}|_0|D^3\phi|_0^4\right].$$

This proves the upper bound in Lemma 5.2 after sending $\delta \rightarrow 0$ while keeping in mind that in this case,

$$|D^n F_2^\delta|_0 \leq |D^n F_2|_0 < \infty \quad \text{and} \quad |D^m F_1^\delta|_0 \leq |D^m F_1|_0 < \infty$$

for $n = 1, \dots, 4$ and $m = 1, 2$. To get the upper bound in Lemma 5.4, we only need to note that in this case $|D^n F_j|_0 \leq C\delta^{1-\delta}$, $n \in \mathbb{N}$, $j = 1, 2$, and then minimize (5.8) w.r.t. δ .

The upper bounds follow in a similar way. We conclude the proof simply by giving the expression corresponding to (5.8),

$$C\left[\Delta t|D^2\phi_t|_0 + \delta \\ + |D^2F_{1,\delta}|_0\left(\Delta t|D^3\phi|^2 + \Delta t^2|D^3F_{2,\delta}|_0|D^3\phi|_0^4 + \Delta t^3|D^3F_{2,\delta}|^2|D^3\phi|_0^6\right)\right].$$

□

Proof of Lemma 5.6. Let

$$w = \phi - t\bar{F}_\delta(D^2\phi),$$

and observe that

$$(5.9) \quad w_t + \bar{F}(D^2w) = -\bar{F}_\delta(D^2\phi) + \bar{F}(D^2\phi) - \bar{F}(D^2\phi) + \bar{F}(D^2w).$$

Since \bar{F} is convex, it is easy to see that $\bar{F}_\delta(X) \geq \bar{F}(X)$, and hence

$$(5.10) \quad -|D\bar{F}|_0\delta \leq -\bar{F}_\delta(D^2\phi) + \bar{F}(D^2\phi) \leq 0.$$

The second difference, $-\bar{F}(D^2\phi) + \bar{F}(D^2w)$, can be written

$$\begin{aligned}
& \int_0^1 \frac{d}{ds} \{ \bar{F}(sD^2w + (1-s)D^2\phi) \} ds \\
&= \sum_{ij} \partial_i \partial_j (w - \phi) \int_0^1 (\partial_{X_{ij}} \bar{F})(sD^2w + (1-s)D^2\phi) ds \\
(5.11) \quad &= -t \sum_{ij} \partial_i \partial_j \{ \bar{F}_\delta(D^2\phi) \} \int_0^1 (\partial_{X_{ij}} \bar{F})(sD^2w + (1-s)D^2\phi) ds.
\end{aligned}$$

We expand $\partial_i \partial_j \{ \bar{F}_\delta(D^2\phi) \}$ and get

$$\begin{aligned}
& \sum_{klmn} (\partial_{X_{kl}} \partial_{X_{mn}} \bar{F}_\delta)(D^2\phi) (\partial_i \partial_k \partial_l \phi) (\partial_j \partial_m \partial_n \phi) \\
&+ \sum_{kl} (\partial_{X_{kl}} \bar{F}_\delta)(D^2\phi) (\partial_i \partial_j \partial_k \partial_l \phi).
\end{aligned}$$

We call the first term $M[\phi]_{ij}$.

Since $\partial_{X_{kl}} \partial_{X_{mn}} \bar{F}_\delta = \partial_{X_{mn}} \partial_{X_{kl}} \bar{F}_\delta$, it follows that M is symmetric,

$$M[\phi]_{ij} = M[\phi]_{ji}.$$

Moreover, since \bar{F} is convex, M is positive semidefinite: For every $\xi \in \mathbb{R}^N$

$$\begin{aligned}
& \sum_i M[\phi]_{ij} \xi_i \xi_j \\
&= \sum_{klmn} (\partial_{X_{kl}} \partial_{X_{mn}} \bar{F}_\delta)(D^2\phi) (\partial_k \partial_l (\sum_i \xi_i \partial_i \phi)) (\partial_m \partial_n (\sum_j \xi_j \partial_j \phi)) \\
&= \sum_{klmn} (\partial_{X_{kl}} \partial_{X_{mn}} \bar{F}_\delta)(D^2\phi) Y_{kl} Y_{mn} \geq 0,
\end{aligned}$$

where $Y_{ij} = \partial_i \partial_j (\sum_k \xi_k \partial_k \phi)$ and where the inequality follows by convexity of \bar{F} .

By the spectral theorem there exists $e^k \in \mathbb{R}^N$ and $\lambda_k \in \mathbb{R}$ for $k = 1, \dots, N$ (depending on ϕ) such that

$$M[\phi] = \sum_k \lambda_k e^k \otimes e^k.$$

Furthermore, since M is positive semidefinite, $\lambda_i \geq 0$ for $i = 1, \dots, N$. Therefore we have

$$\begin{aligned}
& \sum_{ij} M[\phi]_{ij} \int_0^1 (\partial_{X_{ij}} \bar{F})(sD^2w + (1-s)D^2\phi) ds \\
&= \sum_k \lambda_k \int_0^1 \sum_{ij} e_i^k e_j^k (\partial_{X_{ij}} \bar{F})(sD^2w + (1-s)D^2\phi) ds \leq 0,
\end{aligned}$$

where the inequality follows from the fact that \bar{F} is nonincreasing. We conclude that

$$\bar{F}(D^2w) - \bar{F}(D^2\phi) \geq -t |D\bar{F}|_0 |D^2\bar{F}_\delta|_0 |D^4\phi|_0,$$

and hence by (5.9) – (5.11) we get

$$w_t + \bar{F}(D^2w) \geq -|D\bar{F}|_0 \delta - t |D\bar{F}|_0 |D^2\bar{F}_\delta|_0 |D^4\phi|_0.$$

The first part of the Lemma now follows from the comparison principle.

The second part of the Lemma follows from (5.9) – (5.11) and the comparison principle after noting that this time, due to its sign, the $D^2\bar{F}_\delta$ term will be part of the error expression. \square

5.2. Piecewise constant controls. Here we study approximations by piecewise constant controls. Such approximations have been studied e.g. in [25, 22] (see also the references therein). We consider the following simplified version of equation (1.1),

$$(5.12) \quad u_t + \max_i \{-L^i u - f^i(x)\} = 0 \quad \text{in } Q_T,$$

where

$$L^i \phi = \text{tr}[\sigma^i(x)\sigma^{iT}(x)D^2\phi] + b^i(x)D\phi + c^i(x)\phi,$$

and σ, b, c and f satisfy assumption (A1) when α is replaced by i . Note that the coefficients are independent of time. We approximate (5.12) in the following way,

$$(5.13) \quad u^{n+1}(x) = \min_i S_i(\Delta t)u^n(x) \quad \text{in } \{0, 1, \dots, n_T\} \times \mathbb{R}^N,$$

where $S_i(t)\phi(x)$ denotes the solution at (t, x) of the linear equation

$$(5.14) \quad u_t - L^i u - f^i(x) = 0$$

with initial data ϕ at time $t = 0$. As usual, u^n is expected to be an approximation of $u(t_n, x)$, $t_n := n\Delta t$, and we are looking for a bound on the approximation error.

Under assumption (A1) the comparison principle holds for the linear equations (5.14), hence S_i and $\min_i S_i$ are monotone. Furthermore, we have the following consistency relation:

Lemma 5.7. *If (A1) holds, then for any smooth function ϕ we have*

$$\begin{aligned} & \left| \frac{1}{\Delta t} [\min_i S_i(\Delta t) - 1] \phi(t_{n-1}, x) + \Delta t \max_i \{-L^i \phi(t_n, x) - f^i(x)\} \right| \\ & \leq C\Delta t |D^2\phi_t|_0 + C\Delta t \left(\sum_{n=0}^4 |D^n \phi|_0 + 1 \right) + C\Delta t^{1/2} \left(\sum_{n=0}^2 |D^n \phi|_0 + 1 \right). \end{aligned}$$

We have the following error bound:

Proposition 5.8. *Assume (A1). Let u_h denote the solution of (5.1) corresponding to*

$$S_h(t_{n-1}, t_n) = \min_i S_i(\Delta t)$$

and $u_{h,0} = u_0$, and let u be the solution of (5.12) with initial value u_0 . Then

$$-C\Delta t^{\frac{1}{10}} \leq u - u_h \leq 0 \quad \text{in } \Delta t\{0, 1, 2, \dots, n_T\} \times \mathbb{R}^N.$$

Proof. We first observe that $u_h \geq u$ in Q_T . This can be easily seen from the control interpretation of u_h (which we have not provided!) or from the comparison principle since u_h is a supersolution of (5.12) (solutions of (5.14) are supersolutions of (5.12) and so is the min of such solutions). The other bound follows from Lemma 5.7 and Proposition 5.1. \square

Remark 5.6. Assuming more regularity on the coefficients does not lead to any improvement of the bound. The principal contribution to the error comes from the $|D^4\phi|_0$ -term, and this term does not depend on the regularity of the coefficients (only on the L^∞ norm of σ).

Remark 5.7. In [22] Krylov obtains a better rate, namely $1/6$. His approach is different for ours, he works on the dynamic programming principle directly using control techniques.

Proof of Lemma 5.7. Let $\sigma_\delta^i = \sigma^i * \rho_\delta$, and define similarly b_δ^i , c_δ^i , and f_δ^i , and let L_δ^i be the operator L^i corresponding to $\sigma_\delta^i, b_\delta^i, c_\delta^i$. Observe that

$$\begin{aligned} & |\min_i S_i(t)\phi - \phi + t \max_i \{-L_\delta^i \phi - f_\delta^i(x)\}| \\ &= |\min_i (S_i(t)\phi - \phi) - t \min_i \{L_\delta^i \phi + f_\delta^i(x)\}| \\ &\leq \max_i |S_i(t)\phi - \phi + t(-L_\delta^i \phi - f_\delta^i(x))|. \end{aligned}$$

Next, define

$$w^\pm = \phi - t(-L_\delta^i \phi - f_\delta^i(x)) \pm \frac{1}{2}t^2 |L^i L_\delta^i \phi - L^i f_\delta^i|_0 \pm t|(L_\delta^i - L^i)\phi - (f_\delta^i - f^i)|_0,$$

and observe that w^+ is a supersolution of (5.14) while w^- is a subsolution. By the comparison principle and properties of mollifiers we get

$$\begin{aligned} & |S_i(t)\phi - \phi + t(-L_\delta^i \phi - f_\delta^i(x))| \\ &\leq \frac{1}{2}t^2 |L^i L_\delta^i \phi - L^i f_\delta^i|_0 + t\delta C \left(\sum_{n=0}^2 |D^n \phi|_0 + 1 \right). \end{aligned}$$

Furthermore, by properties of mollifiers and the Lipschitz regularity of the coefficients we see that

$$|L^i L_\delta^i \phi + L_i f_i|_0 \leq C \left(\sum_{n=0}^4 |D^n \phi|_0 + \delta^{-1} \sum_{n=0}^2 |D^n \phi|_0 + \delta^{-1} + 1 \right).$$

By combining the above estimates we get

$$\begin{aligned} & |\min_i S_i(t)\phi - \phi + t \max_i \{-L^i \phi - f^i(x)\}| \\ &\leq Ct^2 \left(\sum_{n=0}^4 |D^n \phi|_0 + \delta^{-1} \sum_{n=0}^2 |D^n \phi|_0 + \delta^{-1} + 1 \right) + t\delta C \left(\sum_{n=0}^2 |D^n \phi|_0 + 1 \right), \end{aligned}$$

and the result follows by similar arguments as was given in the proofs of Lemmas 5.2 and 5.4 after optimizing w.r.t. δ . \square

6. REMARKS ON THE HÖLDER CONTINUOUS CASE

In this section we give an extension of the main result Theorem 3.1 to the case when solutions of (1.1) do no longer belong to the space $\mathcal{C}^{0,1}$ but rather belong to the bigger space \mathcal{C}^β for some $\beta \in (0, 1)$.

In the time-dependent case \mathcal{C}^β regularity of the solution is observed typically when assumption (A1) is relaxed in the following way:

(A1') For any $\alpha \in \mathcal{A}$, $a^\alpha = \frac{1}{2}\sigma^\alpha \sigma^{\alpha T}$ for some $N \times P$ matrix σ^α . Moreover, there is a constant K independent of α such that

$$|u_0|_\beta + |\sigma^\alpha|_1 + |b^\alpha|_1 + |c^\alpha|_\beta + |f^\alpha|_\beta \leq K.$$

In other words u_0, c^α, f^α now belongs to \mathcal{C}^β .

Lemma 6.1. *If (A1') holds, then there exists a unique solution $u \in \mathcal{C}^{0,\beta}(\overline{Q}_T)$ of (1.1) and (1.2).*

This standard result is proved e.g. in [19]. We claim that under (A1'), we have the same regularity (the same β) for all equations considered in this paper. We skip the proof of this claim. In the rest of this section, the solutions of the different equations belong to $\mathcal{C}^{0,\beta}(\overline{Q}_T)$ with the same fixed $\beta \in (0, 1]$.

Lower than $\mathcal{C}^{0,1}$ regularity of solutions implies lower convergence rates than obtained in Sections 2 – 5. We will now state the Hölder versions of some these results without proofs. The proofs are not much different from the proofs given above, and moreover, the Hölder case was extensively studied in [1]. We start by the convergence rate for the switching system approximation of Section 2.

Proposition 6.2. *Assume (A1'). If \bar{u} and v are the solutions of (2.2) and (2.1) in $\mathcal{C}^{0,\beta}(\overline{Q}_T)$, then for k small enough,*

$$0 \leq v_i - \bar{u} \leq Ck^{\frac{\beta}{2+\beta}} \quad \text{in } \overline{Q}_T, \quad i \in \mathcal{I},$$

where C only depends on T and K from (A1').

In order to state a \mathcal{C}^β version of Theorem 3.1 we need to modify assumption (S3). The requirement on ϕ_ε should be changed to

$$|\partial_t^{\beta_0} D^{\beta'} \phi_\varepsilon(x, t)| \leq \tilde{K} \varepsilon^{\beta - 2\beta_0 - |\beta'|} \quad \text{in } \overline{Q}_T, \quad \text{for any } \beta_0 \in \mathbb{N}, \beta' \in \mathbb{N}^N.$$

We will denote the modified assumption by (S3'). Now we state the \mathcal{C}^β version of our main result, Theorem 3.1.

Theorem 6.3. *Assume (A1'), (S1), (S2) and that (1.3) has a unique solution $u_h \in C_b(\mathcal{G}_h)$. Let u denote the solution of (1.1)-(1.2), and let h be sufficiently small.*

(a) **(Upper bound)** *If (S3')(i) holds, then there exists a constant C depending only μ, K in (S1), (A1') such that*

$$u - u_h \leq e^{\mu t} |(u_0 - u_{0,h})^+|_0 + C \min_{\varepsilon > 0} \left(\varepsilon^\beta + E_1(\tilde{K}, h, \varepsilon) \right) \quad \text{in } \mathcal{G}_h,$$

where $\tilde{K} = |u|_1$.

(b) **(Lower bound)** *If (S3')(ii) and (A2) holds, then there exists a constant C depending only μ, K in (S1), (A1') such that*

$$u - u_h \geq -e^{\mu t} |(u_0 - u_{0,h})^-|_0 - C \min_{\varepsilon > 0} \left(\varepsilon^{\frac{\beta^2}{2+\beta}} + E_2(\tilde{K}, h, \varepsilon) \right) \quad \text{in } \mathcal{G}_h,$$

where $\tilde{K} = |u|_1$.

Remark 6.1. For the FDMs described in Section 4 we get an upper rate of $\frac{\beta}{2}$ and a lower rate of $\frac{2\beta^2}{4(2+\beta)-2\beta}$ in the \mathcal{C}^β case. Compare with Theorem 4.1.

APPENDIX A. WELL-POSEDNESS, REGULARITY, AND CONTINUOUS DEPENDENCE FOR SWITCHING SYSTEMS

In this section we give well-posedness, regularity, and continuous dependence results for solutions of a very general switching system that has as special cases the scalar HJB equations (1.1), and the switching systems (2.1), (2.3), (3.3).

We consider the following system:

$$(A.1) \quad F_i(x, u, \partial_t u_i, Du_i, D^2 u_i) = 0 \quad \text{in } Q_T, \quad i \in \mathcal{I} := \{1, \dots, M\},$$

with

$$F_i(t, x, r, p_t, p_x, X) = \max \left\{ p_t + \sup_{\alpha \in \mathcal{A}} \inf_{\beta \in \mathcal{B}} \mathcal{L}_i^{\alpha, \beta}(x, r_i, p_x, X); r_i - \mathcal{M}_i r \right\},$$

$$\mathcal{L}_i^{\alpha, \beta}(t, x, s, q, X) = -\text{tr}[a_i^{\alpha, \beta}(t, x)X] - b_i^{\alpha, \beta}(t, x)q - c_i^{\alpha, \beta}(t, x)s - f_i^{\alpha, \beta}(t, x),$$

where \mathcal{M} is defined below (2.1), \mathcal{A}, \mathcal{B} are compact metric spaces, r is a vector $r = (r_1, \dots, r_M)$, and $k > 0$ is a constant (the switching cost). See [14, 7, 28, 17, 16] for more information about such systems.

We make the following assumption:

(A) For any α, β, i , $a_i^{\alpha, \beta} = \frac{1}{2} \sigma_i^{\alpha, \beta} \sigma_i^{\alpha, \beta T}$ for some $N \times P$ matrix $\sigma_i^{\alpha, \beta}$. Furthermore, there is a constant C independent of i, α, β, t , such that

$$|\sigma_i^{\alpha, \beta}(t, \cdot)|_1 + |b_i^{\alpha, \beta}(t, \cdot)|_1 + |c_i^{\alpha, \beta}(t, \cdot)|_1 + |f_i^{\alpha, \beta}(t, \cdot)|_1 \leq \bar{C}.$$

We start by comparison, existence, uniqueness, and L^∞ bounds on the solution and its gradient. Before stating the results, we define $USC(\bar{Q}_T; \mathbb{R}^M)$ and $LSC(\bar{Q}_T; \mathbb{R}^M)$ to be the spaces of upper and lower semi-continuous functions from \bar{Q}_T into \mathbb{R}^M respectively.

Theorem A.1. *Assume (A) holds.*

(i) *If $u \in USC(\bar{Q}_T; \mathbb{R}^M)$ is a subsolution of (A.1) bounded above and $v \in LSC(\bar{Q}_T; \mathbb{R}^M)$ supersolution of (A.1) bounded below, then $u \leq v$ in \bar{Q}_T .*

(ii) *There exists a unique bounded continuous solution u of (A.1).*

(iii) *The solution u of (A.1) belongs to $C^{0,1}(\bar{Q}_T)$, and satisfies for all $t, s \in [0, T]$*

$$e^{-\lambda t} \max_i |u_i(t, \cdot)|_0 \leq \max_i |u_{0,i}|_0 + t \sup_{i, \alpha, \beta} |f_i^{\alpha, \beta}|_0,$$

where $\lambda := \sup_{i, \alpha, \beta} |c_i^{\alpha, \beta+}|_0$,

$$e^{\lambda_0 t} \max_i [u_i(t, \cdot)]_1 \leq \max_i [u_{0,i}]_1 + t \sup_{i, \alpha, \beta, s} \left\{ |u^i|_0 [c_i^{\alpha, \beta}(s, \cdot)]_1 + [f_i^{\alpha, \beta}(s, \cdot)]_1 \right\},$$

where $\lambda_0 := \sup_{i, \alpha, \beta, s} \{ |c_i^{\alpha, \beta+}(s, \cdot)|_0 + [\sigma_i^{\alpha, \beta}(s, \cdot)]_1^2 + [b_i^{\alpha, \beta}(s, \cdot)]_1 \}$, and

$$\max_i |u_i(t, x) - u_i(s, x)| \leq C |t - s|^{1/2},$$

where $C \leq 8M\bar{C} + \sqrt{T}\bar{C}(2M + 1)$ and $M := \sup_{i,t} |u_i(t, \cdot)|_1$.

Before giving the proof we state a key technical lemma.

Lemma A.2. *Let $u \in USC(\bar{Q}_T; \mathbb{R}^M)$ be a bounded above subsolution of (A.1) and $\bar{u} \in LSC(\bar{Q}_T; \mathbb{R}^M)$ be a bounded below supersolution of an other equation (A.1) where the functions $\mathcal{L}_i^{\alpha, \beta}$ are replaced by functions $\bar{\mathcal{L}}_i^{\alpha, \beta}$ satisfying the same assumptions. Let $\phi : [0, T] \times \mathbb{R}^{2N} \rightarrow \mathbb{R}$ be a smooth function bounded from below. We denote by*

$$\psi_i(t, x, y) = u_i(t, x) - \bar{u}_i(t, y) - \phi(t, x, y),$$

and $M = \sup_{i,t,x,y} \psi_i(t, x, y)$. *If there exists a maximum point for M , i.e. a point (i', t_0, x_0, y_0) such that $\psi_{i'}(t_0, x_0, y_0) = M$, then there exists $i_0 \in \mathcal{I}$ such that (i_0, t_0, x_0, y_0) is also a maximum point for M , and, in addition $\bar{u}_{i_0}(t_0, y_0) < \mathcal{M}_{i_0} \bar{u}(t_0, y_0)$.*

Loosely speaking this lemma means that whenever we do doubling of variables for systems of the type (A.1), we can ignore the $u_i - \mathcal{M}_i u$ parts of the equations. So we are more or less back in the scalar case with equations $\partial_t u_{i_0} + \sup_\alpha \inf_\beta \mathcal{L}_{i_0}^{\alpha, \beta} [u_{i_0}] \leq 0$ and $\partial_t \bar{u}_{i_0} + \sup_\alpha \inf_\beta \bar{\mathcal{L}}_{i_0}^{\alpha, \beta} [\bar{u}_{i_0}] \geq 0$. We skip the proof since it is similar to the proof given in [2] for the stationary case.

Proof of Theorem A.1. Comparison, uniqueness, and existence is proved in [17] for the stationary Dirichlet problem for (1.1) on a bounded domain under similar assumptions on the data. To extend the comparison result to a time dependent problem in an unbounded domain, we only need to modify the test function used in [17] in the standard way. (See also the arguments given below). Comparison implies uniqueness, and existence follows from Perron's method. This last argument is similar to the argument given in [17], but easier since we have no boundary conditions other than the initial condition.

Let

$$w(t) := e^{\lambda t} \left\{ \max_i |u_{0,i}|_0 + t \sup_{i, \alpha, \beta} |f_i^{\alpha, \beta}|_0 \right\},$$

then the bound on $|u|_0$ follows from the comparison principle after checking that w ($-w$) is a supersolution (subsolution) of (A.1).

To get the bound on the gradient of u , consider

$$m := \sup_{i, t, x, y \in \mathbb{R}^N} \{u_i(t, x) - u_i(t, y) - \bar{w}(t)|x - y|\},$$

where

$$\bar{w}(t) := e^{\lambda_0 t} \left\{ \max_i [u_{0,i}]_1 + t \sup_{i, \alpha, \beta, s} \left\{ |u_i|_0 [C_i^{\alpha, \beta}(s, \cdot)]_1 + [f_i^{\alpha, \beta}(s, \cdot)]_1 \right\} \right\}.$$

We are done if we can prove that $m \leq 0$. Assume this is not the case, $m > 0$, and for simplicity that this maximum is attained in $\bar{t}, \bar{x}, \bar{y}$. Then there exists a $k > 0$ such that

$$u_i(\bar{t}, \bar{x}) - u_i(\bar{t}, \bar{y}) - \bar{w}(\bar{t})|\bar{x} - \bar{y}| - \bar{t}e^{\lambda_0 \bar{t}}k > 0, \quad i \in \mathcal{I}.$$

Let $\psi_i(t, x, y) := u_i(t, x) - u_i(t, y) - \bar{w}(t)|x - y| - te^{\lambda_0 t}k$, then ψ also has maximum $M > 0$ at some point $(\tilde{t}, \tilde{x}, \tilde{y})$. Since $M > 0$, $\tilde{x} \neq \tilde{y}$ and $\tilde{t} > 0$. Therefore $\bar{w}(t)|x - y| + te^{\lambda_0 t}k$ is a smooth function at $(\tilde{t}, \tilde{x}, \tilde{y})$ and a standard argument using the viscosity sub- and supersolution inequalities for (A.1) at $(\tilde{t}, \tilde{x}, \tilde{y})$ and Lemma A.2 leads to $k \leq 0$. See the proof of Theorem A.3 for a similar argument. This is a contradiction and hence $m \leq 0$.

In the general case when the maximum m need not be attained at some finite point, we must modify the test function in the standard way. We skip the details.

To get the time regularity result, assume that $s < t$ and let u^ε be the solution of (A.1) in $t \in (s, T]$ starting from $u(s, \cdot) * \rho_\varepsilon(x) =: u_0^\varepsilon(x)$. By the comparison principle

$$|u - u^\varepsilon| \leq \sup_{r \in [s, T]} [u(r, \cdot)]_1 \varepsilon \quad \text{in } [s, T] \times \mathbb{R}^N,$$

and easy computations show that

$$w_\varepsilon^\pm(t, x) = e^{\lambda t} \left\{ u_0^\varepsilon(x) \pm (t - s)C_\varepsilon \right\}$$

are subsolution (w^-) and supersolution (w^+) of (A.1) if

$$C_\varepsilon = \bar{C}^2 |D^2 u_0^\varepsilon|_0 + \bar{C} (|Du_0^\varepsilon|_0 + |u_0^\varepsilon|_0 + 1)$$

and \bar{C} is given by (A). Another application of the comparison principle then yields

$$w_\varepsilon^- \leq u^\varepsilon \leq w_\varepsilon^+ \quad \text{in } [s, T] \times \mathbb{R}^N.$$

The result now follows from

$$\begin{aligned} & |u(t, x) - u(s, x)| \\ & \leq |u(t, x) - u^\varepsilon(t, x)| + |u^\varepsilon(t, x) - u_0^\varepsilon(x)| + |u_0^\varepsilon(x) - u(s, x)| \\ & \leq ([u(t, \cdot)]_1 + [u(s, \cdot)]_1)\varepsilon + |t - s|C_\varepsilon, \end{aligned}$$

and a minimization in ε after noting that $C_\varepsilon \leq C(\varepsilon^{-1} + 1)$. \square

We proceed to obtain continuous dependence on the coefficients.

Theorem A.3. *Let u and \bar{u} be solutions of (A.1) with coefficients σ, b, c, f and $\bar{\sigma}, \bar{b}, \bar{c}, \bar{f}$ respectively. If both sets of coefficients satisfy (A1), and $|u|_0 + |\bar{u}|_0 + [u(t, \cdot)]_1 + [\bar{u}(t, \cdot)]_1 \leq M < \infty$ for $t \in [0, T]$, then*

$$\begin{aligned} & e^{-\lambda t} \max_i |u_i(t, \cdot) - \bar{u}_i(t, \cdot)|_0 \leq \max_i |u_i(0, \cdot) - \bar{u}_i(0, \cdot)|_0 \\ & + t^{1/2} K \sup_{i, \alpha, \beta} |\sigma - \bar{\sigma}|_0 + t \sup_{i, \alpha, \beta} \left\{ 2M|b - \bar{b}|_0 + M|c - \bar{c}|_0 + |f - \bar{f}|_0 \right\}, \end{aligned}$$

where $\lambda := \sup_{i, \alpha, \beta} |c^-|_0$ and

$$\begin{aligned} K^2 & \leq 8M^2 + 8MT \sup_{i, \alpha, \beta} \left\{ 2M[\sigma]_1^2 \wedge [\bar{\sigma}]_1^2 \right. \\ & \left. + 2M[b]_1 \wedge [\bar{b}]_1 + M[c]_1 \vee [\bar{c}]_1 + [f]_1 \wedge [\bar{f}]_1 \right\}. \end{aligned}$$

Proof. We only indicate the proof in the case $\lambda = 0$. Define

$$\begin{aligned} \psi^i(t, x, y) & := u_i(t, x) - \bar{u}_i(t, y) - \frac{1}{\delta}|x - y|^2 - \varepsilon(|x|^2 + |y|^2), \\ m & := \sup_{i, t, x, y} \psi^i(t, x, y) - \sup_{i, x, y} (\psi^i(0, x, y))^+, \\ \bar{m} & := \sup_{i, t, x, y} \left\{ \psi^i(t, x, y) - \frac{\sigma m t}{T} \right\}, \end{aligned}$$

where $\sigma \in (0, 1)$. We assume $m > 0$ since otherwise we are done. We will now derive an upper bound on m . To do this we consider \bar{m} . By the assumptions this supremum is attained at some point (i_0, t_0, x_0, y_0) . Since $m > 0$ it follows that $\bar{m} > 0$ and $t_0 > 0$, and by Lemma A.2, the index i_0 may be chosen so that $\bar{u}_{i_0}(t_0, y_0) < \mathcal{M}_{i_0} \bar{u}(t_0, y_0)$. With this in mind, the maximum principle for semi continuous functions [8, 9] and the definition of viscosity solutions imply the following inequality:

$$p_t - \bar{p}_t + \sup_\alpha \inf_\beta \mathcal{L}_{i_0}^{\alpha, \beta}(t_0, x_0, u_{i_0}, p_x, X) - \sup_\alpha \inf_\beta \bar{\mathcal{L}}_{i_0}^{\alpha, \beta}(t_0, y_0, \bar{u}_{i_0}, p_y, Y) \leq 0,$$

where $(p_t, p_x, X) \in \bar{\mathcal{P}}^{2,+} u_{i_0}(x_0)$ and $(\bar{p}_t, p_y, Y) \in \bar{\mathcal{P}}^{2,-} \bar{u}_{i_0}(y_0)$ (see [8, 9] for the notation). Furthermore $p_t - \bar{p}_t = \frac{\sigma m}{T}$, $p_x = \frac{2}{\delta}(x_0 - y_0) + 2\varepsilon x_0$, $p_y = \frac{2}{\delta}(x_0 - y_0) - 2\varepsilon y_0$, and

$$\begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix} \leq \frac{2}{\delta} \begin{pmatrix} I & -I \\ -I & I \end{pmatrix} + 2\varepsilon \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} + \mathcal{O}(\kappa),$$

for some $\kappa > 0$. In the end we will fix σ , δ , and ε and send $\kappa \rightarrow 0$, so we simply ignore the $\mathcal{O}(\kappa)$ -term in the following. The first inequality implies

$$\begin{aligned} \frac{\sigma m}{T} \leq \sup_{i,\alpha,\beta} \left\{ -\operatorname{tr}[\bar{a}(t_0, y_0)Y] + \operatorname{tr}[a(t_0, x_0)X] + \bar{b}(t_0, y_0)p_x - b(t_0, x_0)p_y \right. \\ \left. + \bar{c}(t_0, y_0)\bar{u}(t_0, y_0) - c(t_0, x_0)u(t_0, x_0) + \bar{f}(t_0, y_0) + f(t_0, x_0) \right\}, \end{aligned}$$

Note that Lipschitz regularity of the solutions and a standard argument yields

$$|x_0 - y_0| \leq \delta M.$$

So using Ishii's trick on the 2nd order terms [15, pp. 33,34], and a few other manipulations, we get

$$\begin{aligned} \frac{\sigma m}{T} \leq \sup_{i,\alpha,\beta} \left\{ \frac{2}{\delta} |\sigma(t_0, x_0) - \bar{\sigma}(t_0, y_0)|^2 + 2M |b(t_0, x_0) - \bar{b}(t_0, y_0)| \right. \\ \left. + C\varepsilon(1 + |x_0|^2 + |y_0|^2) \right. \\ \left. + M |c(t_0, x_0) - \bar{c}(t_0, y_0)| + |f(t_0, x_0) - \bar{f}(t_0, y_0)| \right\}. \end{aligned}$$

Some more work leads to an estimate for m depending on T , σ , δ , and ε , and using the definition of m and estimates on $\sup_{i,x,y} \psi_i(0, x, y)$, we obtain a similar upper bound for $u - \bar{u}$. We finish the proof of the upper bound on $u - \bar{u}$ by sending $\sigma \rightarrow 1$, minimizing this expression w.r.t. δ , sending $\varepsilon \rightarrow 0$, and noting that the result still holds if we replace T by any $t \in [0, T]$. The lower bound follows in a similar fashion. \square

Remark A.1. For more details on such manipulations, we refer to [19].

REFERENCES

- [1] G. Barles and E. R. Jakobsen. On the convergence rate of approximation schemes for Hamilton-Jacobi-Bellman equations. *M2AN Math. Model. Numer. Anal.* 36(1):33–54, 2002.
- [2] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for Hamilton-Jacobi-Bellman equations. To appear in *SIAM J. Numer. Anal.*
- [3] G. Barles and P. E. Souganidis. Convergence of approximation schemes for fully nonlinear second order equations. *Asymptotic Anal.* 4(3):271–283, 1991.
- [4] F. Bonnans, E. Ottenwaelter, and H. Zidani. A fast algorithm for the two dimensional HJB equation of stochastic control. *M2AN Math. Model. Numer. Anal.* 38(4):723–735, 2004.
- [5] F. Bonnans and H. Zidani. Consistency of generalized finite difference schemes for the stochastic HJB equation. *SIAM J. Numer. Anal.* 41(3):1008–1021, 2003.
- [6] F. Camilli and M. Falcone. An approximation scheme for the optimal control of diffusion processes. *RAIRO Modél. Math. Anal. Numér.* 29(1): 97–122, 1995.
- [7] I. Capuzzo-Dolcetta and L. C. Evans. Optimal switching for ordinary differential equations. *SIAM J. Control Optim.* 22(1):143–161, 1984.
- [8] M. G. Crandall and H. Ishii. The maximum principle for semicontinuous functions. *Differential Integral Equations*, 3(6):1001–1014, 1990.
- [9] M. G. Crandall, H. Ishii, and P.-L. Lions. User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [10] M. G. Crandall and P.-L. Lions. Two approximations of solutions of Hamilton-Jacobi equations. *Math. Comp.* 43(167):1–19, 1984.
- [11] H. Dong and N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman equations with constant coefficients. To appear in *St. Petersburg Math. J.*
- [12] H. Dong and N. V. Krylov. On the Rate of Convergence of Finite-difference Approximations for Parabolic Equations with C^1 and C^2 Coefficients. Preprint.
- [13] H. J. Kushner and P. Dupuis. *Numerical methods for stochastic control problems in continuous time*. Springer-Verlag, New York, 2001.

- [14] L. C. Evans and A. Friedman. Optimal stochastic switching and the Dirichlet problem for the Bellman equation. *Trans. Amer. Math. Soc.* 253:365-389, 1979.
- [15] H. Ishii. On uniqueness and existence of viscosity solutions of fully nonlinear second-order elliptic PDEs. *Comm. Pure Appl. Math.*, 42(1):15-45, 1989.
- [16] H. Ishii and S. Koike. Viscosity solutions for monotone systems of second-order elliptic PDEs. *Comm. Partial Differential Equations* 16(6-7):1095-1128, 1991.
- [17] H. Ishii and S. Koike. Viscosity solutions of a system of nonlinear second-order elliptic PDEs arising in switching games. *Funkcial. Ekvac.*, 34:143-155, 1991.
- [18] E. R. Jakobsen. On the rate of convergence of approximation schemes for Bellman equations associated with optimal stopping time problems. *Math. Models Methods Appl. Sci. (M3AS)* 13(5):613-644, 2003.
- [19] E. R. Jakobsen and K. H. Karlsen. Continuous dependence estimates for viscosity solutions of fully nonlinear degenerate parabolic equations. *J. Differential Equations* 183:497-525, 2002.
- [20] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman's equations. *St. Petersburg Math. J.*, 9(3):639-650, 1997.
- [21] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman's equations with variable coefficients. *Probab. Theory Relat. Fields*, 117:1-16, 2000.
- [22] N. V. Krylov. Approximating value functions for controlled degenerate diffusion processes by using piece-wise constant policies. *Electron. J. Probab.* 4(2), 1999.
- [23] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman equations with Lipschitz coefficients. To appear.
- [24] P.-L. Lions. Personal communication.
- [25] P.-L. Lions and B. Mercier. Approximation numérique des équations de Hamilton-Jacobi-Bellman. *RAIRO Anal. Numér.* 14(4):369-393, 1980.
- [26] C.-W. Shu. Total-variation-diminishing time discretizations. *SIAM J. Sci. Statist. Comput.* 9(6):1073-1084, 1988.
- [27] P. E. Souganidis. Approximation schemes for viscosity solutions of Hamilton-Jacobi equations. *J. Differential Equations* 59(1):1-43, 1985.
- [28] N. Yamada. Viscosity solutions for a system of elliptic inequalities with bilateral obstacles. *Funkcial. Ekvac.* 30(2-3):417-425, 1987.

(Guy Barles)

LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE
 UNIVERSITY OF TOURS
 PARC DE GRANDMONT
 37200 TOURS, FRANCE

E-mail address: `barles@lmpt.univ-tours.fr`
URL: `http://www.phys.univ-tours.fr/~barles`

(Espen R. Jakobsen)

DEPARTMENT OF MATHEMATICAL SCIENCES
 NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY
 N-7491 TRONDHEIM, NORWAY

E-mail address: `erj@math.ntnu.no`
URL: `http://www.math.ntnu.no/~erj`