

Critical revision of the manuscript for important intellectual content: Gaston, Newcomb, Sharifi, Zein.

Statistical analysis: Gaston, Marozkina, Zein.

Obtained funding: Gaston.

Administrative, technical, or material support: Gaston, Marozkina, Newcomb.

Supervision: Gaston, Newcomb, Sharifi.

Conflict of Interest Disclosures: Dr Gaston reported grants from the National Heart, Lung, and Blood Institute during the conduct of the study. Dr Newcomb reported grants from the National Institutes of Health during the conduct of the study. Dr Sharifi reported being a paid consultant for Pfizer and Celgene. Dr Zein reported grants from National Heart, Lung, and Blood Institute during the conduct of the study. No other disclosures were reported.

Funding/Support: This work was funded by grants 1 P01 HL128192 (Drs Gaston and Marozkina) and KO8HL133381 (Dr Zein).

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

- Baratchian M, McManus JM, Berk M, et al. Sex, androgens and regulation of pulmonary AR, TMPRSS2 and ACE2. *bioRxiv* 2020:2020.04.21.051201.
- Zein J, Gaston B, Bazeley P, et al. *HSD3B1* genotype identifies glucocorticoid responsiveness in severe asthma. *Proc Natl Acad Sci U S A*. 2020;117(4):2187-2193. doi:10.1073/pnas.1918819117
- Mohan SS, Knuiman MW, Divitini ML, et al. Higher serum testosterone and dihydrotestosterone, but not oestradiol, are independently associated with favourable indices of lung function in community-dwelling men. *Clin Endocrinol (Oxf)*. 2015;83(2):268-276. doi:10.1111/cen.12738
- Fuseini H, Yung JA, Cephus JY, et al. Testosterone decreases house dust mite-induced type 2 and IL-17A-mediated airway inflammation. *J Immunol*. 2018;201(7):1843-1854. doi:10.4049/jimmunol.1800293
- Batista RL, Costa EMF, Rodrigues AS, et al. Androgen insensitivity syndrome: a review. *Arch Endocrinol Metab*. 2018;62(2):227-235. doi:10.20945/2359-3997000000031

Estimates and Projections of COVID-19 and Parental Death in the US

The scale of COVID-19 mortality in the United States, including among prime-age adults, merits efforts to continuously track how many children are affected by parental death. Children who lose



Supplemental content

a parent are at elevated risk of traumatic grief, depression, poor educational outcomes, and unintentional death or suicide, and these consequences can persist into adulthood.¹ Sudden parental death, such as that occurring owing to COVID-19, can be particularly traumatizing for children and leave families ill prepared to navigate its consequences. Moreover, COVID-19 losses are occurring at a time of social isolation, institutional

strain, and economic hardship, potentially leaving bereaved children without the supports they need.

Methods | We estimated the expected number of affected children for each COVID-19 death (the parental bereavement multiplier), allowing us to track parental bereavement as the pandemic evolves. We used kinship networks of White and Black individuals in the US estimated through demographic microsimulation to calculate the bereavement multiplier, then used the multiplier to estimate the scope of parental bereavement under various mortality scenarios (eMethods in the Supplement).² We assumed comparable multipliers for other racial groups. First, we used COVID-19 deaths and excess death counts (to address underestimation of mortality and deaths indirectly due to the pandemic) as of February 2021. Second, to facilitate comparison with a typical year, we extracted the incidence rate of parental bereavement under mortality conditions absent COVID-19 from the microsimulations data.³ Finally, we estimated future bereavement under a natural herd immunity scenario. Note that the main results use population-averaged White and Black bereavement multipliers; prior work found small differences by race.³ For current COVID-19 mortality, we also ran supplemental statistics using race-specific mortality and bereavement multipliers. This study uses deidentified, publicly available data and is not considered human subjects research.

Results | Our model suggests that each COVID-19 death leaves 0.078 children aged 0 to 17 parentally bereaved. This represents a 17.5% to 20.2% increase in parental bereavement absent COVID-19. Although the bereavement multiplier is small, it translates into large numbers of children who have lost a parent. As of February 2021, 37 300 children aged 0 to 17 years had lost at least 1 parent due to COVID-19, three-quarters of whom were adolescents (Table). Of these, 20 600 were non-Hispanic White children and 7600 were non-Hispanic Black children. When we rely on excess deaths, we estimate that 43 000 children have lost a parent. A natural herd immunity strategy that results in 1.5 million deaths⁴ demonstrates the potential effect of inaction: 116 900 parentally bereaved children.

Discussion | The number of children experiencing a parent dying of COVID-19 is staggering, with an estimated 37 300 to

Table. Estimated Number of Children Aged 0 to 17 Years Who Will Lose a Parent Owing to the COVID-19 Pandemic Under Various Scenarios

Characteristic	Estimated children experiencing parental loss, median (range) ^a		
	Age 0-17 y	Age 0-9 y	Age 10-17 y
Bereavement multiplier	0.078 (0.059-0.126)	0.021 (0.016-0.054)	0.057 (0.043-0.071)
Mortality owing to the COVID-19 pandemic			
Current mortality estimates from February 2020 to February 2021 ²			
479 000 Recorded COVID-19 deaths	37 337 (28 195-60 119)	9863 (7717-25 923)	27 474 (20 478-34 196)
552 000 Estimated excess deaths ^b	43 027 (32 492-69 281)	11 366 (8893-29 873)	31 661 (23 599-39 408)
Future mortality scenarios			
1 500 000 COVID-19 deaths	116 922 (88 295-188 264)	30 887 (24 167-81 177)	86 035 (64 128-107 086)

^a Estimates are based on the median of 40 simulations with the ranges of simulation results given in parentheses.

^b Excess deaths refer to the difference between the number of observed deaths and the number of expected deaths for the same time period, and thus captures all-cause mortality that is both directly and indirectly due to the pandemic.

43 000 already affected. For comparison, the attacks on September 11, 2001, left 3000 children without a parent.⁵ The burden will grow heavier as the death toll continues to mount. Black children are disproportionately affected, comprising only 14% of children in the US but 20% of those losing a parent to COVID-19.⁶ We note these estimates rely on demographic modeling, not survey or administrative data. Moreover, they do not include bereavement of nonparental primary caregivers.

Sweeping national reforms are needed to address the health, educational, and economic fallout affecting children. Parentally bereaved children will also need targeted support to help with grief, particularly during this period of heightened social isolation. Brief evidence-based interventions may be able to prevent the development of severe psychological problems when delivered widely,¹ although some children will need longer-term support. The establishment of a national child bereavement cohort could identify children who have lost parents, monitor them for early identification of emerging challenges, link them to locally delivered care, and form the basis for a longitudinal study of the long-term effects of mass parental bereavement during a uniquely challenging time of social isolation and economic uncertainty.

Rachel Kidman, PhD

Rachel Margolis, PhD

Emily Smith-Greenaway, PhD

Ashton M. Verdery, PhD

Author Affiliations: Program in Public Health, Stony Brook University, Stony Brook, New York (Kidman); Family, Population, and Preventive Medicine, Renaissance School of Medicine, Stony Brook University, Stony Brook, New York (Kidman); Department of Sociology, University of Western Ontario, London, Ontario, Canada (Margolis); Department of Sociology and Spatial Sciences Institute, University of Southern California, Los Angeles (Smith-Greenaway); Department of Sociology and Criminology, The Pennsylvania State University, University Park (Verdery).

Corresponding Author: Rachel Kidman, PhD, Stony Brook University, 101 Nicolls Rd, Stony Brook, NY 11794 (rachel.kidman@stonybrook.edu).

Accepted for Publication: January 21, 2021.

Published Online: April 5, 2021. doi:10.1001/jamapediatrics.2021.0161

Author Contributions: Dr Verdery had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Kidman, Margolis, Verdery.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Kidman, Verdery.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Verdery.

Obtained funding: Verdery.

Administrative, technical, or material support: Kidman.

Supervision: Verdery.

Conflict of Interest Disclosures: Dr Verdery reported grants from the National Institute on Aging and the Eunice Kennedy Shriver National Institute of Child Health and Human Development during the conduct of the study. No other disclosures were reported.

1. Bergman A-S, Axberg U, Hanson E. When a parent dies: a systematic review of the effects of support programs for parentally bereaved children and their caregivers. *BMC Palliat Care*. 2017;16(1):39. doi:10.1186/s12904-017-0223-y

2. Centers for Disease Control and Prevention. COVID-19 death data and resources. Accessed March 1, 2021. <https://www.cdc.gov/nchs/covid19/mortality-overview.htm>

3. Verdery AM, Smith-Greenaway E, Margolis R, Daw J. Tracking the reach of COVID-19 kin loss with a bereavement multiplier applied to the United States. *Proc Natl Acad Sci U S A*. 2020;117(30):17695-17701. doi:10.1073/pnas.2007476117

4. Fontanet A, Cauchemez S. COVID-19 herd immunity: where are we? *Nat Rev Immunol*. 2020;20(10):583-584. doi:10.1038/s41577-020-00451-5

5. Chemtob CM, Conroy DL, Hochhauser CJ, et al. Children who lost a parent as a result of the terrorist attacks of September 11, 2001: registry construction and population description. *Death Stud*. 2007;31(1):87-100. doi:10.1080/07481180600995263

6. US Census Bureau. 2019 Monthly national population estimates by age, sex, race, hispanic origin, and population universe for the United States: April 1, 2010 to December 1, 2020 (NC-EST2019-ALLDATA). Accessed November 11, 2020. <https://www2.census.gov/programs-surveys/popest/datasets/2010-2019/national/asrh/nc-est2019-alldata-r-file22.csv>

Racial and Ethnic Differences in Marijuana Use in e-Cigarettes Among US Youth in 2017, 2018, and 2020

Marijuana use in e-cigarettes increased among US adolescents during 2017 to 2018.¹ States are increasingly legalizing adult marijuana sales, which could affect ease of access, social acceptance, and youth marijuana use. Marijuana use during adolescence is associated with decline in memory, attention, and learning.² Differences in use by racial and ethnic groups in adults have been reported,³ but little is known about differences among youth. Identifying differences in marijuana use in e-cigarettes by racial and ethnic groups opens avenues for targeted interventions. This study reports the prevalence of ever use of marijuana in e-cigarettes among US youth in 2020 and changes in prevalence from 2017 to 2020 by racial and ethnic groups.

Methods | Self-reported data were collected from the 2017, 2018, and 2020 National Youth Tobacco Surveys (NYTS), annual cross-sectional surveys administered to students in grades 6 to 12 attending public and private schools.⁴ NYTS was paper based in 2017 (N = 20 675; response rate = 71.6%) and 2018 (N = 20 189; response rate = 68.2%) and administered electronically in 2020 (N = 14 531; response rate = 43.6%). Marijuana use in e-cigarettes was not queried in 2019. Persons who ever used marijuana in e-cigarettes responded yes to the question, “Have you ever used marijuana, marijuana concentrates, marijuana waxes, THC [tetrahydrocannabinol], or hash oils in an e-cigarette?” (2017 and 2018) or “Have you ever vaped marijuana or cannabis (including concentrates, waxes, or hash oils)?” (2020). NYTS data collection was approved by the US Centers for Disease Control and Prevention institutional review board. Participation in NYTS was voluntary; written parental consent and student assent were required for NYTS participation.

Weighted analyses provided national estimates and accounted for the complex sampling design. Using 2020 data, correlates of use assessed included sex, race and ethnicity, school level, current (past 30-day) e-cigarette use, and frequency of e-cigarette use. Statistical significance ($P < .05$) of differences in estimates within years and between years (2020 vs 2017) and by covariates were determined using pairwise *t* tests. There were 365 students excluded owing to missing self-reported race/ethnicity. Analyses were conducted using SAS-Callable SUDAAN version 11.0.3 (SAS Institute).

Results | In 2020, 19.5% (weighted) of students (2485 of 13 719) reported ever using marijuana in an e-cigarette (vs 11.1% [1975 of 16 872] in 2017; $P < .001$), including 68% (1119 of 1685) of