

## Estimates of eigenvalues of a boundary value problem with a parameter\*

ALEXEY VLADISLAVOVICH FILINOVSKIY<sup>1,†</sup>

<sup>1</sup> *Department of Fundamental Sciences, N.E. Bauman Moscow State Technical University, 2<sup>nd</sup> Baumanskaya 5, 105 005 Moscow, Russia*

Received October 31, 2013; accepted March 21, 2014

---

**Abstract.** We study an eigenvalue problem for the Laplace operator with a boundary condition containing a parameter. We estimate the rate of convergence of the eigenvalues to the eigenvalues of the Dirichlet problem for large positive values of the parameter.

**AMS subject classifications:** 35J05, 35P15, 49R05

**Key words:** Laplace operator, boundary value problem, large parameter, eigenvalues

---

### 1. Introduction

We consider the eigenvalue problem

$$\Delta u + \lambda u = 0 \quad \text{in } \Omega, \quad (1)$$

$$\frac{\partial u}{\partial \nu} + \alpha \sigma(x) u = 0 \quad \text{on } \Gamma, \quad (2)$$

where  $\Omega \subset R^n$ ,  $n \geq 2$ , is a bounded domain with boundary  $\Gamma = \partial\Omega \in C^2$ . By  $\nu$  we denote the outward unit normal vector to  $\Gamma$ ,  $\alpha$  is a real parameter. The function  $\sigma(x) \in C^1(\Gamma)$  is positive:

$$0 < \sigma_0 \leq \sigma(x) \leq \sigma_1, \quad \sigma_0 = \inf_{x \in \Gamma} \sigma(x) \quad \text{and} \quad \sigma_1 = \sup_{x \in \Gamma} \sigma(x).$$

Problem (1), (2) with  $\sigma(x) = 1$  is known as the Robin (Fourier) problem for  $\alpha > 0$  (see [6, Ch. 7, Par. 7.2]), and the generalized Robin problem for all  $\alpha$  ([5]).

There is a sequence of eigenvalues  $\lambda_1(\alpha) < \lambda_2(\alpha) \leq \dots$  of problem (1) - (2) enumerated according to their multiplicities with

$$\lim_{k \rightarrow \infty} \lambda_k(\alpha) = +\infty.$$

We also consider the sequence of eigenvalues  $0 < \lambda_1^D < \lambda_2^D \leq \dots$  of the Dirichlet eigenvalue problem

$$\Delta u + \lambda u = 0 \quad \text{in } \Omega, \quad (3)$$

$$u = 0 \quad \text{on } \Gamma, \quad (4)$$

---

\*This work was supported by the RFBR Grant No. 11-01-00989.

†Corresponding author. *Email address:* `f1nv@yandex.ru` (A. Filinovskiy)

with

$$\lim_{k \rightarrow \infty} \lambda_k^D = +\infty.$$

Note that the eigenvalues  $\lambda_1(\alpha)$  and  $\lambda_1^D$  are simple and the corresponding eigenfunctions  $u_{1,\alpha}(x)$  and  $u_1^D(x)$  are positive.

In this paper, we estimate  $\lambda_k(\alpha)$  for large values of  $\alpha$ . We now give some known results.

It is easy to see that  $\lambda_k(\alpha) \leq \lambda_k^D$ ,  $k = 1, 2, \dots$ . These inequalities give the upper bound of  $\lambda_k(\alpha)$  for all values of  $\alpha$ . It was announced in ([2, Ch. 6, Par. 2, No. 1]) that for  $n = 2$  and a smooth boundary  $\lim_{\alpha \rightarrow +\infty} \lambda_k(\alpha) = \lambda_k^D$ .

Later the properties of the first eigenvalue  $\lambda_1(\alpha)$  were studied more precisely. Consider the case  $\sigma(x) = 1$ . The following two-sided estimates:

$$\lambda_1^D \left(1 + \frac{\lambda_1^D}{\alpha q_1}\right)^{-1} \leq \lambda_1(\alpha) \leq \lambda_1^D \left(1 + \frac{4\pi}{\alpha |\Gamma|}\right)^{-1}, \quad \alpha > 0,$$

were obtained in [12] for  $n = 2$ . Here  $|\Gamma|$  is the length of  $\Gamma$  and  $q_1$  is the first eigenvalue of the Steklov problem

$$\begin{aligned} \Delta^2 u &= 0 \quad \text{in } \Omega, \\ u &= 0, \quad \Delta u - q \frac{\partial u}{\partial \nu} = 0 \quad \text{on } \Gamma. \end{aligned}$$

In [4], for any  $n \geq 2$  we establish the following asymptotic expansion:

$$\lambda_1(\alpha) = \lambda_1^D - \frac{\int_{\Gamma} \left(\frac{\partial u_1^D}{\partial \nu}\right)^2 ds}{\int_{\Omega} (u_1^D)^2 dx} \alpha^{-1} + o(\alpha^{-1}), \quad \alpha \rightarrow +\infty.$$

The case  $\alpha < 0$  has recently attracted attention (see, for instance, [9]). It was shown in [9] that for a piecewise- $C^1$  boundary

$$\liminf_{\alpha \rightarrow -\infty} \lambda_1(\alpha)/(-\alpha^2) \geq 1.$$

For  $C^1$  boundaries it was proved ([10]) that

$$\lim_{\alpha \rightarrow -\infty} \lambda_1(\alpha)/(-\alpha^2) = 1.$$

The  $C^1$ -condition is optimal. In [9], the authors constructed plane triangle domains for which

$$\lim_{\alpha \rightarrow -\infty} \lambda_1(\alpha)/(-\alpha^2) > 1.$$

In [3], the authors proved that for  $C^1$  boundaries

$$\lim_{\alpha \rightarrow -\infty} \frac{\lambda_k(\alpha)}{-\alpha^2} = 1 \tag{5}$$

for all  $k = 1, 2, \dots$

## 2. Main results

The main result of this paper reads as follows.

**Theorem 1.** *The eigenvalues  $\lambda_k(\alpha)$ ,  $k = 1, 2, \dots$ , satisfy the estimates*

$$0 \leq \lambda_k^D - \lambda_k(\alpha) \leq C_1 \alpha^{-1/2} (\lambda_k^D)^2, \quad \alpha > 0, \quad (6)$$

where the constant  $C_1$  does not depend on  $k$ .

In the following theorem we gather the qualitative properties of eigenvalues of problem (1) - (2) (see also [2, Ch. 6] for i) and [9] for ii) and iii) for  $\sigma(x) = 1$ )

**Theorem 2.** *The eigenvalues have the following properties:*

i)  $\lambda_k(\alpha)$ ,  $k = 1, 2, \dots$ , are continuous functions of  $\alpha$  and

$$\lambda_k(\alpha_1) \leq \lambda_k(\alpha_2), \quad \alpha_1 < \alpha_2; \quad (7)$$

ii)  $\lambda_1(\alpha)$  is a concave function of  $\alpha$ :

$$\lambda_1(\beta\alpha_1 + (1 - \beta)\alpha_2) \geq \beta\lambda_1(\alpha_1) + (1 - \beta)\lambda_1(\alpha_2), \quad 0 < \beta < 1; \quad (8)$$

iii)  $\lambda_1(\alpha)$  is differentiable and

$$\lambda_1'(\alpha) = \frac{\int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx} > 0; \quad (9)$$

iv) the following estimate

$$\liminf_{\alpha \rightarrow -\infty} \frac{\lambda_1'(\alpha)}{-\alpha} \geq \sigma_1^2 \quad (10)$$

holds.

## 3. Operator treatment

In this section, we introduce two linear operators associated with problems (1) - (2) and (3) - (4) to derive the eigenvalue estimates (6).

Consider problem (1) - (2) in the space  $H^1(\Omega)$  ([1, 11]). We define an eigenvalue of problem (1), (2) as a value  $\lambda$  for which there exists the non-zero function  $u \in H^1(\Omega)$  satisfying the integral identity

$$\int_{\Omega} (\nabla u, \nabla v) dx + \alpha \int_{\Gamma} \sigma uv ds = \lambda \int_{\Omega} uv dx \quad (11)$$

for any  $v \in H^1(\Omega)$ . Relation (11) can be rewritten as

$$\int_{\Omega} ((\nabla u, \nabla v) + Muv) dx + \alpha \int_{\Gamma} \sigma uv ds = (\lambda + M) \int_{\Omega} uv dx, \quad M > 0. \quad (12)$$

Let us define an equivalent scalar product in the space  $H^1(\Omega)$  by the formula

$$[u, v]_M = \int_{\Omega} ((\nabla u, \nabla v) + Muv) \, dx, \quad \|u\|_M^2 = [u, u]_M. \tag{13}$$

Now (12) transforms to

$$[u, v]_M + \alpha[Tu, v]_M = (\lambda + M)[Bu, v]_M,$$

where the linear self-adjoint non-negative operators  $T : H^1(\Omega) \rightarrow H^1(\Omega)$  and  $B : H^1(\Omega) \rightarrow H^1(\Omega)$  were defined by the bilinear forms

$$[Tu, v]_M = \int_{\Gamma} \sigma uv \, ds, \quad [Bu, v]_M = \int_{\Omega} uv \, dx, \quad u, v \in H^1(\Omega). \tag{14}$$

Hence we have an equation in the space  $H^1(\Omega)$  with the norm  $\|\cdot\|_M$ :

$$(I + \alpha T)u = (\lambda + M)Bu. \tag{15}$$

Now we use the inequality ([11, Ch. 3, Par. 5, Formula 19])

$$\|v\|_{L_2(\Gamma)}^2 \leq \varepsilon \|\nabla v\|_{L_2(\Omega)}^2 + C_{\varepsilon} \|v\|_{L_2(\Omega)}^2, \tag{16}$$

which is valid for  $v \in H^1(\Omega)$  with an arbitrary  $\varepsilon > 0$ . Using (14), (16), we obtain

$$\begin{aligned} \|Tu\|_M^2 &= [Tu, Tu]_M = \int_{\Gamma} \sigma u Tu \, ds \leq \sigma_1 \|u\|_{L_2(\Gamma)} \|Tu\|_{L_2(\Gamma)} \\ &\leq \sigma_1 \varepsilon \left( \int_{\Omega} \left( |\nabla Tu|^2 + \frac{C_{\varepsilon}}{\varepsilon} (Tu)^2 \right) dx \right)^{1/2} \\ &\quad \times \left( \int_{\Omega} \left( |\nabla u|^2 + \frac{C_{\varepsilon}}{\varepsilon} u^2 \right) dx \right)^{1/2} \leq C_2 \varepsilon \|Tu\|_M \|u\|_M, \end{aligned} \tag{17}$$

where  $\varepsilon > 0$ ,  $M = M_{\varepsilon}$ . It follows from (17) that

$$\|Tu\|_{M_{\varepsilon}} \leq C_2 \varepsilon \|u\|_{M_{\varepsilon}},$$

and for any arbitrary small  $\varepsilon$  we have  $\|\alpha T\|_{H^1(\Omega) \rightarrow H^1(\Omega)} < 1$  for  $|\alpha| < 1/C_2\varepsilon$ . Therefore, the inverse operator  $(I + \alpha T)^{-1}$  is bounded and

$$\|(I + \alpha T)^{-1}\| \leq (1 - |\alpha| \|T\|)^{-1}.$$

Hence, equation (15) is equivalent to

$$(I - (\lambda + M)(I + \alpha T)^{-1}B)u = 0.$$

The operator  $B$  is compact ([11, Ch. 3, Par. 5, Th. 3]) and the operator  $(I + \alpha T)^{-1}B : H^1(\Omega) \rightarrow H^1(\Omega)$  is also compact. Hence the spectrum of problem (15) consists of real eigenvalues  $\lambda_j(\alpha)$ ,  $j = 1, 2, \dots$ , of finite multiplicity with the only limit point at the infinity. From (14), (15) we obtain the inequality

$$\lambda_j(\alpha) \geq -M_{\varepsilon} + (1 - |\alpha| \|T\|) \frac{\|u_{j,\alpha}\|_{M_{\varepsilon}}^2}{\|u_{j,\alpha}\|_{L_2(\Omega)}^2} \geq -M_{\varepsilon}$$

with the corresponding eigenfunction  $u_{j,\alpha}$ . Thus,  $\lambda_j(\alpha) \rightarrow +\infty, j \rightarrow \infty$ .

By the variational principle ([11, Ch. 4, Par. 1, No. 4]) we have

$$\lambda_k(\alpha) = \sup_{v_1, \dots, v_{k-1} \in L_2(\Omega)} \inf_{\substack{v \in H^1(\Omega) \\ (v, v_j)_{L_2(\Omega)} = 0 \\ j = 1, \dots, k-1}} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx}, \tag{18}$$

$$\lambda_k^D = \sup_{v_1, \dots, v_{k-1} \in L_2(\Omega)} \inf_{\substack{v \in \mathring{H}^1(\Omega) \\ (v, v_j)_{L_2(\Omega)} = 0 \\ j = 1, \dots, k-1}} \frac{\int_{\Omega} |\nabla v|^2 dx}{\int_{\Omega} v^2 dx}, \quad k = 1, 2, \dots \tag{19}$$

To prove inequalities (6) we apply the following statement (see [6, Ch. 2, Th. 2.3.1]).

**Theorem 3.** *Let  $T_1$  and  $T_2$  be two linear self-adjoint, compact and positive operators on a separable Hilbert space  $H$ . Assume also that  $\mu_k(T_1)$  and  $\mu_k(T_2)$  are their  $k$ -th respective eigenvalues. Then*

$$|\mu_k(T_1) - \mu_k(T_2)| \leq \|T_1 - T_2\|. \tag{20}$$

Now we give the proof of Theorem 1.

**Proof.** Consider the boundary value problem

$$-\Delta u + u = h \quad \text{in } \Omega, \tag{21}$$

$$\frac{\partial u}{\partial \nu} + \alpha \sigma(x)u = 0 \quad \text{on } \Gamma, \quad \alpha > 0, \tag{22}$$

with  $h \in L_2(\Omega)$ . A weak solution  $u \in H^1(\Omega)$  of problem (21), (22) satisfy the integral identity

$$\int_{\Omega} ((\nabla u, \nabla v) + uv) dx + \alpha \int_{\Gamma} \sigma uv ds = \int_{\Omega} hv dx \tag{23}$$

for all  $v \in H^1(\Omega)$ . Let us define the scalar product in the space  $H^1(\Omega)$  as

$$(u, v)_{H^1(\Omega), \alpha} = \int_{\Omega} ((\nabla u, \nabla v) + uv) dx + \alpha \int_{\Gamma} \sigma uv ds \tag{24}$$

and the corresponding norm by

$$\|u\|_{H^1(\Omega), \alpha}^2 = (u, u)_{H^1(\Omega), \alpha}.$$

Due to (16), scalar product (24) is equivalent to the standard one

$$(u, v)_{H^1(\Omega)} = \int_{\Omega} ((\nabla u, \nabla v) + uv) dx. \tag{25}$$

Using (23), (24), we obtain the relation

$$(u, v)_{H^1(\Omega), \alpha} = (h, v)_{L_2(\Omega)}. \tag{26}$$

Hence, consider the linear functional

$$l_h(v) = (h, v)_{L_2(\Omega)}.$$

This functional is bounded on the space  $H^1(\Omega)$ :

$$|l_h(v)| \leq \|h\|_{L_2(\Omega)} \|v\|_{L_2(\Omega)}. \quad (27)$$

Now, by the Riesz lemma there exists the unique function  $u \in H^1(\Omega)$  satisfying integral identity (23). Applying (26) with  $v = u$ , we obtain

$$\|u\|_{H^1(\Omega), \alpha}^2 \leq \|h\|_{L_2(\Omega)} \|u\|_{H^1(\Omega), \alpha}.$$

Therefore,

$$\|u\|_{L_2(\Omega)} \leq \|u\|_{H^1(\Omega), \alpha} \leq \|h\|_{L_2(\Omega)}, \quad (28)$$

and we can define the bounded linear operator  $A_\alpha : L_2(\Omega) \rightarrow L_2(\Omega)$  such that  $u = A_\alpha h$  and  $\|A_\alpha\| \leq 1$ . Moreover, if the domain  $\Omega$  with  $C^2$  boundary is bounded, then the space  $H^1(\Omega)$  embeds compactly into the space  $L_2(\Omega)$  ([6, Ch. 1, Th. 1.1.1]). It means that the operator  $A_\alpha$  is compact. Note that

$$\begin{aligned} (h, A_\alpha g)_{L_2(\Omega)} &= \int_{\Omega} h A_\alpha g \, dx = \int_{\Omega} h v \, dx \\ &= \int_{\Omega} ((\nabla u, \nabla v) + uv) \, dx + \alpha \int_{\Gamma} \sigma uv \, ds \\ &= \int_{\Omega} u g \, dx = (A_\alpha h, g)_{L_2(\Omega)}, \quad f, g \in L_2(\Omega), \end{aligned} \quad (29)$$

with  $u = A_\alpha h$ ,  $v = A_\alpha g$ ,  $u, v \in H^1(\Omega)$ . Relation (29) means that  $A_\alpha$  is a self-adjoint operator. Now, by relation (29) we have

$$\begin{aligned} (h, A_\alpha h)_{L_2(\Omega)} &= \int_{\Omega} u h \, dx \\ &= \int_{\Omega} (|\nabla u|^2 + u^2) \, dx + \alpha \int_{\Gamma} \sigma u^2 \, ds = \|u\|_{H^1(\Omega), \alpha}^2 > 0, \quad h \neq 0. \end{aligned}$$

Hence, the operator  $A_\alpha$  is positive. Finally,  $A_\alpha$  is a self-adjoint positive compact operator in the Hilbert space  $H = L_2(\Omega)$ . By the well-known theorem ([6, Ch. 1, Th. 1.2.1]),  $A_\alpha$  has a sequence of eigenvalues  $\{\mu_k(\alpha)\}$ ,  $k = 1, 2, \dots$  with finite multiplicities such that  $\mu_k(\alpha) > 0$ ,  $\mu_k(\alpha) \searrow 0$ ,  $k \rightarrow \infty$ . Let us denote by  $u_{k,\alpha}(x) \in L_2(\Omega)$  the eigenfunction satisfying  $A_\alpha u_{k,\alpha} = \mu_k(\alpha) u_{k,\alpha}$ . Thus,

$$\mu_k(\alpha) (u_{k,\alpha}, v)_{H^1(\Omega), \alpha} = (u_{k,\alpha}, v)_{L_2(\Omega)}$$

and

$$\mu_k(\alpha) \left( \int_{\Omega} ((\nabla u_{k,\alpha}, \nabla v) + u_{k,\alpha} v) \, dx + \alpha \int_{\Gamma} \sigma u_{k,\alpha} v \, ds \right) = \int_{\Omega} u_{k,\alpha} v \, dx.$$

It can be seen that

$$\mu_k(\alpha) = \frac{1}{\lambda_k(\alpha) + 1}.$$

Let us note that for  $\alpha > 0$  we have

$$\mu_k(\alpha) \leq \frac{1}{\lambda_1(\alpha) + 1} < 1,$$

so  $\|A_\alpha\| < 1$ .

Furthermore, consider a Dirichlet problem

$$-\Delta y + y = h \quad \text{in } \Omega, \quad (30)$$

$$y = 0 \quad \text{on } \Gamma. \quad (31)$$

For  $h \in L_2(\Omega)$  a weak solution  $y \in \overset{\circ}{H}^1(\Omega)$  of problem (30), (31) satisfies the integral identity

$$\int_{\Omega} ((\nabla y, \nabla v) + yv) dx = \int_{\Omega} hv dx \quad (32)$$

for all  $v \in \overset{\circ}{H}^1(\Omega)$ . Define the scalar product in the space  $\overset{\circ}{H}^1(\Omega)$  by (25). Using (25), (32), we obtain the relation

$$(y, v)_{\overset{\circ}{H}^1(\Omega)} = l_h(v). \quad (33)$$

Now, by (27) and the Riesz lemma there exists the unique function  $y \in \overset{\circ}{H}^1(\Omega)$  satisfying integral identity (32). Using (32) with  $v = y$ , we obtain

$$\|y\|_{\overset{\circ}{H}^1(\Omega)}^2 \leq \|h\|_{L_2(\Omega)} \|y\|_{\overset{\circ}{H}^1(\Omega)}. \quad (34)$$

Therefore,

$$\|y\|_{L_2(\Omega)} \leq \|y\|_{\overset{\circ}{H}^1(\Omega)} \leq \|h\|_{L_2(\Omega)}, \quad (35)$$

and we can define the bounded linear operator  $A^D : L_2(\Omega) \rightarrow L_2(\Omega)$  such that  $u = A^D h$  and  $\|A^D\| \leq 1$ . If the domain  $\Omega$  is bounded, then the space  $\overset{\circ}{H}^1(\Omega)$  embeds compactly into the space  $L_2(\Omega)$  ([6, Ch. 1, Th. 1.1.1]). Hence, the operator  $A^D$  is compact. Note that

$$\begin{aligned} (h, A^D g)_{L_2(\Omega)} &= \int_{\Omega} h A^D g dx = \int_{\Omega} hv dx = \int_{\Omega} ((\nabla y, \nabla v) + yv) dx \\ &= \int_{\Omega} yg dx = (A^D h, g)_{L_2(\Omega)}, \quad f, g \in L_2(\Omega), \end{aligned} \quad (36)$$

with  $y = A^D h$ ,  $v = A^D g$ ,  $y, v \in \overset{\circ}{H}^1(\Omega)$ . Relation (36) means that  $A^D$  is a self-adjoint operator. Now, by (36) we have

$$(h, A^D h)_{L_2(\Omega)} = \int_{\Omega} yh dx = \int_{\Omega} (|\nabla y|^2 + y^2) dx = \|y\|_{\overset{\circ}{H}^1(\Omega)}^2 > 0, \quad h \neq 0.$$

Hence, the operator  $A^D$  is positive. Finally,  $A^D$  is a self-adjoint positive compact operator in the Hilbert space  $H = L_2(\Omega)$ . By ([6, Ch. 1, Th. 1.2.1]), there exists a

sequence of eigenvalues  $\{\mu_k^D\}$ ,  $k = 1, 2, \dots$ , of the operator  $A^D$  with finite multiplicities such that  $\mu_k^D > 0$ ,  $\mu_k^D \searrow 0$ ,  $k \rightarrow \infty$ . Denote by  $y_k(x) \in L_2(\Omega)$  the respective eigenfunction satisfying  $A^D y_k = \mu_k^D y_k$ . Thus,  $\mu_k^D (y_k, v)_{H^1(\Omega)} = (y_k, v)_{L_2(\Omega)}$  and

$$\mu_k^D \int_{\Omega} ((\nabla y_k, \nabla v) + y_k v) dx = \int_{\Omega} y_k v dx.$$

Then,

$$\mu_k^D = \frac{1}{\lambda_k^D + 1}.$$

Note that

$$\mu_k^D \leq \frac{1}{\lambda_1^D + 1} < 1,$$

so  $\|A^D\| < 1$ .

Now we estimate the norm  $\|A_\alpha - A^D\|_{L_2(\Omega) \rightarrow L_2(\Omega)}$  for large positive values of  $\alpha$ .

Let us remind that in domains with  $C^2$ -class boundaries and positive  $\sigma(x) \in C^1(\Gamma)$  the functions  $u = A_\alpha h$  and  $y = A^D h$  are strong solutions and belong to  $H^2(\Omega)$  ([11, Ch. 4, Par. 2, Th. 4]). Moreover, the following estimate

$$\|y\|_{H^2(\Omega)} \leq C_2 \|h\|_{L_2(\Omega)} \tag{37}$$

holds. Now we use estimate (16) with  $\varepsilon = 1$ :

$$\|y\|_{L_2(\Gamma)} \leq C_3 \|y\|_{H^1(\Omega)}. \tag{38}$$

Combining (37) and (38) we derive the inequality

$$\|\nabla y\|_{L_2(\Gamma)} \leq C_4 \|y\|_{H^2(\Omega)}. \tag{39}$$

Since  $\left| \frac{\partial y}{\partial \nu} \right| \leq |\nabla y|$  on  $\Gamma$ , from (37), (39) we obtain the estimate

$$\left\| \frac{\partial y}{\partial \nu} \right\|_{L_2(\Gamma)} \leq C_5 \|h\|_{L_2(\Omega)}. \tag{40}$$

Suppose that  $w = (A^D - A_\alpha) h$ . By (21), (22), (30), (31) the function  $w$  is a solution of the boundary value problem

$$-\Delta w + w = 0 \quad \text{in } \Omega, \tag{41}$$

$$\frac{\partial w}{\partial \nu} + \alpha \sigma w = \frac{\partial y}{\partial \nu} \quad \text{on } \Gamma. \tag{42}$$

Multiplying equation (41) by  $w$  and integrating it over  $\Omega$  with respect to boundary condition (42), we get the relation

$$\int_{\Omega} (|\nabla w|^2 + w^2) dx + \frac{1}{\alpha} \int_{\Gamma} \left( \frac{\partial w}{\partial \nu} \right)^2 \frac{ds}{\sigma} = \frac{1}{\alpha} \int_{\Gamma} \frac{\partial w}{\partial \nu} \frac{\partial y}{\partial \nu} \frac{ds}{\sigma}, \quad \alpha > 0. \tag{43}$$



Then we obtain the inequality

$$\|w\|_{L_2(\Omega)}^2 + \frac{1}{\alpha} \left\| \frac{\partial w}{\partial \nu} \right\|_{L_2(\Gamma)}^2 \leq \frac{C_6}{\alpha} \left\| \frac{\partial w}{\partial \nu} \right\|_{L_2(\Gamma)} \left\| \frac{\partial y}{\partial \nu} \right\|_{L_2(\Gamma)}$$

and, consequently,

$$\|w\|_{L_2(\Omega)}^2 + \frac{1}{\alpha} \left\| \frac{\partial w}{\partial \nu} \right\|_{L_2(\Gamma)}^2 \leq \frac{1}{2\alpha} \left\| \frac{\partial w}{\partial \nu} \right\|_{L_2(\Gamma)}^2 + \frac{C_6^2}{2\alpha} \left\| \frac{\partial y}{\partial \nu} \right\|_{L_2(\Gamma)}^2.$$

Therefore, we have the estimate

$$\|w\|_{L_2(\Omega)} \leq \frac{C_6}{\sqrt{2\alpha}} \left\| \frac{\partial y}{\partial \nu} \right\|_{L_2(\Gamma)}, \quad \alpha > 0. \quad (44)$$

Combining (44) with (40), we get

$$\|w\|_{L_2(\Omega)} \leq C_7 \alpha^{-1/2} \|h\|_{L_2(\Omega)}, \quad \alpha > 0,$$

with the constant  $C_6$  independent of  $\alpha$ . Thus, for all  $h \in L_2(\Omega)$  we have the estimate

$$\|(A^D - A_\alpha)h\|_{L_2(\Omega)} \leq C_7 \alpha^{-1/2} \|h\|_{L_2(\Omega)}$$

and

$$\|A^D - A_\alpha\| \leq C_7 \alpha^{-1/2}, \quad \alpha > 0. \quad (45)$$

Now we apply (20) to the operators  $T_1 = A_\alpha$ ,  $T_2 = A^D$ . Then, by the relations

$$\mu_k(\alpha) = \frac{1}{\lambda_k(\alpha) + 1}, \quad \mu_k^D = \frac{1}{\lambda_k^D + 1},$$

and inequalities (20), (45) we get the estimate

$$\left| \frac{1}{\lambda_k(\alpha) + 1} - \frac{1}{\lambda_k^D + 1} \right| \leq C_7 \alpha^{-1/2}. \quad (46)$$

Therefore,

$$|\lambda_k^D - \lambda_k(\alpha)| \leq C_7 \alpha^{-1/2} (\lambda_k^D + 1) (\lambda_k(\alpha) + 1). \quad (47)$$

and taking into account inequalities (49) (see Section 4), we obtain the estimate

$$0 \leq \lambda_k^D - \lambda_k(\alpha) \leq C_7 \alpha^{-1/2} (\lambda_k^D + 1)^2 \leq C_1 \alpha^{-1/2} (\lambda_k^D)^2. \quad (48)$$

The proof of Theorem 1 is completed.  $\square$

### 4. General properties of eigenvalues

In this Section, we give the proof of Theorem 2.

**Proof.** Due to (18),  $\lambda_k(\cdot)$  is an increasing function. Using (19) and the inclusion  $\overset{\circ}{H}^1(\Omega) \subset H^1(\Omega)$ , we have

$$\begin{aligned} \lambda_k(\alpha) &= \sup_{v_1, \dots, v_{k-1} \in L_2(\Omega)} \inf_{\substack{v \in H^1(\Omega) \\ (v, v_j)_{L_2(\Omega)} = 0 \\ j = 1, \dots, k-1}} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &\leq \sup_{v_1, \dots, v_{k-1} \in L_2(\Omega)} \inf_{\substack{v \in \overset{\circ}{H}^1(\Omega) \\ (v, v_j)_{L_2(\Omega)} = 0 \\ j = 1, \dots, k-1}} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &= \sup_{v_1, \dots, v_{k-1} \in L_2(\Omega)} \inf_{\substack{v \in \overset{\circ}{H}^1(\Omega) \\ (v, v_j)_{L_2(\Omega)} = 0 \\ j = 1, \dots, k-1}} \frac{\int_{\Omega} |\nabla v|^2 dx}{\int_{\Omega} v^2 dx} = \lambda_k^D. \end{aligned} \tag{49}$$

The continuity of  $\lambda_k(\alpha)$  was proved in ([2, Ch. 6, Par. 2, No. 6]).

Inequality (8) can be proved by the following:

$$\begin{aligned} \lambda_1(\beta\alpha_1 + (1 - \beta)\alpha_2) &= \inf_{v \in H^1(\Omega)} \frac{\int_{\Omega} |\nabla v|^2 dx + (\beta\alpha_1 + (1 - \beta)\alpha_2) \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &\geq \beta \inf_{v \in H^1(\Omega)} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha_1 \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &\quad + (1 - \beta) \inf_{v \in H^1(\Omega)} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha_2 \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &= \beta\lambda_1(\alpha_1) + (1 - \beta)\lambda_1(\alpha_2), \quad 0 < \beta < 1. \end{aligned}$$

The eigenvalue  $\lambda_1(\alpha)$  is simple for all  $-\infty < \alpha < \infty$ . The family of self-adjoint operators  $(I + \alpha T)^{-1}B$  in the space  $H^1(\Omega)$  with norm (13) satisfies the conditions of the asymptotic perturbation theorem ([7, Ch. 8, Par. 4, Th. 2.9]). It means that the eigenvalue  $\lambda_1(\alpha)$  is a differentiable function of  $\alpha$ . So

$$\lim_{j \rightarrow \infty} \frac{\lambda_1(\alpha_j) - \lambda_1(\alpha)}{\alpha_j - \alpha} = \lambda_1'(\alpha) \tag{50}$$

for an arbitrary sequence  $\alpha_j \rightarrow \alpha, j \rightarrow \infty, \alpha_j \neq \alpha$ . Let  $\alpha_j \rightarrow \alpha, j \rightarrow \infty$ , and  $\|u_{1,\alpha_j}\|_{L_2(\Omega)} = 1, u_{1,\alpha_j} \geq 0$ . Therefore,  $\|u_{1,\alpha_j}\|_{H^1(\Omega)} \leq C_8$ . By (11), the functions  $u_{1,\alpha_j}$  satisfy

$$\int_{\Omega} (\nabla u_{1,\alpha_j}, \nabla v) dx + \alpha_j \int_{\Gamma} \sigma u_{1,\alpha_j} v ds = \lambda_1(\alpha_j) \int_{\Omega} u_{1,\alpha_j} v dx. \tag{51}$$

Now, we can choose a subsequence  $u_{1,\alpha_j} \rightharpoonup u$  weakly in  $H^1(\Omega)$  and  $\|u_{1,\alpha_j} - u\|_{L_2(\Omega)} \rightarrow 0$ ,  $\|u_{1,\alpha_j} - u\|_{L_2(\Gamma)} \rightarrow 0$ . It means that  $u \geq 0$  and  $\|u\|_{L_2(\Omega)} = 1$ . Due to (51),  $u$  satisfies the integral identity

$$\int_{\Omega} (\nabla u, \nabla v) dx + \alpha \int_{\Gamma} \sigma uv ds = \lambda_1(\alpha) \int_{\Omega} uv dx. \quad (52)$$

Hence, by the uniqueness of the first positive normalized eigenfunction  $u = u_{1,\alpha}$  and

$$\|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Omega)} \rightarrow 0, \quad j \rightarrow \infty. \quad (53)$$

Now, we have

$$\begin{aligned} & \int_{\Omega} |\nabla(u_{1,\alpha_j} - u_{1,\alpha})|^2 dx + \alpha \int_{\Gamma} \sigma(u_{1,\alpha_j} - u_{1,\alpha})^2 ds \\ &= \lambda_1(\alpha) \int_{\Omega} (u_{1,\alpha_j} - u_{1,\alpha})^2 dx \\ &+ (\lambda_1(\alpha_j) - \lambda_1(\alpha)) \int_{\Omega} u_{1,\alpha_j} (u_{1,\alpha_j} - u_{1,\alpha}) dx \\ &- (\alpha_j - \alpha) \int_{\Gamma} \sigma u_{1,\alpha_j} (u_{1,\alpha_j} - u_{1,\alpha}) ds. \end{aligned} \quad (54)$$

It follows from (54) that

$$\begin{aligned} \|u_{1,\alpha_j} - u_{1,\alpha}\|_{H^1(\Omega)}^2 &\leq C_9 \left( |\alpha| \|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Gamma)}^2 \right. \\ &+ (|\lambda_1(\alpha)| + 1) \|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Omega)}^2 \\ &+ |\lambda_1(\alpha_j) - \lambda_1(\alpha)| \|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Omega)} \|u_{1,\alpha_j}\|_{L_2(\Omega)} \\ &\left. + |\alpha_j - \alpha| \|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Gamma)} \|u_{1,\alpha_j}\|_{L_2(\Gamma)} \right). \end{aligned} \quad (55)$$

Applying (50) and (16) with sufficiently small  $\varepsilon$  we obtain

$$\|u_{1,\alpha_j} - u_{1,\alpha}\|_{H^1(\Omega)}^2 \leq C_{10} \left( \|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Omega)}^2 + (\alpha_j - \alpha)^2 \|u_{1,\alpha_j}\|_{H^1(\Omega)}^2 \right). \quad (56)$$

Due to (16), (53) and (56) we get

$$\|u_{1,\alpha_j} - u_{1,\alpha}\|_{L_2(\Gamma)} \leq C_{11} \|u_{1,\alpha_j} - u_{1,\alpha}\|_{H^1(\Omega)} \rightarrow 0, \quad j \rightarrow \infty.$$

Therefore,

$$\int_{\Gamma} \sigma u_{1,\alpha_j}^2 ds \rightarrow \int_{\Gamma} \sigma u_{1,\alpha}^2 ds, \quad j \rightarrow \infty. \quad (57)$$

Now, to obtain (9) we use the inequalities

$$\begin{aligned} \lambda_1(\alpha_j) - \lambda_1(\alpha) &= \lambda_1(\alpha_j) - \inf_{v \in H^1(\Omega)} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} \\ &\geq \lambda_1(\alpha_j) - \frac{\int_{\Omega} |\nabla u_{1,\alpha_j}|^2 dx + \alpha \int_{\Gamma} \sigma u_{1,\alpha_j}^2 ds}{\int_{\Omega} u_{1,\alpha_j}^2 dx} = (\alpha_j - \alpha) \frac{\int_{\Gamma} \sigma u_{1,\alpha_j}^2 ds}{\int_{\Omega} u_{1,\alpha_j}^2 dx} \end{aligned}$$

and

$$\begin{aligned} \lambda_1(\alpha_j) - \lambda_1(\alpha) &= \inf_{v \in H^1(\Omega)} \frac{\int_{\Omega} |\nabla v|^2 dx + \alpha_j \int_{\Gamma} \sigma v^2 ds}{\int_{\Omega} v^2 dx} - \lambda_1(\alpha) \\ &\leq \frac{\int_{\Omega} |\nabla u_{1,\alpha}|^2 dx + \alpha_j \int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx} - \lambda_1(\alpha) = (\alpha_j - \alpha) \frac{\int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx}. \end{aligned}$$

Therefore, for  $\alpha_j > \alpha$

$$\frac{\int_{\Gamma} \sigma u_{1,\alpha_j}^2 ds}{\int_{\Omega} u_{1,\alpha_j}^2 dx} \leq \frac{\lambda_1(\alpha_j) - \lambda_1(\alpha)}{\alpha_j - \alpha} \leq \frac{\int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx}. \tag{58}$$

Finally, it follows from (50), (57) and (58) that

$$\lambda'_1(\alpha) = \frac{\int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx}.$$

By ([11, Ch. 4, Par. 2, Th. 4]),  $u_{1,\alpha} \in H^2(\Omega)$  and it satisfies equation (1) almost everywhere and the boundary condition in the sense of trace (the so-called strong solution). In the case  $\int_{\Gamma} \sigma u_{1,\alpha}^2 ds = 0$ , by (2) we have:

$$u_{1,\alpha} = \frac{\partial u_{1,\alpha}}{\partial \nu} = 0 \quad \text{on } \Gamma.$$

Applying the uniqueness theorem to the Cauchy problem for second-order elliptic equations ([8, Ch. 1, Par. 3, Th. 1.46]), we get  $u_{1,\alpha} = 0$  in  $\Omega$ . This contradiction proves that  $\lambda'_1(\alpha) > 0$  for all  $\alpha$ . Taking into account (9), we have the inequality  $\lambda_1(\alpha) < \lambda_1^D$ .

By combining the result from [10] with (9) we obtain the relations

$$\begin{aligned} \alpha \lambda'_1(\alpha) &= \frac{\alpha \int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx} \leq \frac{\int_{\Omega} |\nabla u_{1,\alpha}|^2 dx + \alpha \int_{\Gamma} \sigma u_{1,\alpha}^2 ds}{\int_{\Omega} u_{1,\alpha}^2 dx} \\ &= \lambda_1(\alpha) = -\alpha^2 \sigma_1^2 (1 + \varrho(\alpha)), \quad \varrho(\alpha) \rightarrow 0, \quad \alpha \rightarrow -\infty. \end{aligned}$$

Hence,

$$\frac{\lambda'_1(\alpha)}{-\alpha} \geq \sigma_1^2 (1 + \varrho(\alpha)), \quad \alpha < 0,$$

and inequality (10) is proved.

This completes the proof of Theorem 2. □

### References

[1] R. A. CASTRO, *Regularity for the solutions of a Robin problem and some applications*, Rev. Colombiana Mat. **42**(2008), 127–144.  
 [2] R. COURANT, D. HILBERT, *Methods of mathematical physics*, Wiley, New York, 1989.  
 [3] D. DANERS, J. B. KENNEDY, *On the asymptotic behaviour of the eigenvalues of a Robin problem*, Differential Integral Equations **23**(2010), 659–669.

- [4] A. V. FILINOVSKIY, *Asymptotic behaviour of the first eigenvalue of a Robin problem*, Differ. Equ. **47**(2011), 1681–1682.
- [5] T. GIORGI, R. G. SMITS, *Monotonicity results for the principal eigenvalue of the generalized Robin problem*, Illinois J. Math. **49**(2005), 1133–1143.
- [6] A. HENROT, *Extremum problems for eigenvalues of elliptic operators*, Birkhäuser, Basel, 2006.
- [7] T. KATO, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1995.
- [8] V. A. KONDRATIEV, E. M. LANDIS, *Qualitative theory of second order linear partial differential equations*, Itogi nauki i tehniki, Ser. Sovrem. Probl. Mat., Fundam. Napravleniya. Moscow **32**(1988), 99–215, in Russian.
- [9] A. A. LACEY, J. R. OCKENDON, J. SABINA, *Multidimensional reaction-diffusion equations with nonlinear boundary conditions*, SIAM J. Appl. Math. **58**(1998), 1622–1647.
- [10] Y. LOU, M. ZHU, *A singularly perturbed linear eigenvalue problem in  $C^1$  domains*, Pacific J. Math. **214**(2004), 323–334.
- [11] V. P. MIKHAĬLOV, *Partial Differential Equations*, Nauka, Moscow, 1983., in Russian.
- [12] R. SPERB, *Untere und obere Schranken für den tiefsten Eigenwert der elastisch gestützten Membran*, Zeitschrift Angew. Math. Phys. **23**(1972), 231–244.