

Estimating Surface Normals in Noisy Point Cloud Data

Niloy J. Mitra
Stanford Graphics Laboratory
Stanford University
CA, 94305
niloy@stanford.edu

An Nguyen
Stanford Graphics Laboratory
Stanford University
CA, 94305
anguyen@cs.stanford.edu

ABSTRACT

In this paper we describe and analyze a method based on local least square fitting for estimating the normals at all sample points of a point cloud data (PCD) set, in the presence of noise. We study the effects of neighborhood size, curvature, sampling density, and noise on the normal estimation when the PCD is sampled from a smooth curve in \mathbb{R}^2 or a smooth surface in \mathbb{R}^3 and noise is added. The analysis allows us to find the optimal neighborhood size using other local information from the PCD. Experimental results are also provided.

Categories and Subject Descriptors

I.3.5 [**Computing Methodologies**]: Computer Graphics Computational Geometry and Object Modeling [Curve, surface, solid, and object representations]

Keywords

normal estimation, noisy data, eigen analysis, neighborhood size estimation

1. INTRODUCTION

Modern range sensing technology enables us to make detailed scans of complex objects generating point cloud data (PCD) consisting of millions of points. The data acquired is usually distorted by noise arising out of various physical measurement processes and limitations of the acquisition technology.

The traditional way to use PCD is to reconstruct the underlying surface model represented by the PCD, for example as a triangle mesh, and then apply well known methods on that underlying manifold model. However, when the size of the PCD is large, such methods may be expensive. To do surface reconstruction on a PCD, one would first need to filter out the noise from the PCD, usually by some smoothing filter [12]. Such a process may remove sharp features,

however, which may be undesirable. A reconstruction algorithm such as those in [2, 4, 8] then computes a mesh that approximates the noise free PCD. Both the smoothing and the surface reconstruction processes may be computationally expensive. For certain applications like rendering or visualization, such a computation is often unnecessary and direct rendering of PCD has been investigated by the graphics community [14, 16].

Alexa et al. [1] and Pauly et al. [16] have proposed to use PCD as a new modeling primitive. Algorithms running directly on PCD often require information about the normal at each of the points. For example, normals are used in rendering PCD, making visibility computation, answering inside-outside queries, etc. Also some curve (or surface) reconstruction algorithms, as in [6], need to have the normal estimates as a part of the input data.

The normal estimation problem has been studied by various communities such as computer graphics, image processing, and mathematics, but mostly in the case of manifold representations of the surface. We would like to estimate the normal at each point in a PCD, given to us only as an unstructured set of points sampled from a smooth curve in \mathbb{R}^2 or a smooth surface in \mathbb{R}^3 and without any additional manifold structure.

Hoppe et al. [11] proposed an algorithm where the normal at each point is estimated as the normal to the fitting plane obtained by applying the total least square method to the k nearest neighbors of the point. This method is robust in the presence of noise due to the inherent low pass filtering. In this algorithm, the value of k is a parameter and is chosen manually based on visual inspection of the computed estimates of the normals, and different trial values of k may be needed before a good selection of k is found. Furthermore, the same value of k is used for normal estimation at all points in the PCD.

We note that the accuracy of the normal estimation using a total least square method depends on (1) the noise in the PCD, (2) the curvature of the underlying manifold, (3) the density and the distribution of the samples, and (4) the neighborhood size used in the estimation process. In this paper, we make precise such dependencies and study the contribution of each of these factors on the normal estimation process. This analysis allows us to find the optimal neighborhood size to be used in the method. The neighborhood size can be computed adaptively at each point based on its local information, given some estimates about the noise, the local sampling density, and bounds on the local curvature. The computational complexity of estimating all

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

SoCG'03, June 8–10, 2003, San Diego, California, USA.
Copyright 2003 ACM 1-58113-663-3/03/0006 ...\$5.00.

normals of a PCD with m points is only $O(m \log m)$.

1.1 Related Work

In this section, we summarize some of the previous works that are related to the computation of the normal vectors of a PCD. Many current surface reconstruction algorithms [2, 4, 8] can either compute the normal as part of the reconstruction, or the normal can be trivially computed once the surface has been reconstructed. As the algorithms require that the input is noise free, a raw PCD with noise needs to go through a smoothing process before these algorithms can be applied.

The work of Hoppe et al. [11] for surface reconstruction suggests a method to compute the normals for the PCD. The normal estimate at each point is done by fitting a least square plane to its k nearest neighbors. The value of k is selected experimentally. The same approach has also been adopted by Pauly et al. [16] for local surface estimation. Higher order surfaces have been used by Welch et al. [15] for local parameterization. However, as pointed out by Amenta et al. [3] such algorithms can fail even in cases with arbitrarily dense set of samples. This problem can be resolved by assuming uniformly distributed samples which prevents errors resulting from biased fits. As noted before, all these algorithms work well even in presence of noise because of the inherent filtering effect. The success of these algorithms depends largely on selecting a suitable value for k , but usually little guidance is given on the selection of this crucial parameter.

1.2 Paper Overview

In section 2, we study the normal estimation for PCD which are samplings of curves in \mathbb{R}^2 , and the effects of different parameters on the error of that estimation process. In section 3, we derive similar results for PCD which come from a surface in \mathbb{R}^3 . In section 4, we provide some simulations to illustrate the results obtained in sections 2 and 3. We also show how to use our theoretical result on practical data. We conclude in section 5.

2. NORMAL ESTIMATION IN \mathbb{R}^2

In this section, we consider the problem of approximating the normals to a point cloud in \mathbb{R}^2 . Given a set of points, which are noisy samples of a smooth curve in \mathbb{R}^2 , we can use the following method to estimate the normal to the curve at each of the sample points. For each point O , we find all the points of the PCD inside a circle of radius r centered at O , then compute the total least square line fitting those points. The normal to the fitting line gives us an approximation to the undirected normal of the curve at O . Note that the orientation of the normals is a global property of the PCD and thus cannot be computed locally. Once all the undirected normals are computed, a standard breadth first search algorithm [11] can be applied to obtain all the normal directions in a consistent way. Through out this paper, we only consider the computation of the undirected normals.

We analyze the error of the approximation when the noise is small and the sampling density is high enough around O . Under these assumptions, which we will make precise later, the computed normal approximates well the true normal. We observe that if r is large, the neighborhood of the point cannot be well approximated by a line in the presence of curvature in the data and we may incur large error. On the

other hand, if r is small, the noise in the data can result in significant estimation error. We aim for the optimal r that strikes a balance between the errors caused by the noise and the local curvature.

2.1 Modeling

Without loss of generality, we assume that O is the origin, and the y -axis is along the normal to the curve at O . We assume that the points of the PCD in a disk of radius r around O come from a segment of the curve (a 1-D topological disk). Under this assumption, the segment of the curve near O is locally a graph of a smooth function $y = g(x)$ defined over some interval R containing the interval $[-r, r]$. We assume that the curve has a bounded curvature in R , and thus there is a constant $\kappa > 0$ such that $|g''(x)| < \kappa \quad \forall x \in R$.

Let $p_i = (x_i, y_i)$ for $1 \leq i \leq k$ be the points of the PCD that lie inside a circle of radius r centered at O . We assume the following probabilistic model for the points p_i . Assume that x_i 's are instances of a random variable X taking values within $[-r, r]$, and $y_i = g(x_i) + n_i$, where the noise terms n_i are independent instances of a random variable N . X and N are assumed to be independent. We assume that the noise N has zero mean and standard deviation σ_n , and takes values in $[-n, n]$.

Using Taylor series, there are numbers ψ_i , $1 \leq i \leq k$ such that $g(x_i) = g''(\psi_i)x_i^2/2$ with $|\psi_i| \leq |x_i| \leq r$. Let $\gamma_i = g''(\psi_i)$, then $|\gamma_i| \leq \kappa$.

Note that if κr is large, even when there is no noise in the PCD, the normal to the best fit line may not be a good approximation to the tangent as shown in Figure 1. Similarly, if σ_n/r is large and the noise is biased, this normal may not be a good approximation even if the original curve is a straight line, see Figure 2. In order to keep the normal approximation error low we assume *a priori* that κr and σ_n/r are sufficiently small.




Figure 1: Curvature causes error in the estimated normal




Figure 2: Noise causes error in the estimated normal

We assume that the data is *evenly distributed*; there is a radius $r_0 > 0$ (possibly dependent on O) so that any neighborhood of size r_0 in R contains at least 2 points of the x_i 's but no more than some small constant number of them. We observe that the number of points k inside any disk of radius r is bounded from above by $\Theta(1)r\rho$, and also is bounded from below by another $\Theta(1)r\rho$, where ρ is the sampling density of the point cloud. Here we use $\Theta(1)$ to denote some small positive constant, and for notational simplicity, different appearances of $\Theta(1)$ may denote different constants. We note that distributions satisfying the (ϵ, δ) sampling condition proposed by Dey et. al. [7] are evenly distributed.

Under the above assumptions, we would like to bound the normal estimation error and study the effects of different parameters. The analysis involves probabilistic arguments to account for the random nature of the noise.

2.2 Total Least Square Line

In this section, we briefly describe the well-known total least square method. Given a set of points p_i , $1 \leq i \leq k$, we would like to find the line $a^T x = c$, with $a^T a = 1$ such that the sum of square distances from the points p_i 's to the line is minimized. Let $f(a, c) = \frac{1}{2k} \sum_{i=1}^k (a^T p_i - c)^2 = \frac{1}{2} a^T \left(\frac{1}{k} \sum_{i=1}^k p_i p_i^T \right) a - c \bar{p}^T a + \frac{1}{2} c^2$ where $\bar{p} = \frac{1}{k} \sum_{i=1}^k p_i$. We would like to find a and c minimizing $f(a, c)$ under the constraint that $a^T a = 1$.

To solve this quadratic optimization problem, we need to solve the following system of equations:

$$\begin{aligned} \frac{\partial}{\partial a} f(a, c) = \lambda a &\Rightarrow \left(\frac{1}{k} \sum_{i=1}^k p_i p_i^T \right) a - c \bar{p} = \lambda a, \\ \frac{\partial}{\partial c} f(a, c) = 0 &\Rightarrow -\bar{p}^T a + c = 0, \end{aligned}$$

where λ is a Lagrangian multiplier. It follows that $c = \bar{p}^T a$, $\left(\frac{1}{k} \sum_{i=1}^k p_i p_i^T - \bar{p} \bar{p}^T \right) a = \lambda a$, and $f(a, c) = \frac{\lambda}{2}$. Thus λ is an eigenvalue of $M = \frac{1}{k} \sum_{i=1}^k p_i p_i^T - \bar{p} \bar{p}^T$ with a as the corresponding eigenvector. It is clear that to minimize $f(a, c)$, λ has to be the minimum eigenvalue of M . The corresponding eigenvector a is the normal to the total least square line and is our normal estimate.

Note that this approach can be generalized to higher dimensional space. The normal to the total least square fitting plane (or hyperplane) of a set of k points p_i , $1 \leq i \leq k$ in \mathbb{R}^d for $d \geq 2$ can be obtained by computing the eigenvector corresponding to the smallest eigenvalue of $M = \frac{1}{k} \sum_{i=1}^k p_i p_i^T - \bar{p} \bar{p}^T$. We observe that M can be written as $M = \frac{1}{k} \sum_{i=1}^k (p_i - \bar{p})(p_i - \bar{p})^T$ and thus it is always symmetric positive semi-definite, and has non-negative eigenvalues and non-negative diagonal.

2.3 Eigen-analysis of M

We can write the 2×2 symmetric matrix M , as defined in the previous section, as $\begin{bmatrix} m_{11} & m_{12} \\ m_{12} & m_{22} \end{bmatrix}$. Note that in absence of noise and curvature, $m_{12} = m_{22} = 0$ which means 0 is the smallest eigenvalue of M with $[0 \ 1]^T$ as the corresponding eigenvector. Under our assumption that the noise and the curvature are small, y_i 's are small, and thus m_{12} and m_{22} are small. Let $\alpha = (|m_{12}| + m_{22})/m_{11}$. We would like to estimate the smallest eigenvalue of M and its corresponding eigenvector when α is small.

Using the *Gershgorin Circle Theorem* [9], there is an eigenvalue λ_1 such that $|m_{11} - \lambda_1| \leq |m_{12}|$, and an eigenvalue λ_2 such that $|m_{22} - \lambda_2| \leq |m_{12}|$. When $\alpha \leq 1/2$, we have that $\lambda_1 \geq m_{11} - |m_{12}| \geq m_{22} + |m_{12}| \geq \lambda_2$. It follows that the two eigenvalues are distinct, and λ_2 is the smallest eigenvalue of M . Let $[v \ 1]^T$ be the eigenvector corresponding to λ_2 , then

$$\begin{aligned} \begin{bmatrix} m_{11} & m_{12} \\ m_{12} & m_{22} \end{bmatrix} \begin{bmatrix} v \\ 1 \end{bmatrix} &= \lambda_2 \begin{bmatrix} v \\ 1 \end{bmatrix}, \\ \begin{bmatrix} m_{11} - \lambda_2 \\ m_{12} \end{bmatrix} v &= - \begin{bmatrix} m_{12} \\ m_{22} - \lambda_2 \end{bmatrix}. \end{aligned}$$

Thus

$$\begin{aligned} v &= - \frac{(m_{11} - \lambda_2)m_{12} + m_{12}(m_{22} - \lambda_2)}{(m_{11} - \lambda_2)^2 + m_{12}^2}, \quad (1) \\ |v| &\leq \frac{|m_{12}|(m_{11} - \lambda_2 + |m_{12}|)}{(m_{11} - \lambda_2)^2}, \\ &\leq \frac{\alpha(1 + \alpha)}{(1 - \alpha)^2}. \end{aligned}$$

Thus, the estimation error, which is the angle between the estimated normal and the true normal (which is $[0 \ 1]^T$ in this case), is less than $\tan^{-1}(\alpha(1 + \alpha)/(1 - \alpha)^2) \approx \alpha$, for small α . Note that we could write the error explicitly in closed form, then bound it. Our approach is more complicated, though as we will show later, it can be extended to obtain the error bound for the 3D case. To bound the estimation error, we need to estimate α .

2.4 Estimating Entries of M

The assumption that the sample points are evenly distributed in the interval $[-r, r]$ implies that, given any number x in that interval, the number of points p_i 's satisfying $|x_i - x| \geq r/4$ is at least $\Theta(1)k$. It follows easily that $m_{11} = \frac{1}{k} \sum_{i=1}^k (x_i - \bar{x})^2 \geq \Theta(1)r^2$. The constant $\Theta(1)$ depends only on the distribution of the random variable X .

For the entries m_{12} and m_{22} , we use $|x_i| \leq r$ and $|y_i| \leq \kappa r^2/2 + n$ to obtain the following trivial bound:

$$\begin{aligned} |m_{12}| &= \left| \frac{1}{k} \sum_{i=1}^k x_i y_i - \frac{1}{k^2} \sum_{i=1}^k x_i \sum_{i=1}^k y_i \right| \\ &\leq 2r(\kappa r^2/2 + n), \\ m_{22} &\leq \frac{1}{k} \sum_{i=1}^k y_i^2 \\ &\leq 2((\kappa r^2/2)^2 + n^2). \end{aligned}$$

Thus,

$$\begin{aligned} \alpha &\leq \Theta(1) \left(\kappa r + \frac{n}{r} + \kappa^2 r^2 + \frac{n^2}{r^2} \right) \\ &\leq \Theta(1) \left(\kappa r + \frac{n}{r} \right). \quad (2) \end{aligned}$$

This bound illustrates the effects of r , κ and n on the error. For large values of r , the error caused by the curvature κr dominates, while for a small neighborhood the term n/r is dominating. Nevertheless, the expression depends on the absolute bound n of the noise N . This bound n can be unnecessarily large or unbounded for many distribution models of N . We would like to use our assumption on the distribution of the noise N to improve our bound on α further.

Note that

$$\begin{aligned}
|m_{12}| &= \left| \frac{1}{k} \sum_{i=1}^k x_i y_i - \frac{1}{k^2} \sum_{i=1}^k x_i \sum_{i=1}^k y_i \right| \\
&\leq \left| \frac{1}{k} \sum_{i=1}^k (\gamma_i x_i^3 / 2 + x_i n_i) \right| \\
&\quad + \left| \frac{1}{k^2} \sum_{i=1}^k x_i \sum_{i=1}^k (\gamma_i x_i^2 / 2 + n_i) \right| \\
&\leq \Theta(1) \kappa r^3 + \left| \frac{1}{k} \sum_{i=1}^k x_i n_i \right| \\
&\quad + \Theta(1) r \left(\kappa r^2 + \left| \frac{1}{k} \sum_{i=1}^k n_i \right| \right).
\end{aligned}$$

Furthermore, under the assumption that X and N are independent, we have $E[x_i n_i] = E[x_i] E[n_i] = 0$ since $E[n_i] = 0$ and $\text{Var}(x_i n_i) = \Theta(1) r^2 \sigma_n^2$ since $\text{Var}(n_i) = \sigma_n^2$. Let $\epsilon > 0$ be some small constant. Using the *Chebyshev Inequality* [13], we can show that the following bound on $|m_{12}|$ holds with probability at least $1 - \epsilon$:

$$\begin{aligned}
|m_{12}| &\leq \Theta(1) \kappa r^3 + \Theta(1) \sqrt{\frac{r^2 \sigma_n^2}{\epsilon k}} + \Theta(1) r \sqrt{\frac{\sigma_n^2}{\epsilon k}} \\
&= \Theta(1) \kappa r^3 + \Theta(1) \sqrt{\frac{r^2 \sigma_n^2}{\epsilon r \rho}} + \Theta(1) r \sqrt{\frac{\sigma_n^2}{\epsilon r \rho}} \\
&\leq \Theta(1) \kappa r^3 + \Theta(1) \sigma_n \sqrt{\frac{r}{\epsilon \rho}}. \tag{3}
\end{aligned}$$

For reasonable noise models, we also have that

$$\begin{aligned}
m_{22} &\leq \frac{1}{k} \sum_{i=1}^k 2(\gamma_i^2 x_i^4 / 4 + n_i^2) \\
&\leq \Theta(1) \kappa^2 r^4 + \Theta(1) \sigma_n^2.
\end{aligned}$$

2.5 Error Bound for the Estimated Normal

From the estimations of the entries of M , we obtain the following bound on α , with probability at least $1 - \epsilon$:

$$\alpha \leq \Theta(1) \kappa r + \Theta(1) \frac{\sigma_n}{\sqrt{\epsilon \rho r^3}} + \Theta(1) \frac{\sigma_n^2}{r^2}. \tag{4}$$

Note that this bound depends on the standard deviation σ_n of the noise N rather than its magnitude bound n .

For a given set of parameters κ , σ_n , ρ , and ϵ , we can find the optimal r that minimizes the right hand side of inequality 4. As this optimal value of r is not easily expressed in closed form, let us consider a few extreme cases.

- When there is no curvature ($\kappa = 0$) we can make the bound on α arbitrarily small by increasing r . For sufficiently large r , the bound is linear in σ_n and it decreases as $r^{-3/2}$.
- When there is no noise, we can make the error bound small by choosing r as small as possible, say $r = r_0$.
- When both noise and curvature are present, the error bound cannot be arbitrarily reduced. When the density ρ of the PCD is sufficiently high, $\alpha \leq \Theta(1) \kappa r +$

$\Theta(1) \sigma_n^2 / r^2$. The error bound is minimized when $r = \Theta(1) \sigma_n^{2/3} \kappa^{-1/3}$, in which case $\alpha \leq \Theta(1) \kappa^{2/3} \sigma_n^{2/3}$. The sufficiently high density condition on ρ can be shown to be $\rho > \Theta(1) \epsilon^{-1} \sigma_n^{-4/3} \kappa^{-1/3}$.

- When there are both noise and curvature, and the density ρ is sufficiently low, $\alpha \leq \Theta(1) \kappa r + \Theta(1) \sigma_n / \sqrt{\epsilon \rho r^3}$. The bound is smallest when $r = \Theta(1) (\sigma_n^2 / (\epsilon \rho \kappa^2))^{1/5}$, in which case, $\alpha \leq \Theta(1) (\kappa^3 \sigma_n^2 / (\epsilon \rho))^{1/5}$. The sufficiently low condition on ρ can be expressed more specifically as $\rho < \Theta(1) \epsilon^{-1} \sigma_n^{-4/3} \kappa^{-1/3}$. We would like to point out that the constant hidden in the $\Theta(1)$ notation in the ‘‘sufficiently low’’ condition is 3/4 of that in the ‘‘sufficiently high’’ condition.

3. NORMAL ESTIMATION IN \mathbb{R}^3

We can extend the results obtained for curves in \mathbb{R}^2 to surfaces in \mathbb{R}^3 . Given a point cloud obtained from a smooth 2-manifold in \mathbb{R}^3 and a point O on the surface, we can estimate the normal to the surface at O as follows: find all the points of the PCD inside a sphere of radius r centered at O , then compute the total least square plane fitting those points. The normal vector to the fitting plane is our estimate of the undirected normal at O .

Given a set of k points p_i , $1 \leq i \leq k$, let $M = \frac{1}{k} \sum_{i=1}^k p_i p_i^T - \bar{p} \bar{p}^T$, where $\bar{p} = \frac{1}{k} \sum_{i=1}^k p_i$. As pointed out in subsection 2.2, the normal to the total least square plane for this set of k points is the eigenvector corresponding to the minimum eigenvalue of the M . We would like to bound the angle between this eigenvector and the true normal to the surface.

3.1 Modeling

We model the PCD in a similar fashion as in the \mathbb{R}^2 case. We assume that O is the origin, the z-axis is the normal to the surface at O , and that the points of the PCD in the sphere of radius r around O are samples of a topological disk on the surface. Under these assumptions, we can represent the surface as the graph of a function $z = g(\underline{x})$ where $\underline{x} = [x, y]^T$. Using Taylor Theorem, we can write $g(\underline{x}) = \frac{1}{2} \underline{x}^T H \underline{x}$ where H is the Hessian of f at *some* point ψ such that $|\psi| \leq |\underline{x}|$.

We assume that the surface has bounded curvature in some neighborhood around O so that there is a $\kappa > 0$ such that the Hessian H of g satisfies $\|H\|_2 \leq \kappa$ in that neighborhood.

Write the points p_i as $p_i = (x_i, y_i, z_i) = (\underline{x}_i, z_i)$. We assume that $z_i = g(\underline{x}_i) + n_i$, where the n_i 's are independent instances of some random variable N with zero mean and standard deviation σ_n . We similarly assume that the points \underline{x}_i are *evenly distributed* in the xy -plane on a disk D of radius r centered at O , i.e. there is a radius r_0 such that any disk of size r_0 inside D contains at least 3 points among the x_i 's but no more than some small constant number of them. We also assume that the noise and the surface curvature are both small.

3.2 Eigen-analysis in \mathbb{R}^3

We write the analogous matrix $M = \begin{bmatrix} m_{11} & m_{12} & m_{13} \\ m_{12} & m_{22} & m_{23} \\ m_{13} & m_{23} & m_{33} \end{bmatrix} = \begin{bmatrix} M_{11} & M_{13} \\ M_{13}^T & m_{33} \end{bmatrix}$. As pointed out in subsection 2.2, M is sym-

metric and positive semi-definite. Under the assumptions that the noise and the curvature are small, and that the points \underline{x}_i are evenly distributed, m_{11} and m_{22} are the two dominant entries in M . We assume, without loss of generality, that $m_{11} \leq m_{22}$. Let $\alpha = (|m_{13}| + |m_{23}| + m_{33}) / (m_{11} - |m_{12}|)$. As in the \mathbb{R}^2 case, we would like to bound the angle between the computed normal and the true normal to the point cloud in term of α .

Denote by $\lambda_1 \leq \lambda_2$ the eigenvalues of the 2×2 symmetric matrix M_{11} . Using again the Gershgorin Circle Theorem, it is easy to see that $m_{11} - |m_{12}| \leq \lambda_1, \lambda_2 \leq m_{22} + |m_{12}|$.

Let λ be the smallest eigenvalue of M . From the Gershgorin Circle Theorem we have $\lambda \leq |m_{13}| + |m_{23}| + m_{33} = \alpha(m_{11} - m_{12}) \leq \alpha\lambda_1$. Let $[\underline{y}^T \ 1]^T$ be the eigenvector of M corresponding with λ . Then, as with Equation 1, we have that:

$$\begin{aligned} \underline{y} &= - \left((M_{11} - \lambda I)^2 + M_{13}M_{13}^T \right)^{-1} \\ &\quad \left((M_{11} - \lambda I)M_{13} + M_{13}(m_{33} - \lambda) \right) \\ &= - (M_{11} - \lambda I)^{-2} \left(I + (M_{11} - \lambda I)^{-2} M_{13}M_{13}^T \right)^{-1} \\ &\quad \left((M_{11} - \lambda I)M_{13} + M_{13}(m_{33} - \lambda) \right), \\ \|\underline{y}\|_2 &\leq \left\| (M_{11} - \lambda I)^{-2} \right\|_2 \times \\ &\quad \left\| \left(I + (M_{11} - \lambda I)^{-2} M_{13}M_{13}^T \right)^{-1} \right\|_2 \times \\ &\quad \left(\|(M_{11} - \lambda I)\|_2 \|M_{13}\|_2 + \|M_{13}\|_2 |m_{33} - \lambda| \right). \end{aligned}$$

Note that

$$\begin{aligned} \|(M_{11} - \lambda I)^{-2} M_{13}M_{13}^T\|_2 &\leq \|(M_{11} - \lambda I)^{-2}\|_2 \|M_{13}\|_2 \|M_{13}^T\|_2 \\ &\leq (\lambda_1 - \lambda)^{-2} (m_{13}^2 + m_{23}^2) \\ &\leq (1 - \alpha)^{-2} \alpha^2. \end{aligned}$$

Thus

$$\begin{aligned} &\left\| \left(I + (M_{11} - \lambda I)^{-2} M_{13}M_{13}^T \right)^{-1} \right\|_2 \\ &\leq \frac{1}{1 - (1 - \alpha)^{-2} \alpha^2} \leq \frac{(1 - \alpha)^2}{1 - 2\alpha}. \end{aligned}$$

It follows that

$$\begin{aligned} \|\underline{y}\|_2 &\leq \frac{1}{(1 - \alpha)^2 \lambda_1^2} \frac{(1 - \alpha)^2}{1 - 2\alpha} (\lambda_2 \alpha \lambda_1 + \alpha \lambda_1 \alpha \lambda_1) \\ &\leq \frac{\alpha(1 + \alpha)}{1 - 2\alpha} \frac{\lambda_2}{\lambda_1}. \end{aligned}$$

When α is small, the right hand side is approximately $(\lambda_2/\lambda_1)\alpha$, and thus the angle between the computed normal and the true normal, $\tan^{-1} \|\underline{y}\|_2$, is approximately bounded by $(\lambda_2/\lambda_1)\alpha \leq ((m_{22} + |m_{12}|)/(m_{11} - |m_{12}|))\alpha$,

3.3 Estimation of the entries of M

As in the \mathbb{R}^2 case, from the assumption that the samples are evenly distributed, we can show that $\Theta(1)r^2 \leq m_{11}, m_{22} \leq r^2$. We can also show that $m_{33} \leq \Theta(1)\kappa^2 r^4 + \Theta(1)\sigma_n^2$. Let ρ be the sampling density of the PCD at O , then $k = \Theta(1)\rho r^2$. Again, let $\epsilon > 0$ be some small positive number. Using the Chebyshev inequality, we can show that $m_{13}, m_{23} \leq \Theta(1)\kappa r^3 + \Theta(1)\sigma_n r / \sqrt{\epsilon k} \leq \Theta(1)\kappa r^3 + \Theta(1)\sigma_n / \sqrt{\epsilon \rho}$ with probability at least $1 - \epsilon$. For the term

m_{12} , we note that $E[x_i y_i] = 0$ and $Var(x_i y_i) = \Theta(1)r^4$, and so, by the Chebyshev inequality, $m_{12} \leq \Theta(1)r / \sqrt{\epsilon \rho}$ with probability at least $1 - \epsilon$.

3.4 Error Bound for the Estimated Normal

Let $\beta = m_{12}/m_{11}$. We restrict our analysis to the cases when β is sufficiently less than 1, say $\beta < 1/2$. This restriction simply means that the points \underline{x}_i 's are not degenerate, i.e. not all of the points \underline{x}_i 's are lying on or near any given line on the xy -plane. With this restriction, it is clear that $(\lambda_2/\lambda_1)\alpha \leq (m_{22}/m_{11})((1 + \beta)/(1 - \beta))\alpha = \Theta(1)\alpha$.

From the estimations of the entries of M , we obtain the following bound with probability at least $1 - \epsilon$:

$$\begin{aligned} \frac{\lambda_2}{\lambda_1} \alpha &\leq \Theta(1)\kappa r + \Theta(1) \frac{\sigma_n}{r^2 \sqrt{\epsilon \rho}} \\ &\quad \Theta(1)\kappa^2 r^2 + \Theta(1) \frac{\sigma_n^2}{r^2} \\ &\leq \Theta(1)\kappa r + \Theta(1) \frac{\sigma_n}{r^2 \sqrt{\epsilon \rho}} + \Theta(1) \frac{\sigma_n^2}{r^2} \end{aligned}$$

This is an approximate bound on the angle between the estimated normal and the true normal. To minimize this error bound, it is clear that we should pick

$$r = \left(\frac{1}{\kappa} \left(c_1 \frac{\sigma_n}{\sqrt{\epsilon \rho}} + c_2 \sigma_n^2 \right) \right)^{1/3}, \quad (5)$$

for some constants c_1, c_2 . The constants c_1 and c_2 are small and they depend only on the distribution of the PCD.

We notice that from the above result, when there is no noise, we should pick the radius r to be as small as possible, say $r = r_0$. When there is no curvature, the radius r should be as large as possible. When the sampling density is high, the optimal value of r that minimizes the error bound is approximately $r = \Theta(1)(\sigma_n^2/\kappa)^{1/3}$. This result is similar to that for curves in \mathbb{R}^2 , and it is not at all intuitive.

4. EXPERIMENTS

In this section, we discuss some simulations to validate our theoretical results. We then show how to use the results in obtaining a good neighborhood size for the normal computation with the least square method.

4.1 Validation

We considered a PCD whose points were noisy samples of the curves $(x, \kappa \operatorname{sgn}(x) x^2/2)$, for $x \in [-1, 1]$ for different values of κ . We estimated the normals to the curves at the origin by applying the least square method on their corresponding PCD. As the y -axis is known to be the true normal to the curves, the angles between the computed normals and the y -axis gives the estimation errors.

To obtain the PCD in our experiments, we let the sampling density ρ be 100 points per unit length, and let x be uniformly distributed in the interval $[-1, 1]$. The y -components of the data were polluted with uniformly random noise in the interval $[-n, n]$, for some value n . The standard deviation σ_n of this noise is $n/\sqrt{3}$.

Figure 3 shows the error as a function of the neighborhood size r when $n = 0.05$ for 3 different values of κ , $\kappa = 0.4, 0.8$, and 1.2. As predicted by Equation 4 for large value of r , the error increases as r increases. In the experiments, it can be seen that the error is proportional to κr for $r > 0.2$. Note

that the PCD we chose generates the worst case behavior of the error.




Figure 3: The normal estimation error increases as r increases for $r > 0.2$.

Figure 4 shows the estimation error as a function of the neighborhood size r for small r when $\kappa = 1.2$ for 3 different values of n , $n = 0.017, 0.033$, and 0.05 . We observe that the error tends to decrease as r increases for $r < 0.08$. This is expected as from Equation 4, the bound on the error is a decreasing function of r when r is small.




Figure 4: The normal estimation error decreases as n decreases and r increases for $r < 0.08$.

The dependency of the error on r for small values of r can be seen more easily in Figure 5, which shows the average of the estimation errors over 50 runs for each r .

4.2 Estimating Neighborhood Size for the Normal Computation

In this part, we used the results obtained in Section 3 to estimate the normals of a PCD. The data points in the PCD were assumed to be noisy samples of a smooth surface in \mathbb{R}^3 . This is the case, for example, for PCD obtained by range scanners. To obtain the neighborhood size for the normal computation using the least square method, we would like to use Equation 5.

We assumed that the standard deviation σ_n of the noise was given to us as part of the input. We estimated the other local parameters in Equation 5, then computed r . Note that this value of r minimizes the bound of the normal computation error, and there is no guarantee that this would minimize the error itself. The constants c_1 and c_2 depend on the




Figure 5: The average error over 50 runs exhibits a clear tendency to decrease as r increases for small r .



Figure 6: Tangent planes on the original bunny

sampling distribution of the PCD. While we could attempt to compute the exact values of c_1 and c_2 , we simply guessed the value c_1 and c_2 . The value of ϵ was fixed at 0.1.

Given a PCD, we estimated the local sampling density as follows. For a given point p in the PCD, we used the approximate nearest neighbor library ANN [5] to find the distance s from p to its k -th nearest neighbor for some small number k , $k = 15$ in our experiments. The local sampling density at p was then approximated as $\rho = k/(\pi s^2)$ samples per unit area.

To estimate the local curvature, we used the method proposed by Gumhold et al. [10]. Let p_j , $1 \leq j \leq k$ be the k nearest sample points around p , and let μ be the average distance from p to all the points p_j . We computed the best fit least square plane for those k points, and let d be the distance from p to that best fit plane. The local curvature at p can then be estimated as $\kappa = 2d/\mu^2$.

Once all the parameters were obtained, we computed the neighborhood size r using Equation 5. Note that the estimated value of r could be used to obtain a good value for k , which can be used to re-estimate the local density and the local curvature. This suggests an iterative scheme in which we repeatedly estimate the local density, the local curvature, and the neighborhood size. In our experiments, we found that 3 iterations were enough to obtain good values for all the quantities.

We still have problems with obtaining good estimates for the constants c_1 and c_2 . Fortunately, we only have to estimate the constants once for a given PCD, and we can use the same constants for many PCD with a similar point distribution. In our experiments, we used the same value for both c_1 and c_2 . This value was chosen so that the computed normals on a small region of the PCD were visually satisfactory.

Figure 6 shows the computed tangent planes for the orig-



Figure 7: Normal estimation errors for the bunny PCD with noise added. The subfigures show the points of the PCD using the pink color whenever the errors are above 10° , 8° , and 5° respectively.

inal Stanford bunny. The planes are drawn as small fixed size square patches¹. We noted that our computed normals are similar to those obtained using the *cocone* method by Amenta et al. [4].

Noisy PCD used in our experiments were obtained by adding noise to the original bunny. The x , y , and z components of the noise were chosen independently and uniformly random in the range $[-0.0005, 0.0005]$. The amplitude of this noise is comparable to the average distance between the sample points and their nearest samples.

We computed the normals of the noisy PCD, and used the angles between those normals and the normals of the original PCD as estimates of the normal computation errors. In Figure 7, we color coded the estimation errors using a convention in which the color of the square patch at a point of the PCD showed the error at that point. The color of a patch is blue when there is no error, and it gets darker as the error increases. When the error is larger than a certain threshold, the patch becomes pink. Figure 7 shows the tangent planes where the thresholds are 10° , 8° , and 5° respectively.

We ran the least square normal estimation algorithm on the bunny with different amounts of noise added to it and observed that the algorithm worked well. We also noted that the normal estimation method based on *cocone* performed poorly in the presence of noise.

5. CONCLUSIONS

We have analyzed the method of least square in estimating the normals to a point cloud data derived either from a smooth curve in \mathbb{R}^2 or a smooth surface in \mathbb{R}^3 , with noise added. In both cases, we provided theoretical bound on the maximum angle between the estimated normal and the true normal of the underlying manifold. This theoretical study allowed us to find an optimal neighborhood size to be used in the least square method.

6. ACKNOWLEDGMENTS

We would like to thank Leonidas Guibas for his suggestions, comments and encouragement. We are grateful to Tamal K. Dey for useful discussions and also for providing the software for evaluating normals using *cocone*. We also thank Marc Levoy, Ron Fedkiw for helpful discussions. We acknowledge the generous support of the Stanford Graduate Fellowship program and of NSF CARGO grant 0138456.

¹The small holes in the bunny are observable due to the fact that the patches do not cover the bunny entirely.

We also want to thank the numerous referees of the previous versions of this paper for their extremely useful suggestions.

7. REFERENCES

- [1] M. Alexa, J. Behr, D. Cohen-Or, S. Fleishman, D. Levin, and C. T. Silva. Point set surfaces. *IEEE Visualization 2001*, pages 21–28, October 2001. ISBN 0-7803-7200-x.
- [2] N. Amenta, M. Bern, and M. Kamvysselis. A new Voronoi-based surface reconstruction algorithm. *Computer Graphics*, 32(Annual Conference Series):415–421, 1998.
- [3] N. Amenta and M. W. Bern. Surface reconstruction by voronoi filtering. In *Symposium on Computational Geometry*, pages 39–48, 1998.
- [4] N. Amenta, S. Choi, T. K. Dey, and N. Leekha. A simple algorithm for homeomorphic surface reconstruction. *International Journal of Computational Geometry and Applications*, 12(1-2):125–141, 2002.
- [5] S. Arya, D. M. Mount, N. S. Netanyahu, R. Silverman, and A. Y. Wu. An optimal algorithm for approximate nearest neighbor searching fixed dimensions. *Journal of the ACM*, 45(6):891–923, 1998.
- [6] J.-D. Boissonnat and F. Cazals. Smooth surface reconstruction via natural neighbour interpolation of distance functions. In *Symposium on Computational Geometry*, pages 223–232, 2000.
- [7] T. K. Dey, J. Giesen, S. Goswami, and W. Zhao. Shape dimension and approximation from samples. In *Proc. 13th ACM-SIAM Sympos, Discrete Algorithms*, pages 772–780, 2002.
- [8] S. Funke and E. Ramos. Smooth-surface reconstruction in near-linear time, 2002.
- [9] G. Golub and C. V. Loan. *Matrix Computations*. The John Hopkins University Press, Baltimore, 1996.
- [10] S. Gumhold, X. Wang, and R. MacLeod. Feature extraction from point clouds. In *10th International Meshing Roundtable, Sandia National Laboratories*, pages 293–305, October 2001.
- [11] H. Hoppe, T. DeRose, T. Duchamp, J. McDonald, and W. Stuetzle. Surface reconstruction from unorganized points. *Computer Graphics*, 26(2):71–78, 1992.
- [12] I. Lee. Curve reconstruction from unorganized points. *Computer Aided Geometric Design*, 17:161–177, 2000.
- [13] A. Leon-Garcia. *Probability and Random Processes for Electrical Engineering*. Addison Wesley, 1994.
- [14] S. Rusinkiewicz and M. Levoy. QSPat: A multiresolution point rendering system for large meshes. In K. Akeley, editor, *Siggraph 2000, Computer Graphics Proceedings*, pages 343–352. ACM Press / ACM SIGGRAPH / Addison Wesley Longman, 2000.
- [15] W. Welch and A. Witkin. Free-form shape design using triangulated surfaces. *Computer Graphics*, 28(Annual Conference Series):247–256, 1994.
- [16] M. Zwicker, M. Pauly, O. Knoll, and M. Gross. Pointshop 3d: An interactive system for point-based surface editing. In *Proc. ACM SIGGRAPH '02, Computer Graphics Proceedings, Annual Conference Series*, 2002.