

Estimating vertebrate, benthic macroinvertebrate, and diatom taxa richness in raftable Pacific Northwest rivers for bioassessment purposes

Robert M. Hughes · Alan T. Herlihy ·
William J. Gerth · Yangdong Pan

Received: 15 November 2010 / Accepted: 8 June 2011 / Published online: 29 June 2011
© Springer Science+Business Media B.V. 2011

Abstract The number of sites sampled must be considered when determining the effort necessary for adequately assessing taxa richness in an ecosystem for bioassessment purposes; however, there have been few studies concerning the number of sites necessary for bioassessment of large rivers. We evaluated the effect of sample size (i.e., number of sites) necessary to collect vertebrate (fish and aquatic amphibians), macroinvertebrate, and diatom taxa from seven large rivers in Oregon and Washington, USA during the summers of 2006–2008. We used Monte Carlo simulation to determine the

number of sites needed to collect 90–95% of the taxa 75–95% of the time from 20 randomly located sites on each river. The river wetted widths varied from 27.8 to 126.0 m, mean substrate size varied from 1 to 10 cm, and mainstem distances sampled varied from 87 to 254 km. We sampled vertebrates at each site (i.e., 50 times the mean wetted channel width) by nearshore-raft electrofishing. We sampled benthic macroinvertebrates nearshore through the use of a 500- μ m mesh kick net at 11 systematic stations. From each site composite sample, we identified a target of 500 macroinvertebrate individuals to the lowest possible taxon, usually genus. We sampled benthic diatoms nearshore at the same 11 stations from a 12-cm² area. At each station, we sucked diatoms from soft substrate into a 60-ml syringe or brushed them off a rock and rinsed them with river water into the same jar. We counted a minimum of 600 valves at 1,000 \times magnification for each site. We collected 120–211 diatom taxa, 98–128 macroinvertebrate taxa, and 14–33 vertebrate species per river. To collect 90–95% of the taxa 75–95% of the time that were collected at 20 sites, it was necessary to sample 11–16 randomly distributed sites for vertebrates, 13–17 sites for macroinvertebrates, and 16–18 sites for diatoms. We conclude that 12–16 randomly distributed sites are needed for cost-efficient sampling of vertebrate richness in the main stems of our study rivers, but 20 sites markedly underestimates the species richness of benthic macroinvertebrates and diatoms in those rivers.

R. M. Hughes (✉)
Department of Fisheries and Wildlife,
Oregon State University,
Corvallis, OR 97333, USA
e-mail: hughes.bob@epa.gov

R. M. Hughes
Amnis Opes Institute,
Corvallis, OR 97333, USA

R. M. Hughes
Departamento de Conservação e Manejo de Vida Silvestre,
Universidade Federal de Minas Gerais,
Belo Horizonte, Minas Gerais, Brazil

A. T. Herlihy · W. J. Gerth
Department of Fisheries and Wildlife, Oregon State
University,
Corvallis, OR 97331, USA

Y. Pan
Department of Environmental Sciences and Management,
Portland State University,
Portland, OR 97207, USA

Keywords Sampling effort · Periphyton · Benthos ·
Fish · Large unwadeable rivers · Oregon · Washington

Introduction

Because of increased state and federal interest in biological assessments of large unwadeable rivers, a number of papers have recently been published documenting the level of sampling effort needed for adequately estimating the taxa richness of a river site, particularly for fish (Cao et al. 2001; Lyons et al. 2001; Hughes et al. 2002; Dußling et al. 2004; Utrup and Fisher 2006; Eros et al. 2008). Sampling effort studies have also been conducted for assessing the site-scale taxonomic richness of benthic macroinvertebrates in rivers (Li et al. 2001; Cao et al. 2002; Fesl 2002; Flotemersch et al. 2006). Fewer studies have assessed the site-scale sampling effort for diatoms (e. g., Weilhoefer and Pan 2007).

In addition to the need for assessing site-scale sampling effort sufficiency, managers and ecologists are interested in determining the number of sites needed for adequately assessing taxa richness at the extent of entire mainstem rivers or their catchments. This is an important issue because too few sites underestimate taxa richness and too many sites increase monitoring costs unnecessarily (Hughes and Peck 2008; Kanno et al. 2009; Ligeiro et al. 2010). Also, studies of entire rivers as landscapes with emergent properties changing with the spatial extent of the observations have received increased attention in recent years (Schlosser 1991; Ward 1998; Fausch et al. 2002). Smith and Jones (2005) estimated that electrofishing 17–49 sites, each 30 times the mean wetted channel width (MWCW), were needed to collect 90% of the fish species in wadeable streams of nine Michigan watersheds. Based on their research in four wadeable Great Plains streams, Fischer and Paukert (2009) concluded that six to ten electrofishing sites of 40 MWCW were needed to detect 90% of the species in segments 20–28 km long. Blocksom et al. (2009) found that 15 electrofishing sites, each 500 m long, were needed for detecting 90% of the species found in seven Ohio River reaches that were 58–153 km long. Discontinuously distributed, patchy or rare species are the major reason for the reported levels of sampling effort needed to assess fish species richness at site, basin or river spatial extents (Angermeier and Smogor 1995; Cao et al. 2001; Hughes et al. 2002; Eros et al. 2008; Kanno et al. 2009).

Possibly because of their high diversity, fewer studies are available concerning the number of sites

needed to adequately assess macroinvertebrate richness in long river reaches or segments. Bartsch et al. (1998) concluded that 18–40 sites, depending on the indicator, were needed for assessing the upper Mississippi River. Raunio and Antilla-Huhtinen (2008) estimated that eight sites were necessary for assessing a large soft-bottomed Finnish river.

We located no rigorous studies of the number of sites need to assess periphyton (diatom) richness in mainstem rivers. Reavie et al. (2010) sampled over 100 sites in each of the Ohio (1,560 km), upper Mississippi (1,400 km), and undammed reaches of the Missouri (2,900 km) Rivers. The Delaware River Basin Commission (2007) samples 25 sites along the 320 km long river. But Miettinen (2007), Raunio and Soinen (2007), Sgro et al. (2007), and Taylor et al. (2007) sampled only 8–11 sites in rivers 115–400 km long.

Following the goals of the U.S. Environmental Protection Agency's Environmental Monitoring and Assessment Program, our objectives were to determine the single-gear sampling effort needed to collect 90–95% of the vertebrate, macroinvertebrate, and diatom taxa collected from intensive surveys of seven raftable river mainstems (each 87–254 km long) 75–95% of the time. Such percentages also have been reported sufficient for assessing fish assemblage condition through use of multimetric indices of biological integrity at individual sites (Reynolds et al. 2003; Hughes and Herlihy 2007; Maret et al. 2007). Those site-scale levels of sampling effort have also been found sufficient for assessing the dissimilarity of benthic macroinvertebrate assemblages among sites (Cao et al. 2002; Flotemersch et al. 2010). Based on our field experience and the literature, we hypothesized that seven to ten sites would be sufficient for collecting 90–95% of the vertebrate species collected from 20 sites 75–95% of the time, but 14–19 sites would be necessary for the more speciose benthic macroinvertebrate and diatom assemblages. We chose 20 sites as the upper limit for these rivers because our sample site lengths of 50 MWCW meant that with 20 samples we actually electrofished approximately half of the total length of each river mainstem. We also hypothesized that the highest and lowest vertebrate species richness would occur in the largest and smallest catchments, respectively, and that the highest and lowest benthic macroinvertebrate and diatom taxa richness would occur in

the rivers with largest and smallest mean substrate sizes, respectively.

Methods

We sampled seven large unwadeable rivers in Oregon and Washington by raft during the summers of 2006–2008 (Fig. 1; Table 1). We selected those rivers because of interests by tribes or federal or state agencies and because they lacked major dams in the mainstems. We randomly selected 20 sites on each river from the 1:100,000 scale NHDPlus digital line graph (USEPA and USGS 2006) to be unequally dispersed but not overlapping (Stevens and Olsen 2004). The rivers included three with largely forested catchments on the wet west side of the Cascade Mountains and four with largely steppe catchments on the dry east side of the Cascades. They varied in size from the Malheur (27.8 m mean width, 1.0 m mean thalweg depth, 7,847 km² catchment area) to the Willamette (126 m mean width, 2.9 m mean thalweg depth, 13,554 km² catchment area). Mean slopes of the rivers varied from 0.03% to 0.19%, percent snag incidence per site varied from 0% to 57.7%, percent catchment urbanized varied from 0.2% to 8.5% per site, and percent catchment agriculture varied from 0.4% to 21.7% per site. The mainstem distances sampled varied from 87 to 254 km for the Chehalis and Willamette, respectively (Fig. 1).

The site-scale sampling design followed that described for rivers in Hughes and Peck (2008) and Peck et al. (In Press). The site length was 50 times the MWCW, which was divided into ten equidistant sub-sites (each 5 MWCW long) and separated by 11 transects. We sampled vertebrate (fish and aquatic amphibian) assemblages by daytime raft electrofishing along alternating shorelines for two sub-sites (10 MWCW) to reduce potential bias from differentially shaded banks and to avoid excessive ferrying; we electrofished the thalweg when rapids or other obstacles necessitated it. One netter collected vertebrates as the rower maneuvered the raft downstream at a slightly greater velocity than the river. The electrofisher was a Smith-Root GPP 2.5 (Smith-Root, Vancouver, Washington, USA) operated at 30–60 pps pulsed DC and 400–1,000 V depending on conductivity. Voucher specimens were preserved in 10% formalin and confirmed at the Oregon State Univer-

sity Museum of Ichthyology (Corvallis, Oregon, USA).

At each of 11 nearshore systematic transects 5 MWCW apart (and also sampled at alternating shorelines every two transects), we sampled benthic macroinvertebrates with a 500- μ m mesh kick net with a 30 \times 30-cm opening and a bag length of 80 cm. The area sampled per transect was 0.09 m². We preserved the 11 macroinvertebrate sub-samples in 95% ethanol and, like the vertebrates, composited sub-samples into a single sample for each site, for a total sample area of 0.99 m². In our Oregon State University laboratory, we identified macroinvertebrates to the lowest practical taxonomic resolution, typically genus—except for Annelids, Arachnids, and Ostracods to class; Maxillipods to family; and Cnidaria, Platyhelminthes, Nemata, and Tardigrada to phylum—with a count goal of 500 individuals. Using an air-lift sampler in the Danube River, Schönbauer (1999) found that macroinvertebrates were distributed across the entire channel, but that diversities and abundances were greatest in the vicinity of protected areas nearshore. Cao et al. (2002) reported that for Oregon macroinvertebrates, ten composited sub-samples and 400–500 individuals were sufficient for distinguishing dissimilar assemblages when using Jaccard or Bray-Curtis similarity estimators.

We collected one benthic diatom sample from the littoral zone near each of the 11 macroinvertebrate transects. At each transect, we collected diatoms from a 12-cm² area through use of a 3.9-cm diameter PVC pipe as a template, and we combined all 11 samples into a single diatom composite sample per site, for a total sample area of 132 cm². In fine sediment locales, we sucked diatoms from the upper 1–2 mm of sediment into a 60-ml syringe and expelled the contents into a 500-ml jar. In coarse sediment areas, we brushed diatoms off rocks with a toothbrush and rinsed the sample with river water into the same jar. After sampling all 11 transects, we thoroughly mixed the contents of the jar, poured 50 ml into a centrifuge tube, and preserved the sample with 2 ml of 37% formalin.

At our Portland State University laboratory, we cleaned diatom valves with concentrated nitric acid using the Microwave Accelerated Reaction System (Model MARS[®] 5, CEM Corporation, Matthews, North Carolina, USA) following a pre-programmed




Fig. 1 River locations and survey design for seven Pacific Northwest rivers, USA

Table 1 Habitat characteristics (medians and ranges) for 20 sites in Pacific Northwest rivers, USA

Variable	Chehalis	Willamette	Umpqua	Sprague	Malheur	John Day	Okanogan
Mean width (m)	47.7	126.2	84.5	30.9	27.8	51.3	79.5
	29–65	88–236	33–134	15–55	18–47	42–72	40–104
Mean thalweg depth (m)	1.8	2.9	2.3	1.2	1.0	1.4	1.3
	0.9–4.8	1.7–18.0	0.6–4.0	0.6–2.5	0.5–1.3	0.9–1.9	0.3–2.3
Mean littoral depth (m)	0.3	1.0	0.2	0.2	0.6	0.2	0.7
	0.2–0.7	0.7–1.5	0.1–1.0	0.2–0.5	0.2–1.1	0.1–0.4	0.4–1.5
Catchment area (km ²)	2,738	13,554	8,637	3,637	7,847	15,214	18,890
	1,586–3,554	5,318–28,912	1,950–10,491	1,346–4,169	6,278–9,053	12,871–17,819	17,392–21,115
Map sinuosity	1.3	1.3	1.7	1.4	1.2	1.5	1.1
	1.0–2.6	1.1–2.6	1.0–3.6	1.1–2.5	1.0–4.1	1.0–17.8	1.0–2.7
Map slope (%)	0.04	0.04	0.08	0.03	0.14	0.19	0.03
	0.03–0.2	0.01–0.1	0.04–0.4	0.01–0.4	0.01–0.3	0.06–0.4	0.02–0.08
Log mean substrate (mm)	1.4	1.6	2.3	0.7	1.2	2.1	0.9
	–2.0–2.7	–0.4–2.9	1.3–3.4	–0.2–2.4	0.3–2.4	1.6–2.4	–0.1–1.5
Littoral sand dominant (%)	9.1	18.2	27.3	18.2	27.3	9.1	9.1
	0.0–27.3	0.0–36.4	9.1–45.5	4.5–63.6	9.1–54.	0.0–45.5	0.0–36.4
% site with snags present	57.7	47.5	20.1	3.9	2.0	0.0	39.0
	18–78	20–88	0–59	0–58	0–28	0–7	26–64
Bank canopy cover (%)	39.2	36.2	29.9	7.0	27.7	2.3	37.7
	14–70	16–58	6–44	0–34	11–54	0–9	13–64
Water temp. (°C)	20.6	19.9	23.1	21.8	21.3	22.5	20.8
	19–25	16–25	22–26	20–27	15–29	20–24	15–24
Max. July air temp. (°C)	25.4	27.2	28.7	28.8	34.0	32.3	31.3
	25–26	27–28	26–29	28–30	33–35	30–34	30–33
Total N (µg/L)	550	355	160	250	465	215	160
	240–920	90–570	130–890	200–320	270–4,830	180–260	120–240
Total P (µg/L)	69.5	59.0	49.0	77.5	283.5	31.0	26.0
	48–113	45–104	21–168	53–118	223–458	23–37	19–52
Sulfate (µeq/L)	70.6	87.4	44.7	15.0	371.9	112.3	733.0
	51–88	14–124	41–174	12–21	158–4,531	85–140	476–801
Chloride (µeq/L)	174.5	82.2	109.0	34.8	142.7	39.1	99.4
	162–209	30–127	85–403	30–46	67–787	27–46	84–122
Conductivity (µS/cm)	104.0	66.5	79.0	113.0	269.5	200.0	301.0
	92–115	43–79	67–173	101–132	131–1,120	150–220	236–332
Riparian disturb./transect	1.4	2.2	1.4	1.4	1.3	0.9	2.2
	0.4–2.3	0.6–3.6	0.8–3.2	0.2–2.4	0.5–2.6	0.3–1.7	1.5–3.0
% catchment urban	7.9	4.3	2.2	0.4	0.3	0.8	1.8
	6.7–8.5	2.2–7.0	0.6–2.8	0.3–0.4	0.2–0.7	0.8–0.9	1.4–2.4
% catchment agriculture	8.6	14.7	5.8	2.2	0.9	0.5	2.0
	7.7–9.1	3.5–21.7	1.9–6.1	1.4–2.3	0.5–3.8	0.4–1.0	0.7–2.6

digestion scheme (temperature, 180°C; pressure, 350 PSI; ramp, 15 min; hold, 15 min). We repeatedly

rinsed the digested diatoms with distilled water until the sample pH approximated 7. We mounted the

cleaned valves in NAPHRAX® to make permanent slides for taxa identification and counted a minimum of 600 valves at 1,000× magnification using a compound microscope with differential interference contrast optics. Diatom taxonomy mainly followed Krammer and Lange-Bertalot (1986, 1988, 1991a, b) and Patrick and Reimer (1966, 1975).

We measured physical habitat from a second raft in the thalweg and at the 11 systematic transects. Depths, dominant substrate, and snag incidence were measured at ten systematic points in the thalweg of each sub-site (100 total measurements per site). We measured additional physical habitat variables at each transect in a 100-m² area. The transect measurements included nearshore depths and substrates, wetted and bank full widths, bank full and incision heights, bank angle, vertebrate and benthos cover, large woody debris count, canopy density, riparian vegetation structure, and human disturbance counts.

We measured water conductivity and temperature at each transect, and at the end of the sample site, a water sample was taken, sealed, iced, and transported to the laboratory for analysis of total nitrogen and phosphorus by persulfate digestion and colorimetry, and sulfate and chloride by ion chromatography (USEPA 1987).

We estimated the minimum number of sites needed to estimate 75%, 90%, and 95% of the total river taxa richness by comparing the taxa richness of all 20 sites sampled against an increasingly greater number of sites, beginning with one randomly selected site, then two randomly selected sites, and so on. We assumed that the taxa richness of all 20 sites provided an adequate estimate of total river richness of the entire mainstem for purposes of our analyses—but they do not represent true taxa richness, especially for diatoms and macroinvertebrates (Cao et al. 2001, 2007; LaVigne et al. 2008b). We analyzed the taxa richness data through use of 1,000-run Monte Carlo analyses for each site number to obtain random samples without replacement of 1–20 study site composites. This technique is unbiased by the initial starting site, which may occur when field sampling until no new taxa are encountered (e.g., Gammon 1976; Lyons et al. 2001) or when subjectively choosing a starting site in a data set (Angermeier and Smogor 1995; Reynolds et al. 2003). For example, for one site, we randomly selected a single site without replacement from the 20 candidates and repeated this 999 times to

determine an average and range of taxa richness for one site. For two sites, we randomly selected two sites without replacement and repeated this 999 times to determine an average and range of taxa richness for two sites. This process was repeated for 1–20 site composites for each river for all three assemblages. The Monte Carlo approach avoids most of the shortcomings of rarefaction and statistical estimators (varying results with different models) as reported by Cao et al. (2001, 2007) and Hughes et al. (2002).

For most purposes, we used the mean of the 1,000 Monte Carlo simulations as the best estimate of the taxa richness in each of the 1–20 sample size composites and constructed individual river taxa accumulation curves for each biotic assemblage. We made box and whisker plots that combined the results of the seven rivers into one taxa accumulation curve per assemblage after normalizing richness among rivers by calculating it as a percent of total river richness. We also reported on the variability in the Monte Carlo analysis by presenting results for the taxa richness in each sample size composite based on the 5th and 95th percentiles of the 1,000 Monte Carlo simulations instead of the mean. These results show a type of 90% confidence interval for the taxa accumulation curves based on the random accumulation of sample sites that was expressed in the Monte Carlo analysis.

We explored the correlations between mean river environmental data and the shape of the individual river taxa accumulation curves using the mean Monte Carlo simulation value for the five sample composite expressed as percent of total river richness. We chose five sites because they represent the steepest part of the accumulation curves but were a large enough sample to be meaningful (markedly fewer sites would have little meaning and using all 20 sites has the same 100% of total value for each river). The environmental data used were means of the 20 sites at each river for field data and whole watershed values for landscape data because we were seeking to produce a single value for each assemblage in each of the seven rivers.

Results

Cumulative taxa richness continued to increase with effort for all three assemblages in all seven rivers

(Figs. 2, 3, and 4), especially for macroinvertebrates and diatoms, suggesting greater taxa richness than we were able to sample adequately with 20 sites.

Diatom taxa richness varied from 120–210 in the John Day and Sprague, respectively, and the Sprague had the highest individual site richness (62 taxa), whereas the John Day, Malheur, and Willamette had the lowest site richness (40 taxa; Table 2). Contrary to what we hypothesized, diatom taxa richness was highest in the two rivers (Sprague and Okanogan) with the smallest mean substrate size (Tables 1 and 2). Maximum sampled richness varied from 98–128 macroinvertebrate taxa for the Okanogan and Umpqua, respectively; but the Umpqua and Chehalis had the lowest and highest richness at individual sites with 23 and 62 taxa, respectively (Table 2). As hypothesized, there was a weak tendency for macroinvertebrate richness to be highest and lowest in the rivers with the largest and smallest mean substrate sizes, respectively (Tables 1 and 2). For vertebrates, maxima varied from 14–33 species in the John Day and Willamette Rivers, respectively; those two rivers also had the lowest and highest species richness at a site (3 and 24 species, respectively). Although the highest vertebrate species richness was associated with the largest catchment area, that was not the case for the lowest




Fig. 2 The mean diatom species richness from 1,000 Monte Carlo simulations for cumulative sample sizes of 1 to 20 samples collected from seven Pacific Northwest rivers, USA




Fig. 3 The mean macroinvertebrate species richness from 1,000 Monte Carlo simulations for cumulative sample sizes of 1 to 20 samples collected from seven Pacific Northwest rivers, USA

richness because the John Day catchment was the third largest (Tables 1 and 2).




Fig. 4 The mean vertebrate species richness (fish and aquatic amphibians) from 1,000 Monte Carlo simulations for cumulative sample sizes of 1 to 20 samples collected from seven Pacific Northwest rivers, USA

Table 2 Means and ranges of summer vertebrate, macroinvertebrate, and diatom richness of seven Pacific Northwest rivers, USA

River	Vertebrate mainstem richness	Vertebrate site richness mean (range)	Macroinvertebrate mainstem richness	Macroinvertebrate site richness mean (range)	Diatom mainstem richness	Diatom site richness mean (range)
Chehalis	25	12 (7–16)	115	46 (31–62)	180	46 (29–65)
Willamette	33	18 (7–24)	118	42 (32–58)	143	40 (31–62)
Umpqua	25	10 (6–16)	128	45 (23–60)	153	46 (37–62)
Sprague	16	10 (7–12)	111	41 (29–51)	211	62 (52–71)
Malheur	19	10 (7–18)	103	35 (28–43)	128	40 (30–54)
John Day	14	6 (3–8)	101	44 (31–52)	120	40 (34–53)
Okanogan	23	9 (5–12)	98	39 (33–48)	208	60 (53–68)

The percent of total river taxa richness increased with number of samples at a faster rate, but with greater variability, for vertebrates than for benthic macroinvertebrates and diatoms, as expected for a less speciose assemblage (Fig. 5). We found that 75% of the time, 90% of all vertebrate species encountered in all 20 sites could be encountered at 11 sites; to collect 95% of the total river vertebrate species, 16 sites would be needed. To collect 90% of the total river

benthic macroinvertebrate taxa 75% of the time, 13 sites were required; to collect 95% of the total river benthic macroinvertebrate taxa 75% of the time, 17 sites were needed. To collect 90% of the benthic diatom taxa collected at 20 sites 75% of the time, 16 sites were required; and to collect 95% of the diatom taxa collected at 20 sites 75% of the time, 18 sites were needed.




Fig. 5 Box and whisker plot of the combined results for the seven rivers showing percent of total river richness versus cumulative sample size based on the mean of 1,000 Monte Carlo simulations for each sample size in each river. The *line in the box* represents the median among the seven sample rivers and the boxes are the 1st and 3rd quartile and the whiskers are the minimum and maximum value. *Horizontal lines* show 75%, 90%, and 95% of total river richness

In examining variability associated with the Monte Carlo simulations and using the 95th percentile of the simulations instead of the mean, we collected 90% of the vertebrate species after accumulating two sites (Fig. 6). At the other extreme with the 5th percentile of the simulations, we collected 90% of the vertebrate species only after compositing 11 sites. For macroinvertebrates, in 95% of the Monte Carlo simulations we collected 90% of the total river taxa by accumulating nine sites, but in 5% of the simulations we collected 90% of the total taxa only after accumulating 15 sites. Similarly, in 95% of the Monte Carlo simulations, we collected 90% of the total river diatom taxa by compositing 11 sites, but in 5% of the simulations we collected 90% of the river diatom taxa only after compositing 15 sites (Fig. 6). These numbers of sites approximated the ranges of the numbers that we hypothesized: seven to ten for vertebrates and 14–19 for benthic macroinvertebrates and diatoms.

Several patterns are evident in comparing taxa richness versus physical and chemical habitat variables (Table 3). Malheur River taxa richness was associated with the greatest number of predictor variables and demonstrated the most significant correlations. No single predictor variable was highly correlated with taxa richness in all rivers for all 3




Fig. 6 Same plot as Fig. 5, but instead of plotting the mean of the 1,000 Monte Carlo simulations both the 5th and 95th percentiles of the 1,000 simulations are shown for each assemblage

assemblages. Greater mean thalweg depth and greater mean littoral depth were associated with lower taxa richness. In most cases increased thalweg substrate size and increased littoral sand tended to decrease taxa richness. Increased air or water temperature was occasionally correlated with increased taxa richness, but so was increased canopy cover (for diatoms). Increased nutrients (total nitrogen, total phosphorus) were correlated with increased vertebrate species richness, but decreased diatom and macroinvertebrate richness. Similarly, vertebrate species richness increased with increased riparian or catchment disturbance in three rivers.

Most correlations were insignificant between mean river environmental variables and the steepness of the taxa accumulation curves as indicated by the percent of total river taxa accumulated after five samples. Because the sample size was only 7 (the seven rivers) and we examined 30 correlations per assemblage, *r* had to be >0.75 to be significant at *p*<0.05 (Harrell 2001; Tabachnick and Fidell 2001). All correlations were insignificant between environmental variables

and the percent of total river macroinvertebrate taxa observed after sampling any five sites. However, there were significant correlations between percent of total river vertebrate species in five samples and watershed road density (*r*=+0.79). There were also significant correlations between percent total river diatom taxa in five samples and bank canopy density (*r*=-0.86), maximum July air temperature (*r*=+0.82), and watershed population density (*r*=-0.79). These associations suggest a relationship between river-scale disturbance and within-river sample taxa homogeneity for vertebrates and diatoms. The more homogeneous the samples are within a river, the fewer the number of samples required to accumulate all the taxa in the river.

Discussion

There is no absolute number of sample sites needed for assessing taxa richness or biological condition in a river. We found that 75% of the time, we needed 11–16 randomly selected sites to assess 90–95% of the vertebrate species collected at 20 sites. Beyond those numbers of sites, each additional site typically only added a single individual of a rarely occurring species. Kanno et al. (2009) also reported that singletons and doubletons governed the recommended sampling distance for stream and river sites. These site numbers are slightly greater than the six to ten sites found appropriate by Fischer and Paukert (2009) for wadeable Great Plains rivers and the nine to ten sites typically used by Ohio EPA for assessing point sources along nonwadeable Ohio rivers (Yoder and Smith 1999). The range in our site numbers encompass the 15 sites reported by Blocksom et al. (2009) for navigation pools on the Ohio River and the 10–56 sites used by Gammon and Simon (2000) on the Wabash River, but their site lengths were shorter than ours and they sampled less heterogeneous rivers. Our baselines of 50 MWCW of site length and 20 sampling sites per river may seem insufficient to some readers. However, LaVigne et al. (2008a, b) documented that they sufficed for detecting significantly more vertebrate species than previous surveys of the Willamette and Malheur Rivers, and we detected a previously unrecognized species in the Umpqua River (Kettratad and Markle 2010).

Table 3 Correlations between taxa richness and physical and chemical habitat variables for seven Pacific Northwest rivers, USA

Variable	Chehalis	Willamette	Umpqua	Sprague	Malheur	J. Day	Okanogan	ALL (seven)
Mean width	0.52*							
Mean thalweg depth	-0.49*	-0.47*			<i>0.52*</i>			0.36***
Mean littoral depth		-0.47*			-0.58*		-0.46*	
Catchment area				0.44*	0.89***			
Map sinuosity	-0.44*		0.60*					
Map slope	0.51*					<i>0.45*</i>		<i>0.50***</i>
Log mean size substrate		<i>-0.49*</i>			-0.68**	-0.44*		<i>0.41***</i>
Littoral sand dominant		-0.43*				<i>0.57*</i>	-0.48*	<i>0.45***</i>
% snags							-0.44*	0.52***
Bank canopy cover	<i>0.58*</i>	<i>0.69**</i>					<i>0.44*</i>	0.44***
Water temp.				-0.44*	0.53*			
Max. July air temp.		<i>0.65*</i>			-0.45*			-0.59***
Total N	<i>-0.56*</i>			0.46*	0.67**			-0.33***
Total P			0.46*		0.88***		-0.60*	0.35***
Sulfate				<i>0.60*</i>	0.83***	-0.49*		-0.35***
Chloride					-0.47*			
Conductivity			-0.46*		0.86***			
Riparian disturbance					0.85***	-0.51*		-0.56***
% urban	<i>-0.46*</i>	-0.53*	<i>-0.54*</i>	-0.51*	0.89***	<i>0.45*</i>		0.48***
% agriculture			-0.47*		0.76***	0.46*		0.71***

The bold text refers to vertebrates, the underlined refers to macroinvertebrates, and the italicized refers to diatoms. Significance: 0.05–0.002*; <0.002–0.0001**; <0.0001***; less significant correlations not shown

Much greater sampling effort than the 11 composite samples from 20 sites that we employed is needed for estimating true taxa richness of benthic macroinvertebrates in large rivers. Similar to our results from each of seven rivers from which we collected 98 to 128 taxa, Fesl (2002) reported a log-normal distribution of Chironomid species that continued increasing even after 120 grab samples and 80 species at nearshore sites in the Danube River. Because of the greater alpha and beta diversity and available microhabitats for macroinvertebrates versus vertebrates, the number of sites out of 20 that are needed to collect 90–95% of the benthic macroinvertebrate taxa 75% of the time is 13–17, respectively. Bartsch et al. (1998)

estimated 18–40 sites were needed for the upper Mississippi River and Raunio and Antilla-Huhtinen (2008) recommended eight sites, but they were studying soft-bottomed systems compared to our typically hard bottomed rivers (Table 1). Collier and Lill (2008) collected 63 macroinvertebrate taxa from the Waikato River, New Zealand, by using a kick net with the same mesh size as ours and by sampling all available habitats, but they only sampled 47 stations (versus our 220 per river) and only counted 200 individuals per sample (versus our 500 per composite sample).

Despite using site-scale sampling and processing methods similar to ours, diatom taxa richness may

have been underestimated by the 8–11 sites in rivers 115–400 km long sampled by Miettinen (2007), Raunio and Soininen (2007), Sgro et al. (2007), and Taylor et al. (2007). Miettinen (2007) did not report taxa richness, but Sgro et al. only reported 80 taxa and a site mean of 20 taxa; however, their study river was affected by metal mining. Raunio and Soininen (2007) and Taylor et al. (2007) recorded 161 and 245 diatom taxa in their studies (which exceeded the taxa richness we recorded), despite sampling half the number of sites that we did. Reavie et al. (2010) used the same site-scale periphyton protocol as we did, plus phytoplankton sampling, at over 100 sites in three great rivers and collected 410 algae taxa (diatoms and soft algae).

Because natural characteristics such as catchment area and substrate are known to affect local fish, macroinvertebrate, and diatom taxa richness in large rivers (e.g., McGarvey and Hughes 2008; Angradi et al. 2009; Biggs and Kilroy 2000), sampling effort may also need to be adjusted for such factors. However, a sample size of only seven rivers is insufficient for rigorously evaluating the relative importance of various natural variables. For example, discharge or water volume may be more important than catchment area (Oberdorff et al. 1995; McGarvey and Hughes 2008). Nonetheless, the John Day River, with the lowest vertebrate species richness, lacks dams and has a greater volume and catchment area than other rivers in this study that drain semi-arid landscapes (Fig. 1; Table 1). Clearly other factors than size and fragmentation are involved. Macroinvertebrate taxa richness followed the expected pattern of being highest in the river with the coarsest substrate. Substrate size has long been considered a key determinant of benthos richness (Hynes 1970; Allan 1995), and it incorporates aspects of current velocity, as well as temperature and oxygen concentration, which are also associated with increased invertebrate richness. In addition, we found that it was important to distribute the sampling effort along the entire mainstems of our study rivers because of taxa additions and subtractions associated with differences in temperature, nutrients, and substrate size between the upper and lower segments of each river. Ligeiro et al. (2010), working in a neotropical catchment, also reported that macroinvertebrate diversity was maximized by sampling many sites in many segments over a large area, and across diverse substrate types. Y. Pan, Portland State University, Portland, OR. unpublished

data concluded that there was a substantial tributary effect on diatom assemblages at sampling sites of some rivers. Lane et al. (2007) and Leland et al. (2001) reported significant micro- and macro-habitat effects on benthic diatoms collected from large river sites in the Ohio and San Joaquin River Basins.

Although beyond the intent of this paper, the correlations between taxa richness and environmental variables (Table 3) stimulate additional thought concerning site-scale and catchment-scale determinants of richness. Most importantly, the lack of a single set of taxa richness predictor variables for any or all three assemblages in any or all seven rivers indicates that there is no key variable for predicting riverine richness or the necessary sampling effort, at least for these rivers and at the spatial resolution at which we studied them. This is one reason we support measuring a large and diverse set of environmental variables (Hughes and Peck 2008). We believe that the greater number of higher correlations between taxa richness and predictor variables in the Malheur River was associated with the stronger disturbance gradient in that river, which also covaried with its size (LaVigne et al. 2008a; Y. Pan, Portland State University, Portland, OR. unpublished data). The negative correlation between taxa richness and mean thalweg and littoral depth may have been an artifact of decreased sampling effectiveness because deeper sites are more difficult to sample effectively (Flotemersch et al. 2010). Increased temperature, nutrients, and anthropogenic disturbance were occasionally highly significantly correlated with increased taxa richness, particularly of vertebrates. As stated above, this is partly a result of those variables covarying with river size because river size is strongly correlated with fish species richness (e.g., Fausch et al. 1984; McGarvey and Hughes 2008). The three rivers (Umpqua, Sprague, and Malheur) for which these correlations were strongest also varied markedly in size from up- to down-river (Table 1). But for all these reasons, one must be cautious when attempting to use simple taxa richness when assessing anthropogenic impacts on rivers (Moya et al. 2011; Stoddard et al. 2008; Whittier et al. 2007).

We conclude that randomization analytical procedures, such as multiple-run Monte Carlo analyses, as recommended by Flotemersch et al. (2010) and Schneck and Melo (2010) are useful for estimating sufficient sampling effort for assessing vertebrate, macroinvertebrate, and diatom taxa richness in rivers.

We encourage others to undertake similar studies of large river sampling effort because such systems are rapidly losing their vertebrate and macroinvertebrate taxa as a result of water projects and water pollution (Allan and Flecker 1993; Dynesius and Nilsson 1994; Hughes et al. 2005; Dudgeon et al. 2006) generated by increased human population and economic growth (Czech et al. 2004; Perkins 2004; Leprieur et al. 2008; Limburg et al. 2011).

Acknowledgments This research was funded by grants to Oregon State University from the U.S. Environmental Protection Agency (RM832827, B. Hill, Project Officer), National Marine Fisheries Service (AB133FO8SE3579, C. Jordan, Project Officer), and U.S. Fish and Wildlife Service (81450-7-528, M. Buettner, Project Officer). Data analyses and manuscript preparation were funded by the Fundacao de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG grant 00011/09) and the Companhia Energetica de Minas Gerais (CEMIG Projeto Peixe Vivo). Vertebrates were collected under permits from the National Marine Fisheries Service (1559), U.S. Fish and Wildlife Service (TE141832), Oregon Department of Fish and Wildlife (OR2008-4575), Washington Department of Fish and Wildlife (WA3424), and Oregon State University Institutional Animal Care and Use Committee (3430). Field work was conducted by H. LaVigne, J. Adams, T. Botsford, R. Emig, A. Farmer, B. Freese, C. Gewecke, L. Genzoli, E. Hughes, and S. Wiedemer. Chemical analyses were conducted by the Cooperative Chemical Analytical Laboratory, Forest Science Department, Oregon State University; vertebrate voucher specimens were confirmed by D. Markle; Fig. 1 and catchment values were produced by C. Johnson; and site physical habitat structure values were produced by C. Seeliger. R. Van Driesche and J. DiGuilio assisted in identifying macroinvertebrates. Critical reviews by T. Eros, V. Vono, J. Flotemersch, and two anonymous reviewers improved the submitted manuscript.

References

- Allan, J. D. (1995). *Stream ecology: structure and function of running waters*. New York: Chapman & Hall.
- Allan, J. D., & Flecker, A. S. (1993). Biodiversity conservation in running waters. *BioScience*, 43, 32–43.
- Angermeier, P. L., & Smogor, R. A. (1995). Estimating number of species and relative abundances in stream-fish communities: effects of sampling effort and discontinuous spatial distributions. *Canadian Journal of Fisheries and Aquatic Sciences*, 52, 936–949.
- Angradi, T. R., Pearson, M. S., Bolgrien, D. W., Jicha, T. M., Taylor, D. L., & Hill, B. H. (2009). Multimetric macroinvertebrate indices for mid-continent US great rivers. *Journal of the North American Benthological Society*, 28, 785–804.
- Bartsch, L. A., Richardson, W. B., & Naimo, T. J. (1998). Sampling benthic macroinvertebrates in a large flood-plain river: considerations of study design, sample size, and cost. *Environmental Monitoring and Assessment*, 52, 425–439.
- Biggs, B. J. F., & Kilroy, C. (2000). *Stream periphyton monitoring manual*. Christchurch: NIWA, New Zealand Ministry for the Environment.
- Blocksom, K., Emery, E., & Thomas, J. (2009). Sampling effort needed to estimate condition and species richness in the Ohio River, USA. *Environmental Monitoring and Assessment*, 155, 157–167.
- Cao, Y., Larsen, D. P., & Hughes, R. M. (2001). Evaluating sampling sufficiency in fish assemblage surveys: a similarity-based approach. *Canadian Journal of Fisheries and Aquatic Sciences*, 58, 1782–1793.
- Cao, Y., Larsen, D. P., Hughes, R. M., Angermeier, P. L., & Patton, T. M. (2002). Sampling effort affects multivariate comparisons of stream assemblages. *Journal of the North American Benthological Society*, 21, 701–714.
- Cao, Y., Hawkins, C. P., Larsen, D. P., & Van Sickle, J. (2007). Effects of sample standardization on mean species detectabilities and estimates of relative differences in species richness among assemblages. *The American Naturalist*, 170, 381–395.
- Collier, K. J., & Lill, A. (2008). Spatial patterns in the composition of shallow-water macroinvertebrate communities of a large New Zealand river. *New Zealand Journal of Marine and Freshwater Research*, 42, 129–141.
- Czech, B., Angermeier, P. L., Daly, H. E., Pister, E. P., & Hughes, R. M. (2004). Fish conservation, sustainable fisheries, and economic growth: no more fish stories. *Fisheries*, 29(8), 36–37.
- Delaware River Basin Commission. (2007). *DRBC Delaware River biomonitoring program 2006–2007 quality assurance project plan*. West Trenton: Delaware River Basin Commission.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Lévêque, C., et al. (2006). Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews*, 81, 163–182.
- Dußling, U., Berg, R., Klinger, H., Wolter, C., & Wilken, R.-D. (2004). Assessing the ecological status of river systems using fish assemblages. In C. Steinberg, W. Calmano, & H. Klapper (Eds.), *Handbuch angewandte limnologie, VIII-7.4, 20. Erg.Lfg. 12/04* (pp. 1–84). Landsberg am Lech: Ecomed.
- Dynesius, M., & Nilsson, C. (1994). Fragmentation and flow regulation of river systems in the northern third of the world. *Science*, 266, 753–762.
- Eros, T., Toth, B., Sevcsik, A., & Schmera, D. (2008). Comparison of fish assemblage diversity in natural and artificial rip-rap habitats in the littoral zone of a large river (River Danube, Hungary). *International Review of Hydrobiology*, 93, 88–105.
- Fausch, K. D., Karr, J. R., & Yant, P. R. (1984) Regional application of an index of biotic integrity based on stream fish communities. *Transactions of the American Fisheries Society*, 113, 39–55.
- Fausch, K. D., Torgersen, C. E., Baxter, C. V., & Li, H. W. (2002). Landscapes to riverscapes: bridging the gap between research and conservation of stream fishes. *BioScience*, 52, 483–498.

- Fesl, C. (2002). Biodiversity and resource use of larval chironomids in relation to environmental factors in a large river. *Freshwater Biology*, 47, 1065–1087.
- Fischer, J. R., & Paukert, C. P. (2009). Effects of sampling effort, assemblage similarity, and habitat heterogeneity on estimates of species richness and relative abundance of stream fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, 66, 277–290.
- Flotemersch, J. E., Blocksom, K., Hutchens, J. J., Jr., & Autrey, B. C. (2006). Development of a standardized large river bioassessment protocol (LR-BP) for macroinvertebrate assemblages. *River Research and Applications*, 22, 775–790.
- Flotemersch, J. E., Stribling, J. B., Hughes, R. M., Reynolds, L., Paul, M. J., & Wolter, C. (2010). Site length for biological assessment of boatable rivers. *River Research & Applications*, 27, 520–535.
- Gammon, J. R. (1976). *The fish populations of the middle 340 km of the Wabash River* (Water Resources Research Center Technical Report 86). West Lafayette: Purdue University.
- Gammon, J. R., & Simon, T. P. (2000). Variation in a great river index of biotic integrity over a 20-year period. *Hydrobiologia*, 422–423, 291–304.
- Harrell, F. E., Jr. (2001). *Regression modeling strategies: with applications to linear models, logistic regression, and survival analysis*. New York: Springer-Verlag.
- Hughes, R. M., & Herlihy, A. T. (2007). Electrofishing distance needed to estimate consistent index of biotic integrity (IBI) scores in raftable Oregon rivers. *Transactions of the American Fisheries Society*, 136, 135–141.
- Hughes, R. M., & Peck, D. V. (2008). Acquiring data for large aquatic resource surveys: the art of compromise among science, logistics, and reality. *Journal of the North American Benthological Society*, 27, 837–859.
- Hughes, R. M., Kaufmann, P. R., Herlihy, A. T., Intelmann, S. S., Corbett, S. C., Arbogast, M. C., et al. (2002). Electrofishing distance needed to estimate fish species richness in raftable Oregon rivers. *North American Journal of Fisheries Management*, 22, 1229–1240.
- Hughes, R. M., Rinne, J. N., & Calamusso, B. (2005). Introduction to historical changes in large river fish assemblages of the Americas. In J. N. Rinne, R. M. Hughes, & B. Calamusso (Eds.), *Historical changes in large river fish assemblages of the Americas* (pp. 1–12). Bethesda: Symposium 45, American Fisheries Society.
- Hynes, H. B. N. (1970). *The ecology of running waters*. Toronto: University of Toronto Press.
- Kanno, Y., Vokoun, J. C., Dauwalter, D. C., Hughes, R. M., Herlihy, A. T., Maret, T. R., et al. (2009). Influence of rare species on electrofishing distance–species richness relationships at stream sites. *Transactions of the American Fisheries Society*, 138, 1240–1251.
- Kettrattad, J., & Markle, D. F. (2010). Redescription of the Tye sucker, *Catostomus tsilicoosensis* (Catostomidae). *Western North American Naturalist*, 70, 273–287.
- Krammer, K., & Lange-Bertalot, H. (1986). *Bacillariophyceae, teil 1. Naviculaceae*. Heidelberg: Spektrum Akademischer Verlag.
- Krammer, K., & Lange-Bertalot, H. (1988). *Bacillariophyceae, teil 2. Epithemiaceae, Bacillariophyceae, Surirellaceae*. Heidelberg: Spektrum Akademischer Verlag.
- Krammer, K., & Lange-Bertalot, H. (1991a). *Bacillariophyceae, teil 3. Centrales, Fragilariaceae, Eunotiaceae, Achnantheaceae*. Heidelberg: Spektrum Akademischer Verlag.
- Krammer, K., & Lange-Bertalot, H. (1991b). *Bacillariophyceae, teil 4. Achnantheaceae, kritische erganzen zu Navicula (lineolate) und Gomphonema*. Heidelberg: Spektrum Akademischer Verlag.
- Lane, C. L., Flotemersch, J. E., Blocksom, K. A., & DeCelles, S. (2007). Effect of sampling method on diatom composition for use in monitoring and assessing large river condition. *River Research and Applications*, 23, 1126–1146.
- LaVigne, H. R., Hughes, R. M., & Herlihy, A. T. (2008). Bioassessments to detect changes in Pacific Northwest river fish assemblages: a Malheur River case study. *Northwest Science*, 82, 251–258.
- LaVigne, H. R., Hughes, R. M., Wildman, R. C., Gregory, S. V., & Herlihy, A. T. (2008). Summer distribution and species richness of non-native fishes in the mainstem Willamette River, Oregon, 1944–2006. *Northwest Science*, 82, 83–93.
- Leland, H. V., Brown, L. R., & Mueller, D. K. (2001). Distribution of algae in the San Joaquin River, California, in relation to nutrient supply, salinity and other environmental factors. *Freshwater Biology*, 46, 1139–1167.
- Leprieur, F., Beauchard, O., Blanchet, S., Oberdorff, T., & Brosse, S. (2008). Fish invasions in the world's river systems: when natural processes are blurred by human activities. *Public Library of Science: Biology*, 6(2), e28. doi:10.1371/journal.pbio.0060028.
- Li, J., Herlihy, A. T., Gerth, W., Kaufmann, P. R., Gregory, S. V., Urquhart, S., et al. (2001). Variability in stream macroinvertebrates at multiple spatial scales. *Freshwater Biology*, 46, 87–97.
- Ligeiro, R., Melo, A. S., & Callisto, M. (2010). Spatial scale and the diversity of macroinvertebrates in a neotropical catchment. *Freshwater Biology*, 55, 424–435.
- Limburg, K. E., Hughes, R. M., Jackson, D. C., & Czech, B. (2011). Population increase, economic growth, and fish conservation: collision course or savvy stewardship. *Fisheries*, 36, 27–34.
- Lyons, J., Piette, R. R., & Niermeyer, K. W. (2001). Development, validation, and application of a fish-based index of biotic integrity for Wisconsin's large warmwater rivers. *Transactions of the American Fisheries Society*, 130, 1077–1094.
- Maret, T. R., Ott, D. S., & Herlihy, A. T. (2007). Electrofishing effort required to estimate biotic condition in southern Idaho rivers. *North American Journal of Fisheries Management*, 27, 1041–1052.
- McGarvey, D. J., & Hughes, R. M. (2008). Longitudinal zonation of Pacific Northwest (U.S.A.) fish assemblages and the species-discharge relationship. *Copeia*, 2008, 311–321.
- Miettinen, J. (2007). River Tenojoki: periphyton survey 2003. In S. Sivonen (Ed.), *Ecological state of the River Tenojoki—periphyton, macrozoobenthos and fish communities* (pp. 6–46). Rovaniemi: University of Lapland Printing Centre.
- Moya, N., Hughes, R. M., Dominguez, E., Gibon, F.-M., Goita, E., & Oberdorff, T. (2011). Macroinvertebrate-based multimetric predictive models for measuring the biotic condition of Bolivian streams. *Ecological Indicators*, 11, 840–847.

- Oberdorff, T., Guegan, J., & Huguency, B. (1995). Global scale patterns of fish species richness in rivers. *Ecography*, *18*, 345–352.
- Patrick, R., & Reimer, C. W. (1966). *The diatoms of the United States. Volume 1* (Monographs of the Academy of Natural Sciences of Philadelphia, no. 13). Philadelphia: Academy of Natural Sciences of Philadelphia.
- Patrick, R., & Reimer, C. W. (1975). *The diatoms of the United States. Volume 2* (Monographs of the Academy of Natural Sciences of Philadelphia, no. 13). Philadelphia: Academy of Natural Sciences of Philadelphia.
- Peck, D. V., Averill, D. K., Herlihy, A. T., Hill, B. H., Hughes, R. M., Kaufmann, P. R., et al. (Eds.). (In Press). *Environmental Monitoring and Assessment Program: surface waters western pilot study—field operations manual for nonwadeable rivers and streams*. Washington, DC: US Environmental Protection Agency.
- Perkins, J. (2004). *Confessions of an economic hit man*. New York: Penguin Books.
- Raunio, J., & Antilla-Huhtinen, M. (2008). Sample size determination for soft-bottom sampling in large rivers and comparison with the chironomid pupal exuvial technique (CPET). *River Research and Applications*, *24*, 835–843.
- Raunio, J., & Soinen, J. (2007). A practical and sensitive approach to large river periphyton monitoring: comparative performance of methods and taxonomic levels. *Boreal Environmental Research*, *12*, 55–63.
- Reavie, E. D., Jicha, T. M., Angradi, T. R., Bolgrien, D. W., & Hill, B. H. (2010). Algal assemblages for large river monitoring: comparison among biovolume, absolute and relative abundance metrics. *Ecological Indicators*, *10*, 167–177.
- Reynolds, L., Herlihy, A. T., Kaufmann, P. R., Gregory, S. V., & Hughes, R. M. (2003). Electrofishing effort requirements for assessing species richness and biotic integrity in Western Oregon streams. *North American Journal of Fisheries Management*, *23*, 450–461.
- Schlosser, I. J. (1991). Stream fish ecology: a landscape perspective. *BioScience*, *41*, 704–712.
- Schneck, F., & Melo, A. S. (2010). Reliable sample sizes for estimating similarity among macroinvertebrate assemblages in tropical streams. *Annals of Limnology – International Journal of Limnology*, *46*, 93–100.
- Schönbauer, B. (1999). Spatio-temporal patterns of macrobenthic invertebrates in a free-flowing section of the River Danube in Austria. *Archiv für Hydrobiologie*, *115*, 375–397.
- Sgro, G. V., Poole, J. B., & Johansen, J. R. (2007). Diatom species composition and ecology of the Animas River watershed, Colorado, USA. *Western North American Naturalist*, *67*, 510–519.
- Smith, K. L., & Jones, M. L. (2005). Watershed-level sampling effort requirements for determining riverine fish species composition. *Canadian Journal of Fisheries and Aquatic Sciences*, *62*, 1580–1588.
- Stevens, D. L., & Olsen, A. R. (2004). Spatially balanced sampling of natural resources. *Journal of the American Statistical Association*, *99*, 262–278.
- Stoddard, J. L., Herlihy, A. T., Peck, D. V., Hughes, R. M., Whittier, T. R., & Tarquinio, E. (2008). A process for creating multi-metric indices for large-scale aquatic surveys. *Journal of the North American Benthological Society*, *27*, 878–891.
- Tabachnick, B. G., & Fidell, L. S. (2001). *Using multivariate statistics* (4th ed.). Boston: Allyn & Bacon.
- Taylor, J. C., Janse van Vuuren, M. S., & Pieterse, A. J. H. (2007). The application and testing of diatom-based indices in the Vaal and Wilge Rivers, South Africa. *Water SA*, *33*(1), 51–59.
- USEPA. (1987). *Handbook of methods for acid deposition studies: laboratory analyses for surface water chemistry. EPA/600/4-87/026*. Washington, DC: U. S. Environmental Protection Agency.
- USEPA and USGS (2006) *NHD Plus*. US Environmental Protection Agency and US Geological Survey, Washington, DC. (available at: <http://www.epa.gov/waters/doc/rad/nhdplus.html>).
- Utrup, N. J., & Fisher, W. L. (2006). Development of a rapid bioassessment protocol for sampling fish in large prairie rivers. *North American Journal of Fisheries Management*, *26*, 714–726.
- Ward, J. V. (1998). Riverine landscapes: biodiversity patterns, disturbance regimes, and aquatic conservation. *Biological Conservation*, *83*, 269–278.
- Weilhoefer, C. L., & Pan, Y. (2007). A comparison of periphyton assemblages generated by two sampling protocols. *Journal of North American Benthological Society*, *26*, 308–318.
- Whittier, T. R., Hughes, R. M., Stoddard, J. L., Lomnický, G. A., Peck, D. V., & Herlihy, A. T. (2007). A structured approach to developing indices of biotic integrity: three examples from western USA streams and rivers. *Transactions of the American Fisheries Society*, *136*, 718–735.
- Yoder, C. O., & Smith, M. A. (1999). Using fish assemblages in a state biological assessment and criteria program: essential concepts and considerations. In T. P. Simon (Ed.), *Assessing the sustainability and biological integrity of water resources using fish communities* (pp. 17–56). Boca Raton: CRC Press.