

*Full Length Research Paper*

# Estimation of genetic parameters for body weight at different ages in Mehraban sheep

Vali Aghaali Gamasae<sup>1</sup>, Seyed Hasan Hafezian<sup>1,\*</sup>, Ahmad Ahmadi<sup>2</sup>, Hasan Baneh<sup>1</sup>, Ayyoob Farhadi<sup>1</sup> and Alborz Mohamadi<sup>3</sup>

<sup>1</sup>Department of Animal Science, Faculty of Animal and Fishery Sciences, Agricultural Science and Natural Resources, University of Sari, Iran.

<sup>2</sup>Department of Animal Science, Faculty of Agriculture, Bu Ali Sina University, Hamadan, Iran.

<sup>3</sup>Department of Animal Science, Faculty of Agriculture, Islamic Azad University, Yasouj branch, Iran.

Accepted 17 May, 2010

The objective of the present study is to estimate genetic parameters of birth weight (BW, n = 3005), weaning weight (WW, n = 2800), 6 months weight (6 MW, n = 2600), 9 months weight (9 MW, n = 1990) and yearling weight (YW, n = 1450) of Mehraban sheep, collected during 1995 - 2007 at Mehraban sheep Breeding Station in Hamedan province, Iran. (Co)variance components and genetic parameters were estimated with univariate and multivariate animal model using restricted maximum likelihood (REML) procedure. Effect of herd, lamb's sex, and year of birth were significant on all traits ( $P < 0.05$ ). The estimates of direct heritability for BW, WW, 6MW, 9MW and YW were  $0.30 \pm 0.05$ ,  $0.30 \pm 0.04$ ,  $0.35 \pm 0.05$ ,  $0.37 \pm 0.04$  and  $0.43 \pm 0.04$  respectively. Maternal heritability estimates for mentioned traits were  $0.17 \pm 0.03$ ,  $0.18 \pm 0.03$ ,  $0.14 \pm 0.03$ ,  $0.12 \pm 0.03$  and  $0.10 \pm 0.02$ , respectively. The estimates of the direct genetic correlation between BW-WW, BW-6MW, BW-9MW, BW-YW, WW-6MW, WW-9MW, WW-YW, 6MW-9MW, 6MW-YW and 9MW-YW were  $0.287 \pm 0.09$ ,  $0.305 \pm 0.09$ ,  $0.249 \pm 0.03$ ,  $0.136 \pm 0.07$ ,  $0.825 \pm 0.34$ ,  $0.713 \pm 0.05$ ,  $0.845 \pm 0.52$ ,  $0.862 \pm 0.06$ ,  $0.596 \pm 0.09$  and  $0.712 \pm 0.02$  respectively. The estimates of the phenotypic correlation between traits were positive and ranged from 0.152 for BW-9MW to 0.835 for 9MW-YW.

**Key words:** Mehraban sheep, heritability, genetic correlation, body weight traits.

## INTRODUCTION

Body weight and growth traits are one of the economically important traits in sheep production, specially in Iran that lamb sale is the main source of income for breeder while other products are in secondary importance. Although mutton is the most important source of protein in Iran, meat production from the sheep does not cover the increasing consumer demand (Rashidi et al., 2008). On the other hand, increase in sheep number for more meat production has been limited by low quality and quantity of forage range. Therefore enhancing meat production should be achieved by selecting the animals that have

maximum genetic merit as future generation parents. To design an efficient improvement program and genetic evaluation system for maximization response to selection for economically important traits, accurate estimates of the genetic parameters and the genetic relationships between the traits are necessary (Safari et al., 2005; Baneh, 2009).

Mehraban sheep is one of Iranian fat-tailed breed that are raised in Hamadan province, western Iran. The region's average annual rainfall is between 300 and 500 mm with mean temperature, 19°C; and average elevation, 1747 m (Bathaei and Pascal, 1998). In this province, the predominant breed is Mehraban, numbering approximately 2.1 million heads (Aghaali Gamasae, 2009). They are well adapted to the cold condition. Though Mehraban sheep have high performance of milk, wool and repro-duction efficiency, mutton is the most important productive trait in this breed. Nowadays, meat is the main source of income for producers.

\*Corresponding author. E-mail: [sh.hafezian@yahoo.com](mailto:sh.hafezian@yahoo.com).

**Abbreviations:** BW, Birth weight; WW, weaning weight; 6 MW, weight at six months of age; 9 MW, weight at nine months of age; YW, yearling weight.

**Table 1.** Descriptive data used in the analysis.

Trait	No of records	Mean	S.D	Minimum	Maximum	C.V (%)
Birth weight	3005	3.38	0.82	1.20	8.01	24.24
Weaning weight	2800	22.21	5.80	9.00	30.00	26.12
6 months weight	2600	35.73	9.29	13.10	59.00	26.00
9 months weight	1990	46.90	9.99	14.04	62.00	21.30
Yearling weight	1450	55.48	7.97	25.29	69.25	14.37

Observations have shown that only a few attempts have been made to estimate genetic parameters in Mehraban sheep with no published result on the genetic parameters estimation of body weight traits using animal models for this breed. In a previous study, the genetic parameters for growth traits in Mehraban sheep were estimated using least-squares procedure analysis by Bathaei and Pascal (1998).

The objective of the present study is to investigate the environmental effects (herd, lamb's sex, birth type and year of birth) as well as estimate the (Co)variance components and genetic parameters for birth weight (BW), weaning weight (WW), weight at six months (6 MW), weight at nine months (9 MW) and yearling weight (YW) in Mehraban sheep using animal model methodology. In addition, genetic and phenotypic correlations between traits were estimated.

## MATERIALS AND METHODS

### Data and management

Data and pedigree information of Mehraban sheep collected at Mehraban sheep Breeding Station in Hamedan Province, over a 12-years period (1995 - 2007), was used in this study. Number of records for birth weight, weaning weight, 6 months weight, 9 months weight and yearling weight were 3005, 2800, 2600, 1990 and 1450, respectively. The structure of data set used in this research are presented in Table 1.

Lambs were ear-tagged and pedigree information (animal code, sire and dam) and birth information (date of birth, lambs' sex, birth type) were registered for each lambs separately at birth time. Lambs were weighed at birth, 3, 6, 9 and 12 months of age. Ewes were exposed to the rams at about 18 months of age. Mating period for all ewes started from 15<sup>th</sup> of September and continued to the beginning of November each year. Lambing commenced from 15<sup>th</sup> of February to April. Mating was controlled and each mating groups including 10 – 15 ewes were set aside to a ram. The lambs were weaned at around 90 days of age. Flocks were grazed during the daytime and housed at night. Lambs were kept indoors and fed manually during winter.

### Statistical models

Investigation of environmental effects was carried out using general linear model (GLM) procedure of SAS software package (SAS, 1996). Statistical model utilized for investigation of studied environmental factors was:

$$y_{ijklm} = \mu + Y_i + S_k + T_l + H_m + e_{ijklm}$$

Where,  $y_{ijklm}$  is each of the records,  $\mu$  the overall mean,  $Y_i$  the effect of  $i^{\text{th}}$  birth year in 12 class (1995-2007),  $S_k$  the effect of  $k^{\text{th}}$  sex in 2 class (male and female),  $T_l$  the effect of  $l^{\text{th}}$  type of birth in 4 class (single, twin, triplet and quadruplet),  $H_m$ , the effect of  $m^{\text{th}}$  herd in 17 class and  $e_{ijklm}$  is the residual effect.

As ewes had different mating times and lambs were given birth to during the lambing period, all lambs weighed at the same day had different age. Therefore the age was used as covariate for phenotype observation of age adjustment on all traits except birth weight.

Co-variance components and corresponding genetic parameters for the various traits were estimated with a univariate and multivariate animal model. For this purpose, data were analyzed by restricted maximum likelihood (REML) method using ASREML (Gilmour et al., 2002). The following model was fitted to the data:

$$y = Xb + Z_1a + Z_2m + e \quad \text{with Cov}(a,m) = 0$$

Where,  $y$  is a vector of observations for different traits;  $b$ ,  $a$ ,  $m$  and  $e$  are vectors of fixed effects, direct additive genetic effects, maternal additive genetic effects and the residual effects, respectively.  $X$ ,  $Z_1$  and  $Z_2$  are corresponding design matrices associating the fixed effects, direct additive genetic effects and maternal additive genetic effects to vector of  $y$ .

Total heritability was calculated according to Willham (1972) as:

$$h^2_T = (\sigma^2_a + 0.5 \sigma^2_m) / \sigma^2_p$$

## RESULTS

### Environmental effects

The results from general linear model analysis for investigation of environmental effects, means and standard error for different traits are presented in Table 2. The effects of sex, year of birth and herd on the all traits were significant ( $p < 0.05$ ). Type of birth was significant on BW, WW, 6MW, 9MW, but had no effects on YW ( $p < 0.05$ ). At all ages, male lambs were significantly heavier than females ( $p < 0.05$ ).

### Heritability estimates

The estimates of variance component, direct heritability and maternal heritability for all traits are given in Table 3. Total heritability estimates of all traits were moderate,

**Table 2.** Least squares means  $\pm$  S.E. of body weight traits.

Fixed effects	Traits				
	BW	WW	6MW	9MW	YW
Sex	*	*	*	*	*
Male	3.59 <sup>a</sup> $\pm$ 0.32	23.03 <sup>a</sup> $\pm$ 0.09	39.94 <sup>a</sup> $\pm$ 0.25	49.19 <sup>a</sup> $\pm$ 0.35	57.20 <sup>a</sup> $\pm$ 0.85
Female	3.43 <sup>b</sup> $\pm$ 0.34	22.26 <sup>b</sup> $\pm$ 0.55	37.60 <sup>b</sup> $\pm$ 0.46	45.73 <sup>b</sup> $\pm$ 0.62	54.72 <sup>b</sup> $\pm$ 0.23
Type of birth	*	*	*	*	ns
Single	4.46 <sup>a</sup> $\pm$ 0.55	23.05 <sup>a</sup> $\pm$ 0.41	53.78 <sup>a</sup> $\pm$ 0.02	55.81 <sup>a</sup> $\pm$ 0.45	57.53 <sup>a</sup> $\pm$ 0.24
Twin	3.02 <sup>c</sup> $\pm$ 0.02	22.03 <sup>b</sup> $\pm$ 0.80	42.72 <sup>b</sup> $\pm$ 0.51	51.40 <sup>b</sup> $\pm$ 0.23	57.44 <sup>a</sup> $\pm$ 0.67
Triplet	3.74 <sup>b</sup> $\pm$ 0.81	21.48 <sup>b</sup> $\pm$ 0.96	38.55 <sup>c</sup> $\pm$ 0.12	48.01 <sup>c</sup> $\pm$ 0.78	55.33 <sup>a</sup> $\pm$ 0.53
Quadruplet	2.76 <sup>d</sup> $\pm$ 0.04	-	37.49 <sup>c</sup> $\pm$ 0.09	44.56 <sup>d</sup> $\pm$ 0.29	53.03 <sup>a</sup> $\pm$ 0.42
Year of birth	*	*	*	*	*
Herd	*	*	*	*	*

Means with similar letters within a column do not differ from another at  $p < 0.05$ . BW: birth weight, WW: weaning weight, 6MW: 6 months weight, 9MW: 9 months weight, YW: yearling weight, \*  $p < 0.05$ .

**Table 3.** Estimates of variance components, genetic, maternal and total heritability for body weight traits.

Trait	BW	WW	6MW	9MW	YW
$\sigma_a^2$	0.06422	4.215	5.521	8.961	12.61
$\sigma_m^2$	0.03602	2.456	2.234	2.906	2.960
$\sigma_e^2$	0.1109	7.301	8.246	12.352	13.73
$\sigma_p^2$	0.2112	13.97	16	24.22	29.30
$h_a^2$	0.30 $\pm$ 0.05	0.30 $\pm$ 0.04	0.35 $\pm$ 0.05	0.37 $\pm$ 0.04	0.43 $\pm$ 0.04
$h_m^2$	0.17 $\pm$ 0.03	0.18 $\pm$ 0.03	0.14 $\pm$ 0.03	0.12 $\pm$ 0.03	0.10 $\pm$ 0.02
$h_T^2$	0.38	0.39	0.42	0.43	0.48

BW: birth weight, WW: weaning weight, 6MW: 6 months weight, 9MW: 9 months weight, YW: yearling weight,  $\sigma_a^2$ : direct additive genetic variance,  $\sigma_m^2$ : maternal additive genetic variance,  $\sigma_e^2$ : residual variance,  $\sigma_p^2$ : phenotypic variance,  $h_a^2$ : direct heritability,  $h_m^2$ : maternal heritability,  $h_T^2$ : total heritability.

and ranged from 0.38 for BW to 0.48 for YW. Direct heritability for BW, WW, 6MW, 9MW and YW were estimated as 0.30, 0.30, 0.35, 0.37 and 0.43, respectively. Maternal heritability was lower than direct heritability for all traits. This parameter for BW, WW, 6MW, 9MW and YW were estimated as 0.17, 0.18, 0.14, 0.12, and 0.10, respectively.

### Correlation estimates

The estimates of phenotypic correlations, direct genetic correlations and maternal genetic correlations between body weight traits are shown in Table 4. All phenotypic correlation estimates were positive and ranged from 0.152 for BW-9MW to 0.835 for 9MW-YW. Direct genetic correlation estimates except for BW-YW and 9MW-YW were generally higher than those of phenotypic correlation and varied from 0.136 for BW-YW to 0.862 for 6MW-9MW. The maternal additive genetic correlations between

all studied traits were positive and ranged from 0.160 for 9MW-YW to 0.840 for 6MW-YW.

### DISCUSSION

The studied environmental factors showed significant sources of variation on growth traits in Mehraban sheep. Male lambs were heavier than female at all ages. Differences in sexual chromosomes, physiological characteristics and endocrinal system (type and measure of hormone secretion especially sexual hormones) can lead to significant influences of sex factor. Also, effect of lamb's sex on body weight traits at different ages have been reported in various breeds of sheep (Baneh and Hafezian, 2009; Rashidi et al., 2008; Mokhtari et al., 2008; Kalantar, 2003; Vatankhah and Talebi, 2008b; Dixit et al., 2001).

Birth type was significant on all traits except YW. Order of body weight traits for single and other type of birth were single > twin > triplet > quadruplet. The significant

**Table 4.** Estimates of phenotypic, direct genetic and maternal correlations between traits.

Trait 1	Trait 2	$r_p$	$r_d$	$r_m$
BW	WW	0.272±0.18	0.287±0.09	0.412±0.05
BW	6MW	0.273±0.04	0.305±0.09	0.345±0.11
BW	9MW	0.152±0.01	0.249±0.03	0.180±0.13
BW	YW	0.191±0.02	0.136±0.07	0.180±0.08
WW	6MW	0.705±0.20	0.825±0.34	0.812±0.02
WW	9MW	0.552±0.01	0.713±0.05	0.748±0.13
WW	YW	0.452±0.08	0.845±0.52	0.702±0.05
6MW	9MW	0.707±0.05	0.862±0.06	0.745±0.06
6MW	YW	0.405±0.02	0.596±0.09	0.840±0.03
9MW	YW	0.835±0.05	0.712±0.02	0.160±0.08

BW: Birth weight, WW: weaning weight, 6MW: 6 months weight, 9MW: 9 months weight, YW: yearling weight,  $r_p$ : phenotypic correlation,  $r_d$ : direct genetic correlation,  $r_m$ : maternal genetic correlation.

effect of birth type on body weight can be explained by limited uterine space during pregnancy, nutrition of dam especially during last pregnancy and competition for milk suckling between multiple birth lambs during birth to weaning.

In addition, high correlation between pre-weaning weight with 6MW and 9MW is probably the reason for significant effect of type of birth on post-weaning traits in this study.

Significant effect of birth type on body weight has been reported in other breeds such as Zandi (Kalantar, 2003), Merino (Dixit et al., 2001), Lori-Bakhtiari (Vatankhah and Talebi, 2008a, b) and Kermani (Rashidi et al., 2008). However, Mokhtari et al. (2008) reported that birth type has no significant effect on post-weaning weights in Kermani sheep.

Also, significant influences of herd and birth year on studied traits can be explained by differences in management, food availability, diseases, climatic condition (rate of rainfall, humidity and temperature that had effect on the quality and quantity of pasture forages) and raising systems in different years and different herds. Our results on birth year showed that significant effect were similar with those reported by Dixit et al. (2001), Ozcan et al. (2005), Abegaz et al. (2005), Rashidi et al. (2007), Kalantar (2003), Baneh and Hafezian (2009). In addition, the same results were shown by the researchers who investigated the effects of herd on body weight (Yazdi et al., 1998; Nourian, 2000; Naser et al., 2001; Baneh and Hafezian, 2009).

Direct heritability estimates increased with age from 0.30 at birth to 0.43 at yearling, but the estimates of maternal heritability reduced with increase in age. The direct additive heritability estimate (0.30) of BW in the present study is within the range reported from 0.04 in Kermani (Rashidi et al., 2008) and Romanov (Maria et al., 1993) breeds to 0.46 in Menz breed (Gizaw et al., 2007). The estimate of direct additive heritability of birth weight in the present study was similar to that reported by Hanford et al. (2002), for Columbia breed (0.27), Hanford

et al. (2005), for Rambouillet breed (0.27), Miraei-Ashtiani et al. (2007), for Sangesari breed (0.33) and Vatankhah and Talebi (2008 b), for Lori-Bakhtiari (0.31).

Our estimate of maternal heritability (0.17) for BW was higher than those reported for Dorper (Naser et al., 2000), Sabi (Matika et al., 2003), Horro (Abegaz et al., 2005) and Merino (Ozcan et al., 2005) breeds. This estimate is in agreement with the findings of Mousa et al. (1999) (0.17 for Compound), Bromley et al. (2000) (0.18 and 0.19 for Rambouillet and Targhee respectively), Hanford et al. (2005) (0.19 for Rambouillet) and Maxa et al. (2007) (0.17 and 0.19 for Shropshire and Danish Texel respectively). However, estimates reported by Tosh and Kemp (1994), Larsgard et al. (1998), El Fadili et al. (2000) and Miraei-Ashtiani et al. (2007) were higher than the present estimate.

Estimate of direct heritability for WW (0.30) obtained in the present study is within the range of those published in the literature, which varied from 0.07 in Polypay (Notter, 1997) and Elsenburg Dormer (Van Wyk et al., 2003) to 0.48 in Menz sheep (Gizaw et al., 2007) and is in agreement with the finding of Rashidi et al. (2008) in Kermani sheep. Lower direct heritability estimates were reported by Tosh and Kemp (1994) in Romanov sheep, Snyman et al. (1996) in Merino sheep, Larsgard et al. (1998) in Norwegian sheep, Ligda et al. (2000) in Choise sheep, Ozcan et al. (2005) in Turkish Merino sheep, Abegaz et al. (2005) in Horro sheep, Hanford et al. (2006) in Polypay sheep, Vatankhah and Talebi (2008a, b) in Lori-Bakhtiari sheep. However, estimates reported by Maria et al. (1993) in Romanov, Tosh and Kemp (1994) in Hampshire and Gizaw et al. (2007) in Menz sheep were higher than our estimates.

Also the obtained maternal heritability for WW (0.18) in this research is in agreement with those reported by Tosh and Kemp (1994) (0.19 for Hampshire), Larsgard et al. (1998) (0.17 for Norwegian) and Vatankhah and Talebi (2008b) (0.16 for Lori-Bakhtiari), However published results of estimates in Chios (Ligda et al., 2000), Columbia

(Hanford et al., 2002), Sabi (Matika et al., 2003) and Turkish Merino (Ozcan et al., 2005) were lower than our estimates.

Direct heritability estimated for 6MW (0.35) was lower than those reported by Gizaw et al. (2007) and Miraei-Ashtiani et al. (2007), but was greater than the estimates reported by Snyman et al. (1996), Abegaz et al. (2005), Vatankhah and Talebi (2008a, b). Estimate of maternal heritability for 6MW (0.14) obtained in the present study was close to those published by Snyman et al. (1996) for Merino sheep (0.10) and Miraei-Ashtiani et al. (2007) for Sangesari sheep (0.11); however, our estimate was greater than those reported by Abegaz et al. (2005), Vatankhah and Talebi (2008b).

Estimate of direct additive heritability for 9MW (0.37) in the present study is relatively high and is in the range reported by other researchers. The range of direct heritability estimate for 9MW in literature vary from 0.03 (Mokhtari et al., 2008) to 0.59 (Snyman et al., 1995). In addition, our estimate was higher than that of 0.08 reported by Miraei-Ashtiani et al. (2007) in Sangesari lambs.

Estimate of direct heritability for YW (0.43) obtained in the present study was within the range reported from 0.10 by Miraei-Ashtiani et al. (2007) to 0.58 by Snyman et al. (1995) in literature, and was higher than those published by Bahreini Behzadi et al. (2007) in Kermani sheep (0.14), by Ozcan et al. (2005) in Turkish Merino sheep (0.25) and by Abegaz et al. (2005) in Horro sheep (0.33), but lower than those of 0.56 reported by Gizaw et al. (2007) in Menz sheep.

Our estimate of maternal heritability for 9MW was 0.12. Abegaz et al. (2005) reported an estimate of 0.06 for maternal heritability for body weight at 8 months of ages in Horro sheep. Maternal heritability estimated for YW in this study (0.10) was higher than those reported by Ozcan et al. (2005) for Turkish Merino sheep (0.03) and Matika et al. (2003) for Sabi sheep (0.00).

Estimates of direct genetic correlation between traits in this study were positive and varied from 0.136 for BW-YW to 0.862 for 6MW-9MW. Our direct genetic correlation estimates between BW-WW and BW-6MW were lower than those reported by Miraei-Ashtiani et al. (2007) in Sangesari sheep (0.46 for BW-WW and 0.50 for BW-6MW). Also genetic correlation estimated for BW-9MW (0.249) and BW-YW (0.136) were low and these magnitudes were lower than that reported by Mokhtari et al. (2008) in Kermani sheep. Estimates of direct genetic correlation for WW-6MW, WW-9MW and WW-YW were relatively high and were 0.825, 0.713 and 0.845, respectively. The results published by Mokhtari et al. (2008) in Kermani sheep were higher than the present estimates. Direct genetic correlation between post-weaning traits varied from 0.596 for 6MW-YW to 0.862 for 6MW-9MW.

All phenotypic correlation estimates among studied traits were positive and changed from 0.152 for BW-9MW to 0.835 for 9MW-YW. In addition, these estimates except for BW-YW and 9MW-YW were lower than those

of the genetic correlation estimates. Phenotypic correlations between weaning and post-weaning traits were positive and relatively high estimates.

The positive maternal genetic correlations obtained between all investigated traits varied from 0.160 (for 9MW-YW) to 0.840 (for 6MW-YW).

The results of the present investigation showed that due to positive genetic and phenotypic correlation, selection for any of the studied traits can result in an increase of phenotypic magnitudes and genetic potentials for other body weight traits.

## REFERENCES

- Abegaz S, VanWyk JB, Olivier JJ (2005). Model comparisons and genetic and environmental parameter estimates of growth and the Kleiber ratio in Horro sheep, S. Afr. J. Anim. Sci. 35: 30-40.
- Aghaali Gamasae V (2009). Estimation of genetic parameter for body weight in Mehraban breed sheep. M.Sc. Thesis, University of Mazandara, Sari, Iran, p. 104.
- Bahreini Behzadi MR, Shahroudi FE, Van Vleck LD (2007). Estimates of genetic parameters for growth traits in Kermani sheep. J. Anim. Breed. Genet. 124: 296-301.
- Baneh H (2009). Estimation of genetic parameter for body weight in Ghezel breed sheep. M.Sc. Thesis, University of Mazandara, Sari, Iran, p. 74.
- Baneh H, Hafezian SH (2009). Effects of environmental factors on growth traits in Ghezel sheep. Afr. J. Biotechnol. 12: 2903-2907.
- Bathaei SS, Pascal L (1998). Genetic and phenotypic aspects of the growth curve characteristics in mehraban Iranian fat-tailed sheep small. Rumin. Res. 29: 261-269.
- Bromley CM, Snowden GD, Van Vleck LD (2000). Genetic parameters among eight, prolificacy and wool traits of Columbia, Polypay Rambouillet and Targhee sheep. J. Anim. Sci. 78: 846-858.
- Dixit SP, Dhilon JS, Sing G (2001). Genetic and non-genetic parameter estimates for growth traits of Bharat Merino lambs. Small Rumin. Res. 42: 101-104.
- El Fadili M, Michaux C, Detilleux J, Leroy PL (2000). Genetic parameters for growth traits of the Moroccan Timahdit breed of sheep. Small Rumin. Res. 37: 203-208.
- Gilmour AR, Gogel BJ, Cullis BR, Welham SJ, Thompson R (2002). ASREML User Guide, Release 1.0. VSN International Ltd., UK.
- Gizaw S, Lemma S, Komen H, Johan AM (2007). Estimates of genetic parameters and genetic trends for live weight and fleece traits in Menz sheep. Small Rumin. Res. 70:145-153.
- Hanford KJ, Van Vleck LD, Snowden GD (2002). Estimates of genetic parameters and genetic change for reproduction, weight, and wool characteristic of Columbia sheep. J. Anim. Sci. 80: 3086-3098.
- Hanford KJ, Van Vleck LD, Snowden GD (2005). Estimates of genetic parameters and genetic change for reproduction, weight, and wool characteristics of Rambouillet sheep. Small Rumin. Res. 57: 175-186.
- Hanford KJ, Van Vleck LD, Snowden GD (2006). Estimates of genetic parameters and genetic trend for reproduction, weight, and wool characteristics of Polypay sheep. Livest. Prod. Sci. 102: 72-82.
- Kalantar M (2003). Evaluation of some environmental effect on Growth traits in Zandi sheep. Agric. Res. 4: 49-58.
- Larsgard AG, Olesen I (1998). Genetic parameters for direct and maternal effects on weights and ultrasonic muscle and fat depth of lambs. Livest. Prod. Sci. 55: 273-278.
- Ligda CH, Gabriilidis G, Papadopoulos Th, Georgoudis A (2000). Investigation of direct and maternal genetic effects on birth and weaning weight of Chios lambs. Livest. Prod. Sci. 67: 75-80.
- Maria GA, Boldman KG, Van Vleck LD (1993). Estimates of variances due to direct and maternal effects for growth traits of Romanov sheep. J. Anim. Sci. 71: 845-849.
- Matika O, Van Wyk JB, Erasmus GJ, Baker RL (2003). Genetic parameter estimates in Sabi sheep. Livest. Prod. Sci. 79: 17-28.
- Maxa J, Norberg E, Berg P, Milerski M (2007). Genetic parameters for

- body weight, longissimus muscle depth and fat depth for Suffolk sheep in the Czech Republic. *Small Rumin. Res.* 72: 87-91.
- Miraei-Ashtiani SR, Seyedalian AR, Moradi Shahrabak M (2007). Variance components and heritabilities for body weight traits in Sangsari sheep, using univariate and multivariate animal models. *Small Rumin. Res.* 73: 109-114.
- Mokhtari MS, Rashidi A, Mohammadi Y (2008). Estimation of genetic parameters for post-weaning traits of Kermani sheep. 80: 22-27.
- Mousa E, Van Vleck LD, Leymaster KA (1999). Genetic parameters for growth traits for a composite terminal sire breed of sheep. *J. Anim. Sci.* 77: 1659-1665.
- Neser FWC, Erasmus GJ, Van Wyk JB (2000). Genetic parameter estimates for pre-weaning growth traits in Dorper sheep. *S. Afr. J. Anim. Sci.* 30: 99-101.
- Neser FWC, Erasmus GJ, Van Wyk JB (2001). Genetic parameter estimates for pre-weaning weight traits in Dorper sheep. *Small Rumin. Res.* 40: 197-202.
- Notter DR (1997). Genetic parameters for growth traits in Suffolk and Polypay sheep. *Livest. Prod. Sci.* 55: 205-213.
- Nourian E (2000). Estimation genetic parameters for pre-weaning in Ghezel sheep. M.Sci. Thesis. University of Tarbyat Modares, Iran, p. 96.
- Ozcan M, Ekiz B, Yilmaz A, Ceyhan A (2005). Genetic parameter estimates for lamb growth traits and greasy fleece weight at first shearing in Turkish Merino sheep. *Small Rumin. Res.* 56: 215-222.
- Rashidi A, Mokhtari MS, Safi Jahanshahi A, Mohammad Abadi MR (2008). Genetic parameter estimates of pre-weaning growth traits in Kermani sheep. *Small Rumin. Res.* 74: 165-171.
- Safari E, Fogarty NM, Gilmour AR (2005). A review of genetic parameter estimates for wool, growth, meat and reproduction traits in sheep. *Livest. Prod. Sci.* 92: 271-289.
- SAS (1996). Release 6.11, SAS Institute Inc., Cary, NC, USA.
- Snyman MA, Erasmus GJ, Van Wyk JB, Oliver JJ (1995). Direct and maternal co-variance components and heritability estimates for body weight at different ages and fleece traits in Afrino Sheep. *Livest. Prod. Sci.* 44: 229-235.
- Snyman MA, Olivier JJ, Olivier WJ (1996). Variance components and genetic parameters for body weight and fleece traits of Merino sheep in an arid environment. *S. Afr. J. Anim. Sci.* 26: 11-14.
- Tosh JJ, Kemp RA (1994). Estimation of variance components for lamb weights in three sheep populations. *J. Anim. Sci.* 72: 1184-1190.
- Van Wyk JB, Fair MD, Cloete SWP (2003). Revised models and genetic parameter estimates for production and reproduction traits in the Elsenburg Dormer sheep stud. *S. Afr. J. Anim. Sci.* 30: 213-222.
- Vatankhah M Talebi MA (2008a). Genetic parameters of body weight and fat-tail measurements in lambs. *Small Rumin Res.* 75: 1-6.
- Vatankhah M, Talebi MA (2008b). Heritability estimates and correlations between production and reproductive traits in Lori-Bakhtiari sheep in Iran. *S. Afr. J. Anim. Sci.* 38: 110-118.
- Yazdi MH, Engstrom G, Nasholm A, Johansson K, Jorjani H, Liljedahl LE (1997). Genetic parameters for lamb weight at different ages and wool production in Baluchi sheep. *J. Anim. Prod.* 65: 247-255.