



Published in final edited form as:

Menopause. 2013 June ; 20(6): . doi:10.1097/GME.0b013e31828392c4.

Estrogen Alone and Joint Symptoms in the Women's Health Initiative Randomized Trial

Rowan T. Chlebowski, MD, PhD¹, Dominic J. Cirillo, MD, PhD², Charles B. Eaton, MD³, Marcia L. Stefanick, Ph.D⁴, Mary Pettinger, MS⁵, Laura D. Carbone, MD, MS⁶, Karen C. Johnson, MD, MPH⁷, Michael S Simon, MD, MPH⁸, Nancy F. Woods, PhD, RN, FAAN⁹, and Jean Wactawski-Wende, PhD¹⁰

¹Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA

²University of Iowa Carver College of Medicine, Division of General Internal Medicine

³The Memorial Hospital of Rhode Island, Pawtucket, RI

⁴Stanford Prevention Research Center, Dept of Medicine, Stanford University

⁵Fred Hutchinson Cancer Research Center, Seattle, WA

⁶University of Tennessee Health Sciences, Center Department of Medicine, Memphis, TN

⁷University of Tennessee Health Science Center Department of Preventive Medicine Memphis, TN

⁸Karmanos Cancer Institute, Department of Oncology, Wayne State University, Detroit, MI

⁹University of Washington

¹⁰The State University of New York, Buffalo, NY

Abstract

Objectives—While joint symptoms are commonly reported after menopause, observational studies examining exogenous estrogen influence on joint symptoms provide mixed results. Against this background, estrogen alone effects on joint symptoms were examined in post hoc analyses in the Women's Health Initiative randomized, placebo-controlled clinical trial.

Methods—10,739 postmenopausal women with prior hysterectomy were randomized to receive daily oral conjugated equine estrogen (0.625 mg/d) or matching placebo. The frequency and severity of joint pain and joint swelling were assessed by questionnaire at entry and year 1 from all participants and in a random 9.9% subsample (n=1062) following years 3 and 6. Logistic regression models were used to compare frequency and severity of symptoms by randomization group. Sensitivity analyses evaluated adherence influence on symptoms.

Results—At baseline, joint pain and swelling were closely comparable in the randomization groups (about 77% with joint pain and 40% with joint swelling). After one year, joint pain

Correspondence to: Rowan T. Chlebowski, MD, PhD, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, 1124 W. Carson Street, Torrance, CA, 90502; phone: 310-222-2219; Fax: 310-320-2564; rowanchlebowski@gmail.com.

Conflicts of Interest: RTC has been a consultant for AstraZeneca, Novartis, Amgen, and received funding support from Amgen, and served on speaker's bureaus for AstraZeneca, Novartis, Amgen and Celgene. Dr. Woods reports board membership for Procter and Gamble. All other authors declare that they have no conflicts of interest.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

frequency was significantly lower in the estrogen alone compared to the placebo group (76.3% vs 79.2%, $P=0.001$) as was joint pain severity and the difference in pain between randomization groups persisted through year 3. However, joint swelling frequency was higher in the estrogen alone group (42.1% vs 39.7%, $P=0.02$). Adherence adjusted analyses strengthen the estrogen association with reduced joint pain but attenuated the estrogen association with increased joint swelling.

Conclusions—The current findings suggest that estrogen alone use in postmenopausal women results in a modest but sustained reduction in the frequency of joint pain.

Keywords

Estrogen; joint pain; joint swelling; Women's Health Initiative; postmenopausal women; randomized clinical trial

Introduction

While joint symptoms are commonly reported by women after menopause,^{1, 2} a determinant role for estrogen in the process is not established.^{3, 4} While some observational studies examining relationships between exogenous estrogen use and joint symptoms report a favorable effect,^{2, 5-8} negative studies have been reported⁹⁻¹² and no clear association has emerged.^{3, 13}

The issue of hormonal influence on joint symptoms was examined previously in the WHI trial evaluating estrogen alone use in women with prior hysterectomy where analyses compared women with no joint pain to those with moderate or severe pain. At one year, compared to placebo, a marginal effect of estrogen alone on joint pain was seen ($P<0.04$). In analyses conducted in adherent women at the end of intervention, about 5% more women in the estrogen group were free of joint pain ($P=0.001$).¹⁴ The current analyses in this WHI trial expands those observations. Joint symptoms are now evaluated for severity and serially assessed alone with joint swelling frequency and severity. The findings are presented for women assigned estrogen alone and compared to those assigned placebo at baseline and after one year (all participants) and after 3 and 6 years (in a randomly identified 9.9% subset with joint symptoms assessments) in both intent-to-treat analyses and analyses adjusted for adherence.

The study objective was to determine whether estrogen alone use favorably influences the incidence or severity of joint pain or joint swelling in postmenopausal women. The Women's Health Initiative randomized placebo-controlled clinical trial evaluating estrogen alone use in postmenopausal women with prior hysterectomy provides an opportunity to evaluate this association in a rigorous manner.

Methods

WHI estrogen alone trial

The study design and conduct of the WHI trial evaluating estrogen alone has been reported elsewhere.^{15, 16} Postmenopausal women between 50-79 years old who had previous hysterectomy with life expectancy ≥ 3 years and no prior breast cancer were entered into the randomized, double-blind, placebo-controlled trial at 40 US clinical centers. Women using hormones at baseline required a three month washout period before study entry.

A total of 10,739 postmenopausal women were randomized using a permuted-blocked algorithm to receive daily oral conjugated equine estrogen (0.625 mg/d) or matching

placebo. The influence of estrogen alone on primary disease outcomes has been reported.¹⁵⁻¹⁸ Women participating in the estrogen alone trial were invited to join an additional randomized, placebo-controlled trial evaluating daily calcium (1000 mg) plus vitamin D (400 IU) supplementation at their first or second annual follow-up visit. The influence of calcium plus vitamin D supplementation on major primary study endpoints has also been previously reported.¹⁹⁻²¹ The influence of estrogen alone on joint symptoms is the focus of the current report.

The WHI estrogen alone clinical trial had institutional review board approval from all participating institutions and written informed consent was obtained from all participants. Statistical analyses and data management was conducted at the WHI Clinical Coordinating Center.

Data Collection

At entry, information on demographics, family and medical history and lifestyle factors were obtained by self-reported questionnaires. Medication and supplement use was assessed by interviewer-administered questionnaire. A written protocol, central training of clinic staff and quality assurance visits by the WHI Coordinating Center ensured uniform data collection across centers.

Physical activity was assessed by questionnaire with information used to generate metabolic equivalent (MET) values.²² Measurements of height and weight were made by clinical personnel to permit body mass index (BMI) determination. Total daily calcium and vitamin D intake were defined as the sum of the dietary intake (assessed with a modified Block food frequency questionnaire)²³ and the self-reported intake from supplement and prescription medication.

Joint symptoms were assessed by questionnaire at entry into the trial and at the first annual visit from all participants. Additionally, joint symptoms were assessed in a random 9.9% subsample of participants at years 3 and 6 after entry in the estrogen alone randomized clinical trial. The sampling was done on the entire clinical trial population (n=68,132) with a 6-fold higher odds at selection for non-White participants. Joint pain and joint symptoms were separately assessed and categorized by presence (yes/no) and severity (mild, moderate, severe) among those with each symptom. The joint pain and swelling severity scores were calculated as an average from: 0 (none), one (mild), two (moderate) and three (severe).

Information on other clinical outcomes was collected at annual clinic visits and semi-annual contacts. Annual clinic visits included counting or weighing returned pills as an adherence measure.

Patients entered between 1993-1998. The estrogen alone and placebo intervention ended on February 29, 2004 after 7.1 years mean follow-up. Thus, all presented data through year 3 and almost all data through year 6 reflect findings during active intervention. Of the 10,739 estrogen alone trial participants, 6,176 were randomized into the calcium and vitamin D supplement trial receiving either calcium carbonate (1,000 mg as elemental calcium) with vitamin D₃ (400 IU) or matching placebo daily.

Statistical Analysis

Analyses of joint symptoms utilized all available data at each time point. The frequency and severity of joint symptoms (pain and swelling) were compared by randomization group assignment (active vs placebo). A logistic regression model was used to compare the frequency of having any symptoms compared to having no symptoms in analyses both unadjusted and adjusted for age and race/ethnicity. Similarly, the average symptom score

where a response of “none” equals zero and “severe” equals 3 was compared in unadjusted and adjusted linear regression models incorporating age, BMI, and WHI calcium and vitamin D supplementation trial participation. Score differences between baseline and follow-up were computed the same way. To help place the joint score differences in clinical context regression models examined the year-to-year change in joint pain and swelling in the placebo group.

A sensitivity analyses examined estrogen influence in participants who were adherent to study medication use by censoring follow-up 6 months after a participant became non-adherent. Non-adherence was defined as using < 80% of study pills or initiating non-protocol hormone therapy.

All analyses were done with SAS version 9.1.3. All P-values are two-sided and P-values of 0.05 or less were regarded as significant. The WHI study is registered with clinicaltrials.gov, NCT000000611.

Results

Most baseline clinical and demographic characteristics were comparable in the two randomization groups including age, body mass index, non-steroidal anti-inflammatory drug (NSAID) use, physical activity, self-reported history of rheumatoid arthritis and previous hormonal exposure. In addition, total calcium and vitamin D intakes at baseline, reflecting both dietary intake and supplement use, were similar in the two randomization groups. Finally, participation in the WHI calcium and vitamin D supplementation trial was also balanced between the randomization groups (Table 1), the latter relevant for the year 3 and year 6 results.

By the termination of study drug intervention, after 7.1 years mean follow-up, 53.8% of participants had stopped study drugs with similar frequency noted between randomization groups. In addition, 5.7% percent of estrogen alone group women and 9.1% of placebo group women had started hormone therapy outside the trial.¹⁵

Joint pain and swelling at entry into the estrogen alone trial were closely comparable in the two randomization groups with about 77% of participants reporting some joint pain and about 40% reporting joint swelling. After one year, women randomized to estrogen alone compared to placebo had joint pain significantly less frequently, (76.3% vs 79.2%, respectively P=0.001) and had significantly lower joint pain scores (1.16 ± 0.87 vs 1.22 ± 0.88 , mean \pm SD, P<0.001, respectively). In contrast, joint swelling frequency was higher in the estrogen alone group (42.1% vs 39.7%, P=0.02) as was swelling severity score (0.58 vs 0.52, P<0.001).

Serial analyses of joint symptoms also found differences between randomization groups. At year 3 in the subset of participants with joint symptom assessment, women randomized to estrogen continued to have joint pain less frequently compared to women randomized to placebo (74.2% vs 79.8%, respectively, P=0.03). In addition, women in the estrogen alone group had no change in pain score from baseline (0.01 ± 0.81) while those in the placebo group had an increase in pain score (0.15 ± 0.94 , P=0.01). Similar findings for joint pain continue through year 6 but did not achieve statistical significance. Joint swelling continued to be significantly higher through year 6 in the estrogen group (P=0.04) while the change in the joint swelling severity score from baseline was no longer statistically significant (P=0.31) (Table 2).

To put the joint pain score and joint swelling score differences in the estrogen compared to the placebo group at 1 year of 0.06 and 0.08, respectively, in clinical context, regression

models were used to assess year-to-year change in these parameters in the placebo group. The mean joint pain score increased by 0.0003 for a one year increase in age and the mean joint swelling score decreased by 0.001 for a one year increase in age.

The influence of adherence on joint symptom findings was examined by censoring joint symptoms reported six months after participants first became non-adherent (took < 80% of study drugs or started non-study hormone therapy). A stronger association between estrogen use and lower frequency of joint pain was seen, especially for findings after 1 and 3 years follow-up. At year 3, 72.5% of adherent women randomized to estrogen alone had joint pain compared to 81.7% of adherent women randomized to placebo (P=0.006) (Table 3). In contrast to joint pain where adherence adjusted analyses provided stronger incidence of favorable influence of estrogen alone, adherence adjusted analyses attenuated the estrogen effect on joint swelling. At year 3, the difference in joint swelling between estrogen alone and placebo group adherent participants was no longer statistically significant (P=0.31) (Table 3). Analyses in year 6 for both joint pain and joint swelling were hindered by limited number of adherent participants.

Discussion

In the present post hoc analyses in a randomized clinical trial setting, statistically significantly fewer women in the estrogen alone compared to the placebo group had joint pain after one and three years. In analyses adjusted for adherence, stronger favorable associations with estrogen use and reduced joint pain are seen. Thus, in a randomized clinical trial, estrogen alone use in postmenopausal women results in a modest but sustained and statistically significant reduction in joint pain. In contrast, joint swelling was more common in estrogen alone group participants but the findings were attenuated in adherence adjusted analyses.

The current report expands on prior findings in this trial¹⁴ by including information on joint pain severity, joint swelling, joint symptom severity and adding serial and adherence analyses. The statistically significant reduction in joint pain frequency in intention-to-treat analyses after 1 year on study currently reported in the estrogen alone group included all participants while the prior analyses excluded women with mild joint pain.¹⁴ To our review, no other randomized trial has described estrogen alone influence on joint symptoms. While the reduction in joint pain score with estrogen alone use were modest, they far exceeded the year-to-year increase in joint pain score seen in placebo group participants.

The apparent opposite effects of estrogen alone on joint pain (reduction) and joint swelling (increase) appears contradictory but may be related to the performance of the self-reported joint symptom measures. Self-reported joint pain has reasonable correlation with clinical and radiographic osteoarthritis measures.^{24, 25} However, the relation between self-reported joint swelling and articular change has been questioned.²⁶ Importantly, analyses adjusted for adherence strengthened the estrogen alone association with reduced joint pain but attenuated the estrogen association with increased joint swelling.

Supportive findings for a favorable influence of estrogen alone use on joint pain come from other prior analyses in this WHI randomized trial.^{27, 28} Women with prior hysterectomy randomized to estrogen alone had fewer cases of rheumatoid arthritis (25 cases of 5,076 vs 37 cases of 5,195, for estrogen alone vs placebo, respectively) but the difference was not statistically significant (HR 0.69, 95% CI 0.41-1.14, P=0.149).²⁷ Estrogen alone users in the trial were also found to have significantly fewer hip and knee joint replacements (222 cases of 5,076 vs 269 cases of 5,195 for estrogen alone vs placebo, respectively, HR 0.84, 95% CI 0.70-1.00, P=0.05).²⁸ Given that arthroplasty due to osteoarthritis is generally indicated

when pain can no longer be managed with pain medications, the findings support an association between estrogen alone use and lower frequency of joint symptoms.

In contrast to the findings in the WHI estrogen alone trial, in the WHI estrogen plus progestin trial in women with an intact uterus, there was no association seen between combined hormone therapy use and arthroplasty frequency seen.²⁸ This difference between the two WHI hormone trials reinforces the message that current findings are based on one regimen, conjugated equine estrogen, 0.625 mg/d alone in women with prior hysterectomy, and the results cannot be extrapolated to other hormone regimens or treatment durations.

Biological plausibility for an estrogen and joint pain association is provided by clinical studies of estradiol and its metabolites and osteoarthritis risk. In a study of 842 pre and perimenopausal women, women with radiographically defined osteoarthritis had estradiol concentrations in the lowest tertile (OR 1.88, 95% CI 1.07-3.51) compared to women without osteoarthritis.²⁹ More recently, significantly lower free estradiol levels were seen in both premenopausal and postmenopausal women with osteoarthritis compared to levels in healthy women.³⁰

Findings from observational studies examining relationships among joint problems including arthritis and menopausal hormone therapy have been mixed and their heterogeneity with respect to outcome measures and study populations have precluded pooling.³ Nonetheless, in a recent review, while insufficient information to support strong conclusions was acknowledged, nonetheless, evidence was felt supportive of an effect of endogenous and exogenous estrogen on joint health.³¹ While further study is warranted, the current results, seen in a randomized clinical trial setting, support a moderate effect of exogenous estrogen in mitigating joint pain. Any consideration of estrogen use for this purpose must incorporate available information on the identified risks and benefits of menopausal hormone therapy including the admonition to use the lowest dose for the shortest duration consistent with the intended therapeutic goal.^{15, 16}

Likely mechanisms mediating estrogen influence on joint pain include reduction in inflammation markers and reduction in cartilage turnover, as potential contributors to arthritis risk^{32, 33} seen in both preclinical³⁴ and clinical settings.^{35, 36} In addition, if future studies could confirm associations among cartilage turnover, joint pain and estrogen levels, a clinical model for more rapidly identifying potential intervention strategies for joint problems could result. Finally, both preclinical and clinical studies suggest estrogens may modulate pain processing pathways.¹

The current findings are of most relevance to women with limiting climacteric symptoms near the beginning of menopause who have had a prior hysterectomy and are considering estrogen alone use. Recent follow-up^{16, 18} and subgroup analyses³⁷ from this WHI randomized trial evaluating estrogen alone indicate a favorable benefit/risk balance for estrogen use for about 5 years.¹⁶ A modest, favorable effect on joint symptoms represents one additional factor for women contemplating estrogen alone use in this setting to consider.

These findings also may inform understanding of aromatase inhibitor-associated joint symptoms. Aromatase inhibitors substantially lower circulatory estrogen levels³⁸ and increase arthralgias.³⁹⁻⁴¹ The effect of exogenous estrogen to reduce joint pain frequency supports the concept that such arthralgias, at least in part, may be influenced by circulatory estrogen levels. Given the uncertainty regarding the potential influence of exogenous hormones on breast cancer recurrence,^{42, 43} estrogen alone should not be used to treat joint symptoms arising from aromatase inhibitor use in women with resected breast cancer.

Study strengths include the size of the large well characterized, ethnically diverse study population, serial joint symptom determination within the context of a randomized clinical trial using a quantitative instrument which was prospectively applied. However, joint symptoms were not primary study endpoints and the findings emerge from post hoc analyses. In addition, the joint pain and joint swelling scales used have not been compared to other instruments or formally validated.

Conclusion

Current study findings suggest that estrogen alone use in postmenopausal women with prior hysterectomy results in a modest but sustained and statistically significant reduction in joint pain.

Acknowledgments

Role of the Sponsor: The Women's Health Institute (WHI) Project Office at the National Heart, Lung, and Blood Institute (NHLBI), the Sponsor, reviewed and approved the final manuscript but had no other role in the preparation of this report. Decisions concerning study design, data collection and analysis, interpretation of the results, the preparation of the manuscript, or the decision to submit the manuscript for publication resided with committees comprised of WHI investigators that included NHLBI representatives.

WHI INVESTIGATORS

Program Office: (National Heart, Lung, and Blood Institute, Bethesda, Maryland)

Barbara Alving, Jacques Rossouw, Linda Pottern, Shari Ludlam, Joan McGowan, Nancy Geller, Leslie Ford.

Clinical Coordinating Center: (Fred Hutchinson Cancer Research Center, Seattle, WA)

Ross Prentice, Garnet Anderson, Andrea LaCroix, Ruth Patterson, Anne McTiernan, Barbara Cochrane, Julie Hunt, Lesley Tinker, Charles Kooperberg, Martin McIntosh, C. Y. Wang, Chu Chen, Deborah Bowen, Alan Kristal, Janet Stanford, Nicole Urban, Noel Weiss, Emily White; (Wake Forest University School of Medicine, Winston-Salem, NC) Sally Shumaker, Ronald Prineas, Michelle Naughton; (Medical Research Laboratories, Highland Heights, KY) Evan Stein, Peter Laskarzewski; (San Francisco Coordinating Center, San Francisco, CA) Steven R. Cummings, Michael Nevitt, Lisa Palermo; (University of Minnesota, Minneapolis, MN) Lisa Harnack; (Fisher BioServices, Rockville, MD) Frank Cammarata, Steve Lindenfesler; (University of Washington, Seattle, WA) Bruce Psaty, Susan Heckbert.

Clinical Centers: (Albert Einstein College of Medicine, Bronx, NY) Sylvia Wassertheil-Smoller, William Frishman, Judith Wylie-Rosett, David Barad, Ruth Freeman; (Baylor College of Medicine, Houston, TX) Jennifer Hays, Ronald Young, Jill Anderson, Sandy Lithgow, Paul Bray; (Brigham and Women's Hospital, Harvard Medical School, Boston, MA) JoAnn Manson, J. Michael Gaziano, Claudia Chae, Kathryn Rexrode, Caren Solomon (Brown University, Providence, RI) Annlouise R. Assaf, Carol Wheeler, Charles Eaton, Michelle Cyr; (Emory University, Atlanta, GA) Lawrence Phillips, Margaret Pedersen, Ora Strickland, Margaret Huber, Vivian Porter; (Fred Hutchinson Cancer Research Center, Seattle, WA) Shirley A.A. Beresford, Vicky M. Taylor, Nancy F. Woods, Maureen Henderson, Robyn Andersen; (George Washington University, Washington, DC) Judith Hsia, Nancy Gaba, Joao Ascensao; (Harbor-UCLA Research and Education Institute, Torrance, CA) Rowan Chlebowski, Robert Detrano, Anita Nelson, Michele Geller; (Kaiser Permanente Center for Health Research, Portland, OR) Evelyn Whitlock, Victor Stevens, Njeri Karanja; (Kaiser Permanente Division of Research, Oakland, CA) Bette Caan, Stephen Sidney, Geri Bailey Jane Hirata; (Medical College of Wisconsin, Milwaukee, WI) Jane Morley Kotchen, Vanessa Barnabei, Theodore A. Kotchen, Mary Ann C. Gilligan, Joan Neuner; (MedStar Research Institute/Howard University, Washington, DC) Barbara V. Howard, Lucile Adams-Campbell, Lawrence Lessin, Monique Rainford, Gabriel Uwaifo; (Northwestern University, Chicago/Evanston, IL) Linda Van Horn, Philip Greenland, Janardan Khandekar, Kiang Liu, Carol Rosenberg; (Rush University Medical Center, Chicago, IL) Henry Black, Lynda Powell, Ellen Mason; Martha Gulati; (Stanford Prevention Research Center, Stanford, CA) Marcia L. Stefanick, Mark A. Hlatky, Bertha Chen, Randall S. Stafford, Sally Mackey; (State University of New York at Stony Brook, Stony Brook, NY) Dorothy Lane, Iris Granek, William Lawson, Gabriel San Roman, Catherine Messina; (The Ohio State University, Columbus, OH) Rebecca Jackson, Randall Harris, Electra Paskett, W. Jerry Mysiw, Michael Blumenfeld; (University of Alabama at Birmingham, Birmingham, AL) Cora E. Lewis, Albert Oberman, James M. Shikany, Monika Safford, Mona Fouad; (University of Arizona, Tucson/Phoenix, AZ) Cyndi Thomson, Tamsen Bassford, Marcia Ko, Ana Maria Lopez, Cheryl Ritenbaugh; (University at Buffalo, Buffalo, NY) Jean Wactawski-Wende, Maurizio Trevisan, Ellen Smit, Susan Graham, June Chang; (University of

California at Davis, Sacramento, CA) John Robbins, S. Yasmeen; (University of California at Irvine, CA) F. Allan Hubbell, Gail Frank, Nathan Wong, Nancy Greep, Bradley Monk; (University of California at Los Angeles, Los Angeles, CA) Howard Judd, David Heber, Robert Elashoff; (University of California at San Diego, LaJolla/Chula Vista, CA) Robert D. Langer, Michael H. Criqui, Gregory T. Talavera, Cedric F. Garland, Matthew A. Allison; (University of Cincinnati, Cincinnati, OH) Margery Gass, Suzanne Wernke; (University of Florida, Gainesville/Jacksonville, FL) Marian Limacher, Michael Perri, Andrew Kaunitz, R. Stan Williams, Yvonne Brinson;¹ (University of Hawaii, Honolulu, HI) J. David Curb, Helen Petrovitch, Beatriz Rodriguez, Kamal Masaki, Santosh Sharma; (University of Iowa, Iowa City/Davenport, IA) Robert Wallace, James Torner, Susan Johnson, Linda Snetselaar, Jennifer Robinson; (University of Massachusetts/Fallon Clinic, Worcester, MA) Judith Ockene, Milagros Rosal, Ira Ockene, Robert Yood, Patricia Aronson; (University of Medicine and Dentistry of New Jersey, Newark, NJ) Norman Lasser, Baljinder Singh, Vera Lasser, John Kostis, Peter McGovern; (University of Miami, Miami, FL) Mary Jo O'Sullivan, Linda Parker, Timothy DeSantis, Diann Fernandez, Pat Caralis; (University of Minnesota, Minneapolis, MN) Karen L. Margolis, Richard H. Grimm, Mary F. Perron, Cynthia Bjerk, Sarah Kempainen; (University of Nevada, Reno, NV) Robert Brunner, William Graettinger, Vicki Oujevolk, Michael Bloch; (University of North Carolina, Chapel Hill, NC) Gerardo Heiss, Pamela Haines, David Ontjes, Carla Sueta, Ellen Wells; (University of Pittsburgh, Pittsburgh, PA) Lewis Kuller, Jane Cauley, N. Carole Milas; (University of Tennessee Health Science Center, Memphis, TN) Karen C. Johnson, Suzanne Satterfield, Raymond W. Ke, Stephanie Connelly, Fran Tylavsky; (University of Texas Health Science Center, San Antonio, TX) Robert Brzyski, Robert Schenken, Jose Trabal, Mercedes Rodriguez-Sifuentes, Charles Mouton; (University of Wisconsin, Madison, WI) Gloria E. Sarto, Douglas Laube, Patrick McBride, Julie Mares-Perlman, Barbara Loevinger; (Wake Forest University School of Medicine, Winston-Salem, NC) Denise Bonds, Greg Burke, Robin Crouse, Mara Vitolins, Scott Washburn; (Wayne State University School of Medicine/Hutzel Hospital, Detroit, MI) Susan Hendrix, Michael Simon, Gene McNeeley.

Former Principal Investigators and Project Officers:

Baylor College of Medicine, John Foreyt, Ph.D., Emory University, Dallas Hall, M.D., George Washington University, Valery Miller, M.D., Kaiser, Oakland, Robert Hiatt, M.D., Kaiser, Portland, Barbara Valanis, Dr.Ph., National Cancer Institute, Bethesda, Maryland, Carolyn Clifford², University of California, Irvine, Frank Meyskens, Jr., M.D., University of Cincinnati, James Liu, M.D., Nelson Watts, University of Miami, Marianna Baum, Ph.D., University of Minnesota, Richard Grimm, M.D., University of Nevada, Sandra Daugherty, M.D.², University of North Carolina, Chapel Hill, David Sheps, M.D., Barbara Hulka, M.D., University of Tennessee, Memphis, William Applegate, M.D., University of Wisconsin, Catherine Allen, Ph.D.²

Funding/Support: The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26, 42129-32, and 44221.

References

1. Magliano M. Menopausal arthralgia: Fact or fiction. *Maturitas*. 2010; 67:29–33. [PubMed: 20537472]
2. Szoek CE, Circuttini FM, Guthrie JR, Dennerstein L. The relationship of reports of aches and joint pains to the menopausal transition: a longitudinal study. *Climacteric*. 2008; 11:55–62. [PubMed: 18202965]
3. de Klerk BM, Schiphof D, Groeneveld FPMJ, et al. Limited evidence for a protective effect of unopposed oestrogen therapy for osteoarthritis of the hip: a systemic review. *Rheumatology (Oxford)*. 2009a; 48:104–12. [PubMed: 19000994]
4. Roman-Blas JA, Castaneda S, Largo R, Herrero-Beaumont G. Osteoarthritis associated with estrogen deficiency. *Arthritis Research and Therapy*. 2009; 11:241. [PubMed: 19804619]
5. Szoek CE, Circuttini FM, Guthrie JR, Clark MS, Dennerstein L. Factors affecting the prevalence of osteoarthritis in healthy middle-aged women: data from the longitudinal Melbourne Women's Midlife Health Project. *Bone*. 2006; 39:1149–55. [PubMed: 16844440]
6. Nevitt MC, Cummings SR, Lane NE, et al. For the Study of Osteoporotic Fractures Research Group. Association of estrogen replacement therapy with the risk of osteoarthritis of the hip in elderly white women. *Arch Intern Med*. 1996; 156:2073–80. [PubMed: 8862099]
7. Wluka AE, Davis SR, Bailey M, Stuckey SL, Cicuttini FM. Users of oestrogen replacement therapy have more knee cartilage than non-users. *Ann Rheum Dis*. 2001; 60:332–6. [PubMed: 11247861]

²deceased

8. Richette P, Corvol M, Bardin T. Estrogens, cartilage, and osteoarthritis. *Joint Bone Spine*. 2003; 70:257–62. [PubMed: 12951307]
9. Hannan MT, Felson DT, Anderson JJ, Naimark A, Kannel WB. Estrogen use and radiographic osteoarthritis of the knee in women: The Framingham Osteoarthritis Study. *Arthritis Rheum*. 1990; 33:525–32. [PubMed: 2328031]
10. Zhang Y, McAlindon TE, Hannan MT, et al. Estrogen replacement therapy and worsening of radiographic knee osteoarthritis: The Framingham Study. *Arthritis Rheum*. 1998; 41:1867–73. [PubMed: 9778229]
11. Erb A, Brenner H, Gunther KP, Sturmer T. Hormone replacement therapy and patterns of osteoarthritis: baseline data from the Ulm Osteoarthritis Study. *Ann Rheum Dis*. 2000; 59:105–9. [PubMed: 10666164]
12. Carbone LD, Nevitt MC, Wildy K, et al. The relationship of anti-resorptive drug use to structural findings and symptoms of knee osteoarthritis. *Arthritis and Rheum*. 2004; 50:3516–3525. [PubMed: 15529367]
13. de Klerk BM, Schiphof D, Groeneveld FPMJ, et al. No clear association between female hormonal aspects and osteoarthritis of the hand, hip and knee: a systematic review. *Rheumatology*. 2009; 48:1160–1165. [PubMed: 19608726]
14. Brunner RL, Aragaki A, Barnabei V, et al. Menopausal symptoms experience before and after stopping estrogen therapy in the Women’s Health Initiative randomized, placebo-controlled trial. *Menopause*. 2010; 17:946–54. [PubMed: 20505547]
15. Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: The Women’s Health Initiative randomized controlled trial. *JAMA*. 2004; 291:1701–1712. [PubMed: 15082697]
16. LaCroix AZ, Chlebowski RT, Manson JE, et al. Health risks and benefits 5 years after stopping randomized treatment with conjugated equine estrogens in postmenopausal women with prior hysterectomy. *JAMA*. 2011; 305:1305–14. [PubMed: 21467283]
17. Stefanick ML, Anderson GL, Margolis KL, et al. Effects of conjugated equine estrogens on breast cancer and mammography screening in postmenopausal women with hysterectomy. *JAMA*. 2006; 295:1647–1657. [PubMed: 16609086]
18. Anderson GL, Chlebowski RT, Aragaki AK, et al. Conjugated equine oestrogen and breast cancer incidence and mortality in postmenopausal women with hysterectomy: extended follow-up of the Women’s Health Initiative randomized placebo-controlled trial. *Lancet Oncol*. 2012; 13:476–86. [PubMed: 22401913]
19. Jackson RD, Wactawski-Wende J, LaCroix AZ, et al. Effects of conjugated equine estrogen on risk of fractures and BMD in postmenopausal women with hysterectomy: results from the Women’s Health Initiative randomized trial. *J Bone Miner Res*. 2006; 21:817–28. [PubMed: 16753012]
20. Wactawski-Wende J, Kotchen JM, Anderson GL, et al. Calcium plus vitamin D supplementation and the risk of colorectal cancer. *N Engl J Med*. 2006; 354:684–696. [PubMed: 16481636]
21. Chlebowski RT, Johnson KC, Kooperberg C, et al. Calcium plus vitamin D supplementation and the risk of breast cancer. *J Natl Cancer Inst*. 2008; 100:1604–1616.
22. McTiernan A, Kooperberg C, White E, et al. Recreational physical activity and the risk of breast cancer in postmenopausal women. *JAMA*. 2003; 290:1331–1336. [PubMed: 12966124]
23. Patterson RE, Kristal AR, Tinker LF, et al. Measurement characteristics of the Women’s Health Initiative food frequency questionnaire. *Ann Epidemiol*. 1999; 9:178–87. [PubMed: 10192650]
24. Kirk KM, Bellamy N, O’Gorman LE, et al. The validity and heritability of self-report osteoarthritis in an Australian older twin sample. *Twin Res*. 2002; 5:98–106. [PubMed: 11931687]
25. March LM, Schwarz JM, Carfrae BH, Bagge E. Clinical validation of self-reported osteoarthritis. *Osteoarthritis Cartilage*. 1998; 6:87–93. [PubMed: 9692063]
26. Taal E, Abdel-Nasser AM, Rasker JJ, Wiegman O. A self-report Thompson articular index: what does it measure? *Clin Rheumatol*. 1998; 17:125–9.
27. Walitt B, Pettinger M, Weinstein A, et al. Effects of postmenopausal hormone therapy on rheumatoid arthritis: the Women’s Health Initiative randomized controlled trials. *Arthritis Rheum*. 2008; 59:320–310.

28. Crillo DJ, Wallace RB, Wu L, Yood RA. Effect of hormone therapy on risk of hip and knee joint replacement in the Women's Health Initiative. *Arthritis Rheum.* 2006; 54:3194–204. [PubMed: 17009251]
29. Sowers MR, McConnell D, Jannausch M, Buyuktur AG, Hochberg M, Jamadar DA. Estradiol and its metabolites and their association with knee osteoarthritis. *Arthritis Rheum.* 2006; 54:2481–7. [PubMed: 16871545]
30. Gao W, Zeng C, Cai D, Liu B, Wen X, Chen Y. Serum concentrations of selected endogenous estrogen and estrogen metabolites in pre- and post-menopausal Chinese women with osteoarthritis. *J Endocrinol Invest.* 2010; 33:644–9. [PubMed: 20339312]
31. Tanamas SK, Wijethilake P, Wluka AE, et al. Sex hormones and structural changes in osteoarthritis: a systematic review. *Maturitas.* 2011; 69:141–56. [PubMed: 21481553]
32. Jochems C, Islander U, Erlandsson M, et al. Osteoporosis in experimental postmenopausal polyarthritis: the relative contributions of estrogen deficiency and inflammation. *Arthritis Res Ther.* 2005; 7:R837–43. [PubMed: 15987485]
33. Hoegh-Andersen P, Tanko LB, Andersen TL, et al. Ovariectomized rats as a model of postmenopausal osteoarthritis: validation and application. *Arthritis Res Ther.* 2004; 6:R169–R180. [PubMed: 15059281]
34. Ham KD, Loeser RF, Lindgren BR, Carlson CS. Effects of long term estrogen replacement therapy on osteoarthritis severity in cynomolgus monkeys. *Arthritis Rheum.* 2002; 46:1956–64. [PubMed: 12124881]
35. Mouritzen U, Christgau S, Lehmann HJ, Tanko LB, Christiansen C. CartiLaps. A novel marker of cartilage degradation. The Influence of age, gender, menopause, hormone replacement therapy and bone mass index. *Ann Rheum Dis.* 2003; 62:332–336. [PubMed: 12634232]
36. Josse RG. Roles for estrogen in bone loss and arthralgia during aromatase inhibitor treatment. *Curr Opin Oncol.* 2007; 19:S1–S8.
37. Rossouw JE, Prentice RL, Manson JE, et al. Postmenopausal hormone therapy and risk of cardiovascular disease by age and years since menopause. *JAMA.* 2007; 297:1465–77. [PubMed: 17405972]
38. Geisler J, Haynes B, Anker G, Dowsett M, Lonning PE. Influence of letrozole and anastrozole on total body aromatization and plasma estrogen levels in postmenopausal breast cancer patients evaluated in a randomized, cross-over study. *J Clin Oncol.* 2002; 20:751–7. [PubMed: 11821457]
39. The Arimidex, Tamoxifen, Alone or in Combination Trialists Group. Buzdar AM, Howell A, Cuzick J, et al. Comprehensive side-effect profile of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: long-term safety analysis of the ATAC trial. *Lancet Oncol.* 2006; 7:633–43. [PubMed: 16887480]
40. Felson DT, Cummings SR. Aromatase inhibitors and the syndrome of arthralgia with estrogen deprivation. *Arthritis Rheum.* 2005; 52:2594–8. [PubMed: 16142740]
41. Chlebowski RT. Aromatase inhibitor-associated arthralgias. *J Clin Oncol.* 2009; 27:4932–4. [PubMed: 19752332]
42. Chlebowski RT, Col N. Menopausal hormone therapy after breast cancer. *Lancet.* 2004; 363:410–411. [PubMed: 14962515]
43. Col NF, Kim JA, Chlebowski RT. Menopausal hormone therapy in breast cancer survivors: A meta-analysis and critical appraisal of the evidence. *Breast Cancer Res.* 2005; 42:861–872.

TABLE 1
Descriptive characteristics of participants at baseline by randomization assignment

	Conjugated equine estrogens alone (n = 5,310)	Placebo (n = 5,429)
Age at screening		
50-59 y	1,637 (30.8)	1,673 (30.8)
60-69 y	2,387 (45.0)	2,465 (45.4)
70-79 y	1,286 (24.2)	1,291 (23.8)
Race/ethnicity		
White	4,007 (75.5)	4,075 (75.1)
Black	782 (14.7)	835 (15.4)
Hispanic	322 (6.1)	333 (6.1)
American Indian	41 (0.8)	34 (0.6)
Asian/Pacific Islander	86 (1.6)	78 (1.4)
Unknown	72 (1.4)	74 (1.4)
Education		
None to some high school	535 (10.2)	518 (9.6)
High school diploma/GED	1,233 (23.5)	1,188 (22.1)
School after high school	2,271 (43.2)	2,350 (43.7)
College degree or higher	1,216 (23.1)	1,327 (24.7)
Age at menarche		
11 y	1,215 (23.0)	1,280 (23.7)
12-13 y	2,805 (53.1)	2,853 (52.8)
14 y	1,259 (23.8)	1,274 (23.6)
Body mass index		
<25 kg/m ²	1,110 (21.0)	1,096 (20.3)
25-29.9 kg/m ²	1,795 (34.0)	1,912 (35.5)
30 kg/m ²	2,376 (45.0)	2,383 (44.2)
Physical activity		
0 MET/wk	1,081 (22.2)	1,043 (21.3)
1-3.5 MET/wk	887 (18.2)	930 (19.0)
3.6-8.0 MET/wk	983 (20.1)	983 (20.0)
8.1-16.5 MET/wk	981 (20.1)	945 (19.3)
>16.5 MET/wk	948 (19.4)	1,003 (20.5)
Alcohol use		
Nondrinker	718 (13.7)	737 (13.7)
Past drinker	1,277 (24.3)	1,270 (23.6)
<1 drink/mo	767 (14.6)	766 (14.2)
<1 drink/wk	1,001 (19.1)	1,049 (19.5)

	Conjugated equine estrogens alone (n = 5,310)	Placebo (n = 5,429)
1-6 drinks/wk	1,027 (19.6)	1,091 (20.2)
7 drinks/wk	457 (8.7)	475 (8.8)
Smoking		
Never smoked	2,723 (51.9)	2,705 (50.4)
Past smoker	1,986 (37.8)	2,089 (38.9)
Current smoker	542 (10.3)	571 (10.6)
Self-reported history of rheumatoid arthritis		
No	4,679 (93.5)	4,784 (94.0)
Yes	327 (6.5)	304 (6.0)
Nonsteroidal anti-inflammatory drug use		
No	4,987 (93.9)	5,100 (93.9)
Yes	323 (6.1)	329 (6.1)
Daily total calcium (supplements + diet), mean (SD), mg	983.5 (643.5)	994.0 (652.2)
Daily total calcium (supplements + diet)		
<800 mg	2,420 (47.7)	2,469 (47.7)
800-1,199 mg	1,190 (23.4)	1,229 (23.8)
1,200 mg	1,466 (28.9)	1,476 (28.5)
Daily vitamin D (supplements + diet), mean (SD), IU	354.5 (257.4)	355.1 (256.6)
Daily vitamin D (supplements + diet)		
<200 IU	1,953 (38.5)	1,969 (38.1)
200-399 IU	1,190 (23.4)	1,212 (23.4)
400-599 IU	1,018 (20.1)	1,090 (21.1)
600 IU	915 (18.0)	903 (17.5)
CaD trial participation		
None	2,236 (42.1)	2,327 (42.9)
CaD randomization	1,531 (28.8)	1,540 (28.4)
Placebo randomization	1,543 (29.1)	1,562 (28.8)
Estrogen-alone use		
Nonuser	2,872 (54.1)	2,891 (53.3)
<5 y	1,317 (24.8)	1,368 (25.2)
5 y	1,121 (21.1)	1,170 (21.6)
Estrogen plus progestin		
Nonuser	5,093 (95.9)	5,178 (95.4)
<5 y	144 (2.7)	158 (2.9)
5 y	73 (1.4)	93 (1.7)

Data are presented as n (%), unless stated otherwise.

MET, metabolic equivalent; CaD, calcium and vitamin D.

TABLE 2
Joint symptoms at baseline and follow-up by randomization assignment in the estrogen-alone trial

	Baseline		Year 1		Year 3		Year 6	
	CEE	Placebo	CEE	Placebo	CEE	Placebo	CEE	Placebo
Joint pain								
n	5,248	5,377	4,780	4,893	523	539	465	477
None, n (%)	1,196 (22.8)	1,249 (23.2)	1,135 (23.7)	1,018 (20.8)	135 (25.8)	109 (20.2)	97 (20.9)	80 (16.8)
Any, n (%)	4,052 (77.2)	4,128 (76.8)	3,645 (76.3)	3,875 (79.2)	388 (74.2)	430 (79.8)	368 (79.1)	397 (83.2)
Severity, n (%)								
Mild	2,381 (45.4)	2,435 (45.3)	2,139 (44.8)	2,236 (45.7)	197 (37.7)	217 (40.3)	189 (40.7)	203 (42.6)
Moderate	1,235 (23.5)	1,271 (23.6)	1,136 (23.8)	1,191 (24.3)	136 (26.0)	150 (27.8)	124 (26.7)	141 (29.6)
Severe	436 (8.3)	422 (7.9)	370 (7.7)	448 (9.2)	55 (10.5)	63 (11.7)	55 (11.8)	53 (11.1)
Severity score, mean (SD)	1.17 (0.87)	1.16 (0.87)	1.16 (0.87)	1.22 (0.88)	1.21 (0.95)	1.31 (0.92)	1.29 (0.93)	1.35 (0.89)
Change in score from baseline, mean (SD)			-0.00 (0.79)	0.06 (0.80)	0.01 (0.81)	0.15 (0.94)	0.09 (0.90)	0.18 (0.95)
Joint swelling								
n	5,254	5,378	4,788	4,890	521	537	468	473
None, n (%)	3,136 (59.7)	3,283 (61.0)	2,772 (57.9)	2,949 (60.3)	299 (57.4)	341 (63.5)	244 (52.1)	282 (59.6)
Any, n (%)	2,118 (49.3)	1,562 (29.0)	2,016 (42.1)	1,941 (39.7)	222 (42.6)	196 (36.5)	224 (47.9)	191 (40.4)
Severity, n (%)								

	Baseline		Year 1		Year 3		Year 6	
	CEE bo	Place bo	CEE bo	Place bo	CEE bo	Place bo	CEE bo	Place bo
Mild	1,535 (29.2)	1,562 (29.0)	1,399 (29.2)	1,431 (29.3)	151 (29.0)	130 (24.2)	153 (32.7)	129 (27.3)
Moderate	481 (9.2)	427 (7.9)	483 (10.1)	408 (8.3)	51 (9.8)	48 (8.9)	54 (11.5)	47 (9.9)
Severe	102 (1.9)	106 (2.0)	134 (2.8)	102 (2.1)	20 (3.8)	18 (3.4)	11 (3.6)	15 (3.2)
Severity score, mean (SD)	0.53 (0.74)	0.51 (0.73)	0.58 (0.78)	0.52 (0.74)	0.60 (0.82)	0.52 (0.79)	0.67 (0.82)	0.57 (0.80)
Change in score from baseline, mean (SD)			0.06 (0.75)	0.02 (0.70)	0.10 (0.81)	0.05 (0.80)	0.17 (0.79)	0.09 (0.89)
				0.03			0.31	0.13
				<0.001			0.11	0.06
		0.14		0.001			0.23	0.15

Statistical tests comparing hormone therapy randomization: unadjusted *P* values from either a logistic regression model (none, any) or a linear regression model (average), CEE, conjugated equine estrogens.

TABLE 3
Joint symptoms at baseline and follow-up by randomization assignment among adherent women in the estrogen-alone trial

	Baseline		Year 1		Year 3		Year 6	
	CEE	Placebo	CEE	Placebo	CEE	Placebo	CEE	Placebo
Joint pain								
n	5,248	5,377	4,125	4,314	298	312	155	183
None, n (%)	1,196 (22.8)	1,249 (23.2)	997 (24.2)	905 (21.0)	82 (27.5)	57 (18.3)	38 (24.5)	30 (16.4)
Any, n (%)	4,052 (77.2)	4,128 (76.8)	3,128 (75.8)	3,409 (79.0)	216 (72.5)	255 (81.7)	117 (75.5)	153 (83.6)
Severity, n (%)								
Mild	2,381 (45.4)	2,435 (45.3)	1,877 (45.5)	1,970 (45.7)	119 (39.9)	134 (43.0)	74 (47.7)	94 (51.4)
Moderate	1,235 (23.5)	1,271 (23.6)	945 (22.9)	1,047 (24.3)	71 (23.8)	86 (27.6)	31 (20.0)	45 (24.6)
Severe	436 (8.3)	422 (7.9)	306 (7.4)	392 (9.1)	26 (8.7)	35 (11.2)	12 (7.7)	14 (7.7)
Severity score, mean (SD)	1.17 (0.87)	1.16 (0.87)	1.14 (0.87)	1.21 (0.88)	1.14 (0.92)	1.32 (0.90)	1.11 (0.86)	1.24 (0.82)
Change in score from baseline, mean (SD)			-0.01 (0.78)	0.06 (0.79)	0.01 (0.74)	0.13 (0.96)	0.12 (0.81)	0.12 (0.87)
Joint swelling								
n	5,254	5,378	4,134	4,307	299	301	156	171
None, n (%)	3,136 (59.7)	3,283 (61.0)	2,403 (58.1)	2,613 (60.7)	175 (58.5)	194 (62.6)	92 (59.0)	113 (62.4)
Any, n (%)	2,118 (49.3)	1,562 (29.0)	1,731 (41.9)	1,694 (39.3)	124 (41.5)	116 (937.4)	64 (41.0)	68 (37.6)
Severity, n (%)								

	Baseline			Year 1			Year 3			Year 6		
	CEE	Placebo	P	CEE	Placebo	P	CEE	Placebo	P	CEE	Placebo	P
Mild	1,535 (29.2)	1,562 (29.0)	0.14	1,217 (29.4)	1,257 (29.2)	0.007	99 (33.1)	80 (25.8)	0.16	48 (30.8)	40 (22.1)	0.23
Moderate	481 (9.2)	427 (7.9)		409 (9.9)	350 (8.1)		21 (7.0)	28 (9.0)		22 (12.2)	13 (8.3)	
Severe	102 (1.9)	106 (2.0)		105 (92.5)	87 (2.0)		4 (1.3)	8 (2.6)		6 (3.3)	3 (1.9)	
Severity score, mean (SD)	0.53 (0.74)	0.51 (0.73)	0.08	0.57 (0.77)	0.52 (0.73)	0.001	0.51 (0.69)	0.52 (0.77)	0.94	0.53 (0.73)	0.56 (0.83)	0.71
Change in score from baseline, mean (SD)				0.05 (0.74)	0.02 (0.69)	0.06	0.08 (0.72)	0.05 (0.76)	0.70	0.18 (0.69)	0.16 (0.83)	0.76

Participants were excluded if they became nonadherent (using <80% of study pills or initiating nonprotocol hormone therapy) before the follow-up visit.
CEE, conjugated equine estrogens.