

Ethanol-induced oxidative stress: basic knowledge

Mario Comporti · Cinzia Signorini · Silvia Leoncini ·
Concetta Gardi · Lucia Ciccoli · Anna Giardini ·
Daniela Vecchio · Beatrice Arezzini

Received: 30 July 2009 / Accepted: 9 November 2009 / Published online: 24 December 2009
© Springer-Verlag 2009

Abstract After a general introduction, the main pathways of ethanol metabolism (alcohol dehydrogenase, catalase, coupling of catalase with NADPH oxidase and microsomal ethanol-oxidizing system) are shortly reviewed. The cytochrome P₄₅₀ isoform (CYP2E1) specifically involved in ethanol oxidation is discussed. The acetaldehyde metabolism and the shift of the NAD/NADH ratio in the cellular environment (reductive stress) are stressed. The toxic effects of acetaldehyde are mentioned. The ethanol-induced oxidative stress: the increased MDA formation by incubated liver preparations, the absorption of conjugated dienes in mitochondrial and microsomal lipids and the decrease in the most unsaturated fatty acids in liver cell membranes are discussed. The formation of carbon-centered (1-hydroxyethyl) and oxygen-centered (hydroxyl) radicals during the metabolism of ethanol is considered: the generation of hydroxyethyl radicals, which occurs likely during the process of univalent reduction of dioxygen, is highlighted and is carried out by ferric cytochrome P₄₅₀ oxy-complex (P₄₅₀-Fe³⁺O₂⁻) formed during the reduction of heme-oxygen. The ethanol-induced lipid peroxidation has been evaluated, and it has been shown that plasma F₂-isoprostanes are increased in ethanol toxicity.

Keywords CYP2E1 isoform · Ethanol metabolism · Hydroxyethyl radicals · Liver-free non-protein bound iron · Oxidative stress · Plasma isoprostanes

Introduction

Ethanol unlike many other hepatotoxic chemicals is not a foreign substance for living organisms; it occurs, in fact, in small amounts in mammalian tissues [82] and is conceivably formed by the alcohol dehydrogenase-catalyzed reduction of acetaldehyde derived from the decarboxylation of the intermediary metabolite pyruvate [83]. A similar reaction also occurs in biological fermentation (such as that yielding ethyl alcohol in wine and other alcoholic beverages) in which glucose is fermented by yeast. Moreover, significant amounts of ethanol are normally formed in the gastrointestinal tract, absorbed by the portal vein and metabolized in the liver [64]; most alcohol is of microbial origin, the other portion probably arising from the acetaldehyde formed by the normal pathways of degradation of threonine, deoxyribose phosphate and β -alanine. Furthermore, ethanol is rapidly and most entirely converted to the key intermediate, acetate, which can enter a wide-spread variety of metabolic pathways. This “more physiological” aspect of ethanol when compared with other hepatotoxic drugs complicates to a large extent the study of the mechanisms involved in the pathogenetic effects of alcohol. The great interest on ethylism from both a clinical and a sociological point of views resulted in an impressive number of studies on the effects of ethyl alcohol. However, the results have not been even univocal because the models used (alcohol dosage, acute or chronic administration, animals studied, different susceptibility to ethanol consumption, sex, nutritional status, etc.) have been extremely variable. Even the choice of the model itself for the study is questionable: in fact, if the acute (single large dose of alcohol) intoxication is a model more suitable for the study of the direct effects of ethanol at the cellular, subcellular and molecular level, it is obviously away from the

M. Comporti (✉) · C. Signorini · S. Leoncini · C. Gardi ·
L. Ciccoli · A. Giardini · D. Vecchio · B. Arezzini
Department of Pathophysiology,
Experimental Medicine and Public Health,
University of Siena, via A. Moro, 53100 Siena, Italy
e-mail: comporti@unisi.it

conditions of the alcoholic; the chronic intoxication, on the other hand, is more adherent to such conditions but is subjected to a number of variables (of nutritional order in particular) that can contribute to a great extent to the development of the lesions, so that it is more difficult to discern the net role played by alcohol. Since the aim of the present review is to focus on the mechanisms of ethanol-induced liver injury and on the role of oxidative stress, particular attention will be paid to the results obtained with acute ethanol intoxication.

Ethanol metabolism and consequent alterations of the overall metabolism of the hepatic cell

The pathophysiological bases of the hepatic alterations produced by ethanol start obviously from its metabolism. Ethanol is quickly absorbed in the gastrointestinal tract and is almost completely oxidized to carbon dioxide and water. Minor amounts only are excreted, unmetabolized, in urine, breath and sweat. The rate of alcohol metabolism varies between 50 and 180 mg/h/kg body wt [118] in adult men, while in rats and mice, notwithstanding the natural aversion to alcohol, a two to threefold higher rate has been reported [67].

Alcohol is mainly metabolized in the liver, and the metabolic pathways are summarized in Fig. 1. The most important pathway is represented by alcohol dehydrogenase, a zinc containing metalloenzyme [121] widely distributed in nature, localized in the soluble cytoplasm and NAD^+ -dependent. It catalyzes the oxidation of ethanol to acetaldehyde (Fig. 1A). It is generally accepted that ethanol is also metabolized by catalase, a heme-containing enzyme widely distributed in nature and particularly expressed in the peroxisomes. Catalase, which normally catalyzes the decomposition of H_2O_2 to H_2O and O_2 (catalytic reaction), can also catalyze the reduction of H_2O_2 to H_2O if electron donors are present (peroxidatic reaction); generally such reaction is catalyzed by peroxidase, but catalase can also catalyze peroxidative type reactions, when hydrogen peroxide is produced and a source of electron donors is available. Ethanol, which acts as an electron donor, is so oxidized to acetaldehyde (Fig. 1B). Hydrogen peroxide necessary for the reaction can be produced by oxidases, some of which also occur in the peroxisomes [14].

Ethanol oxidation can also result from the coupling of the reactions catalyzed by NADPH oxidase (microsomal) and xanthine oxidase with catalase (Fig. 1C, D, respectively). Which is the real importance of catalase in ethanol metabolism is not clear; it seems to be the most important pathway for the metabolism of methanol [1], while it would be of minor importance for the oxidation of ethanol, at least under normal conditions.




Fig. 1 Metabolic pathways of ethanol

Finally, Lieber and De Carli [69] and Lieber et al. [78] have described a third metabolic pathway for ethanol oxidation occurring in the endoplasmic reticulum of the hepatocyte and named “microsomal ethanol-oxidizing system” (MEOS) (Fig. 1E). It utilizes the terms of the enzymatic system of “drug metabolism” (mixed function oxidase system, driven by the microsomal electron transport chain). MEOS with NADPH as a cofactor, oxidizes, it too, ethanol to acetaldehyde (Fig. 1E). A long debate followed the report of MEOS as a separate and individual ethanol-oxidizing system: some authors [19, 115] claimed that microsomal ethanol oxidation could result from the coupling of the activities of NADPH oxidase, the real microsomal enzyme, with catalase or with catalase and alcohol dehydrogenase (present as contaminant in the microsomal fraction) [57]. However, the group of Lieber [117] and others [87] showed that (a) MEOS is active even in acatalasemic animals; (b) a reconstituted system consisting of the essential terms of drug metabolism, i.e., cytochrome P_{450} , NADPH-cytochrome *c* reductase and synthetic phospholipids, is able to oxidize, besides benzphetamine (characteristic substrate of drug metabolism), ethanol and other alcohols too and that MEOS is adaptively increased after chronic ethanol consumption, like other drug metabolizing activities. The induction of drug metabolizing enzymes due to alcohol [56, 73, 108, 109] and the fact that ethanol in vitro inhibits in a competitive way the same enzymes, could explain, at least in part, the increased tolerance of alcoholics to sedatives when sober and the enhanced sensitivity to sedatives when inebriated. The quantitative contribution of microsomal ethanol oxidation to the overall ethanol metabolism is another much

debated question. According to some estimations [115] this route does not play a significant role in the metabolism of alcohol *in vivo*; however, according to the calculation of [71] about ¼ of the ethanol ingested could be metabolized by MEOS in normal conditions, while, after prolonged ethanol consumption, the role of MEOS could be considerably greater. Today, it is well known that the isoform of cytochrome P₄₅₀ involved in ethanol oxidation, CYP2E1 (see in the following paragraphs), has a much higher K_m for ethanol (8–10 mmol/l) compared to that of alcohol dehydrogenase (0.2–2 mmol/l); however, chronic alcohol consumption increases the activity of CYP2E1 by several times [3].

Acetaldehyde metabolism

As reported in the previous paragraphs, all the ethanol-metabolizing pathways lead to the formation of acetaldehyde. The latter can be oxidized to acetate mainly by enzymes, aldehydes dehydrogenases, which are mitochondrial, NAD⁺-dependent enzymes. Thus, the two main steps of ethanol metabolism, namely, the NAD⁺-linked oxidation of ethanol to acetaldehyde and the NAD⁺-linked oxidation of acetaldehyde to acetate, cause an increased concentration of the reduced pyridine coenzyme, NADH. Therefore, a decrease in the NAD⁺/NADH ratio occurs in the liver cell and such an imbalance would represent, according to a classic view of the problem [68, 74, 75], the central event in the explanation of the various metabolic alterations produced directly in the hepatic cell by ethanol oxidation, including triglyceride accumulation (the importance of the reductive stress environment created by ethanol oxidation will be outlined in the following paragraphs).

As is known, the reoxidation of NADH formed in the soluble cytoplasm occurs through shuttle mechanisms that transfer reducing equivalents from cytoplasmic NADH to mitochondrial electron transport chain. The latter becomes quickly saturated by reducing equivalents originating from ethanol and thus can slow down the citric acid cycle that, under normal conditions, supplies itself reducing equivalents to the respiratory chain. The slackening of the citric acid cycle leads to various metabolic consequences, the major part of which derive from decreased oxidation of acetate, which can originate, as mentioned in the previous paragraphs, directly by ethanol, even if a consistent part of the latter would be metabolized in peripheral tissues [80]. The metabolic unbalance derived from the decreased acetate oxidation is primarily reflected on fatty acid oxidation, fatty acid synthesis, ketone bodies formation and cholesterol metabolism. Decreased fatty acid oxidation and increased fatty acid synthesis have been demonstrated since a long time ago [40, 74, 95, 99, 126]. It was therefore suggested that ethanol, acting itself as a fuel for

mitochondria, could replace in a “competitive way” fatty acids that, under normal conditions, represent the major source of energy for the liver cell.

Today, it is known that the ethanol-induced stimulation of hepatic triglyceride synthesis depends, to a great extent, upon the increased expression of the enzymes involved in fatty acid synthesis that are regulated by the transcription factor sterol regulatory element-binding protein (SREBP)-1 [123], located in the endoplasmic reticulum. Upon ethanol feeding, SREBP-1 is proteolytically cleaved to the active form that translocates to the nucleus, inducing the expression of the genes coding for lipogenic enzymes.

Pathogenetic mechanisms of the hepatic lesions produced by ethanol

As it is known, the long lasted debate whether nutritional deficiencies associated to alcoholism are responsible of liver injury [15, 49] or whether ethanol per se (or some proximal metabolite) exerts toxic effects on hepatic cell has been largely resolved by the introduction of a novel model for experimental studies based on liquid diets in which ethanol (36% of total calories) replaces in a isocaloric way part of carbohydrates. With these diets, considered nutritionally adequate, the alcohol assumption in the rat was greatly increased and hepatic lesions (hepatic steatosis) were obtained in both rats and humans [76, 77]. These lesions therefore apparently were independent of nutritional deficiencies. Subsequently, with the use of primates as experimental animals, a sequential production of fatty liver, inflammatory reactions, necrosis, fibrosis and sclerosis has been obtained [78] upon prolonged ethanol feeding.

The ethanol-induced ultrastructural changes of the liver cell involve both the mitochondria and the endoplasmic reticulum. The mitochondrial changes consist of enlargement, swelling, shortening and disorganization of the *cristae*, decreased number or absence of matrix granules and intramitochondrial crystalline inclusions. The presence of bizarre shapes and *giant mitochondria* [61, 63] with an increased matrix density was also reported [56, 66, 72, 97]. The changes involving the endoplasmic reticulum [109] consist of a marked proliferation of the smooth membranes with abundant vesicular structures. The alterations of the endoplasmic reticulum are, according to [56], the first morphological changes detectable in the liver cell after ethanol feeding to rats; they were observed, in fact, after few days of alcohol administration, while mitochondrial changes appeared some days later. Autophagic vacuoles, containing altered mitochondria, and residual dense bodies were also described. The presence of hyaline Mallory bodies was repeatedly documented. An increased number of peroxisomes were also reported [12], which is consistent with the increased catalase activity.

The morphological changes of mitochondria have a functional counterpart, consisting in a decrease in cytochrome *a*, *a*₃ and *b* [21, 62, 110], of the enzymatic activities of the Krebs cycle [21, 110] and in general of the respiratory control [45, 59]. Many of these alterations are stable and can be in turn responsible for the changes in lipid metabolism produced by ethanol oxidation itself. Ethanol-induced mitochondrial changes also involve the autochthonous synthesis of some mitochondrial proteins, thus producing an unbalance between proteins synthesized in mitochondria and those synthesized in the endoplasmic reticulum. Such unbalance could influence the mitochondria biosynthesis.

The toxic effects of acetaldehyde

That ethanol-induced liver changes are in part mediated by the proximal metabolite acetaldehyde that has been proposed since a long time ago and reconsidered when it has been observed [21] that many of the earlier mentioned ethanol-induced mitochondrial changes are reproduced by acetaldehyde. In particular, the addition of acetaldehyde to isolated mitochondria at concentrations of the same order as those occurring in the liver actively oxidizing ethanol (1–3 mM), inhibits mitochondrial respiration at the level of complex I (NADH-ubiquinone oxido-reductase) and the coupling site I of oxidative phosphorylation [21, 22]. Moreover, acetaldehyde inhibits, at the same concentrations, fatty acid oxidation, and such inhibition seems to be due to the inhibition of β -oxidation, citric acid cycle and oxidative phosphorylation [24]. The inhibition of the oxidation of NAD⁺-dependent substrates [21] and the inhibition of fatty acid oxidation [24] do not seem to depend from a competition of acetaldehyde with such substrates for NAD⁺. It has been suggested [23, 25], on the other hand, that acetaldehyde could interact with sulphhydryl groups involved in oxidative phosphorylation particularly with those essential for the complex I (NAD⁺ ubiquinone-oxydoreductase) which is the site of the respiratory chain mainly affected by both acetaldehyde and ethanol. Thus, the block of –SH groups would be responsible for the alterations. The reaction of acetaldehyde and other alkanals with cysteine has been actually known since many decades (formation of thiazolidine-carboxylic acids) and other low molecular weight thiols (reduced glutathione, GSH, in particular) can also be involved in this reaction. In effect, ethanol toxicity is always accompanied by a decrease in hepatic GSH content. If, as we will see later, lipid peroxidation is going to develop in ethanol hepatotoxicity, much more reactive aldehydes (alkenals and 4-hydroxyalkenals) will be formed and a much higher reactivity toward –SH groups and other nucleophiles has to be expected.

Finally, the acetaldehyde-produced alterations of the mitochondrial functions impair the mitochondrial acetaldehyde metabolism [51], thus producing worsening of the damage by a vicious cycle.

Ethanol-induced oxidative stress

It has been known since a long time ago that prior administration for a variety of substances known as antioxidants affords a marked protection against the liver damage induced by CCl₄ and other hepatotoxins [44, 54]. It was subsequently demonstrated [29, 31, 36] that CCl₄ both administered in vivo or added in vitro to liver preparations markedly increases the peroxidation of liver lipids, as measured by malonil dialdehyde (MDA) formation. Such a pro-oxidant effect of CCl₄ was immediately confirmed by Recknagel and Ghoshal [100, 101]. The theoretical background for these studies was suggested by [18] and [125], who proposed that in the hepatic cell, CCl₄ could undergo a homolytic cleavage yielding free radicals (CCl₃, CCl₃OO[•], etc.). The latter could rapidly interact with neighboring molecules, such as proteins, nucleic acid, thiols and membrane unsaturated fatty acids. The latter interaction would set into motion lipid peroxidation that seriously affects membrane structure and function. Since antioxidant pretreatment [37] was also found to be effective in ethanol-induced liver damage, it was suggested [38] that the liver injury produced by CCl₄ or ethanol could have a common pathogenetic mechanism, namely, oxidative stress¹ and peroxidation of liver lipids. It was subsequently found [30, 32] that, as in the case of CCl₄, the MDA production by incubated liver homogenates is greatly enhanced during 1, 2, 4, 6 and 12 h after acute ethanol administration. The increase in liver lipid peroxidation precedes the accumulation of triglycerides in the liver, thus excluding that the enhanced MDA formation could result from the increased hepatic fat content; the latter possibility is also ruled out by the fact that no increased MDA production occurs 24 h after ethanol dosing when the hepatic triglyceride level is maximum. Furthermore, it was shown [32] that ethanol added in vitro to liver homogenates has a pro-oxidant effect although to a lesser extent than carbon tetrachloride. The effect is specific for the liver tissue, since ethanol, either in vitro or in vivo, has no effect on the peroxidation of brain [48] or other tissue homogenates [114].

It was proposed [32] that ethanol or its metabolites stress the “peroxidative balance” of the liver cell [16] toward autoxidation, either acting as pro-oxidant or lowering the cellular antioxidant level. Direct evidence for increased

¹ Oxidative stress is generally considered as a disturbance in the pro-oxidant/antioxidant balance in favor of the former, leading to potential damage [47].



hepatic lipoperoxidation *in vivo* after acute ethanol intoxication was forwarded by Kalish and Di Luzio [58], who showed that the peroxide content was increased in the liver in ethanol-treated rats. Hashimoto and Recknagel [50], on the contrary, found no evidence of conjugated diene absorption characteristic of peroxidized lipids [17] in the lipids of any subcellular fraction at any time after acute ethanol intoxication. On the basis of these results, it was concluded that in the case of ethanol-induced liver injury, there is no direct evidence for the *in vivo* occurrence of hepatic lipoperoxidation. Di Luzio [39] questioned the above results and showed that the absorption of conjugated dienes can be detected in the mitochondrial but not in the microsomal lipids of ethanol-treated rats. On the other hand, Corongiu et al. [35] demonstrated the absorption of conjugated dienes in microsomal lipids of ethanol-treated animals by the second-derivative spectroscopy.

An approach to the problem with different technical procedures was then devised. Since the end result of lipoperoxidation is a decrease in the most highly unsaturated fatty acids, which are the major peroxidizable substrates, a decrease in their content in the lipids of isolated subcellular fractions could indicate, among other possibilities, that a peroxidative breakdown of these moieties had actually occurred *in vivo*. As a matter of fact, a progressive decrease in the arachidonic acid content of liver microsomal phospholipids was observed [33] shortly after carbon tetrachloride intoxication; it was also observed, in contrast with liver phospholipids, that hepatic triglycerides do not show any change in arachidonic acid content after poisoning, again suggesting that lipid peroxidation involves structural lipids rather than the lipids accumulating in the liver as a result of the intoxication. A clear decrease in the arachidonic and docosahexaenoic acid content of liver mitochondrial lipids from acutely ethanol-treated rats was actually found [34]. In contrast with the mitochondrial changes, ethanol did not induce a decrease in the most unsaturated components of the fatty acid pattern of liver microsomal phospholipids [34].

A decrease in arachidonic as well as an increase in linoleic acid content of liver mitochondrial lipids was also observed by French et al. [43] after chronic ethanol administration, but these changes were mainly attributed to alterations in the activity of the chain elongation desaturation system.

Implication of oxidative stress in ethanol toxicity would imply that either ethanol is converted, during its metabolism, to a free radical intermediate or that ethanol or its metabolites react with some nucleophile in an antioxidant molecule, thus blocking the molecule and decreasing the antioxidant potential. The latter possibility has been shown above (reaction of acetaldehyde with –SH groups of cysteine or GSH), but the loss of GSH is by far smaller than

that occurring with many other GSH depletors (bromobenzene, allyl alcohol, etc. [28]) and cannot account by itself for the induction of lipid peroxidation. The former possibility—the formation of a free radical during ethanol metabolism—was postulated by Slater [113] since many years ago. Ethanol may enter free radical reaction relatively easily [111], through the interaction with some endogenous radical; the latter could give rise to a homolytic cleavage of ethanol yielding a reducing ethoxy radical ($\text{CH}_3\text{-CH}_2\text{O}^\cdot$), which in the presence of some oxidant would be converted to acetaldehyde:



(from [113]).

Several endogenous radicals are known to be involved in the NADPH-cytochrome P_{450} chain; ethanol may interact at this site during its metabolism in MEOS. Also, in the scheme proposed for the action of catalase-free radical intermediates of the hydrogen donor are formed; if ethanol is the donor, free radical intermediates from ethanol can result.

More recent studies have conclusively shown that ethoxy radical is really generated during ethanol oxidation and that an oxidative stress is imposed on the liver cell as a result of ethanol metabolism [94]. Several sources of such an oxidative stress have been described. Ethanol oxidation results in the production of free radicals, which can derive from both oxygen and ethanol itself. Oxygen radicals can originate as follows: microsomal NADPH-cytochrome *c* reductase and cytochrome P_{450} (components of MEOS) can generate $\text{O}_2^{\cdot-}$ and H_2O_2 [52, 53, 65, 93, 124]; the same oxygen species can be produced by aldehydes oxidase and xanthine oxidase [85], both involved in the metabolism of ethanol-derived acetaldehyde; $\text{O}_2^{\cdot-}$ and H_2O_2 can also be generated by microsomal NADPH oxidase, which has been shown to be increased after acute [120] or chronic [70, 104, 119, 122] ethanol administration; during NADPH oxidation liver microsomes produce significant amount of OH^\cdot (being H_2O_2 the precursor), which in turn appears to be required for ethanol oxidation [20, 26, 55].

With regard to ethanol-derived radicals, it has been shown [4, 5] that ethanol is activated to a free radical intermediate by the ethanol inducible form of cytochrome P_{450} , i.e., the specific isoenzymatic form involved in MEOS, CYP2E1. With the use of electron spin resonance (ESR) spectroscopy in combination with the spin trapping agent 4-pyridyl-1-oxo-*t*-butyl nitron (4-POBN), it has been demonstrated [4, 5] that rat liver microsomes incubated with ethanol and NADPH can produce a free radical intermediate, identified as 1-hydroxyethyl radical. Free radical intermediates are also produced by liver microsomes during the metabolism of various aliphatic alcohols (1-propanol,

2-propanol, 1-butanol, 2-butanol and 1-pentanol), indicating the existence of a common activating pathway for these compounds [5, 7]. The formation of radical intermediates has been confirmed in the whole animal in vivo with the use of 4-POBN [8, 60, 102, 103]. The generation of ethanol radicals would occur during the process of univalent reduction of dioxygen and possibly would be carried out by ferric cytochrome P₄₅₀ oxy-complex (P₄₅₀-Fe³⁺O₂⁻) [10, 11] formed during the reduction of heme-oxygen. In such a state, cytochrome would be sufficiently reactive to abstract a proton from the 1-carbon of ethanol, yielding a carbon-centered radical and H₂O₂ [116]. Alternatively, hydroxyethyl radicals could be produced by addition to ethanol of OH[•] radicals generated by liver microsomes [81]. However, generated hydroxyethyl radicals bind to microsomal protein [9], particularly CYP2E1, and probably play an important role in the induction of lipid peroxidation [6, 42]. The binding of alcohol radical to protein represents another mechanism of hepatic protein alkylation in addition to that operated by acetaldehyde [41] and known to contribute to overall liver cell damage. Furthermore, the hydroxyethyl radical-derived protein adducts are immunogenic and give rise to antibodies different from those generated by acetaldehyde-derived protein adducts [88]. Chronic alcohol feeding of rats leads to the production of antibodies that recognize hydroxyethyl rat serum albumin but do not recognize rat serum albumin [88]. Moreover, sera of alcoholic cirrhotic patients contain IgG and IgA antibodies that recognize proteins modified by hydroxyethyl radicals [27]. Such antibodies may play an important role in the immunologic reactions triggered by ethanol and due to antibodies against liver cells found in the serum of patients with alcoholic liver injury [46].

Ethanol-induced oxidative stress as measured by F₂-isoprostane determination

A great advance in the study of oxidative stress has been represented by the demonstration [89, 90, 106] of the formation of a series of prostaglandin F₂-like compounds, named F₂-isoprostanes, which originate in vitro and in vivo from the peroxidation of phospholipids bound arachidonic acid. Since F₂-isoprostanes, which are initially formed in situ on phospholipids [91], are released into the blood compartment and since these prostanoids are much less reactive than other lipid peroxidation products such as lipoperoxides and aldehydes, they can be found more easily in plasma and urine. Elevated levels of F₂-isoprostanes have been found in various human pathologies [105]. Therefore, we reconsidered the whole problem of ethanol-induced oxidative stress with this methodological approach, which is nowadays considered as the most sophisticated and reliable technique to evaluate oxidative

stress, at least when the determinations are carried out by gas-chromatography mass spectrometry. Plasma F₂-isoprostanes are increased in ethanol toxicity ([92, 86, 13, 96], Comporti et al. unpublished work).

Final considerations

Owing to the fact that this review is concerned with the mechanisms inducing oxidative stress upon ethanol exposure (particularly acute exposure), no attention has been paid to the development of alcoholic liver disease and its progression to liver fibrosis. Thus, no mention has been done of the impairment of endogenous antioxidant defences, accumulation of unfolded proteins and endoplasmic reticulum stress [84, 112], AMP-dependent protein Kinase (AMPK) and adiponectin-regulated hepatic lipid metabolism [107], translocation of gut derived endotoxins to portal circulation [98], activation of Kupffer cells and release of proinflammatory cytokines (TNF α in particular), role of immune reactions in alcoholic-induced inflammation and progression to liver fibrosis through activation of hepatic stellate cells.

Extensive information about the above topics can be found in several reviews [2, 3, 79].

References

1. Aebi H, Koblet H, von Wartburg JP (1957) Ueber den mechanismus der biologischen methanoloxydation. *Elv Physiol Pharmacol Acta* 15:384–399
2. Albano E (2006) Alcohol, oxidative stress and free radical damage. *Proc Nutr Soc* 65:278–290
3. Albano E (2008) New concepts in the pathogenesis of alcoholic liver disease. *Expert Rev Gastroenterol Hepatol* 2:749–759
4. Albano E, Tomasi A, Gloria-Gatti L, Poli G, Vannini V, Dianzani MU (1987) Free radical metabolism of alcohols by rat liver microsomes. *Free Radic Res Commun* 3:243–249
5. Albano E, Tomasi A, Gloria-Gatti L, Dianzani MU (1988) Spin trapping of free radical species produced during the microsomal metabolism of ethanol. *Chem Biol Interact* 65:223–234
6. Albano E, Poli G, Tomasi A, Gloria-Gatti L, Dianzani MU (1988) Study of the mechanisms responsible for ethanol-induced oxidative damage using isolated rat hepatocytes. In: Hayaishi O, Niki E, Kondo M, Yoshikawa T (eds) *Medical biochemical and chemical aspects of free radicals*. Elsevier Biomedical, Amsterdam, pp 1389–1392
7. Albano E, Tomasi A, Persson J, Terelius Y, Gloria-Gatti L, Ingelman-Sundberg M, Dianzani MU (1991) Role of ethanol-inducible cytochrome P₄₅₀ (P₄₅₀IIE1) in catalyzing the free radical activation of aliphatic alcohols. *Biochem Pharmacol* 41:1895–1902
8. Albano E, Tomasi A, Parola M, Comoglio A, Ingelman-Sundberg M, Dianzani MU (1993) Mechanisms responsible for free radical formation during ethanol metabolism and their role in causing oxidative injury by alcohol. In: Corongiu F, Banni S, Dessi MA, Rice Evans C (eds) *Free radicals and antioxidants nutrition*. Richelieu Press, London, pp 77–96

9. Albano E, Parola M, Comoglio A, Dianzani MU (1993) Evidence for the covalent binding of hydroxyethyl radicals to rat microsomal proteins. *Alcohol Alcohol* 28:453–459
10. Albano E, Tomasi A, Ingelman-Sundberg M (1994) Spin trapping of alcohol-derived radicals in microsomes and reconstituted systems by electron spin resonance. *Methods Enzymol* 233:117–127
11. Albano E, French SW, Ingelman-Sundberg M (1999) Hydroxyethyl radicals in ethanol hepatotoxicity. *Front Biosci* 4:533–540
12. Albot G, Parturier-Albot M, Nicolaisen P (1968) Histology, ultrastructure and histobiology of subacute alcoholic hepatitis and cirrhotic alcoholic hepatitis. *Rev Intern Hepatol* 18:117–154
13. Aleynik SI, Leo MA, Aleynik MK, Lieber CS (1998) Increased circulating products of lipid peroxidation in patients with alcoholic liver disease. *Alcohol Clin Exp Res* 22:192–196
14. Baudhuin P (1969) Liver peroxisomes, cytology and function. *Ann NY Acad Sci* 168:214–228
15. Best CH, Hartroft WS, Lucas CC, Ridout JH (1949) Liver damage produced by feeding alcohol or sugar and its prevention by choline. *Br Med J* 11:1001–1006
16. Bieri JG, Anderson AA (1960) Peroxidation of lipids in tissue homogenates as related to vitamin E. *Arch Biochem Biophys* 90:105–110
17. Bolland JL, Koch HP (1945) The course of autoxidation reactions in polyisoprenes and allied compounds. Part IX. The primary thermal oxidation product of ethyl linoleate. *J Chem Soc* 445–447
18. Butler TC (1961) Reduction of carbon tetrachloride in vivo and reduction of carbon tetrachloride and chloroform in vitro by tissues and tissue constituents. *J Pharmacol Exp Ther* 134:311–319
19. Carter EA, Isselbacher KJ (1971) The role of microsomes in the hepatic metabolism of ethanol. *Ann NY Acad Sci* 179:282–294
20. Cederbaum AI, Dicker E (1983) Inhibition of microsomal oxidation of alcohols and of hydroxyl-radical-scavenging by the iron-chelating agent desferrioxamine. *Biochem J* 210:107–113
21. Cederbaum AI, Rubin E (1975) Molecular injury to mitochondria produced by ethanol and acetaldehyde. *Fed Proc* 34:2045–2051
22. Cederbaum AI, Lieber CS, Rubin E (1974) Effects of chronic ethanol treatment on mitochondrial functions. *Arch Biochem Biophys* 165:560–569
23. Cederbaum AI, Lieber CS, Rubin E (1974) The effects of acetaldehyde on mitochondrial function. *Arch Biochem Biophys* 161:26–39
24. Cederbaum AI, Lieber CS, Rubin E (1975) Effect of acetaldehyde on fatty acid oxidation and ketogenesis by hepatic mitochondria. *Arch Biochem Biophys* 169:29–41
25. Cederbaum AI, Lieber CS, Rubin E (1976) Effect of chronic ethanol consumption and acetaldehyde on partial reactions of oxidative phosphorylation and CO₂ production from citric acid cycle intermediates. *Arch Biochem Biophys* 176:525–538
26. Cederbaum AI, Dicker E, Cohen G (1978) Effect of hydroxyl radical scavengers on microsomal oxidation of alcohols and associated microsomal reactions. *Biochemistry* 17:3058–3064
27. Clot P, Bellomo G, Tebone M, Aricòt S, Albano E (1995) Detection of antibodies against proteins modified by hydroxyethyl free radicals in patients with alcoholic cirrhosis. *Gastroenterology* 108:201–207
28. Comporti M (1987) Glutathione depleting agents and lipid peroxidation. *Chem Phys Lipids* 45:143–169
29. Comporti M, Saccocci C (1965) Stimulation of the lipid peroxidation of rat liver homogenates by low concentration of CCl₄ in vitro. *Boll Soc Ital Biol Sper* 41:1066–1069
30. Comporti M, Di Luzio NR (1966) Effect of ethanol administration on the peroxidation of liver lipids. *Physiologist* 9:157 (abstract)
31. Comporti M, Saccocci C, Dianzani MU (1965) Effect on CCl₄ in vitro and in vivo on lipid peroxidation of rat liver homogenates and subcellular fractions. *Enzymologia* 29:185–204
32. Comporti M, Hartman A, Di Luzio NR (1967) Effect of in vivo and in vitro ethanol administration on liver lipid peroxidation. *Lab Invest* 16:616–624
33. Comporti M, Burdino E, Ugazio G (1971) Changes in fatty acid pattern of liver microsomal phospholipids in rats treated with carbon tetrachloride. *Ital J Biochem* 20:156–165
34. Comporti M, Burdino E, Raja F (1971) Fatty acid composition of mitochondrial and microsomal lipids of rat liver after acute ethanol intoxication. *Life Sci* 10:855–866
35. Corongiu FP, Lai M, Milia A (1983) Carbon tetrachloride, bromotrichloromethane and ethanol acute intoxication. *Biochem J* 212:625–631
36. Dianzani MU, Baccino FM, Comporti M (1966) The direct effect of carbon tetrachloride on subcellular particles. *Lab Invest* 15:149–156
37. Di Luzio NR (1964) Prevention of the acute ethanol-induced fatty liver by the simultaneous administration of antioxidants. *Life Sci* 3:113–118
38. Di Luzio NR (1966) A mechanism of the acute ethanol-induced fatty liver and modification of liver injury by antioxidants. *Lab Invest* 15:50–63
39. Di Luzio NR (1968) Role of lipid peroxidation and antioxidants in ethanol-induced lipid alterations. *Exp Mol Pathol* 8:394–402
40. Di Luzio NR, Poggi M (1963) Abnormal lipid tolerance and hyperlipemia in acute ethanol-treated rats. *Life Sci* 2:751–758
41. Donohur TM, Tuma DJ, Sorrell MF (1983) Acetaldehyde adducts with proteins: bindings of ¹⁴C-acetaldehyde to serum albumin. *Arch Biochem Biophys* 220:239–246
42. Ekström G, Ingelman-Sundberg M (1989) Rat liver microsomal NADPH-supported oxidase activity and lipid peroxidation dependent or ethanol-inducible cytochrome P₄₅₀ (P₄₅₀ II E1). *Biochem Pharmacol* 38:1313–1319
43. French SW, Ihrig TJ, Morin RJ (1970) Lipid composition of RBC ghosts, liver mitochondria and microsomes of ethanol-fed rats. *Q J Stud Alcohol* 31:801–809
44. Gallagher CH (1962) The effect of antioxidants on poisoning by carbon tetrachloride. *Austral J Exp Biol Med Sci* 40:241–254
45. Gordon ER (1973) Mitochondrial functions in an ethanol-induced fatty liver. *J Biol Chem* 248:8271–8280
46. Grossley IR, Neuberger J, Davis M, Williams R, Eddleston ALWF (1986) Ethanol metabolism in generation of new antigenic determinations on liver cells. *Gut* 27:186–189
47. Halliwell B, Gutteridge JMC (1999) Free radicals in biology and medicine, 3rd edn. Oxford Science Publication, Oxford
48. Hartman A, Comporti M, Di Luzio NR (1967) Peroxidation of lipids in the pathogenesis of acute ethanol induced liver injury. *Gastroenterology* 52:316 (abstract)
49. Hartroft WS, Porta EA (1968) Alcohol, diet, and experimental hepatic injury. *Can J Physiol Pharmacol* 46:463–473
50. Hashimoto S, Recknagel RO (1968) No chemical evidence of hepatic lipid peroxidation in acute ethanol toxicity. *Exp Mol Pathol* 8:225–242
51. Hasumura Y, Teschke R, Lieber CS (1975) Acetaldehyde oxidation by hepatic mitochondria: decrease after chronic ethanol consumption. *Science* 198:727–729
52. Hildebrandt AG, Roots I (1975) Reduced nicotinamide adenine dinucleotide phosphate (NADPH)-dependent formation and breakdown of hydrogen peroxide during mixed function oxidation reactions in liver microsomes. *Arch Biochem Biophys* 171:385–397

53. Hildebrandt AG, Speck M, Roots I (1973) Possible control of hydrogen peroxide production and degradation in microsomes during mixed function oxidase reaction. *Biochem Biophys Res Commun* 54:968–975
54. Hove EL (1948) Interrelation between alpha-tocopherol and protein metabolism; the protective effect of vitamin E and certain nitrogenous compounds against CCl₄ poisoning in rats. *Arch Biochem* 17:467–473
55. Ingelman-Sundberg M, Johanson I (1984) Mechanisms of radical formation and ethanol oxidation by ethanol inducible and other forms of rabbit liver microsomal cytochromes P₄₅₀. *J Biol Chem* 249:6447–6458
56. Iseri OA, Lieber CS, Gottlieb LS (1966) The ultrastructure of fatty liver induced by prolonged ethanol ingestion. *Am J Pathol* 48:535–555
57. Isselbacher KJ, Carter EA (1970) Ethanol oxidation by liver microsomes: evidence against a separate and distinct enzyme system. *Biochem Biophys Res Commun* 39:530–535
58. Kalish GH, Di Luzio NR (1966) Peroxidation of liver lipids in the pathogenesis of the ethanol-induced fatty liver. *Science* 152:1390–1392
59. Kiessling KH, Pilström L (1968) Effect of ethanol on rat liver. V. Morphological and functional changes after prolonged consumption of various alcoholic beverages. *Q J Stud Alcohol* 29:819–827
60. Knecht KT, Bradford BU, Mason RP, Thurman GR (1990) In vivo formation of radical metabolite of ethanol. *Mol Pharmacol* 38:26–30
61. Koch OR, Porta EA, Hartroft WS (1968) A new experimental approach in the study of chronic alcoholism. III. Role of alcohol versus sucrose or fat derived calories in hepatic damage. *Lab Invest* 18:379–386
62. Koch OR, Boveris A, Sirotzky de Favelukes S, Schwarck de Tarlowsky M, Stoppani AOM (1977) Biochemical lesions of liver mitochondria from rats after chronic alcohol consumption. *Exp Mol Pathol* 27:213–220
63. Koch OR, Roatta De Conti L, Pacheco Bolaños L, Stoppani AOM (1978) Ultrastructural and biochemical aspects of liver mitochondria during recovery from ethanol-induced alterations. *Am J Pathol* 90:325–344
64. Krebs HA, Perkins JR (1970) The physiological role of liver alcohol dehydrogenase. *Biochem J* 118:635–644
65. Lai CS, Grover TA, Piette LH (1979) Hydroxyl radical production in a purified NADPH-cytochrome *c* (P₄₅₀ reductase system). *Arch Biochem Biophys* 193:373–378
66. Lane BP, Lieber CS (1966) Ultrastructural alterations in human hepatocytes following ingestion of ethanol with adequate diets. *Am J Pathol* 49:593–603
67. Lester D, Keokosky WZ, Felzenberg F (1968) Effect of pyrazoles and other compounds on alcohol metabolism. *Q J Stud Alcohol* 29:449–454
68. Lieber CS, Davidson CS (1962) Some metabolic effects of ethyl alcohol. *Am J Med* 33:319–327
69. Lieber CS, De Carli LM (1968) Ethanol oxidation by hepatic microsomes: adaptive increase after ethanol feeding. *Science* 162:917–918
70. Lieber CS, De Carli LM (1970) Reduced nicotinamide-adenine dinucleotide phosphate oxidase: activity enhanced by ethanol consumption. *Science* 170:78–80
71. Lieber CS, De Carli LM (1972) The role of the hepatic microsomal ethanol oxidizing system (MEOS) for ethanol metabolism in vivo. *J Pharmacol Exp Ther* 181:278–287
72. Lieber CS, De Carli LM (1974) An experimental model of alcohol feeding and liver injury in the baboon. *J Med Primatol* 3:153–163
73. Lieber CS, Rubin E (1968) Alcoholic fatty liver in man on a high protein and low fat diet. *Am J Med* 44:200–206
74. Lieber CS, Schmid R (1961) The effect of ethanol on fatty acid metabolism: stimulation of hepatic fatty acid synthesis in vitro. *J Clin Invest* 40:394–399
75. Lieber CS, De Carli LM, Schmid R (1959) Effect of ethanol on fatty acid metabolism in liver slices. *Biochem Biophys Res Commun* 1:302–306
76. Lieber CS, Jones DP, De Carli LM (1965) Effects of prolonged ethanol intake: production of fatty liver despite adequate diets. *J Clin Invest* 44:1009–1021
77. Lieber CS, Lefèvre A, Spritz N, Feinman L, De Carli LM (1967) Difference in hepatic metabolism of long- and medium-chain fatty acids: the role of fatty acid chain length in the production of the alcoholic fatty liver. *J Clin Invest* 46:1451–1460
78. Lieber CS, Teschke R, Hasumura Y, De Carli LM (1975) Difference in hepatic and metabolic changes after acute and chronic alcohol consumption. *Fed Proc* 34:2060–2074
79. Lu Y, Cederbaum AL (2008) CYP2E1 and oxidative liver injury by alcohol. *Free Radic Biol Med* 44:723–738
80. Lundquist F, Tygstrup N, Winkler K, Mellemegaard K, Munck-Petersen S (1962) Ethanol metabolism and production of free acetate in the human liver. *J Clin Invest* 41:955–961
84. Malhotra JD, Kaufman RJ (2007) Endoplasmic reticulum stress and oxidative stress: a vicious circle or a double-edged sword? *Antioxid Redox Signal* 9:2277–2293
85. Massey V, Strickland S, Mayhew SG, Hawell LG, Engel PC, Matthews RG, Schuman M, Sullivan PA (1969) The production of superoxide anion radicals in the reaction of reduced flavins and flavoproteins with molecular oxygen. *Biochem Biophys Res Commun* 36:891–897
81. McCay PB, Reinke LA, Rau JM (1992) Hydroxyl radicals are generated by hepatic microsomes during NADPH oxidation relationship to ethanol metabolism. *Free Radic Res Commun* 15:335–346
82. McManus IR, Contag AO, Olson RE (1960) Characterization of endogenous ethanol in the mammal. *Science* 131:102–103
83. McManus IR, Contag AO, Olson RE (1966) Studies on the identification and origin of ethanol in mammalian tissues. *J Biol Chem* 241:349–356
86. Meagher EA, Barry OP, Burke A, Lucey MR, Lawson JA, Rokach J, FitzGerald GA (1999) Alcohol-induced generation of lipid peroxidation products in humans. *J Clin Invest* 104:805–813
87. Mezey E, Potter JJ, Reed WD (1973) Ethanol oxidation by a component of liver microsomes rich in cytochrome P₄₅₀. *J Biol Chem* 248:1183–1187
88. Moncada C, Torres V, Varghese G, Albano E, Israel Y (1994) Ethanol-derived immunoreactive species formed by free radical mechanisms. *Mol Pharmacol* 46:786–791
89. Morrow JD, Harris TM, Roberts LJ (1990) Noncyclooxygenase oxidative formation of a series of novel prostaglandins: analytical ramifications for measurement of eicosanoids. *Anal Biochem* 184:1–10
90. Morrow JD, Hill KE, Burk RF, Nammour TM, Badr KF, Roberts LJ (1990) A series of prostaglandin F₂-like compounds are produced in vivo in humans by a non-cyclooxygenase, free radical-catalyzed mechanism. *Proc Natl Acad Sci USA* 87:9383–9387
91. Morrow JD, Awad JA, Boss HJ, Blair IA, Roberts LJ (1992) Non-cyclooxygenase derived prostanoids (F₂-isoprostanes) are formed in situ on phospholipids. *Proc Natl Acad Sci USA* 89:10721–10725
92. Nanji AA, Khwaja S, Tahan SR, Sadrzadeh SM (1994) Plasma levels of a novel noncyclooxygenase-derived prostanoid (8-isoprostane) correlate with severity of liver injury in

- experimental alcoholic liver disease. *J Pharmacol Exp Ther* 269:1280–1285
93. Nordblom GD, Coon MJ (1977) Hydrogen peroxide formation and stoichiometry of hydroxylation reactions catalyzed by highly purified liver microsomal cytochrome P₄₅₀. *Arch Biochem Biophys* 180:343–347
 94. Nordmann R, Ribière C, Rouach H (1992) Implication of free radical mechanisms in ethanol-induced cellular injury. *Free Radic Biol Med* 12:219–240
 95. Ontko JA (1973) Effects of ethanol on the metabolism of free fatty acids in isolated liver cells. *J Lipid Res* 14:78–86
 96. Pemberton PW, Smith A, Warnes TW (2005) Non-invasive monitoring of oxidant stress in alcoholic liver disease. *Scand J Gastroenterol* 40:1102–1108
 97. Porta EA, Bergman BJ, Stein AA (1965) Acute alcoholic hepatitis. *Am J Pathol* 46:657–689
 98. Rao RK, Serth A, Sheth P (2004) Recent advances in alcoholic liver disease I. Role of intestinal permeability and endotoxemia in alcoholic liver disease. *Am J Physiol Gastrointest Liver Physiol* 286:G881–G884
 99. Reboucas G, Isselbacher KJ (1961) Studies on the pathogenesis of the ethanol-induced fatty liver. I. Synthesis and oxidation of fatty acids by the liver. *J Clin Invest* 40:1355–1362
 100. Recknagel RO, Ghoshal AK (1966) Quantitative estimation of peroxidative degeneration of rat liver microsomal and mitochondrial lipids after carbon tetrachloride poisoning. *Exp Mol Pathol* 5:413–426
 101. Recknagel RO, Ghoshal AK (1966) Lipoperoxidation as a vector in carbon tetrachloride hepatotoxicity. *Lab Invest* 15:132–148
 102. Reinke LA, Lai EK, Du Bose CM, McCay PB (1987) Reactive free radical generation in vivo in heart and liver of ethanol-fed rats: correlation with radical formation in vitro. *Proc Natl Acad Sci USA* 84:9223–9227
 103. Reinke LA, Kotake Y, McCay PB, Janzen EG (1991) Spin-trapping studies of hepatic free radicals formed following the acute administration of ethanol to rats: in vivo detection of 1-hydroxyethyl radicals with PBN. *Free Radic Biol Med* 11:31–39
 104. Reitz RC (1975) A possible mechanism for the peroxidation of lipids due to chronic ethanol ingestion. *Biochem Biophys Acta* 380:145–154
 105. Roberts LJ II, Morrow JD (2000) Measurement of isoprostanes to assess oxidant stress status in vivo. In: Yoshikawa T, Toyokuni S, Yamamoto Y (eds) *Free radicals in chemistry, biology and medicine*. Nashville, TN, pp 329–340
 106. Roberts LJ II, Morrow JD (2002) Products of the isoprostane pathway: unique bioactive compounds and markers of lipid peroxidation. *Cell Mol Life Sci* 59:808–820
 107. Rogers CQ, Ajmo JM, You M (2008) Adiponectin and alcoholic fatty liver disease. *IUBMB Life* 60:790–797
 108. Rubin E, Lieber CS (1968) Hepatic microsomal enzymes in man and rat: induction and inhibition by ethanol. *Science* 162:690–691
 109. Rubin E, Hutterer F, Lieber CS (1968) Ethanol increases hepatic smooth endoplasmic reticulum and drug-metabolizing enzymes. *Science* 159:1469–1470
 110. Rubin E, Beattie DS, Lieber CS (1970) Effects of ethanol on the biogenesis of mitochondrial membranes and associated mitochondrial functions. *Lab Invest* 23:620–627
 111. Schönberg A (1968) *Preparative organic photochemistry*. Springer, Berlin
 112. Schröder M (2008) Endoplasmic reticulum stress responses. *Cell Mol Life Sci* 65:862–894
 113. Slater TF (1972) *Free radical mechanisms in tissue injury*. Pion Limited, London
 114. Takada A, Ikegami F, Okamura Y, Asumura Y, Kanayama R, Takeuchi J (1970) Effect of alcohol on the liver of rats. III. The role of lipid peroxidation and sulphhydryl compounds in ethanol-induced liver injury. *Lab Invest* 23:421–428
 115. Tephly TR, Tinelli F, Watkins WD (1969) Alcohol metabolism: role of microsomal oxidation in vivo. *Science* 166:627–628
 116. Terelius Y, Lindros KO, Albano E, Ingelman-Sundberg M (1993) Isozyme-specificity of cytochrome P₄₅₀-mediated hepatotoxicity. In: Ruckpaul K, Rein H (eds) *Frontiers in biotransformation*, vol 8. Akademie Verlag, Berlin, pp 186–232
 117. Teschke R, Hasumura Y, Joly J-G, Ishii H, Lieber CS (1972) Microsomal ethanol-oxidizing system (MEOS): purification and properties of a rat liver system free of catalase and alcohol dehydrogenase. *Biochem Biophys Res Commun* 49:1187–1193
 118. Thompson GN (1956) *Alcoholism*. Thomas C.C. Ed, Springfield
 119. Thurman RG (1973) Induction of hepatic microsomal reduced nicotinamide adenine dinucleotide phosphate-dependent production of hydrogen peroxide by chronic prior treatment with ethanol. *Mol Pharmacol* 9:670–675
 120. Valenzuela A, Fernandez N, Fernandez V, Ugarte G, Videla LA, Guerra R, Villanueva A (1980) Effect of acute ethanol ingestion on lipoperoxidation and on the activity of the enzymes related to peroxide metabolism in rat liver. *FASEB Lett* 111:11–13
 121. Vallee BL (1960) Metal and enzyme interactions: correlation of composition, function and structure. In: Boyer PD, Lardy H, Myrbäck K (eds) *The enzymes*. Academic Press, New York, pp 225–276
 122. Videla L, Bernstein J, Isdrael I (1973) Metabolic alterations produced in the liver by chronic ethanol administration: increased oxidative capacity. *Biochem J* 134:507–514
 124. White RE, Coon MJ (1980) Oxygen activation by cytochrome P₄₅₀. *Annu Rev Biochem* 49:325–356
 125. Wirtschafter ZT, Cronyn MW (1964) Free radical mechanism for solvent toxicity. *Arch Environ Health* 9:186–191
 126. Wooles WR (1966) Depressed hepatic fatty acid oxidation as a factor in the etiology of acute ethanol-induced fatty liver. *Life Sci* 5:267–276
 123. You M, Crabb DW (2005) Recent advances in alcoholic liver disease. II. Minireview: molecular mechanisms of alcoholic fatty liver. *Am J Physiol Gastrointest Liver Physiol* 287:G4–G63