

# Etiology of Severe Non-malaria Febrile Illness in Northern Tanzania: A Prospective Cohort Study

John A. Crump<sup>1,2,3,4,5,6\*</sup>, Anne B. Morrissey<sup>1</sup>, William L. Nicholson<sup>7</sup>, Robert F. Massung<sup>7</sup>, Robyn A. Stoddard<sup>8</sup>, Renee L. Galloway<sup>8</sup>, Eng Eong Ooi<sup>9</sup>, Venance P. Maro<sup>4,5</sup>, Wilbrod Saganda<sup>10</sup>, Grace D. Kinabo<sup>4,5</sup>, Charles Muiruri<sup>1,3</sup>, John A. Bartlett<sup>1,3,4,5</sup>

**1** Division of Infectious Diseases and International Health, Department of Medicine, Duke University Medical Center, Durham, North Carolina, United States of America, **2** Department of Pathology, Duke University Medical Center, Durham, North Carolina, United States of America, **3** Duke Global Health Institute, Duke University, Durham, North Carolina, United States of America, **4** Kilimanjaro Christian Medical Centre, Moshi, Tanzania, **5** Kilimanjaro Christian Medical College, Tumaini University, Moshi, Tanzania, **6** Centre for International Health, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand, **7** Rickettsial Zoonoses Branch, Division of Vector-Borne Diseases, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America, **8** Bacterial Special Pathogens Branch, Division of High-Consequence Pathogens and Pathology, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America, **9** Emerging Infectious Diseases Signature Research Program, Duke-National University of Singapore Graduate Medical School, Singapore, Singapore, **10** Mawenzi Regional Hospital, Moshi, Tanzania

## Abstract

**Introduction:** The syndrome of fever is a commonly presenting complaint among persons seeking healthcare in low-resource areas, yet the public health community has not approached fever in a comprehensive manner. In many areas, malaria is over-diagnosed, and patients without malaria have poor outcomes.

**Methods and Findings:** We prospectively studied a cohort of 870 pediatric and adult febrile admissions to two hospitals in northern Tanzania over the period of one year using conventional standard diagnostic tests to establish fever etiology. Malaria was the clinical diagnosis for 528 (60.7%), but was the actual cause of fever in only 14 (1.6%). By contrast, bacterial, mycobacterial, and fungal bloodstream infections accounted for 85 (9.8%), 14 (1.6%), and 25 (2.9%) febrile admissions, respectively. Acute bacterial zoonoses were identified among 118 (26.2%) of febrile admissions; 16 (13.6%) had brucellosis, 40 (33.9%) leptospirosis, 24 (20.3%) had Q fever, 36 (30.5%) had spotted fever group rickettsioses, and 2 (1.8%) had typhus group rickettsioses. In addition, 55 (7.9%) participants had a confirmed acute arbovirus infection, all due to chikungunya. No patient had a bacterial zoonosis or an arbovirus infection included in the admission differential diagnosis.

**Conclusions:** Malaria was uncommon and over-diagnosed, whereas invasive infections were underappreciated. Bacterial zoonoses and arbovirus infections were highly prevalent yet overlooked. An integrated approach to the syndrome of fever in resource-limited areas is needed to improve patient outcomes and to rationally target disease control efforts.

**Citation:** Crump JA, Morrissey AB, Nicholson WL, Massung RF, Stoddard RA, et al. (2013) Etiology of Severe Non-malaria Febrile Illness in Northern Tanzania: A Prospective Cohort Study. *PLoS Negl Trop Dis* 7(7): e2324. doi:10.1371/journal.pntd.0002324

**Editor:** Mathieu Picardeau, Institut Pasteur, France

**Received:** March 22, 2013; **Accepted:** June 9, 2013; **Published:** July 18, 2013

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

**Funding:** This research was supported by an International Studies on AIDS Associated Co-infections (ISAAC) award, a United States National Institutes of Health (NIH) funded program (U01 AI062563). Investigator support was received from NIH awards ISAAC (JAC ABM VPM GDK CM JAB); AIDS International Training and Research Program (D43 PA-03-018 to JAC VPM GDK CM JAB); the Duke Clinical Trials Unit and Clinical Research Sites (U01 AI069484 to JAC VPM GDK CM JAB); the Duke Center for AIDS Research (P30 AI 64518 to JAB); the Center for HIV/AIDS Vaccine Immunology (U01 AI067854 to JAC, JAB); and the joint NIH-NSF Ecology of Infectious Disease program and the UK Economic and Social Research Council and Biotechnology and Biological Sciences Research Council (R01TW009237 to JAC, VPM, WS, CM). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: john.crump@duke.edu

## Introduction

Fever without a localized cause is one of the most common presenting complaints among persons seeking healthcare in many low- and middle-income countries [1,2]. However, unlike the syndromes of pneumonia and diarrhea that feature in global disease burden estimates and have well coordinated programs integrating efforts across the range of responsible pathogens to avert morbidity and mortality, there has been a lack of a coordinated approach for febrile illness. While illness and death due to some specific infections causing fever, such as malaria [3] and increasingly bacterial sepsis are well quantified [4–6], others

such as a range of zoonoses and viral infections are uncounted and consequently may be underappreciated.

The various etiologies of febrile illnesses are difficult to distinguish from one another clinically [7,8]. As clinical laboratory services are often limited in areas where febrile conditions are particularly common [9,10], clinicians may have few diagnostic tools to establish an etiologic diagnosis. Therefore, clinical management is often driven by syndrome-based guidelines employing empiric treatment [11–13]. In the absence of systematically collected data on fever etiology, considerable mismatch between clinical diagnosis, clinical management, and actual etiology may occur resulting in poor patient outcomes [14]. It is

## Author Summary

The syndrome of fever is caused by a large number of infectious diseases. Malaria is thought to have been declining in the tropics since 2004. Increasing use of malaria diagnostic tests reveal a growing proportion of patients with fever who do not have malaria. While malaria diagnostic tests may be available, healthcare workers have few tools to diagnose causes of fever other than malaria. In order to identify major causes of fever other than malaria in northern Tanzania, we studied 870 patients with fever who were sufficiently ill to require admission to hospital. Malaria was uncommon and over-diagnosed, whereas invasive infections, including bloodstream infections, were underappreciated. Infections associated with animals such as brucellosis, leptospirosis, Q fever, and spotted fever group rickettsioses as well as viral infections transmitted by mosquitoes were common yet overlooked. We recommend that research on the syndrome of fever in resource-limited areas should focus on a wide range of potential causes. Animal-associated infections should be prioritized in patient management and disease control.

increasingly recognized that malaria is over-diagnosed in many areas [14,15]. To address this problem, the World Health Organization (WHO) malaria treatment guidelines moved away from clinical diagnosis of malaria to treatment based on the results of a malaria diagnostic test such as a blood smear or a malaria rapid diagnostic test. With more widespread availability of diagnostic tests to exclude malaria and apparent declines in malaria worldwide [3], clinicians in resource-limited areas are faced with a growing proportion of febrile patients who do not have malaria and few tools to guide subsequent management.

We sought to describe comprehensively the causes of febrile illness in northern Tanzania among patients sufficiently ill to require hospitalization. Febrile patients admitted to two hospitals were evaluated for a wide range of infectious etiologies using conventional standard diagnostic techniques.

## Methods

### Ethics statement

This study was approved by the Kilimanjaro Christian Medical Centre (KCMC) Research Ethics Committee, the Tanzania National Institutes for Medical Research National Research Ethics Coordinating Committee, and Institutional Review Boards of Duke University Medical Center and the CDC. All minors had written informed consent given from a parent or guardian and all adult participants provided their own written informed consent.

### Setting

Moshi (population, >144 000) is the administrative center of the Kilimanjaro Region (population, >1.4 million) in northern Tanzania and is situated at an elevation of 890 m above mean sea level. The climate is characterized by a long rainy period (March–May) and a short rainy period (October–December) [16]. Malaria transmission intensity is low [17]. KCMC is a consultant referral hospital with 458 inpatient beds serving several regions in northern Tanzania, and Mawenzi Regional Hospital (MRH), with 300 beds, is the Kilimanjaro Regional hospital. Together KCMC and MRH serve as the main providers of hospital care in the Moshi area. In 2008, KCMC admitted 22,099 patients and MRH admitted 21,763 patients.

## Study design

A study team that was independent of the hospital clinical team identified participants among infants and children admitted to KCMC from 17 September 2007 through 25 August 2008, and among adolescents and adult admitted to KCMC and MRH in Moshi, Tanzania, from 17 September 2007 through 31 August 2008. The methods of these studies have been described in detail elsewhere [7,8]. In brief, all admitted patients were screened for eligibility by study team members as soon as possible after admission and no later than 24 hours after admission. Infants and children aged from  $\geq 2$  months to  $< 13$  years, with a history of fever in the past 48 h or an axillary temperature  $\geq 37.5^{\circ}\text{C}$  or a rectal temperature of  $\geq 38.0^{\circ}\text{C}$ , and adolescents and adults aged  $\geq 13$  years and with oral temperatures of  $\geq 38.0^{\circ}\text{C}$  were invited to participate in the study. Patients admitted with known malignancy, renal failure, hepatic failure, bone marrow aplasia, trauma or surgery were excluded.

A standardized clinical history and physical examination were performed on consenting patients by a trained clinical officer who was a member of the study team and who worked in parallel with the hospital admitting team. Provisional diagnoses by the hospital clinical team made independently of the study team were recorded and coded using the International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10) codes. Following cleansing of the skin with isopropyl alcohol and povidone iodine, blood was drawn from adults and adolescents for aerobic blood culture (5 mL) and for mycobacterial blood culture (5 mL) and from pediatric patients for a single aerobic blood culture (4 mL). In addition, blood was drawn for complete blood count, examination for blood parasites, and HIV antibody testing. Acute serum, plasma, and whole blood were archived on all participants. For patients found to be HIV seropositive, CD4-positive T lymphocyte count (CD4 cell count) and serum cryptococcal antigen level were also measured. HIV-seronegative patients were screened for the presence of acute HIV infection by polymerase chain reaction (PCR) for HIV-1 RNA. Urine was collected as soon as possible after admission for detection of urine antimicrobial activity and for antigen detection. A discharge form was completed at the time of discharge from the hospital that captured whether the patient died in hospital, the in-hospital management, and the discharge diagnoses coded using ICD-10 codes. The results of study investigations done in Moshi were provided immediately to the hospital clinical team to inform patient management. The results of investigations done at reference laboratories were provided to the hospital clinical team as they became available. The hospital clinical team was responsible for all aspects of patient management, following clinical judgment and use of locally adapted and developed treatment guidelines. All participants were asked to return to a study clinic 4–6 weeks after enrollment to provide a convalescent serum sample. To promote high levels of follow up, the study team provided a follow up appointment card prior to hospital discharge, made reminder telephone calls to participants during the week prior to the appointment, reimbursed travel expenses of returning participants, and when necessary a field worker made home visits.

### Laboratory evaluations

Laboratory evaluations were selected to reflect a range of infectious diseases that might occur in northern Tanzania. Priority was given to laboratory evaluations for infectious diseases that might require specific management.

**Malaria.** Thick and thin blood films stained with Giemsa were examined for blood parasites by oil immersion microscopy. Parasite density was determined by standard methods [18].

**Bacteria and fungal bloodstream infections.** Blood culture bottles were assessed for volume adequacy comparing the weight before and after inoculation with blood. Adequate volume was defined as the recommended volume  $\pm 20\%$ . BacT/ALERT standard aerobic and mycobacterial bottles were loaded into the BacT/ALERT 3D Microbial Detection system (BioMérieux), where they were incubated for 5 and 42 days, respectively. Standard methods were used for identifying bloodstream isolates [7,8].

**Serum antigen testing.** Cryptococcal antigen level was measured using the Latex Cryptococcal Antigen Detection System assay (Immuno-Mycologies).

**Urine antigen testing.** Urine was tested for all participants for *Legionella pneumophila* serogroup 1 antigen using the Binax NOW *Legionella* urinary antigen test, and for adolescents and adults using the *Streptococcus pneumoniae* using the Binax NOW *S. pneumoniae* antigen test (Binax). Urine was tested for *Histoplasma capsulatum* antigen using the MVista *H. capsulatum* quantitative antigen enzyme immunoassay (Miravista Diagnostics) [19,20].

**Leptospirosis.** Leptospirosis laboratory diagnosis was made using the standard microscopic agglutination test (MAT) performed at the CDC. Live leptospiral cell suspensions representing 20 serovars and 17 serogroups described elsewhere [21] were incubated with serially diluted serum specimens. Resulting agglutination titers were read using darkfield microscopy. The reported titer was the highest dilution of serum that agglutinated at least 50% of the cells for each serovar tested [22]. Confirmed leptospirosis was defined as a  $\geq 4$ -fold rise in the agglutination titer between acute and convalescent serum samples [23].

**Brucellosis.** Brucellosis serology was performed using the standard microagglutination test (MAT) performed at the CDC. Standardized *Brucella abortus* strain 1119-3 killed antigen (National Veterinary Services Laboratory, Ames, IA) was used for MAT at a 1:25 working dilution described elsewhere [24]. Results were read on a Scienceware Plate Reader (Bel-Art Products, Wayne, NJ). Minor modifications were made to the CDC's standard MAT, including the use of U-bottom plates, incubation at 26°C, and discontinued use of staining techniques [25]. Confirmed brucellosis was defined as a  $\geq 4$ -fold rise in the agglutination titer between acute and convalescent serum samples.

**Q fever.** Convalescent-phase serum samples were screened using *C. burnetii* immunoglobulin (Ig) G enzyme-linked immunosorbent assay (ELISA) against Phase II antigen (Inverness Medical Innovations). For samples that were either positive or equivocal by ELISA, paired serum samples were tested by indirect immunofluorescence antibody (IFA) IgG assay to *C. burnetii* (Nine Mile strain) Phase I and Phase II antigens. A fourfold or greater increase in IFA reciprocal titer to Phase II antigen defined acute Q fever [26].

**Spotted fever group and typhus group rickettsioses.** Serum samples were tested for SFGR and TGR by IgG IFA to *R. conorii* (Moroccan strain) and to *R. typhi* (Wilmington strain), respectively. Among paired serum samples, a fourfold or greater increase in IFA titer to *R. conorii* and *R. typhi* defined acute SFGR and TGR infections, respectively [26].

**Arboviruses.** RNA was extracted from serum samples using the QIAamp Viral RNA Mini kit (QIAGEN, Hilden, Germany). Reverse transcription was performed using Invitrogen Superscript III First Strand Synthesis System (Life Technologies, Carlsbad, CA). Real-time PCRs for flavivirus, DENV, and CHIKV were carried out with the LightCycler 480 SYBR Green I Master kit (Roche Diagnostics, Penzberg, Germany) in a total reaction volume of 20  $\mu$ L containing 2  $\mu$ L of cDNA using primers published elsewhere [27–29]. Confirmed acute CHIKV, DENV,

and flavivirus infections were defined as a positive PCR result for CHIKV, DENV, and flavivirus viral RNA, respectively [30].

**HIV.** HIV-1 antibody testing was done on whole blood using both the Capillus HIV-1/HIV-2 (Trinity Biotech) and Determine HIV-1/HIV-2 (Abbott Laboratories) rapid HIV antibody tests. The Capillus test was replaced with the SD Bioline HIV-1/HIV-2 test (version 3.0; Standard Diagnostics) on 4 March 2008 after a change in Tanzania Ministry of Health HIV testing guidelines. If rapid tests were discordant, the sample was tested using enzyme-linked immunosorbent assay (ELISA; Vironostika Uni-Form II plus O Ab; bioMérieux). If the ELISA was negative, no further testing was done. If the ELISA was positive, a Western blot (Genetic Systems HIV-1 Western blot kit; Bio-Rad) was done to confirm the result [31]. HIV-1 RNA PCR was done using the Abbott m2000 system RealTime HIV-1 assay (Abbott Laboratories) [32,33].

### Statistic analysis

Data were entered using the Cardiff Teleform system (Cardiff Inc., Vista, Ca., USA) into an Access database (Microsoft Corp., Va., USA). When a diagnostic test was not applied to the whole cohort due lack of availability of an acute or convalescent sample, the proportion of confirmed cases in the tested group was extrapolated to the untested group by assuming that prevalence was the same in the tested group as in the untested group. Statistical analyses were performed with SAS version 9.1 software (SAS Inc, Cary, NC).

## Results

### Participant characteristics


Figure 1 summarizes participant screening, enrollment, and diagnostic testing. Of 870 febrile admissions to two hospitals in northern Tanzania enrolled in the study 484 (55.6%) were female. Of participants, 467 (53.7%) were infants and children with a median (range) age of 2 years (2 months - 13 years); the remainder adolescents and adults with a median (range) age of 38 (14–96) years. Fifty seven (12.2%) infants and children were HIV-infected compared with 157 (39.0%) adolescents and adults. Among infants and children 34 (7.3%) of 464 with hospital outcome data died; 2 (5.9%) of those who died had invasive infections. Among adolescents and adults, 41 (10.3%) of 398 with hospital outcome data died; 11 (26.8%) of those who died had invasive infections. In hospital deaths could not be attributed to etiologies requiring serologic diagnosis due to the requirement for testing a convalescent serum sample.

### Proportions of febrile admissions attributed to specific etiologies

Table 1 shows the number of patients with acute and convalescent samples available for testing for each etiologic agent or group of etiologic agents. Not all tests could be applied to all participants because of limited volumes of sample for some participants, and by the lack of availability of convalescent serum for participants who died before the follow up visit or who did not return. The number of confirmed cases in each group is also shown. The proportion of febrile admissions attributed to each etiology is calculated. A complete sample set was available for 243–467 (52.0–100.0%) infants and children and for 207–403 (51.4–100.0%) adolescents and adults.

### Etiology of fever among infants and children

Of 467 infants and children enrolled, malaria was the clinical diagnosis for 282 (60.4%), but was the actual cause of fever in 6



**Figure 1. Study flow diagram.** KCMC: Kilimanjaro Christian Medical Centre; MRH: Mawenzi Regional Hospital; MAT: microagglutination test; IFA: immunofluorescence assay; NAAT: nucleic acid amplification test. doi:10.1371/journal.pntd.0002324.g001

(1.3%). Bacterial and fungal bloodstream infections described in detail elsewhere [8] accounted for 16 (3.4%) and 4 (0.9%) febrile admissions, respectively, and were underrepresented on admission differential diagnoses. Bacterial zoonoses were identified among 49 (20.2%) of febrile admissions; 5 (2.0%) had brucellosis, 19 (7.7%) leptospirosis, 7 (2.6%) had Q fever, 18 (7.4%) had spotted fever group rickettsioses, and none had typhus group rickettsioses. In addition, 34 (10.2%) of participants had a confirmed acute arbovirus infection, all due to chikungunya (Table 1). No patient had a bacterial zoonoses or an arbovirus infection included in the admission differential diagnosis.

**Etiology of fever among adolescents and adults**

Of 403 adolescents and adults enrolled, malaria was the clinical diagnosis for 254 (63.0%), but was the actual cause of fever in 8 (2.0%). Bacterial, mycobacterial, and fungal bloodstream infections described in detail elsewhere [7] accounted for 69 (17.1%), 14 (3.5%), and 21 (5.2%) febrile admissions, respectively, and were underrepresented on admission differential diagnoses. Bacterial zoonoses were identified among 69 (33.3%) of febrile admissions; 11 (5.3%) had brucellosis, 21 (10.1%) leptospirosis, 17 (7.9%) had Q fever, 18 (8.7%) had spotted fever group rickettsioses, and 2 (1.0%) had typhus group rickettsioses. In addition, 21 (5.7%) of participants had a confirmed acute arbovirus infection, all due to chikungunya (Table 1). No patient had a bacterial zoonosis or an arbovirus infection included in the admission differential diagnosis.

**Etiology of fever overall**

Among all 870 participants, malaria was the clinical diagnosis for 528 (60.7%), but was the actual cause of fever in 14 (1.6%). By contrast, bacterial, mycobacterial, and fungal bloodstream

infections accounted for 85 (9.8%), 14 (1.6%), and 25 (2.9%) febrile admissions, respectively, and were underrepresented on admission differential diagnoses. Bacterial zoonoses were identified among 118 (26.2%) of febrile admissions; 16 (13.6%) had brucellosis, 40 (33.9%) leptospirosis, 24 (20.3%) had Q fever, 36 (30.5%) had spotted fever group rickettsioses, and 2 (1.8%) had typhus group rickettsioses. In addition, 55 (7.9%) of participants had a confirmed acute arbovirus infection, all due to chikungunya (Table 1). No patient had a bacterial zoonoses or an arbovirus infection included in the admission differential diagnosis. The proportional etiology of febrile illness among study participants after extrapolating to the untested group is summarized in Figure 2.

**Discussion**

We demonstrate among hospitalized febrile patients in northern Tanzania that malaria is uncommon and over-diagnosed, while invasive bacterial, mycobacterial, and fungal infections are underappreciated. At the same time, the bacterial zoonoses leptospirosis, Q fever, and spotted fever rickettsioses, and to a lesser extent brucellosis, and the arbovirus infection chikungunya are common yet unrecognized causes of fever. Our findings point to important mismatches between clinical diagnosis and management with actual diagnoses that have major implications for patient care, disease control and prevention, and for judicious use of antimalarial medications.

While the problem of malaria over-diagnosis has been appreciated for some time [14,15], studies that comprehensively describe the causes of severe non-malaria fever requiring hospital admission beyond bloodstream infections have been lacking. The

**Table 1.** Calculation of the proportion of hospitalized infants and children, and adolescents and adults, with specific etiologies of febrile illness, northern Tanzania, 2007–8.

Etiology	Infants and children			Adults and adolescents			All		
	n confirmed cases	n tested	(%)	n confirmed cases	n tested	(%)	n confirmed cases	n tested	(%)
<b>Bloodstream infections</b>									
Bacterial	16	467	(3.4)	69	403	(17.1)	85	870	(9.8)
Mycobacterial	0	467	(0.0)	14	403	(3.5)	14	870	(1.6)
Fungal	4	467	(0.9)	21	403	(5.2)	25	870	(2.9)
Malaria	6	467	(1.3)	8	403	(2.0)	14	870	(1.6)
Subtotal	26	467	(5.6)	112	403	(27.8)	138	870	(15.9)
<b>Bacterial zoonoses</b>									
Brucellosis	5	246	(2.0)	11	207	(5.3)	16	453	(3.5)
Leptospirosis	19	246	(7.7)	21	207	(10.1)	40	453	(8.8)
Q fever	7	268	(2.6)	17	215	(7.9)	24	482	(5.0)
Spotted fever group rickettsioses	18	243	(7.4)	18	207	(8.7)	36	450	(8.0)
Typhus group rickettsioses	0	243	(0.0)	2	207	(1.0)	2	450	(0.4)
Subtotal	49	243	(20.2)	69	207	(33.3)	118	450	(26.2)
<b>Arboviruses</b>									
Chikungunya	34	332	(10.2)	21	368	(5.7)	55	700	(7.9)
Flaviviruses	0	332	(0.0)	0	368	(0.0)	0	700	(0.0)
Subtotal	34	332	(10.2)	21	368	(5.7)	55	700	(7.9)
No diagnosis			(64.0)			(33.2)			(50.1)

Due to changing denominators for individual diagnostic tests, the proportion with no diagnosis is calculated as the proportion without a positive result from any test. Bloodstream infections are those diagnosed predominantly by blood culture, including organisms such as *Salmonella enterica*, *Streptococcus pneumoniae*, *Cryptococcus neoformans*, and *Mycobacterium tuberculosis*. Bacterial zoonoses, including brucellosis, leptospirosis, Q fever, and rickettsioses were diagnosed predominantly by serology, based on a 4-fold or greater rise in antibody titer between an acute and convalescent sample.  
doi:10.1371/journal.pntd.0002324.t001



over-diagnosis of malaria results in inappropriate use of antimalarial medications and may be associated with higher case fatality rates among patients treated for malaria who do not have the infection [14,15,34]. While the underlying causes of the over-diagnosis of malaria are complex [35], the lack of epidemiologic information about the importance of alternative infections and guidance on their management is likely to play a role. Our findings confirm the potential benefits of making reliable malaria diagnostic tests available at healthcare facilities and using the results as the basis for prescription of antimalarial medications [36]. When adopted, such an approach to malaria treatment would support the judicious use of antimalarials and would define the population of patients with nonmalaria fever.

We found that the bacterial zoonoses, leptospirosis, Q fever, and spotted fever group rickettsioses, and to a lesser extent brucellosis, are major causes of febrile illness among patients sufficiently unwell to require hospitalization. That a group of neglected bacterial zoonoses are of major clinical and public health importance in sub-Saharan Africa is a new and paradigm-changing finding. For clinical practice, with the exception of leptospirosis that may be effectively treated with commonly prescribed antibacterials, patients with brucellosis, Q fever, and the rickettsioses are likely to leave hospital without specific treatment. In northern Tanzania where many rely on livestock for their health and economic wellbeing, *Leptospira*, *Brucella*, and *Coxiella* spp. also indirectly affect human health through their impact on animal fertility, growth, and survival. The control and prevention of the neglected bacterial zoonoses is likely to involve interventions that require the collaboration of human health

experts with the animal and environmental health disciplines, an approach that is underdeveloped in many parts of the world.

Clinical guidelines for management of febrile patients in low resource areas focus on the identification and treatment of malaria and bacterial sepsis [11–13]. Our findings suggest that there is a need to identify and incorporate guidance on when to use a tetracycline for treatment of Q fever or rickettsial infection and when to consider treatment for brucellosis. We have previously demonstrated that features of the clinical history and physical examination do not perform well for identifying fever etiology [7,8,21,26,30]. Therefore, improvements to treatment algorithms for febrile patients are likely to require the development and incorporation of reliable diagnostic tests that provide timely diagnostic information to clinicians [37]. Unfortunately, many rapid diagnostic tests for infections related to fever management other than malaria and HIV suffer from poor performance characteristics [38,39].

Lack of coordination among groups working on the various etiologies of febrile illness in low-resource areas has meant that sentinel studies that could provide much more comprehensive information on a wide range of responsible organisms instead have focused on only one or a small handful of etiologies. For example, a clinical trial evaluating the impact of pneumococcal conjugate vaccine on rates of *Streptococcus pneumoniae* bacteremia in a community has the potential to identify and report all bloodstream infections. Similarly, a study designed to estimate the incidence of typhoid fever to inform vaccine policy could collect acute serum along with the blood culture and, with subsequent collection of convalescent serum, would have the ability to estimate the



**Figure 2. Laboratory confirmed causes of febrile illness among infants and children (panel A) and adolescents and adults (panel B) hospitalized in northern Tanzania, 2007–8\*.** \*In instances that diagnostic test results were not available for all participants, the proportion positive from Table 1 was applied to the whole study population. Pie graphs do not account for concurrent infections. A complete listing of specific bacterial, mycobacterial, and fungal bloodstream infections is available elsewhere [7,8].  
doi:10.1371/journal.pntd.0002324.g002

incidence of leptospirosis and a range of other etiologic agents using conventional serologic methods [40]. However, resources for research have tended to be targeted to specific pathogens and researchers have struggled to leverage additional resources to address a broader range of organisms. Sentinel site studies seeking to understand the infectious causes of febrile illness in low-resource settings have utilized blood culture to highlight the importance of invasive bacterial and fungal infections [4,41]. Expanding laboratory evaluations to include serologic and molecular approaches to diagnosing infections requiring specific antimicrobial management such as the bacterial zoonoses brucellosis, leptospirosis, Q fever, and the rickettsioses adds considerable value [40]. Detection of infections of public health importance such as those caused by the arboviruses dengue, Rift Valley fever, and yellow fever can inform national control programs. Since considerable etiologic overlap exists between the syndromes of fever, acute respiratory tract infection, and diarrhea [42,43], addressing these simultaneously in integrated sentinel studies would inform enhancements in empiric treatment guidelines and improvements in the accuracy of syndrome-based disease burden estimates.

Our study had a number of limitations. While we examined a wide range of etiologies of fever, a large proportion of patients were undiagnosed suggesting that we failed to identify potentially important infections. The undiagnosed group is being investigated further using pathogen discovery approaches. Some of the diagnostic tests used in our study are less than 100% sensitive and specific and we did not test for every known pathogen. As a consequence, we probably underestimated the prevalence of some infections while misclassifying others that were falsely positive. Because a number of our diagnostic tests relied on the demonstration of a four-fold rise in antibody titer between the acute and convalescent serum sample, not all enrolled patients returned for collection of convalescent serum to have diagnoses confirmed. It follows that calculation and comparison of case fatality rate was not possible since those who died before the convalescent visit could not be confirmed cases. Incomplete diagnostic information meant that we had to extrapolate prevalence from the tested population to the untested population, potentially introducing bias. Similarly, instances of apparent infection with multiple agents were not accounted for in presentation of pie graphs. Inclusion of a well control group would have allowed the calculation of attributable fractions for individual pathogens, something that should be considered for future febrile illness research, especially in areas where malaria is endemic. Since considerable geographic variation in fever etiology is known to occur, the generalizability of our findings is uncertain.

What is needed to support an integrated approach to the syndrome of fever in resource-limited areas? First, fever should be recognized alongside pneumonia and diarrhea as a major clinical syndrome of public health importance. Achieving this is likely to require leadership from international institutions of public health and reappraisal of the way that the febrile illnesses are approached

in burden of disease estimates. This could include estimating total morbidity and mortality from the syndrome of fever as a first step before attributing the associated illnesses and deaths to specific etiologies, much as is done for the other major syndromes [44,45]. Second, efforts are needed to bring together the diverse groups and disciplines currently working on the febrile illnesses to quantify the morbidity and mortality attributable to each major etiologic agent. Such integration could be facilitated by support for research efforts that study the syndrome of fever comprehensively as well as its etiologies individually, an approach that has been modeled by studies addressing the syndromes of pediatric pneumonia and diarrhea in developing countries [46,47]. Third, improved diagnostic services are urgently needed to establish disease burden estimates and patient management for the febrile illnesses in resource-limited areas [10]. Conventional diagnostic tests for some infections, such as leptospirosis, are complex. For example, the collection of both acute and convalescent serum samples may be required, and testing services may be available at only a few national or supra-national reference laboratories. Assays relying on convalescent samples cannot be used to estimate case fatality rates [21,26]. Conversely, simple, rapid tests applied to acute samples may have poor performance characteristics [38]. Finally, clinical studies, including clinical trials, are needed to test and improve clinical management algorithms for febrile patients. The goal should be to target antimicrobial therapy to those who need it and to avoid inappropriate use among patients who will not benefit. In this way, patient outcomes can be improved, health resources can be conserved, and disease prevention and control efforts for febrile conditions can be rationally resourced.

## Supporting Information

**Text S1 STROBE checklist.**  
(DOC)

## Acknowledgments

We acknowledge the Hubert-Yeargan Center for Global Health at Duke University for critical infrastructure support for the Kilimanjaro Christian Medical Centre-Duke University Collaboration. **Disclaimers:** The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention. Use of trade names and commercial sources is for identification only and does not imply endorsement by the US Department of Health and Human Services or the Centers for Disease Control and Prevention.

## Author Contributions

Conceived and designed the experiments: JAC JAB. Performed the experiments: JAC ABM WLN RFM RAS RLG EEO VPM WS GDK JAB. Analyzed the data: JAC. Contributed reagents/materials/analysis tools: WLN RFM RAS RLG EEO. Wrote the paper: JAC ABM WLN RFM RAS RLG EEO VPM WS GDK CM JAB. Managed the research program: CM. Sought and obtained funding: JAB.

## References

- Petit PL, van Ginneken JK (1995) Analysis of hospital records in four African countries, 1975–1990, with emphasis on infectious diseases. *J Trop Med Hyg* 98: 217–227.
- Feikin DR, Olack B, Bigogo GM, Audi A, Cosmas L, et al. (2011) The burden of common infectious syndromes at the clinic and household level from population-based surveillance in rural and urban Kenya. *PLoS ONE* 6: e16085.

3. Murray CJ, Rosenfeld LC, Lim SS, Andrews KG, Foreman KJ, et al. (2012) Global malaria mortality between 1980 and 2010: a systematic analysis. *Lancet* 379: 413–431.
4. Reddy EA, Shaw AV, Crump JA (2010) Community acquired bloodstream infections in Africa: a systematic review and meta-analysis. *Lancet Infect Dis* 10: 417–432.
5. Nadjim B, Amos B, Mtove G, Ostermann J, Chonya S, et al. (2010) WHO guidelines for antimicrobial treatment in children admitted to hospital in an area of intense *Plasmodium falciparum* transmission: prospective study. *BMJ* 340: c1350.
6. Nadjim B, Mtove G, Amos B, Walker NF, Diefenthal H, et al. (2012) Severe febrile illness in adult hospital admissions in Tanzania: a prospective study in an area of high malaria transmission. *Trans R Soc Trop Med Hyg* 106: 688–695.
7. Crump JA, Ramadhani HO, Morrissey AB, Saganda W, Mwako MS, et al. (2011) Invasive bacterial and fungal infections among hospitalized HIV-infected and HIV-uninfected adults and adolescents in northern Tanzania. *Clin Infect Dis* 52: 341–348.
8. Crump JA, Ramadhani HO, Morrissey AB, Msuya LJ, Yang L-Y, et al. (2011) Invasive bacterial and fungal infections among hospitalized HIV-infected and HIV-uninfected children and infants in northern Tanzania. *Trop Med Int Health* 16: 830–837.
9. Archibald LK, Reller LB (2001) Clinical microbiology in developing countries. *Emerg Infect Dis* 7: 302–305.
10. Petti CA, Polage CR, Quinn TC, Ronald AR, Sande MA (2006) Laboratory medicine in Africa: a barrier to effective health care. *Clin Infect Dis* 42: 377–382.
11. World Health Organization (2011) IMAI district clinician manual: hospital care for adolescents and adults: guidelines for the management of illnesses with limited resources. Geneva: World Health Organization.
12. World Health Organization (2005) Pocket book of hospital care for children: guidelines for the management of common illnesses with limited resources. Geneva: World Health Organization. 378 p.
13. World Health Organisation Division of Diarrhoea and Acute Respiratory Disease Control, United Nations Children's Fund (1995) Integrated management of the sick child. *Bull World Health Organ* 73: 735–740.
14. Reyburn H, Mbatia R, Drakeley C, Carneiro I, Mwakasungula E, et al. (2004) Overdiagnosis of malaria in patients with severe febrile illness in Tanzania: a prospective study. *BMJ* 329: 1212–1215.
15. Amexo M, Tolhurst R, Barnish G, Bates I (2004) Malaria misdiagnosis: effects on the poor and vulnerable. *Lancet* 364: 1896–1898.
16. National Bureau of Statistics, ORC Macro (2005) Tanzania Demographic and Health Survey 2004–05. Dar es Salaam. 381 p.
17. Hay SI, Guerra CA, Gething PW, Patil AP, Tatem AJ, et al. (2009) A world malaria map: *Plasmodium falciparum* endemnicity in 2007. *PLoS Med* 6: e1000048.
18. Greenwood BM, Armstrong JRM (1991) Comparison of two simple methods for determining malaria parasite density. *Trans R Soc Trop Med Hyg* 85: 186–188.
19. Connolly PA, Durkin MM, LeMonte AM, Hackett EJ, Wheat JL (2007) Detection of *Histoplasma* antigen by a quantitative enzyme immunoassay. *Clin Vaccine Immunol* 14: 1587–1591.
20. Lofgren SM, Kirsch EJ, Maro VP, Morrissey AB, Msuya LJ, et al. (2012) Histoplasmosis among hospitalized febrile patients in northern Tanzania. *Trans R Soc Trop Med Hyg* 106: 504–507.
21. Biggs HM, Bui DM, Galloway RL, Stoddard RA, Shadomy SV, et al. (2011) Leptospirosis among hospitalized febrile patients in northern Tanzania. *Am J Trop Med Hyg* 85: 275–281.
22. Dikken H, Kmety E (1978) Serological typing methods of leptospire. In: Bergan T, Norris JR, editors. *Methods in microbiology*. London: Academic Press. pp. 259–307.
23. Centers for Disease Control and Prevention (1997) Case definitions for infectious conditions under public health surveillance. *MMWR Recomm Rep* 46 (RR-10): 1–55.
24. Bouley AJ, Biggs HM, Stoddard RA, Morrissey AB, Bartlett JA, et al. (2012) Brucellosis among hospitalized febrile patients in northern Tanzania. *Am J Trop Med Hyg* 87: 1105–1111.
25. Brown SL, Klein GC, McKinney FT, Jones WL (1981) Safranin O-stained antigen microagglutination test for detection of *Brucella* antibodies. *J Clin Microbiol* 13: 398–400.
26. Prabhu M, Nicholson WL, Roche AJ, Kersh GJ, Fitzpatrick KA, et al. (2011) Q fever, spotted fever group and typhus group rickettsioses among hospitalized febrile patients in northern Tanzania. *Clin Infect Dis* 53: e8–e15.
27. Lai YL, Chung YK, Tan HC, Yap HF, Yap G, et al. (2007) Cost-effective real-time reverse transcriptase PCR (RT-PCR) to screen for dengue virus followed by rapid single-tube multiplex RT-PCT for serotyping the virus. *J Clin Microbiol* 45: 935–941.
28. Pastorino B, Bessaud M, Grandadam M, Murri S, Tolou HJ, et al. (2005) Development of a TaqMan RT-PCR assay without RNA extraction for the detection and quantification of African chikungunya viruses. *J Virol Methods* 124: 65–71.
29. Scaramozzino N, Crance JM, Jouan A, DeBriel DA, Stoll F, et al. (2001) Comparison of flavivirus universal primer pairs and development of a rapid, highly sensitive heminested reverse transcriptase-PCR assay for detection of flaviviruses targeting to a conserved region of the NS5 gene sequences. *J Clin Microbiol* 39: 1922–1927.
30. Hertz JT, Munishi OM, Ooi EE, Howe S, Lim WY, et al. (2012) Chikungunya and dengue fever among hospitalized febrile patients in northern Tanzania. *Am J Trop Med Hyg* 86: 171–177.
31. Mayhood MK, Afwamba IA, Odhiambo CO, Ndanu E, Thielman NM, et al. (2008) Validation, performance under field conditions, and cost-effectiveness of Capillus HIV-1/HIV-2 and Determine HIV-1/2 rapid human immunodeficiency virus antibody assays using sequential and parallel testing algorithms in Tanzania. *J Clin Microbiol* 46: 3946–3951.
32. Scott LE, Crump JA, Msuya E, Morrissey AB, Venter WF, et al. (2011) Abbott RealTime HIV-1 *m2000t* viral load testing: manual extraction versus the automated *m2000sp* extraction. *J Virol Methods* 172: 78–80.
33. Crump JA, Scott LE, Msuya E, Morrissey AB, Kimaro EE, et al. (2009) Evaluation of the Abbott *m2000t* RealTime HIV-1 assay with manual sample preparation compared with the ROCHE COBAS AmpliPrep/AMPLICOR HIV-1 MONITOR 1.5 using specimens from East Africa. *J Virol Methods* 162: 218–222.
34. Clifton DC, Ramadhani HO, Msuya LJ, Njau BN, Kinabo GD, et al. (2012) Predicting death for paediatric inpatients where malaria is uncommon. *Arch Dis Child* 97: 889–894.
35. Chandler CIR, Jones C, Boniface G, Juma K, Reyburn H, et al. (2008) Guidelines and mindlines: why do clinical staff over-diagnose malaria in Tanzania? A qualitative study. *Malaria J* 7: 53.
36. World Health Organization (2010) Guidelines for the treatment of malaria, 2nd edition. Geneva: World Health Organization. 194 p.
37. Baltzell K, Elfving K, Shakely D, Ali AS, Msellem M, et al. (2013) Febrile illness management in children under five years of age: a qualitative pilot study on primary health care workers' practices in Zanzibar. *Malar J* 12: 37.
38. Bajani MD, Ashford DA, Bragg SL, Woods CW, Aye T, et al. (2003) Evaluation of four commercially available rapid serologic tests for diagnosis of leptospirosis. *J Clin Microbiol* 41: 803–809.
39. Olsen SJ, Pruckler J, Bibb W, Thanh NT, Trinh TM, et al. (2004) Evaluation of rapid diagnostic tests for typhoid fever. *J Clin Microbiol* 42: 1885–1889.
40. Crump JA, Youssef FG, Luby SP, Wasfy MO, Rangel JM, et al. (2003) Estimating the incidence of typhoid fever and other febrile illnesses in developing countries. *Emerg Infect Dis* 9: 539–544.
41. Berkley JA, Lowe BS, Mwangi I, Williams T, Banui E, et al. (2005) Bacteremia among children admitted to a rural hospital in Kenya. *N Engl J Med* 352: 39–47.
42. Graham SM, English M (2009) Non-typhoidal salmonellae: a management challenge for children with community-acquired invasive disease in tropical African countries. *Lancet* 373: 267–269.
43. O'Dempsey TJ, McArdle TF, Laurence BE, Lamont AC, Todd JE, et al. (1993) Overlap in the clinical features of pneumonia and malaria in African children. *Trans R Soc Trop Med Hyg* 87: 662–665.
44. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, et al. (2012) Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2197–2223.
45. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, et al. (2012) Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2095–2128.
46. Levine OS, O'Brien KL, Deloria-Knoll M, Murdoch DR, Feikin DR, et al. (2012) The Pneumonia Etiology Research for Child Health Project: a 21st century childhood pneumonia etiology study. *Clin Infect Dis* 54: S93–101.
47. Levine MM, Kotloff KL, Nataro JP, Muhsen K (2012) The Global Enteric Multicenter Study (GEMS): impetus, rationale, and genesis. *Clin Infect Dis* 55: S215–224.