

Evading the Grossman-Nir bound with $\Delta I = 3/2$ new physics

Xiao-Gang He,^{a,b} Xiao-Dong Ma,^a Jusak Tandean^{a,b} and German Valencia^c

^a*Department of Physics, National Taiwan University,
No. 1, Sec. 4, Roosevelt Rd., Taipei 106, Taiwan*

^b*Physics Division, National Center for Theoretical Sciences,
No. 101, Sec. 2, Kuang Fu Rd., Hsinchu 300, Taiwan*

^c*School of Physics and Astronomy, Monash University,
Wellington Road, Clayton, VIC-3800, Australia*

E-mail: hexg@phys.ntu.edu.tw, maxid@phys.ntu.edu.tw,
jtandean@phys.ntu.edu.tw, german.valencia@monash.edu

ABSTRACT: Rare kaon decays with missing energy, $K \rightarrow \pi + E_{\text{miss}}$, have received considerable attention because their rates can be calculated quite precisely within the standard model (SM), where the missing energy is carried away by an undetected neutrino-antineutrino pair. Beyond the SM, clean theoretical predictions can also be made regarding these processes. One such prediction is the so-called Grossman-Nir (GN) bound, which states that the branching fractions of the K_L and K^+ modes must satisfy the relation $\mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}}) \lesssim 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}})$ and applies within and beyond the SM, as long as the hadronic transitions change isospin by $\Delta I = 1/2$. In this paper we extend the study of these modes to include new-physics scenarios where the missing energy is due to unobserved lepton-number-violating neutrino pairs, invisible light new scalars, or pairs of such scalars. The new interactions are assumed to arise above the electroweak scale and described by an effective field theory. We explore the possibility of violating the GN bound through $\Delta I = 3/2$ contributions to the $K \rightarrow \pi$ transitions within these scenarios and find that large violations are only possible in the case where the missing energy is due to an invisible light new scalar.

KEYWORDS: Beyond Standard Model, Chiral Lagrangians, Effective Field Theories, Kaon Physics

ARXIV EPRINT: [2005.02942](https://arxiv.org/abs/2005.02942)

Contents

1	Introduction	1
2	GN-bound violation via EFT operators for $K \rightarrow \pi 2\nu$	4
2.1	EFT operators for $K \rightarrow \pi 2\nu$	4
2.2	Evaluation of hadronic matrix elements at low energies	6
2.3	Numerical analysis	9
3	GN-bound violation via EFT operators for $K \rightarrow \pi S$ or πSS	12
3.1	Operators and matching	12
3.2	Numerical analysis	16
3.2.1	$K \rightarrow \pi S$	17
3.2.2	$K \rightarrow \pi SS$	19
4	Summary and conclusions	20
A	Isospin decomposition of quark parts of dim-9 operators	22
B	RG running of dim-6 four-quark operators for $K \rightarrow \pi S(S)$	23
C	Chiral structure and hadronization of quark operators for $K \rightarrow \pi S(S)$	24

1 Introduction

The rare kaon decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ are known as the “golden modes” of kaon physics because they can be predicted quite precisely within the standard model (SM) and are potentially sensitive to the presence of new physics (NP) beyond it [1, 2]. Consequently they have received a great deal of theoretical and experimental attention. The SM predictions for their branching fractions have been known for many years [3–10], and their current values are [11–13] $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{SM}} = (3.0 \pm 0.2) \times 10^{-11}$ and $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} = (8.5 \pm 0.5) \times 10^{-11}$, which suffer mostly from parametric uncertainties. These numbers, and their counterparts in many NP scenarios, respect a particularly interesting prediction, the so-called Grossman-Nir bound [14], which states that $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \lesssim 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$. A key assumption behind this statement is that the $K \rightarrow \pi$ transition proceeds from an interaction which changes isospin by $\Delta I = 1/2$.

On the experimental side, the KOTO collaboration at J-PARC in 2018 set an upper limit on the neutral mode based on the data collected in 2015 [15]: $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{KOTO15}} < 3.0 \times 10^{-9}$ at 90% confidence level (CL). For the charged mode, the combined BNL E787/E949 experiments had earlier yielded $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{E949}} = (1.73_{-1.05}^{+1.15}) \times 10^{-10}$ [16, 17]. Last year the NA62 collaboration [18] at CERN reported the preliminary limit

$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{NA62}} < 1.85 \times 10^{-10}$ at 90% CL [19]. All of these results are in agreement with the SM but leave open a window for NP. This possibility is very intriguing, especially in light of the recent preliminary observation by KOTO of three candidate events in the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ signal region [20], with a single event sensitivity of 6.9×10^{-10} having been achieved. If interpreted as signal, they imply a decay rate about two orders of magnitude higher than the SM prediction and in conflict with the experimentally established GN bound $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{GN}} < 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{NA62}} = 8.0 \times 10^{-10}$ at 90% CL.

It is interesting that there are NP scenarios which can overcome this requisite. For instance, as was first pointed out in ref. [21], the branching fraction of $K_L \rightarrow \pi^0 \mathcal{X}$, with \mathcal{X} being an invisible particle with mass $m_{\mathcal{X}}$ chosen to be around the pion mass, can exceed the aforementioned cap of 8.0×10^{-10} because quests for the charged channel $K^+ \rightarrow \pi^+ \mathcal{X}$ do not cover the $m_{\mathcal{X}} \sim m_{\pi}$ region to avoid the sizable $K^+ \rightarrow \pi^+ \pi^0$ background [17, 18]. This kinematic loophole has been exploited in recent attempts [22–28] to account for KOTO’s anomalous events. As another example, imposing the condition $m_{K^+} - m_{\pi^+} < m_{\mathcal{X}} < m_{K^0} - m_{\pi^0}$ renders the $K^+ \rightarrow \pi^+ \mathcal{X}$ channel closed (here \mathcal{X} can be more than one invisible particle), whereas $K_L \rightarrow \pi^0 \mathcal{X}$ with a big rate can still happen [29]. For other $m_{\mathcal{X}}$ ranges, the stringent restriction on $K^+ \rightarrow \pi^+ \mathcal{X}$ can be evaded or weakened if \mathcal{X} is a particle with an average decay length bigger than the KOTO detector size but less than its E949 and NA62 counterparts [22–26]. Most of these cases, while still fulfilling the GN relation, only appear to contradict it by enhancing $\mathcal{B}(K_L \rightarrow \pi^0 \mathcal{X})$ substantially above $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{GN}}^{\text{max}}$. In a framework of effective field theory with only SM fields, the introduction of (predominantly) $\Delta I = 1/2$ interactions which change lepton flavor/number or possess new sources of CP violation also would not bring about a disruption of the GN bound [30, 31]. On the other hand, it has been proposed that in the presence of additional light particles mediating these decays the GN bound could be violated [32].

In terms of the branching-fraction ratio $r_{\mathcal{B}} = \mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) / \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, the GN bound has a theoretical model-independent maximum of $r_{\mathcal{B}}^{\text{GN}} = 4.3$, which differs much from the central value $r_{\mathcal{B}}^{\text{SM}} = 0.36$ of the SM prediction. A key ingredient in the derivation of the GN inequality is the assumption that the $K \rightarrow \pi$ transitions are mediated by a two-quark $s \leftrightarrow d$ operator, which necessarily carries isospin 1/2 and leads to a ratio of amplitudes for the neutral and charged modes given by $\mathcal{A}_{K^0 \rightarrow \pi^0}^{\Delta I=1/2} / \mathcal{A}_{K^+ \rightarrow \pi^+}^{\Delta I=1/2} = -1/\sqrt{2}$. A true violation of the GN relation requires the $K \rightarrow \pi$ transitions to occur via a $\Delta I = 3/2$ interaction as well, and this possibility has recently been investigated in refs. [31, 33]. A pure $\Delta I = 3/2$ operator would result in an amplitude ratio $\mathcal{A}_{K^0 \rightarrow \pi^0}^{\Delta I=3/2} / \mathcal{A}_{K^+ \rightarrow \pi^+}^{\Delta I=3/2} = \sqrt{2}$ and thus translate into $r_{\mathcal{B}}^{\Delta I=3/2} \lesssim 17$ [31].

In this paper we present a systematic study on how to overcome the GN bound with quark-level operators in the context of effective field theory (EFT). Given that the measurements of interest look for $K \rightarrow \pi \mathcal{X}$ with \mathcal{X} standing for one or more particles carrying away missing energy (E_{miss}), we will consider for \mathcal{X} several different possibilities: a neutrino-antineutrino pair ($\nu \bar{\nu}$), a pair of neutrinos ($\nu \nu$) or antineutrinos ($\bar{\nu} \bar{\nu}$), an invisible light new scalar boson (S), and a pair of these scalars (SS). Since an operator directly giving rise to $\Delta I = 3/2$ $K \rightarrow \pi \mathcal{X}$ transitions has to contain at least four light-quark fields, the minimal mass dimension of such an operator is seven, eight, nine, and ten for $\mathcal{X} = S, SS$,

$\nu\nu$ or $\bar{\nu}\bar{\nu}$, and $\nu\bar{\nu}$, respectively. Being unobserved in the experiments, the neutrinos may have different flavors and can also be replaced by new invisible light fermions. We suppose that the new particles are invisible because they are sufficiently long-lived to escape detection, decay invisibly, or are stable. We will look at a complete set of operators with the lowest dimension necessary for each of these cases, assuming that the interactions of the new light particles with the quarks can be described by an EFT approach valid above the electroweak-symmetry breaking scale.

Before embarking in a detailed study, it is instructive to present a simple dimensional-analysis estimate for the scale Λ_{NP} of NP that is needed in order to produce a rate of $K \rightarrow \pi\mathcal{X}$ that is comparable in size to the SM rate. To this end, it is useful to recall the effective Hamiltonian responsible for $K \rightarrow \pi\nu\bar{\nu}$ in the SM,

$$\mathcal{H}_{\text{SM}} = \frac{G_{\text{F}}}{\sqrt{2}} \frac{g^2 V_{ts}^* V_{td} X(x_t)}{16\pi^2} \bar{s}\gamma_{\mu}(1 - \gamma_5)d \sum_{\ell} \bar{\nu}_{\ell}\gamma^{\mu}(1 - \gamma_5)\nu_{\ell} + \text{H.c.} \quad (1.1)$$

in the conventional notation [13], where $X(x_t) \simeq 1.5$ from top-quark loops. In contrast, a generic NP operator which induces a $\Delta I = 3/2$ transition in the process $K \rightarrow \pi\mathcal{X}$ can be written as

$$\mathcal{L}_{\text{NP}} = \frac{1}{\Lambda_{\text{NP}}^{2+N_{\mathcal{X}}}} \mathcal{C} \bar{s}\Gamma_1 u \bar{u}\Gamma_2 d \mathcal{X} + \text{H.c.}, \quad (1.2)$$

where $N_{\mathcal{X}}$ tells the mass dimension of \mathcal{X} (so $N_{\nu\nu} = 3$, $N_S = 1$, $N_{SS} = 2$, etc.), the coefficient \mathcal{C} is a constant, and $\Gamma_{1,2}$ represent gamma matrices. At the amplitude level, the \mathcal{L}_{NP} contribution to $K \rightarrow \pi\mathcal{X}$ relative to the SM top-quark contribution is then

$$\frac{\mathcal{A}_{\text{NP}}}{\mathcal{A}_{\text{SM}}} \sim 3.8 \times 10^5 \mathcal{C} \left(\frac{v}{\Lambda_{\text{NP}}} \right)^2 \left(\frac{m_K}{\Lambda_{\text{NP}}} \right)^{N_{\mathcal{X}}}, \quad (1.3)$$

where the large numerical factor reflects the one-loop and CKM-angle suppression of the SM coefficient, $v = 2^{-1/4} G_{\text{F}}^{-1/2} = 246 \text{ GeV}$ indicates the electroweak scale, and the relevant hadronic scale is taken to be the kaon mass, m_K . If the NP is defined to enter \mathcal{L}_{NP} with \mathcal{C} of order one (so any loop or mixing-angle suppression factor is absorbed into Λ_{NP}), the result in eq. (1.3) implies that $\Lambda_{\text{NP}} \sim 2.2 \text{ TeV}$, 275 GeV , 78 GeV for $N_{\mathcal{X}} = 1, 2, 3$, respectively, correspond to NP effects at the same level as the SM contribution.

The EFT that we employ in this paper only makes sense for $\Lambda_{\text{NP}} > v$. Otherwise, the organization of effective operators in terms of their dimensionality breaks down. This suggests that scenarios with $\mathcal{X} = S$ could amplify the rate of $K_L \rightarrow \pi^0\mathcal{X}$ to the level implied by the three KOTO events; scenarios with $\mathcal{X} = SS$ may increase the corresponding rate compared to its SM counterpart by factors of 2; and that scenarios in which $\mathcal{X} = \nu\nu$ or $\bar{\nu}\bar{\nu}$ or $\nu\bar{\nu}$ can only modify the $K \rightarrow \pi + E_{\text{miss}}$ rates marginally.

We will explore all these possibilities in detail to quantify how the underlying NP interactions influence these kaon decays. To handle a low-energy process involving hadrons, it is necessary to hadronize the quark-level operators at the mass scale where it takes place. For our investigation, this entails computing the effects of QCD renormalization group (RG) running from the electroweak (EW) scale to the chiral-symmetry breaking scale and

subsequently matching the resulting operators to a low-energy chiral Lagrangian suitable to describe the $K \rightarrow \pi + E_{\text{miss}}$ transitions. In all the cases, we present numerical results illustrating the range of values which the ratio $r_{\mathcal{B}}$ can take when the NP scale is under a couple of TeV.

The arrangement of the rest of the paper is as follows. In section 2 we study how the GN bound can be violated through $K \rightarrow \pi\nu\nu, \pi\bar{\nu}\bar{\nu}$ caused by $\Delta I = 3/2$ interactions in the EFT framework where the operators respect the SM gauge symmetries (SMEFT). In section 3 we extend the SMEFT with the addition of a SM-gauge-singlet scalar S and carry out a similar analysis with $K \rightarrow \pi S, \pi S S$. In section 4, we draw our conclusions. We relegate some extra details to appendices, including the RG running of the four-quark parts of the operators pertaining to $K \rightarrow \pi S(S)$ and their low-energy chiral bosonization.

2 GN-bound violation via EFT operators for $K \rightarrow \pi 2\nu$

2.1 EFT operators for $K \rightarrow \pi 2\nu$

In this section we restrict the fields of the EFT to only those in the minimal SM. Accordingly, in the kaon decays of concern the missing energy is carried away by a pair of SM neutrinos (2ν). It may have no or nonzero lepton number depending on whether the underlying interaction is lepton-number conserving or violating, respectively. If the small contributions to $K \rightarrow \pi 2\nu$ from long-distance physics [1, 30, 34–36] are neglected,¹ the only possible way to break the GN bound significantly in this case is through the inclusion of $\Delta I = 3/2$ operators having nonleptonic parts with at least four quark fields [31, 33]. It follows that the lowest dimension of quark-neutrino operators with $\Delta I = 3/2$ components is nine.

In the SMEFT treatment the effective operators constructed must be singlets under the SM gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$, while in the low-energy effective field theory (LEFT) the operators are to be singlets only under the strong and electromagnetic gauge group $SU(3)_C \times U(1)_{\text{em}}$. This means that there are in general more requirements on the SMEFT operators than on the LEFT ones, which makes the number of quark operators relatively less in the former case. Since furthermore there is still no discovery of new particles beyond the SM, we will rely on the SMEFT to perform our examination.

The fundamental fields (with their SM gauge group assignments) available to construct the pertinent operators are the $U(1)_Y$ gauge field $B(1, 1, 0)$, the $SU(2)_L$ gauge field $W(1, 3, 0)$, the $SU(3)_C$ gluon field $G(8, 1, 0)$, the Higgs doublet $H(1, 2, 1/2)$, the left-handed quark doublet $Q(3, 2, 1/6)$, the right-handed quark fields $u(3, 1, 2/3)$ and $d(3, 1, -1/3)$, the left-handed lepton doublet $L(1, 2, -1/2)$, and the right-handed charged lepton field $e(1, 1, -1)$. All the quarks and leptons come in three families. In the SMEFT approach, operators with four quarks and two neutrinos are necessarily of dimension nine (dim-9) or higher.

¹We also ignore isospin-breaking effects due to the u - and d -quark mass difference and electroweak radiative corrections.

If lepton number is conserved in the decays of concern,² the responsible operator necessarily involves a pair of L and \bar{L} , which supplies the $\nu\bar{\nu}$ pair in $K \rightarrow \pi\nu\bar{\nu}$. One can always utilize some appropriate Fierz relations to arrange the lepton fields such that they appear in the operators in the form $\bar{L}\gamma_\mu L$ or $\bar{L}\mathcal{D}_\mu\gamma_\rho L$, with \mathcal{D}_μ being a covariant derivative. To join $\bar{L}\gamma_\mu L$ with four quark fields to form an operator that is a singlet under the SM gauge group, the quark portion needs to have a Lorentz index to contract with the one in the lepton current. The possible lowest-dimension quark parts are $(\bar{Q}\gamma^\mu Q, \bar{q}\gamma^\mu q)(\bar{Q}q, \bar{q}Q)$, where q may be u or d as appropriate, but they have odd numbers of Q and hence are not $SU(2)_L$ singlets yet. These quark combinations can be made singlets by incorporating the Higgs field H , and so the full quark-lepton operators have dimension ten (dim-10). Additionally, one can insert \mathcal{D}^μ in $(\bar{Q}q, \bar{q}Q)(\bar{Q}q, \bar{q}Q)$ to form singlets, and the resulting operators are also of dim-10. If the lepton bilinear is $\bar{L}\mathcal{D}_\mu\gamma_\rho L$ instead, the lowest-dimensional possibilities of the four-quark portion are $\bar{q}\gamma^\mu q\bar{q}\gamma^\rho q$, $g^{\mu\rho}\bar{Q}q\bar{q}Q$, $\bar{Q}\sigma^{\mu\rho}q\bar{q}Q$, $\bar{Q}q\bar{q}\sigma^{\mu\rho}Q$, and $\bar{Q}\gamma^\mu Q\bar{Q}\gamma^\rho Q$. Again the singlet quark-lepton operators constructed are of dim-10.

If lepton-number violation is allowed, the situation is different, as we have $K \rightarrow \pi\nu\nu, \pi\nu\bar{\nu}$. The lepton bilinear of the lowest dimension can be organized in the form $\bar{L}L^C$ ($\bar{L}\sigma_{\mu\rho}L^C$) or its Hermitian conjugate, the superscript C signifying charge conjugation. One can attach the bilinear to $\bar{Q}q\bar{q}Q$ or $\bar{q}Q\bar{q}Q$ ($\bar{Q}\sigma^{\mu\rho}q\bar{q}Q$, $\bar{Q}q\bar{q}\sigma^{\mu\rho}Q$, or $\bar{q}\sigma^{\mu\rho}Q\bar{q}Q$) to form a SM-gauge singlet. Thus, the resulting operators have dim-9, which is less than that of the lepton-number-conserving ones mentioned in the previous paragraph. Hereafter in this section, we concentrate on the dim-9 operators and later briefly comment on the dim-10 case.

It has been shown that all dim-9 SMEFT operators do not preserve lepton and/or baryon numbers [37]. We will not be interested in the baryon-number violating ones, as our aim is to study how dim-9 operators give rise to $K \rightarrow \pi\nu\nu, \pi\nu\bar{\nu}$ and can violate the GN bound. In the following we enumerate all of those containing one strange quark, s , and two neutrinos or antineutrinos. Upon imposing the SM gauge symmetries and applying Fierz transformations, we find the independent operators that can induce $K \rightarrow \pi\nu\bar{\nu}$ to be³

$$\begin{aligned}
 \mathcal{O}_1^{opxy,\alpha\beta} &= \epsilon_{ij}\delta_{kl}(\bar{Q}_o^k\gamma_\mu Q_p^j)(\bar{d}_x\gamma^\mu u_y)\bar{L}_{\{\alpha}^{ci}L_{\beta\}}^l, \\
 \tilde{\mathcal{O}}_1^{opxy,\alpha\beta} &= \epsilon_{ij}\delta_{kl}(\bar{Q}_o^k\gamma_\mu Q_p^j)[\bar{d}_x\gamma^\mu u_y]\bar{L}_{\{\alpha}^{ci}L_{\beta\}}^l, \\
 \mathcal{O}_2^{opxy,\alpha\beta} &= \epsilon_{ij}\delta_{kl}(\bar{Q}_o^k\gamma_\mu Q_p^j)(\bar{d}_x\gamma_\rho u_y)\bar{L}_{[\alpha}^{ci}\sigma^{\mu\rho}L_{\beta]}^l, \\
 \tilde{\mathcal{O}}_2^{opxy,\alpha\beta} &= \epsilon_{ij}\delta_{kl}(\bar{Q}_o^k\gamma_\mu Q_p^j)[\bar{d}_x\gamma_\rho u_y]\bar{L}_{[\alpha}^{ci}\sigma^{\mu\rho}L_{\beta]}^l, \\
 \mathcal{O}_3^{opxy,\alpha\beta} &= \epsilon_{ik}\epsilon_{jl}(\bar{d}_o Q_p^i)(\bar{d}_x Q_y^j)\bar{L}_{\{\alpha}^{ck}L_{\beta\}}^l, \\
 \tilde{\mathcal{O}}_3^{opxy,\alpha\beta} &= \epsilon_{ik}\epsilon_{jl}(\bar{d}_o Q_p^i)[\bar{d}_x Q_y^j]\bar{L}_{\{\alpha}^{ck}L_{\beta\}}^l, \\
 \mathcal{O}_4^{opxy,\alpha\beta} &= \epsilon_{ik}\epsilon_{jl}(\bar{d}_o Q_p^i)(\bar{d}_x\sigma_{\mu\omega}Q_y^j)\bar{L}_{[\alpha}^{ck}\sigma^{\mu\omega}L_{\beta]}^l, \\
 \tilde{\mathcal{O}}_4^{opxy,\alpha\beta} &= \epsilon_{ik}\epsilon_{jl}(\bar{d}_o\sigma_{\mu\rho}Q_p^i)(\bar{d}_x\sigma_\omega^\rho Q_y^j)\bar{L}_{[\alpha}^{ck}\sigma^{\mu\omega}L_{\beta]}^l,
 \end{aligned} \tag{2.1}$$

²In this and the next paragraph, we suppress the family labels of the SM fermion fields. Generally these processes may change quark and/or lepton flavors.

³The factorization of the quark and lepton components is guaranteed by the application of Fierz transformations.

where o, p, x, y (α, β) denote quark (lepton) family indices, summation over the $SU(2)_L$ indices $i, j, k, l = 1, 2$ is implicit, and the leptonic scalar (tensor) currents have been arranged to be symmetric (antisymmetric) in their family indices with the convention $A_{\{\alpha\beta\}} = (A_\alpha B_\beta + A_\beta B_\alpha)/2$ and $A_{[\alpha\beta]} = (A_\alpha B_\beta - A_\beta B_\alpha)/2$. The two brackets $(,)$ and $[,]$ in the quark bilinears distinguish the two different ways of color contraction in the products of four quark fields to form color invariants: $\bar{q}_1^m q_2^m \bar{q}_3^n q_4^n = (\bar{q}_1 q_2)(\bar{q}_3 q_4)$ and $\bar{q}_1^m q_2^n \bar{q}_3^n q_4^m = (\bar{q}_1 q_2)[\bar{q}_3 q_4]$, with the color labels $m, n = 1, 2, 3$ being summed over. Each of the operators is accompanied by an unknown Wilson coefficient C_i , so that $C_1^{opxy, \alpha\beta}$ belongs to $\mathcal{O}_{1,2,3,4}^{opxy, \alpha\beta}$, etc. The Hermitian conjugates of $\mathcal{O}_{1,2,3,4}^{opxy, \alpha\beta}$ and $\tilde{\mathcal{O}}_{1,2,3,4}^{opxy, \alpha\beta}$ contribute to $K \rightarrow \pi\nu\nu$.

As already stated, we assume that the SMEFT operators in eq. (2.1) arise from NP above the EW scale. Consequently, to address their potential impact on $K \rightarrow \pi\nu\nu, \pi\bar{\nu}\bar{\nu}$, we will first take into account the QCD effects causing the coefficients to evolve from the NP scale down to the hadronic scale, which we select to be the conventional chiral-symmetry breaking scale $\Lambda_\chi = 4\pi F_\pi \simeq 1.2 \text{ GeV}$ with F_π being the pion decay constant. Subsequently, we will rely on chiral perturbation theory [38–40], in conjunction with spurion techniques [41–43], to derive the meson-neutrino operators which contribute to the amplitudes for the kaon decays. Based on the chiral power-counting arguments in this procedure, the operators in eq. (2.1) with the leptonic scalar density are of momentum order p^0 , whereas those with the leptonic tensor current are of order p^2 . This implies that the latter operators yield contributions to the amplitudes which are suppressed relative to those of the former by the factor $p_K p_\pi / \Lambda_\chi^2 \sim 0.05$. Therefore, upon singling out the operators in eq. (2.1) pertaining to $K \rightarrow \pi\nu\nu, \pi\bar{\nu}\bar{\nu}$ and neglecting those with the leptonic tensor current, in what follows we can focus on

$$\begin{aligned}
 \mathcal{O}_1^{usdu} &= (\bar{u}_L \gamma_\mu s_L)(\bar{d}_R \gamma^\mu u_R)J, & \tilde{\mathcal{O}}_1^{usdu} &= (\bar{u}_L \gamma_\mu s_L)[\bar{d}_R \gamma^\mu u_R]J, \\
 \mathcal{O}_1^{udsu} &= (\bar{u}_L \gamma_\mu d_L)(\bar{s}_R \gamma^\mu u_R)J, & \tilde{\mathcal{O}}_1^{udsu} &= (\bar{u}_L \gamma_\mu d_L)[\bar{s}_R \gamma^\mu u_R]J, \\
 \mathcal{O}_3^{dds} &= (\bar{d}_R d_L)(\bar{d}_R s_L)J, & \tilde{\mathcal{O}}_3^{dds} &= (\bar{d}_R d_L)[\bar{d}_R s_L]J, \\
 \mathcal{O}_3^{d ds} &= (\bar{d}_R d_L)(\bar{s}_R d_L)J, & \tilde{\mathcal{O}}_3^{d ds} &= (\bar{d}_R d_L)[\bar{s}_R d_L]J,
 \end{aligned} \tag{2.2}$$

where $f_{L,R} = (1 \mp \gamma_5)f/2$ and the neutrino part is expressed as $J = (\bar{\nu}_\alpha^c \nu_\beta)/(1 + \delta_{\alpha\beta})$.

2.2 Evaluation of hadronic matrix elements at low energies

In treating the kaon decay amplitudes, the contributions of the operators generated by NP above the EW scale are to be evaluated at the low energy of interest. This entails dealing with the QCD RG running of the Wilson coefficients by resumming the large logarithms due to the ratio of the EW scale to the chiral-symmetry breaking scale, Λ_χ , the running between the NP and EW scales having been neglected. We can use the one-loop QCD running results of ref. [43],

$$\begin{aligned}
 \mu \frac{d}{d\mu} \begin{pmatrix} C_1^{uxyu} \\ \tilde{C}_1^{uxyu} \end{pmatrix} &= -\frac{\alpha_s}{2\pi} \begin{pmatrix} -\frac{3}{N} & 0 \\ 3 & 6C_F \end{pmatrix} \begin{pmatrix} C_1^{uxyu} \\ \tilde{C}_1^{uxyu} \end{pmatrix}, \\
 \mu \frac{d}{d\mu} \begin{pmatrix} C_3^{ddxy} \\ \tilde{C}_3^{ddxy} \end{pmatrix} &= -\frac{\alpha_s}{2\pi} \begin{pmatrix} \frac{2}{N} + 6C_F - 4 & \frac{2}{N} - 4C_F + 2 \\ \frac{4}{N} - 2 & -\frac{2}{N} - 2C_F - 2 \end{pmatrix} \begin{pmatrix} C_3^{ddxy} \\ \tilde{C}_3^{ddxy} \end{pmatrix},
 \end{aligned} \tag{2.3}$$

where in the superscripts $xy = sd$ or ds , the color number $N = 3$, and $C_F = (N^2 - 1)/(2N) = 4/3$ is the second Casimir invariant of the color group $SU(3)_C$. Taking the EW scale to be the W -boson mass m_W , from the solutions to these RG equations [43] we arrive at

$$\begin{aligned}
 C_1^{uxyu}(\Lambda_\chi) &= 0.88 C_1^{uxyu}(m_W), \\
 \tilde{C}_1^{uxyu}(\Lambda_\chi) &= 2.74 \tilde{C}_1^{uxyu}(m_W) + 0.62 C_1^{uxyu}(m_W), \\
 C_3^{ddxy}(\Lambda_\chi) &= 1.82 C_3^{ddxy}(m_W) - 0.34 \tilde{C}_3^{ddxy}(m_W), \\
 \tilde{C}_3^{ddxy}(\Lambda_\chi) &= 0.52 \tilde{C}_3^{ddxy}(m_W) - 0.08 C_3^{ddxy}(m_W).
 \end{aligned} \tag{2.4}$$

In the matching to chiral perturbation theory (χ PT), the neutrino bilinear in a dim-9 operator behaves as a fixed external source. Thus, we only have to work with the quark portion of the operator. Suppose the latter has been decomposed into a sum of irreducible representations of the chiral group $SU(3)_L \times SU(3)_R$ under which the quarks transform as

$$q_{L,a} \rightarrow \hat{L}_{ap} q_{L,p}, \quad \overline{q_{R,c}} \rightarrow \overline{q_{R,p}} \hat{R}_{pc}^\dagger, \quad q_{R,a} \rightarrow \hat{R}_{ap} q_{R,p}, \quad \overline{q_{L,c}} \rightarrow \overline{q_{L,p}} \hat{L}_{pc}^\dagger, \tag{2.5}$$

where the indices $a, c, p = 1, 2, 3$ refer to the flavor space, summation over p is implicit, $\hat{L} \in SU(3)_L$, and $\hat{R} \in SU(3)_R$. Given that an irreducible representation has the general form

$$O_q = T_{cd,ab} (\overline{q_{\chi_1,c}} \Gamma_1 q_{\chi_2,a}) (\overline{q_{\chi_3,d}} \Gamma_2 q_{\chi_4,b}), \tag{2.6}$$

where the $T_{cd,ab}$ represent pure numbers which depend on the irrep under consideration, the flavor indices $a, b, c, d = 1, 2, 3$ are summed over, $\chi_{1,2,3,4} = L, R$, and $\Gamma_{1,2}$ designate combinations of Dirac matrices, then promoting T to be a spurion field that transforms properly together with the chiral transformations of quarks would render O_q chirally invariant.

On the χ PT side, we introduce the standard matrices for the lightest octet of pseudoscalar mesons,

$$\Sigma = \xi^2, \quad \xi = \exp\left(\frac{i\Pi}{\sqrt{2}F_0}\right), \quad \Pi = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\sqrt{\frac{2}{3}}\eta \end{pmatrix}, \tag{2.7}$$

where $F_0 = F_\pi/1.0627 \approx 87$ MeV is the meson decay constant in the chiral limit. Under chiral transformations the Σ and ξ matrices transform as

$$\Sigma \rightarrow \hat{L}\Sigma\hat{R}^\dagger, \quad \xi \rightarrow \hat{L}\xi\hat{U}^\dagger = \hat{U}\xi\hat{R}^\dagger, \tag{2.8}$$

and so $\hat{U} \in SU(3)_V$ depends on the meson fields. From the second formula, in terms of matrix elements we have

$$\xi_{ab} \rightarrow \hat{L}_{ap}(\xi\hat{U}^\dagger)_{pb} = (\hat{U}\xi)_{ap}(\hat{R}^\dagger)_{pb}, \quad (\xi^\dagger)_{ab} \rightarrow \hat{R}_{ap}(\xi^\dagger\hat{U}^\dagger)_{pb} = (\hat{U}\xi^\dagger)_{ap}(\hat{L}^\dagger)_{pb}. \tag{2.9}$$

How to construct the leading-order (LO) mesonic interactions from the quark operators has been prescribed previously in the literature [41–43]. Accordingly, eq. (2.9) implies that the matching to χ PT involves the substitutions [43]

$$\overline{q_{L,a}} \Rightarrow \xi_{\rho a}^\dagger, \quad q_{L,a} \Rightarrow \xi_{a\rho}, \quad \overline{q_{R,a}} \Rightarrow \xi_{\rho a}, \quad q_{R,a} \Rightarrow \xi_{a\rho}^\dagger, \tag{2.10}$$

where the free indices ρ are to be contracted when forming an operator with $T_{cd,ab}$.

For a given quark operator, the first step in the matching is to decompose it according to the irreducible representations (irreps) of the chiral group $SU(3)_L \times SU(3)_R$. One can easily see that \mathcal{O}_1^{usdu} , $\tilde{\mathcal{O}}_1^{usdu}$, \mathcal{O}_1^{udsu} , and $\tilde{\mathcal{O}}_1^{udsu}$ in eq. (2.2) belong to the $8_L \times 8_R$ irrep of the chiral group, while \mathcal{O}_3^{ddds} , $\tilde{\mathcal{O}}_3^{ddds}$, $\mathcal{O}_3^{d dsd}$, and $\tilde{\mathcal{O}}_3^{d dsd}$ belong to the $6_L \times \bar{6}_R$. Then, after applying eq. (2.10), one associates with the mesonic counterpart of each irrep a low energy constant g_{irrep} , which encodes nonperturbative QCD effects and is to be determined usually by fitting to data or a model calculation.

For example, the LEC associated with $\mathcal{O}_1^{usdu} = (\bar{u}_L \gamma_\mu s_L)(\bar{d}_R \gamma^\mu u_R)J$, which transforms as the $8_L \times 8_R$ under the chiral group, can be called $g_{8 \times 8}$, and the leading-order chiral realization of \mathcal{O}_1^{usdu} is derived via the procedure

$$\begin{aligned} \mathcal{O}_1^{usdu} &\Rightarrow \frac{F_0^4}{4} g_{8 \times 8} \xi_{\vartheta 1}^\dagger \xi_{3 \varrho} \xi_{\varsigma 2} \xi_{1 \varphi}^\dagger \delta_{\vartheta \varphi} \delta_{\varrho \varsigma} J = \frac{F_0^4}{4} g_{8 \times 8} (\xi \xi)_{32} (\xi^\dagger \xi^\dagger)_{11} J = \frac{F_0^4}{4} g_{8 \times 8} \Sigma_{32} \Sigma_{11}^\dagger J \\ &\Rightarrow \frac{g_{8 \times 8}}{4} F_0^2 \left(\frac{3}{\sqrt{2}} \pi^0 \bar{K}^0 - \pi^+ K^- \right) J + \frac{i g_{8 \times 8}}{4 \sqrt{2}} F_0^3 \bar{K}^0 J + \dots, \end{aligned} \quad (2.11)$$

where $F_0^4/4$ is a normalization factor [42], other contractions among $\varepsilon, \varrho, \varsigma, \varphi$ in the first line vanish due to the unitarity of ξ , and the ellipsis stands for terms with the η field and more than two meson fields. This result is independent of the Lorentz and color structures of \mathcal{O}_1^{usdu} and follows from its transformation properties as an irrep of the chiral group. Evidently, $g_{8 \times 8}$ has mass dimension two. This example can be understood from the perspective of bosonization of each quark bilinear: Fierz-transforming \mathcal{Q}_1^{usdu} yields $-2\bar{u}_L^m u_R^n \bar{s}_L^m d_R^n$ which is summed over the color labels m, n and has a lower-order chiral realization than $(\bar{u}_L \gamma_\mu s_L)(\bar{d}_R \gamma^\mu u_R)$ being naively taken to correspond to the higher order $(\partial^\mu \Sigma \Sigma^\dagger)_{31} (\Sigma^\dagger \partial_\mu \Sigma)_{12}$ due to the derivatives. If the operator comprised instead solely left-handed quarks, $(\bar{u}_L \gamma_\mu s_L)(\bar{d}_L \gamma^\mu u_L)$, its chiral realization would have to be of the form $(\partial^\mu \Sigma \Sigma^\dagger)_{op} (\partial_\mu \Sigma \Sigma^\dagger)_{xy}$ because no scalar density can be constructed with such quarks.

In table 1 we provide the chiral realization of the quark component of each of the operators in eq. (2.2) according to their chiral irreps. In the last column, we display the leading-order contributions to the $K \rightarrow \pi$ transitions in terms of the mesonic operators

$$\mathcal{Q}_{1/2}^S = F_0^2 \left(K^+ \pi^- - \frac{1}{\sqrt{2}} K^0 \pi^0 \right), \quad \mathcal{Q}_{3/2}^S = F_0^2 \left(K^+ \pi^- + \sqrt{2} K^0 \pi^0 \right), \quad (2.12)$$

or their Hermitian conjugates, which correspond to definite isospin changes $\Delta I = 1/2$ and $3/2$, respectively.⁴ It is worth noting that in the combination $5\mathcal{Q}_{1/2}^S - 2\mathcal{Q}_{3/2}^S$, which occurs in all of the lines in table 1 and hence also implicitly in eq. (2.11), the size of the $K^0 \pi^0$ term relative to the $K^+ \pi^-$ one is three times that in $\mathcal{Q}_{1/2}^S$ alone. It follows that every one of these operators can potentially break the GN relation.

The table also shows that each operator with a tilde and its counterpart without it have the same chiral realization but their LECs are different. This is attributable to the fact that the chiral realization of a quark-level operator relies only on its representation

⁴For completeness, in appendix A we give the decomposition of the quark part of each of the operators in eq. (2.2) in terms of their $\Delta I = 1/2, 3/2$ components.

Operator	Chiral irrep	Chiral realization	Contributions to $K \rightarrow \pi$
\mathcal{O}_1^{usdu}	$8_L \times 8_R$	$\frac{1}{4}F_0^2 g_{8 \times 8} \Sigma_{32} \Sigma_{11}^\dagger$	$-\frac{1}{12}g_{8 \times 8} \left(5\mathcal{Q}_{1/2}^{S^\dagger} - 2\mathcal{Q}_{3/2}^{S^\dagger}\right)$
$\tilde{\mathcal{O}}_1^{usdu}$	$8_L \times 8_R$	$\frac{1}{4}F_0^2 \tilde{g}_{8 \times 8} \Sigma_{32} \Sigma_{11}^\dagger$	$-\frac{1}{12}\tilde{g}_{8 \times 8} \left(5\mathcal{Q}_{1/2}^{S^\dagger} - 2\mathcal{Q}_{3/2}^{S^\dagger}\right)$
\mathcal{O}_1^{udsu}	$8_L \times 8_R$	$\frac{1}{4}F_0^2 g_{8 \times 8} \Sigma_{23} \Sigma_{11}^\dagger$	$-\frac{1}{12}g_{8 \times 8} \left(5\mathcal{Q}_{1/2}^S - 2\mathcal{Q}_{3/2}^S\right)$
$\tilde{\mathcal{O}}_1^{udsu}$	$8_L \times 8_R$	$\frac{1}{4}F_0^2 \tilde{g}_{8 \times 8} \Sigma_{23} \Sigma_{11}^\dagger$	$-\frac{1}{12}\tilde{g}_{8 \times 8} \left(5\mathcal{Q}_{1/2}^S - 2\mathcal{Q}_{3/2}^S\right)$
\mathcal{O}_3^{dds}	$6_L \times \bar{6}_R$	$\frac{1}{4}F_0^2 g_{6 \times \bar{6}} \Sigma_{22} \Sigma_{32}$	$-\frac{1}{12}g_{6 \times \bar{6}} \left(5\mathcal{Q}_{1/2}^{S^\dagger} - 2\mathcal{Q}_{3/2}^{S^\dagger}\right)$
$\tilde{\mathcal{O}}_3^{dds}$	$6_L \times \bar{6}_R$	$\frac{1}{4}F_0^2 \tilde{g}_{6 \times \bar{6}} \Sigma_{22} \Sigma_{32}$	$-\frac{1}{12}\tilde{g}_{6 \times \bar{6}} \left(5\mathcal{Q}_{1/2}^{S^\dagger} - 2\mathcal{Q}_{3/2}^{S^\dagger}\right)$
\mathcal{O}_3^{ddsd}	$6_L \times \bar{6}_R$	$\frac{1}{4}F_0^2 g_{6 \times \bar{6}} \Sigma_{23} \Sigma_{22}$	$-\frac{1}{12}g_{6 \times \bar{6}} \left(5\mathcal{Q}_{1/2}^S - 2\mathcal{Q}_{3/2}^S\right)$
$\tilde{\mathcal{O}}_3^{ddsd}$	$6_L \times \bar{6}_R$	$\frac{1}{4}F_0^2 \tilde{g}_{6 \times \bar{6}} \Sigma_{23} \Sigma_{22}$	$-\frac{1}{12}\tilde{g}_{6 \times \bar{6}} \left(5\mathcal{Q}_{1/2}^S - 2\mathcal{Q}_{3/2}^S\right)$

Table 1. The chiral representation and realization of the four-quark portion of each of the effective dim-9 operators in eq. (2.2) relevant to the $K \rightarrow \pi$ processes. In the last column, $\mathcal{Q}_{1/2}^S$ and $\mathcal{Q}_{3/2}^S$ are the mesonic operators defined in eq. (2.12) and correspond to $\Delta I = 1/2$ and $3/2$ transitions, respectively.

under the chiral group, whereas the LEC encodes QCD effects. Numerically, we adopt the values of the LECs extracted from ref. [42] which employed χ PT to connect the matrix elements of $\pi^+ \rightarrow \pi^-$ transitions to kaon-mixing matrix elements for which lattice QCD results were available. Thus, we have

$$\begin{aligned}
 g_{8 \times 8} &= -2.9 \text{ GeV}^2, & \tilde{g}_{8 \times 8} &= -12.4 \text{ GeV}^2, \\
 g_{6 \times \bar{6}} &= 2.7 \text{ GeV}^2, & \tilde{g}_{6 \times \bar{6}} &= -0.91 \text{ GeV}^2.
 \end{aligned} \tag{2.13}$$

2.3 Numerical analysis

From the results of the preceding subsection, we can write down the effective interactions pertinent to $K^+ \rightarrow \pi^+ \nu \nu, \pi^+ \bar{\nu} \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \nu, \pi^0 \bar{\nu} \bar{\nu}$, namely

$$\begin{aligned}
 C_1^{usdu} \mathcal{O}_1^{usdu} + \text{H.c.} &\Rightarrow \frac{g_{8 \times 8}}{4} F_0^2 \left[\frac{3}{2} \left(C_1^{usdu} J + C_1^{usdu*} J^\dagger \right) \pi^0 K_L - C_1^{usdu*} J^\dagger \pi^- K^+ \right], \\
 C_1^{udsu} \mathcal{O}_1^{udsu} + \text{H.c.} &\Rightarrow \frac{g_{8 \times 8}}{4} F_0^2 \left[\frac{3}{2} \left(C_1^{udsu} J + C_1^{udsu*} J^\dagger \right) \pi^0 K_L - C_1^{udsu} J \pi^- K^+ \right], \\
 C_3^{dds} \mathcal{O}_3^{dds} + \text{H.c.} &\Rightarrow \frac{g_{6 \times \bar{6}}}{4} F_0^2 \left[\frac{3}{2} \left(C_3^{dds} J + C_3^{dds*} J^\dagger \right) \pi^0 K_L - C_3^{dds*} J^\dagger \pi^- K^+ \right], \\
 C_3^{ddsd} \mathcal{O}_3^{ddsd} + \text{H.c.} &\Rightarrow \frac{g_{6 \times \bar{6}}}{4} F_0^2 \left[\frac{3}{2} \left(C_3^{ddsd} J + C_3^{ddsd*} J^\dagger \right) \pi^0 K_L - C_3^{ddsd} J \pi^- K^+ \right],
 \end{aligned} \tag{2.14}$$

and analogous expressions for the tilded operators, where $J^\dagger = (\bar{\nu}_\alpha \nu_\beta^c)/(1 + \delta_{\alpha\beta})$. With these, we arrive at the decay amplitudes

$$\begin{aligned} \mathcal{A}_{K_L \rightarrow \pi^0 \nu_\alpha \nu_\beta} &= \frac{3}{8} F_0^2 (C_A^* + C_B^*) \bar{\nu}_\alpha \nu_\beta^c, & \mathcal{A}_{K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta} &= -\frac{1}{4} F_0^2 C_B^* \bar{\nu}_\alpha \nu_\beta^c, \\ \mathcal{A}_{K_L \rightarrow \pi^0 \bar{\nu}_\alpha \bar{\nu}_\beta} &= \frac{3}{8} F_0^2 (C_A + C_B) \bar{\nu}_\alpha^c \nu_\beta, & \mathcal{A}_{K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta} &= -\frac{1}{4} F_0^2 C_A \bar{\nu}_\alpha^c \nu_\beta, \end{aligned} \quad (2.15)$$

involving the effective coupling constants

$$\begin{aligned} C_A &= g_{8 \times 8} C_1^{udsu}(\Lambda_\chi) + \tilde{g}_{8 \times 8} \tilde{C}_1^{udsu}(\Lambda_\chi) + g_{6 \times 6} C_3^{ddsd}(\Lambda_\chi) + \tilde{g}_{6 \times 6} \tilde{C}_3^{ddsd}(\Lambda_\chi), \\ C_B &= g_{8 \times 8} C_1^{usdu}(\Lambda_\chi) + \tilde{g}_{8 \times 8} \tilde{C}_1^{usdu}(\Lambda_\chi) + g_{6 \times 6} C_3^{ddds}(\Lambda_\chi) + \tilde{g}_{6 \times 6} \tilde{C}_3^{ddds}(\Lambda_\chi), \end{aligned} \quad (2.16)$$

which implicitly carry the neutrino family labels α and β . From eq. (2.15), we obtain the spin-summed absolute squares

$$\begin{aligned} \sum_{\text{spins}} |\mathcal{A}_{K_L \rightarrow \pi^0 \nu_\alpha \nu_\beta}|^2 &= \frac{9}{32} F_0^4 |C_A + C_B|^2 \hat{s}, & \sum_{\text{spins}} |\mathcal{A}_{K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta}|^2 &= \frac{1}{8} F_0^4 |C_B|^2 \hat{s}, \\ \sum_{\text{spins}} |\mathcal{A}_{K_L \rightarrow \pi^0 \bar{\nu}_\alpha \bar{\nu}_\beta}|^2 &= \frac{9}{32} F_0^4 |C_A + C_B|^2 \hat{s}, & \sum_{\text{spins}} |\mathcal{A}_{K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta}|^2 &= \frac{1}{8} F_0^4 |C_A|^2 \hat{s}, \end{aligned} \quad (2.17)$$

where $\hat{s} = (p_1 + p_2)^2$. The $\nu_\alpha \nu_\beta$ and $\bar{\nu}_\alpha \bar{\nu}_\beta$ channels having no interference with each other, their branching fractions add up to

$$\begin{aligned} \mathcal{B}(K_L \rightarrow \pi^0 \nu_\alpha \nu_\beta) + \mathcal{B}(K_L \rightarrow \pi^0 \bar{\nu}_\alpha \bar{\nu}_\beta) &= \frac{9 F_0^4 |C_A + C_B|^2}{16 (1 + \delta_{\alpha\beta})} \frac{\tau_{K_L}}{2 m_{K^0}} \int d\Pi_3 \hat{s} \\ &= 1.42 \times 10^{-9} \frac{|\hat{C}_A + \hat{C}_B|^2}{1 + \delta_{\alpha\beta}}, \\ \mathcal{B}(K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta) + \mathcal{B}(K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta) &= \frac{F_0^4 (|C_A|^2 + |C_B|^2)}{8 (1 + \delta_{\alpha\beta})} \frac{\tau_{K^+}}{2 m_{K^+}} \int d\Pi_3 \hat{s} \\ &= 6.98 \times 10^{-11} \frac{|\hat{C}_A|^2 + |\hat{C}_B|^2}{1 + \delta_{\alpha\beta}}, \end{aligned} \quad (2.18)$$

where τ_{K_L} and τ_{K^+} stand for the measured K_L and K^+ lifetimes [13], the factor $1/(1 + \delta_{\alpha\beta})$ in each equation accounts for the identical particles in the final state if $\alpha = \beta$, and $d\Pi_3$ denotes the three-body phase-space factor. Also, in eq. (2.18) we have defined the dimensionless parameters

$$\begin{aligned} \hat{C}_A &= (50 \text{ GeV})^5 \left[C_1^{udsu}(m_W) + 3.3 \tilde{C}_1^{udsu}(m_W) - 0.55 C_3^{ddsd}(m_W) + 0.15 \tilde{C}_3^{ddsd}(m_W) \right], \\ \hat{C}_B &= (50 \text{ GeV})^5 \left[C_1^{usdu}(m_W) + 3.3 \tilde{C}_1^{usdu}(m_W) - 0.55 C_3^{ddds}(m_W) + 0.15 \tilde{C}_3^{ddds}(m_W) \right], \end{aligned} \quad (2.19)$$

which incorporate eq. (2.4) for the RG-running effects on the parameters between $\mu = \Lambda_\chi$ and $\mu = m_W$ and eq. (2.13) for the LEC values.

Assuming that the operators involve only one pair of α and $\beta \neq \alpha$, since these modes do not interfere with the SM ones, we can combine eq. (2.18) with their SM counterparts to find

$$\begin{aligned} \mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}}) &= \mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{SM}} + \mathcal{B}(K_L \rightarrow \pi^0 \nu_\alpha \nu_\beta) + \mathcal{B}(K_L \rightarrow \pi^0 \bar{\nu}_\alpha \bar{\nu}_\beta), \\ \mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}}) &= \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} + \mathcal{B}(K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta) + \mathcal{B}(K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta). \end{aligned} \quad (2.20)$$

Accordingly, the definition of $r_{\mathcal{B}}$ should be modified, and we now have

$$r_{\mathcal{B}} = \frac{\mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}})}{\mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}})} = r_{\mathcal{B}}^{\text{SM}} + \frac{(r_{\mathcal{B}}^{\text{NP}} - r_{\mathcal{B}}^{\text{SM}})\epsilon}{1 + \epsilon} \leq 0.36 + \frac{40.3\epsilon}{1 + \epsilon}, \quad (2.21)$$

where

$$\begin{aligned} r_{\mathcal{B}}^{\text{NP}} &= \frac{\mathcal{B}(K_L \rightarrow \pi^0 \nu_\alpha \nu_\beta) + \mathcal{B}(K_L \rightarrow \pi^0 \bar{\nu}_\alpha \bar{\nu}_\beta)}{\mathcal{B}(K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta) + \mathcal{B}(K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta)} = \frac{20.3 |\hat{C}_A + \hat{C}_B|^2}{|\hat{C}_A|^2 + |\hat{C}_B|^2} \leq 40.6, \\ \epsilon &= \frac{\mathcal{B}(K^+ \rightarrow \pi^+ \nu_\alpha \nu_\beta) + \mathcal{B}(K^+ \rightarrow \pi^+ \bar{\nu}_\alpha \bar{\nu}_\beta)}{\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}}}. \end{aligned} \quad (2.22)$$

The most recent measurement $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{NA62}} < 1.85 \times 10^{-10}$ at 90% CL [19] entails that $\epsilon \lesssim 1$ and consequently $r_{\mathcal{B}}$ is capped to be about 20.5. This can accommodate KOTO's anomalous events [20].

Imposing on eq. (2.20) the KOTO 15 [15] and NA62 [19] limits and further assuming that the only nonvanishing coefficients are $C_1^{udsu}(m_W) = C_1^{rusdu}(m_W) = \Lambda_{\text{NP}}^{-5}$, we have

$$\begin{aligned} \mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}}) &= 3.0 \times 10^{-11} + 5.7 \times 10^{-9} \left(\frac{50 \text{ GeV}}{\Lambda_{\text{NP}}} \right)^{10} \leq 3.0 \times 10^{-9}, \\ \mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}}) &= 8.5 \times 10^{-11} + 1.4 \times 10^{-10} \left(\frac{50 \text{ GeV}}{\Lambda_{\text{NP}}} \right)^{10} \leq 1.85 \times 10^{-10}, \end{aligned} \quad (2.23)$$

where the first number in each line is the corresponding SM central value [11–13]. The stronger of the empirical limits, from KOTO in the first line, translates into $\Lambda_{\text{NP}} \gtrsim 53 \text{ GeV}$. In figure 1 we illustrate the dependence of $\mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}})$ and $\mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}})$ in eq. (2.23) on Λ_{NP} . Also plotted are the corresponding SM predictions and limits from KOTO [15] and NA62 [19]. We see that the NP needs to have an effective scale $\Lambda_{\text{NP}} = \mathcal{O}(60 \text{ GeV})$ if it is to be responsible for the KOTO anomaly. The preferred Λ_{NP} is significantly below the EW scale $\sim v \simeq 246 \text{ GeV}$, which implies that for $\Lambda_{\text{NP}} \sim v$ the dim-9 operators would have negligible impact on $K \rightarrow \pi + E_{\text{miss}}$ and thus respect the GN bound.

If we repeat the above steps with the dim-10 operators discussed earlier, a scaling factor of v/Λ_{NP} will accompany them. As mentioned in subsection 2.1, they conserve lepton number and hence, if not flavor-violating, interfere with the SM contribution. For $\Lambda_{\text{NP}} < v$, this scaling factor helps raise Λ_{NP} slightly, but the latter still will not be very close to the EW scale, $\sim v$, if the NP presence in $K \rightarrow \pi + E_{\text{miss}}$ is to be within the current experimental sensitivity reaches. Hence for $\Lambda_{\text{NP}} \gtrsim v$ the dim-10 operators would also have very little influence on these modes.




Figure 1. The branching fractions of $K \rightarrow \pi\nu\nu, \pi\bar{\nu}\bar{\nu}$ arising from the dim-9 operators versus the NP scale Λ_{NP} for the example described in the text. Also displayed are the corresponding SM predictions for $K \rightarrow \pi\nu\bar{\nu}$ (red and blue horizontal bands) and upper limits from KOTO [15] and NA62 [19] (blue and red horizontal thin lines). The light-blue region is excluded by the KOTO bound. The blue dot corresponds to KOTO’s three events.

3 GN-bound violation via EFT operators for $K \rightarrow \pi S$ or πSS

In the preceding section, the problem of the NP scale Λ_{NP} being too low can be ascribed to the high dimension of the SMEFT operators. As sketched in section 1, any NP that can induce $K_L \rightarrow \pi^0 \mathcal{X}$ with a rate exceeding the SM expectation without conflicting with the K^+ data must have an effective scale Λ_{NP} which goes roughly as $(23000 \text{ TeV}^2 m_K^N)^{1/(2+N)}$, where m_K is the kaon mass and N the mass dimension of the field content of \mathcal{X} . Therefore, one way to increase Λ_{NP} is by reducing the dimension of the operators, which is $6+N$. If in the dim-9 operators examined above the neutrino pair, which has $N = 3$, is replaced with a scalar field S , which has $N = 1$, the dimension of the operators can be decreased by 2 and in turn Λ_{NP} can be raised to the TeV level. If $\mathcal{X} = SS$ instead, the scale will become $\Lambda_{\text{NP}} = \mathcal{O}(v)$. In this section, we explore how SMEFT four-quark operators supplemented with S , which we take to be real and a singlet under the SM gauge group, can give rise to $K \rightarrow \pi S$ transitions which break the GN bound. Moreover, we apply a similar treatment to the $K \rightarrow \pi SS$ case.

3.1 Operators and matching

As is clear from the last paragraph, SMEFT operators that directly give rise to $K \rightarrow \pi S$ at leading order have to be of dimension seven (dim-7). Since S is a SM-gauge singlet, the quark portions of these operators are none other than the SMEFT dimension-six (dim-6) four-quark operators [44, 45] which contribute to nonleptonic kaon decays. In the first column of table 2, we list these dim-6 operators in the Warsaw basis [45]. In the middle column, we exhibit the relevant operators with one s -quark field in the LEFT approach [46]. The third column contains the results for the Wilson coefficients at the electroweak scale

SMEFT operators	LEFT operators	Matching at the EW scale
$Q_{qq}^{(1)} = (\bar{Q}\gamma_\mu Q)(\bar{Q}\gamma_\mu Q)$ $Q_{qq}^{(3)} = (\bar{Q}\tau^I\gamma_\mu Q)(\bar{Q}\tau^I\gamma_\mu Q)$	$O_{ddsd}^{V,LL} = (\bar{d}_L\gamma_\mu d_L)(\bar{s}_L\gamma^\mu d_L)$ $O_{uusd}^{V1,LL} = (\bar{u}_L\gamma_\mu u_L)(\bar{s}_L\gamma^\mu d_L)$ $O_{uusd}^{V8,LL} = (\bar{u}_L T^A\gamma_\mu u_L)(\bar{s}_L T^A\gamma^\mu d_L)$	$C_{ddsd}^{V,LL} = C_{qq}^{(1),1121} + C_{qq}^{(3),1121}$ $C_{uusd}^{V1,LL} = C_{qq}^{(1),1121} - \frac{1}{3}C_{qq}^{(3),1121}$ $C_{uusd}^{V8,LL} = 4C_{qq}^{(3),1121}$
$Q_{dd} = (\bar{d}\gamma_\mu d)(\bar{d}\gamma^\mu d)$ $Q_{ud}^{(1)} = (\bar{u}\gamma_\mu u)(\bar{d}\gamma^\mu d)$ $Q_{ud}^{(8)} = (\bar{u}T^A\gamma_\mu u)(\bar{d}T^A\gamma^\mu d)$	$O_{ddsd}^{V,RR} = (\bar{d}_R\gamma_\mu d_R)(\bar{s}_R\gamma^\mu d_R)$ $O_{uusd}^{V1,RR} = (\bar{u}_R\gamma_\mu u_R)(\bar{s}_R\gamma^\mu d_R)$ $O_{uusd}^{V8,RR} = (\bar{u}_R T^A\gamma_\mu u_R)(\bar{s}_R T^A\gamma^\mu d_R)$	$C_{ddsd}^{V,RR} = C_{dd}^{1121}$ $C_{uusd}^{V1,RR} = C_{ud}^{(1),1121}$ $C_{uusd}^{V8,RR} = C_{ud}^{(8),1121}$
$Q_{qu}^{(1)} = (\bar{Q}\gamma_\mu Q)(\bar{u}\gamma^\mu u)$ $Q_{qu}^{(8)} = (\bar{Q}T^A\gamma_\mu Q)(\bar{u}T^A\gamma^\mu u)$ $Q_{qd}^{(1)} = (\bar{Q}\gamma_\mu Q)(\bar{d}\gamma^\mu d)$ $Q_{qd}^{(8)} = (\bar{Q}T^A\gamma_\mu Q)(\bar{d}T^A\gamma^\mu d)$	$O_{sduu}^{V1,LR} = (\bar{s}_L\gamma_\mu d_L)(\bar{u}_R\gamma^\mu u_R)$ $O_{sduu}^{V8,LR} = (\bar{s}_L T^A\gamma_\mu d_L)(\bar{u}_R T^A\gamma^\mu u_R)$ $O_{sddd}^{V1,LR} = (\bar{s}_L\gamma_\mu d_L)(\bar{d}_R\gamma^\mu d_R)$ $O_{sddd}^{V8,LR} = (\bar{s}_L T^A\gamma_\mu d_L)(\bar{d}_R T^A\gamma^\mu d_R)$	$C_{sduu}^{V1,LR} = C_{qu}^{(1),2111}$ $C_{sduu}^{V8,LR} = C_{qu}^{(8),2111}$ $C_{sddd}^{V1,LR} = C_{qd}^{(1),2111}$ $C_{sddd}^{V8,LR} = C_{qd}^{(8),2111}$
$Q_{qd}^{(1)} = (\bar{Q}\gamma_\mu Q)(\bar{d}\gamma^\mu d)$ $Q_{qd}^{(8)} = (\bar{Q}T^A\gamma_\mu Q)(\bar{d}T^A\gamma^\mu d)$	$O_{uusd}^{V1,LR} = (\bar{u}_L\gamma_\mu u_L)(\bar{s}_R\gamma^\mu d_R)$ $O_{uusd}^{V8,LR} = (\bar{u}_L T^A\gamma_\mu u_L)(\bar{s}_R T^A\gamma^\mu d_R)$ $O_{ddsd}^{V1,LR} = (\bar{d}_L\gamma_\mu d_L)(\bar{s}_R\gamma^\mu d_R)$ $O_{ddsd}^{V8,LR} = (\bar{d}_L T^A\gamma_\mu d_L)(\bar{s}_R T^A\gamma^\mu d_R)$	$C_{uusd}^{V1,LR} = C_{qd}^{(1),1121}$ $C_{uusd}^{V8,LR} = C_{qd}^{(8),1121}$ $C_{ddsd}^{V1,LR} = C_{qd}^{(1),1121}$ $C_{ddsd}^{V8,LR} = C_{qd}^{(8),1121}$
$Q_{quqd}^{(1)} = \epsilon_{ij}(\bar{Q}^i u)(\bar{Q}^j d)$ $Q_{quqd}^{(8)} = \epsilon_{ij}(\bar{Q}^i T^A u)(\bar{Q}^j T^A d)$	$O_{uusd}^{S1,RR} = (\bar{u}_L u_R)(\bar{s}_L d_R)$ $O_{uusd}^{S8,RR} = (\bar{u}_L T^A u_R)(\bar{s}_L T^A d_R)$ $O_{udsu}^{S1,RR} = (\bar{u}_L d_R)(\bar{s}_L u_R)$ $O_{udsu}^{S8,RR} = (\bar{u}_L T^A d_R)(\bar{s}_L T^A u_R)$	$C_{uusd}^{S1,RR} = C_{quqd}^{(1),1121}$ $C_{uusd}^{S8,RR} = C_{quqd}^{(8),1121}$ $C_{uusd}^{S1,RR} = -C_{quqd}^{(1),2111}$ $C_{uusd}^{S8,RR} = -C_{quqd}^{(8),2111}$
$Q_{quqd}^{(1)\dagger} = \epsilon_{ij}(\bar{u}Q^i)(\bar{d}Q^j)$ $Q_{quqd}^{(8)\dagger} = \epsilon_{ij}(\bar{u}T^A Q^i)(\bar{d}T^A Q^j)$	$O_{uusd}^{S1,LL} = (\bar{u}_R u_L)(\bar{s}_R d_L)$ $O_{uusd}^{S8,LL} = (\bar{u}_R T^A u_L)(\bar{s}_R T^A d_L)$ $O_{udsu}^{S1,LL} = (\bar{u}_R d_L)(\bar{s}_R u_L)$ $O_{udsu}^{S8,LL} = (\bar{u}_R T^A d_L)(\bar{s}_R T^A u_L)$	$C_{uusd}^{S1,LL} = C_{quqd}^{(1),1112*}$ $C_{uusd}^{S8,LL} = C_{quqd}^{(8),1112*}$ $C_{udsu}^{S1,LL} = -C_{quqd}^{(1),1112*}$ $C_{udsu}^{S8,LL} = -C_{quqd}^{(8),1112*}$

Table 2. Columns 1 and 2: the SMEFT and LEFT four-quark operators contributing to $K \rightarrow \pi S(S)$. Column 3: the results for the Wilson coefficients at the electroweak scale from the matching of the former operators onto the latter.

from the matching of the SMEFT operators onto the LEFT operators [46]. For the $K \rightarrow \pi S$ and $K \rightarrow \pi SS$ transitions, we just multiply all those operators by an S field and a pair of them, respectively, and the matched Wilson coefficients should be understood as new parameters associated with the operators.

Similarly to what was done in section 2, to examine the impact of each of the dim-6 LEFT operators in table 2, we begin by decomposing their four-quark combinations in

terms of the irreducible representations of the chiral group, $SU(3)_L \times SU(3)_R$. Subsequently, for each of the irreps we derive the chiral realization, as prescribed in subsection 2.2, and complement it with a low-energy constant. Finally, the resulting meson operators are expressed as combinations of their isospin components. The first column of table 3 lists the LEFT operators in table 2 according to their irreps. In the second column we collect the mesonic operators pertinent to the $K \rightarrow \pi$ processes. The ones corresponding to the four-quark operators with purely left-handed or right-handed quarks are written in terms of the $\Delta I = 1/2$ and $3/2$ combinations

$$\begin{aligned} \mathcal{Q}_{1/2}^V &= F_0^2 \left(\partial_\mu K^+ \partial^\mu \pi^- - \frac{1}{\sqrt{2}} \partial_\mu K^0 \partial^\mu \pi^0 \right), \\ \mathcal{Q}_{3/2}^V &= F_0^2 \left(\partial_\mu K^+ \partial^\mu \pi^- + \sqrt{2} \partial_\mu K^0 \partial^\mu \pi^0 \right), \end{aligned} \quad (3.1)$$

respectively. The other entries in the second column involve $\mathcal{Q}_{1/2}^S$ and $\mathcal{Q}_{3/2}^S$ which were already defined in eq. (2.12). More details on the bosonization of the irreps are relegated to appendix C. In this table, we also see that there are more LECs than in table 1. For $g_{1 \times 8}$ and $g_{1 \times 27}$, which are dimensionless, we adopt

$$g_{8 \times 1} = 3.65, \quad g_{27 \times 1} = 0.303 \quad (3.2)$$

from ref. [47], whereas $g_{8 \times 8}$, $\tilde{g}_{8 \times 8}$, $g_{6 \times \bar{6}}$, and $\tilde{g}_{6 \times \bar{6}}$ are already given in eq. (2.13). Moreover, the parity invariance of the QCD suggests that we can set

$$\begin{aligned} g_{1 \times 8} &= g_{8 \times 1}, & g_{1 \times 27} &= g_{27 \times 1}, \\ g_{\bar{6} \times 6} &= g_{6 \times \bar{6}}, & g_{\bar{3} \times 3} &= g_{3 \times \bar{3}} \end{aligned} \quad (3.3)$$

and assume analogous relations for the corresponding LECs with a tilde. For the value of $g_{3 \times \bar{3}}$, there is no estimation yet in literature, and so one can resort to the vacuum saturation approximation (VSA) which yields $g_{3 \times \bar{3}} = g_{6 \times \bar{6}} = B_0^2 \simeq 4 \text{ GeV}^2$ with $B_0 = m_\pi^2 / (m_u + m_d) = m_K^2 / (m_u + m_s)$ and quark masses at a renormalization scale of 1 GeV. Evidently the VSA number for $g_{3 \times \bar{3}}$ is not too far from $g_{6 \times \bar{6}} = 3.2 \text{ GeV}^2$ in eq. (2.13). Additionally, one can implement simple scaling to estimate $\tilde{g}_{3 \times \bar{3}} = g_{3 \times \bar{3}} \tilde{g}_{6 \times \bar{6}} / g_{6 \times \bar{6}} \simeq 1.4 \text{ GeV}^2$.

It is worth pointing out that, unlike those in table 1, the operators in table 3 individually either respect the GN inequality or are dominated by portions which do, partly due to the LEC values employed above. The first two sections of the table contain operators in the latter category as they have $\mathcal{Q}_{1/2}^V$ parts with $g_{8 \times 1} \simeq 12g_{27 \times 1}$. The operators in the remaining sections of the table belong to the first category, such as $\mathcal{O}_{sduu}^{V1,LR}$, which by itself does not affect $K_L \rightarrow \pi^0$ transitions due to its chiral realization being proportional to $2\mathcal{Q}_{1/2}^S + \mathcal{Q}_{3/2}^S = 3F_0^2 K^+ \pi^-$, and $\mathcal{O}_{sddd}^{V1,LR}$, which generates $\mathcal{Q}_{1/2}^S + 2\mathcal{Q}_{3/2}^S \propto K^+ \pi^- + K^0 \pi^0 / \sqrt{2}$ whose two terms have the same relative size as their counterparts in $\mathcal{Q}_{1/2}^S$, albeit with the opposite relative sign. Nevertheless, there are countless combinations of the various oper-

Chiral irrep	Contributions to $K \rightarrow \pi$
$O_{dds}^{V,LL} = O_{dds}^{V,LL} _{27 \times 1} + O_{dds}^{V,LL} _{8 \times 1}$ $O_{uus}^{V1,LL} = O_{uus}^{V1,LL} _{27 \times 1} + O_{uus}^{V1,LL} _{8 \times 1}$ $O_{uus}^{V8,LL} = O_{uus}^{V8,LL} _{27 \times 1} + O_{uus}^{V8,LL} _{8 \times 1}$	$\frac{1}{18}g_{27 \times 1} \left(2Q_{1/2}^V - 5Q_{3/2}^V \right) + \frac{1}{6}g_{8 \times 1} Q_{1/2}^V$ $\frac{1}{18}g_{27 \times 1} \left(Q_{1/2}^V + 5Q_{3/2}^V \right) - \frac{1}{3}g_{8 \times 1} Q_{1/2}^V$ $\frac{1}{54}g_{27 \times 1} \left(Q_{1/2}^V + 5Q_{3/2}^V \right) + \frac{11}{36}g_{8 \times 1} Q_{1/2}^V$
$O_{dds}^{V,RR} = O_{dds}^{V,RR} _{1 \times 27} + O_{dds}^{V,RR} _{1 \times 8}$ $O_{uus}^{V1,RR} = O_{uus}^{V1,RR} _{1 \times 27} + O_{uus}^{V1,RR} _{1 \times 8}$ $O_{uus}^{V8,RR} = O_{uus}^{V8,RR} _{1 \times 27} + O_{uus}^{V8,RR} _{1 \times 8}$	$\frac{1}{18}g_{1 \times 27} \left(2Q_{1/2}^V - 5Q_{3/2}^V \right) + \frac{1}{6}g_{1 \times 8} Q_{1/2}^V$ $\frac{1}{18}g_{1 \times 27} \left(Q_{1/2}^V + 5Q_{3/2}^V \right) - \frac{1}{3}g_{1 \times 8} Q_{1/2}^V$ $\frac{1}{54}g_{1 \times 27} \left(Q_{1/2}^V + 5Q_{3/2}^V \right) + \frac{11}{36}g_{1 \times 8} Q_{1/2}^V$
$O_{sdu}^{V1,LR} = O_{sdu}^{V1,LR} _{8 \times 8} + O_{sdu}^{V1,LR} _{8 \times 1}$ $O_{sdu}^{V8,LR} = O_{sdu}^{V8,LR} _{8 \times 8} + O_{sdu}^{V8,LR} _{8 \times 1}$ $O_{sdd}^{V1,LR} = O_{sdd}^{V1,LR} _{8 \times 8} + O_{sdd}^{V1,LR} _{8 \times 1}$ $O_{sdd}^{V8,LR} = O_{sdd}^{V8,LR} _{8 \times 8} + O_{sdd}^{V8,LR} _{8 \times 1}$	$\frac{1}{6}g_{8 \times 8} \left(2Q_{1/2}^S + Q_{3/2}^S \right)$ $-\frac{1}{36}(g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \left(2Q_{1/2}^S + Q_{3/2}^S \right)$ $-\frac{1}{12}g_{8 \times 8} \left(Q_{1/2}^S + 2Q_{3/2}^S \right)$ $\frac{1}{72}(g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \left(Q_{1/2}^S + 2Q_{3/2}^S \right)$
$O_{uus}^{V1,LR} = O_{uus}^{V1,LR} _{8 \times 8} + O_{uus}^{V1,LR} _{1 \times 8}$ $O_{uus}^{V8,LR} = O_{uus}^{V8,LR} _{8 \times 8} + O_{uus}^{V8,LR} _{1 \times 8}$ $O_{dds}^{V1,LR} = O_{dds}^{V1,LR} _{8 \times 8} + O_{dds}^{V1,LR} _{1 \times 8}$ $O_{dds}^{V8,LR} = O_{dds}^{V8,LR} _{8 \times 8} + O_{dds}^{V8,LR} _{1 \times 8}$	$\frac{1}{6}g_{8 \times 8} \left(2Q_{1/2}^S + Q_{3/2}^S \right)$ $-\frac{1}{36}(g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \left(2Q_{1/2}^S + Q_{3/2}^S \right)$ $-\frac{1}{12}g_{8 \times 8} \left(Q_{1/2}^S + 2Q_{3/2}^S \right)$ $\frac{1}{72}(g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \left(Q_{1/2}^S + 2Q_{3/2}^S \right)$
$O_{uus}^{S1,RR} = O_{uus}^{S1,RR} _{\bar{6} \times 6} + O_{uus}^{S1,RR} _{3 \times \bar{3}}$ $O_{uus}^{S8,RR} = O_{uus}^{S8,RR} _{\bar{6} \times 6} + O_{uus}^{S8,RR} _{3 \times \bar{3}}$ $O_{uds}^{S1,RR} = O_{uds}^{S1,RR} _{\bar{6} \times 6} + O_{uds}^{S1,RR} _{3 \times \bar{3}}$ $O_{uds}^{S8,RR} = O_{uds}^{S8,RR} _{\bar{6} \times 6} + O_{uds}^{S8,RR} _{3 \times \bar{3}}$	$-\frac{1}{24}g_{\bar{6} \times 6} \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) + \frac{1}{8}g_{3 \times \bar{3}} Q_{1/2}^S$ $\frac{1}{144}(g_{\bar{6} \times 6} - 3\tilde{g}_{\bar{6} \times 6}) \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) - \frac{1}{48}(g_{3 \times \bar{3}} - 3\tilde{g}_{3 \times \bar{3}}) Q_{1/2}^S$ $-\frac{1}{24}g_{\bar{6} \times 6} \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) - \frac{1}{8}g_{3 \times \bar{3}} Q_{1/2}^S$ $\frac{1}{144}(g_{\bar{6} \times 6} - 3\tilde{g}_{\bar{6} \times 6}) \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) + \frac{1}{48}(g_{3 \times \bar{3}} - 3\tilde{g}_{3 \times \bar{3}}) Q_{1/2}^S$
$O_{uus}^{S1,LL} = O_{uus}^{S1,LL} _{6 \times \bar{6}} + O_{uus}^{S1,LL} _{\bar{3} \times 3}$ $O_{uus}^{S8,LL} = O_{uus}^{S8,LL} _{6 \times \bar{6}} + O_{uus}^{S8,LL} _{\bar{3} \times 3}$ $O_{uds}^{S1,LL} = O_{uds}^{S1,LL} _{6 \times \bar{6}} + O_{uds}^{S1,LL} _{\bar{3} \times 3}$ $O_{uds}^{S8,LL} = O_{uds}^{S8,LL} _{6 \times \bar{6}} + O_{uds}^{S8,LL} _{\bar{3} \times 3}$	$-\frac{1}{24}g_{6 \times \bar{6}} \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) + \frac{1}{8}g_{\bar{3} \times 3} Q_{1/2}^S$ $\frac{1}{144}(g_{6 \times \bar{6}} - 3\tilde{g}_{6 \times \bar{6}}) \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) + \frac{1}{48}(g_{\bar{3} \times 3} - 3\tilde{g}_{\bar{3} \times 3}) Q_{1/2}^S$ $-\frac{1}{24}g_{6 \times \bar{6}} \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) - \frac{1}{8}g_{\bar{3} \times 3} Q_{1/2}^S$ $\frac{1}{144}(g_{6 \times \bar{6}} - 3\tilde{g}_{6 \times \bar{6}}) \left(5Q_{1/2}^S + 4Q_{3/2}^S \right) + \frac{1}{48}(g_{\bar{3} \times 3} - 3\tilde{g}_{\bar{3} \times 3}) Q_{1/2}^S$

Table 3. The chiral representations and realizations of the LEFT four-quark operators contributing to $K \rightarrow \pi S(S)$. In the second column, $Q_{1/2}^{V,S}$ and $Q_{3/2}^{V,S}$ are the mesonic operators defined in eqs. (2.12) and (3.1) and correspond, respectively, to $\Delta I = 1/2$ and $3/2$ transitions. For $O_{q_1 q_2 q_3 q_4}^{V1,LR}$ and $O_{q_1 q_2 q_3 q_4}^{V8,LR}$ in rows 7–14, the contributions of the 8×1 and 1×8 terms are chirally subleading compared to their 8×8 counterparts and therefore dropped from the second column.

ators which could bring about the breaking of the GN bound. Simple instances include

$$\begin{aligned} \mathcal{O}_{ddsd}^{V,LL} - \frac{6}{11} \mathcal{O}_{uusd}^{V8,LL} &\Rightarrow \frac{10}{99} g_{27 \times 1} \left(\mathcal{Q}_{1/2}^V - \frac{13}{4} \mathcal{Q}_{3/2}^V \right) = \frac{-5}{22} g_{27 \times 1} F_0^2 \left(\partial_\mu K^+ \partial^\mu \pi^- + \frac{5\sqrt{2}}{3} \partial_\mu K^0 \partial^\mu \pi^0 \right), \\ \mathcal{O}_{sduu}^{V1,LR} + 3 \mathcal{O}_{sddd}^{V1,LR} &\Rightarrow \frac{1}{12} g_{8 \times 8} \left(\mathcal{Q}_{1/2}^S - 4 \mathcal{Q}_{3/2}^S \right) = -\frac{1}{4} g_{8 \times 8} F_0^2 \left(K^+ \pi^- + \frac{3}{\sqrt{2}} K^0 \pi^0 \right). \end{aligned} \quad (3.4)$$

We conclude that judicious choices for the coefficients of the operators would need to be made in order to evade the bound in a significant manner.⁵ We will encounter more examples later on.

3.2 Numerical analysis

Summing the mesonic operators in table 3 multiplied by their respective Wilson coefficients leads to the effective Lagrangian $\mathcal{L}_{K\pi S}$ responsible for $K \rightarrow \pi S$. We can express it as

$$\begin{aligned} \mathcal{L}_{K\pi S} &= F_0 \left(a_1 K^+ \pi^- - \frac{b_1}{\sqrt{2}} K^0 \pi^0 \right) S + \frac{1}{F_0} \left(a_2 \partial_\mu K^+ \partial^\mu \pi^- - \frac{b_2}{\sqrt{2}} \partial_\mu K^0 \partial^\mu \pi^0 \right) S + \text{H.c.} \\ &\supset F_0 [a_1 K^+ \pi^- - (\text{Re } b_1) K_L \pi^0] S + \frac{1}{F_0} [a_2 \partial_\mu K^+ \partial^\mu \pi^- - (\text{Re } b_2) \partial_\mu K_L \partial^\mu \pi^0] S, \end{aligned} \quad (3.5)$$

where $a_{1,2}$ and $b_{1,2}$ are dimensionless constants comprising linear combinations of the Wilson coefficients Cs, namely

$$\begin{aligned} a_1 &= \frac{1}{24} F_0 \left[6 \left(2\mathcal{C}_{sduu}^{V1,LR} - \mathcal{C}_{sddd}^{V1,LR} + 2\mathcal{C}_{uusd}^{V1,LR} - \mathcal{C}_{ddsd}^{V1,LR} \right) g_{8 \times 8} \right. \\ &\quad - \left(2\mathcal{C}_{sduu}^{V8,LR} - \mathcal{C}_{sddd}^{V8,LR} + 2\mathcal{C}_{uusd}^{V8,LR} - \mathcal{C}_{ddsd}^{V8,LR} \right) (g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \\ &\quad - 9 \left(\mathcal{C}_{uusd}^{S1,LL} + \mathcal{C}_{udsu}^{S1,LL} + \mathcal{C}_{uusd}^{S1,RR} + \mathcal{C}_{udsu}^{S1,RR} \right) g_{6 \times \bar{6}} \\ &\quad + \frac{3}{2} \left(\mathcal{C}_{uusd}^{S8,LL} + \mathcal{C}_{udsu}^{S8,LL} + \mathcal{C}_{uusd}^{S8,RR} + \mathcal{C}_{udsu}^{S8,RR} \right) (g_{6 \times \bar{6}} - 3\tilde{g}_{6 \times \bar{6}}) \\ &\quad + 3 \left(\mathcal{C}_{uusd}^{S1,LL} - \mathcal{C}_{udsu}^{S1,LL} + \mathcal{C}_{uusd}^{S1,RR} - \mathcal{C}_{udsu}^{S1,RR} \right) g_{3 \times \bar{3}} \\ &\quad \left. - \frac{1}{2} \left(\mathcal{C}_{uusd}^{S8,LL} - \mathcal{C}_{udsu}^{S8,LL} + \mathcal{C}_{uusd}^{S8,RR} - \mathcal{C}_{udsu}^{S8,RR} \right) (g_{3 \times \bar{3}} - 3\tilde{g}_{3 \times \bar{3}}) \right]_{\Lambda_\chi}, \end{aligned} \quad (3.6)$$

$$\begin{aligned} b_1 &= \frac{1}{24} F_0 \left[6 \left(\mathcal{C}_{sddd}^{V1,LR} + \mathcal{C}_{ddsd}^{V1,LR} \right) g_{8 \times 8} - \left(\mathcal{C}_{sddd}^{V8,LR} + \mathcal{C}_{ddsd}^{V8,LR} \right) (g_{8 \times 8} - 3\tilde{g}_{8 \times 8}) \right. \\ &\quad + 3 \left(\mathcal{C}_{uusd}^{S1,LL} + \mathcal{C}_{udsu}^{S1,LL} + \mathcal{C}_{uusd}^{S1,RR} + \mathcal{C}_{udsu}^{S1,RR} \right) g_{6 \times \bar{6}} \\ &\quad \left. - \frac{1}{2} \left(\mathcal{C}_{uusd}^{S8,LL} + \mathcal{C}_{udsu}^{S8,LL} + \mathcal{C}_{uusd}^{S8,RR} + \mathcal{C}_{udsu}^{S8,RR} \right) (g_{6 \times \bar{6}} - 3\tilde{g}_{6 \times \bar{6}}) \right] \end{aligned}$$

⁵This conclusion is in line with what has been argued qualitatively in [32], namely that for heavy mediators four-quark operators, especially those of the types in the first two sections of table 3, might not be able to cause $K \rightarrow \pi S$ decays to violate the GN bound by more than a factor of a few. As our examples demonstrate, preference for the $\Delta I = 3/2$ components of the operators is required to achieve substantial violations.

$$\begin{aligned}
 & + 3 \left(\mathbf{C}_{uusd}^{S1,LL} - \mathbf{C}_{udsu}^{S1,LL} + \mathbf{C}_{uusd}^{S1,RR} - \mathbf{C}_{udsu}^{S1,RR} \right) g_{3 \times \bar{3}} \\
 & - \frac{1}{2} \left(\mathbf{C}_{uusd}^{S8,LL} - \mathbf{C}_{udsu}^{S8,LL} + \mathbf{C}_{uusd}^{S8,RR} - \mathbf{C}_{udsu}^{S8,RR} \right) g_{3 \times \bar{3}} \Big]_{\Lambda_\chi}, \tag{3.7}
 \end{aligned}$$

$$\begin{aligned}
 a_2 = \frac{1}{36} F_0^3 \Big[& 6 \left(\mathbf{C}_{ddsd}^{V,LL} + \mathbf{C}_{ddsd}^{V,RR} - 2\mathbf{C}_{uusd}^{V1,LL} - 2\mathbf{C}_{uusd}^{V1,RR} \right) (g_{8 \times 1} - g_{27 \times 1}) \\
 & + \left(\mathbf{C}_{uusd}^{V8,LL} + \mathbf{C}_{uusd}^{V8,RR} \right) (11g_{8 \times 1} + 4g_{27 \times 1}) \Big]_{\Lambda_\chi}, \tag{3.8}
 \end{aligned}$$

$$\begin{aligned}
 b_2 = \frac{1}{36} F_0^3 \Big[& 6 \left(\mathbf{C}_{ddsd}^{V,LL} + \mathbf{C}_{ddsd}^{V,RR} \right) (g_{8 \times 1} + 4g_{27 \times 1}) - 6 \left(\mathbf{C}_{uusd}^{V1,LL} + \mathbf{C}_{uusd}^{V1,RR} \right) (2g_{8 \times 1} + 3g_{27 \times 1}) \\
 & + \left(\mathbf{C}_{uusd}^{V8,LL} + \mathbf{C}_{uusd}^{V8,RR} \right) (11g_{8 \times 1} - 6g_{27 \times 1}) \Big]_{\Lambda_\chi}, \tag{3.9}
 \end{aligned}$$

the subscript Λ_χ indicating that the \mathbf{C} s on the right-hand sides are evaluated at $\mu = \Lambda_\chi$. These coefficients scale as Λ_{NP}^{-3} .

For $K \rightarrow \pi SS$, the interaction Lagrangian $\mathcal{L}_{K\pi SS}$ has an expression similar to $S\mathcal{L}_{K\pi S}$, namely

$$\mathcal{L}_{K\pi SS} = \left(\hat{a}_1 K^+ \pi^- - \frac{\hat{b}_1}{\sqrt{2}} K^0 \pi^0 \right) S^2 + \left(\hat{a}_2 \partial_\mu K^+ \partial^\mu \pi^- - \frac{\hat{b}_2}{\sqrt{2}} \partial_\mu K^0 \partial^\mu \pi^0 \right) \frac{S^2}{F_0^2} + \text{H.c.} \tag{3.10}$$

The dimensionless parameters $\hat{a}_{1,2}$ and $\hat{b}_{1,2}$ are the same in form as $a_{1,2}F_0$ and $b_{1,2}F_0$, respectively, but with the Wilson coefficients now denoted by $\hat{\mathbf{C}}$ s, which scale as Λ_{NP}^{-4} because the underlying quark-level operators are of dimension eight.

3.2.1 $K \rightarrow \pi S$

From $\mathcal{L}_{K\pi S}$, we obtain the amplitudes for $K \rightarrow \pi S$ to be

$$\begin{aligned}
 \mathcal{A}_{K^+ \rightarrow \pi^+ S} &= a_1 F_0 + a_2 \frac{m_{K^+}^2 + m_{\pi^+}^2 - m_S^2}{2F_0}, \\
 \mathcal{A}_{K_L \rightarrow \pi^0 S} &= -\text{Re } b_1 F_0 - \text{Re } b_2 \frac{m_{K^0}^2 + m_{\pi^0}^2 - m_S^2}{2F_0}, \tag{3.11}
 \end{aligned}$$

and hence the branching fractions

$$\begin{aligned}
 \mathcal{B}(K^+ \rightarrow \pi^+ S) &= \tau_{K^+} \frac{\sqrt{(m_{K^+}^2 - m_{\pi^+}^2)^2 - (2m_{K^+}^2 + 2m_{\pi^+}^2 - m_S^2)m_S^2}}{16\pi m_{K^+}^3} \\
 &\quad \times \left| a_1 + a_2 \frac{m_{K^+}^2 + m_{\pi^+}^2 - m_S^2}{2F_0^2} \right|^2 F_0^2, \tag{3.12}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{B}(K_L \rightarrow \pi^0 S) &= \tau_{K_L} \frac{\sqrt{(m_{K^0}^2 - m_{\pi^0}^2)^2 - (2m_{K^0}^2 + 2m_{\pi^0}^2 - m_S^2)m_S^2}}{16\pi m_{K^0}^3} \\
 &\quad \times \left| \text{Re } b_1 + \text{Re } b_2 \frac{m_{K^0}^2 + m_{\pi^0}^2 - m_S^2}{2F_0^2} \right|^2 F_0^2. \tag{3.13}
 \end{aligned}$$

To account for the KOTO anomaly, one could consider various possibilities. For illustration, we look at a scenario in which the only contributing operators are those with purely right-handed quarks: $\mathcal{O}_{ddsd}^{V,RR}$ and $\mathcal{O}_{uusd}^{(V1,V8),RR}$. This implies that $a_1 = b_1 = 0$ and

$$\begin{aligned} a_2 &= \frac{F_0^3}{6} \left[\left(\mathcal{C}_{ddsd}^{V,RR} - 2\mathcal{C}_{uusd}^{V1,RR} + \frac{11}{6}\mathcal{C}_{uusd}^{V8,RR} \right) g_{8\times 1} - \left(\mathcal{C}_{ddsd}^{V,RR} - 2\mathcal{C}_{uusd}^{V1,RR} - \frac{2}{3}\mathcal{C}_{uusd}^{V8,RR} \right) g_{27\times 1} \right]_{\Lambda_\chi}, \\ b_2 &= \frac{F_0^3}{6} \left[\left(\mathcal{C}_{ddsd}^{V,RR} - 2\mathcal{C}_{uusd}^{V1,RR} + \frac{11}{6}\mathcal{C}_{uusd}^{V8,RR} \right) g_{8\times 1} + \left(4\mathcal{C}_{ddsd}^{V,RR} - 3\mathcal{C}_{uusd}^{V1,RR} - \mathcal{C}_{uusd}^{V8,RR} \right) g_{27\times 1} \right]_{\Lambda_\chi}. \end{aligned} \quad (3.14)$$

Moreover, we select $\mathcal{C}_{ddsd}^{V,RR} = 2\mathcal{C}_{uusd}^{V1,RR} - (11/6)\mathcal{C}_{uusd}^{V8,RR}$ to make the $g_{8\times 1}$ terms above vanish, changing eq. (3.14) to

$$a_2 = \frac{5F_0^3}{12} g_{27\times 1} \mathcal{C}_{uusd}^{V8,RR}(\Lambda_\chi), \quad b_2 = \frac{25F_0^3}{18} g_{27\times 1} \left(\frac{3}{5} \mathcal{C}_{uusd}^{V1,RR}(\Lambda_\chi) - \mathcal{C}_{uusd}^{V8,RR}(\Lambda_\chi) \right), \quad (3.15)$$

with which, for $m_S = 0$, we arrive at

$$\tilde{r}_B^{\text{NP}} = \frac{\mathcal{B}(K_L \rightarrow \pi^0 S)}{\mathcal{B}(K^+ \rightarrow \pi^+ S)} = 4.13 \frac{(17.6 \text{ Re } b_2)^2}{|17.4 a_2|^2} = 47 \left| \frac{\text{Re} \left[\mathcal{C}_{uusd}^{V8,RR}(\Lambda_\chi) - 0.6 \mathcal{C}_{uusd}^{V1,RR}(\Lambda_\chi) \right]}{\mathcal{C}_{uusd}^{V8,RR}(\Lambda_\chi)} \right|^2. \quad (3.16)$$

It is worth remarking that the potential enlargement of \tilde{r}_B^{NP} in this equation can be expected from the fact that it arises from the quark operators $\mathcal{O}_{ddsd}^{V,RR}$ and $\mathcal{O}_{uusd}^{(V1,V8),RR}$ which, as rows 4–6 in table 3 show, in the absence of the $g_{8\times 1}$ portions, generate the combinations $2\mathcal{Q}_{1/2}^V - 5\mathcal{Q}_{3/2}^V$ and $\mathcal{Q}_{1/2}^V + 5\mathcal{Q}_{3/2}^V$ of the mesonic operators defined in eq. (3.1) and hence all contain significant $\Delta I = 3/2$ components. Clearly, a much amplified \tilde{r}_B^{NP} can be easily realized with some more tuning of the parameters in eq. (3.16).

To be more precise in our numerical treatment, we again must take into account the QCD RG running of the coefficients from the EW scale, which we choose to be the W -boson mass m_W as before, down to the chiral-symmetry breaking scale Λ_χ . The pertinent one-loop RG equations are available in ref. [48], from which we collect the formulas in appendix B. We use them to get

$$\begin{aligned} \mathcal{C}_{uusd}^{V1,RR}(\Lambda_\chi) &= 1.07 \mathcal{C}_{uusd}^{V1,RR}(m_W) - 0.19 \mathcal{C}_{uusd}^{V8,RR}(m_W), \\ \mathcal{C}_{uusd}^{V8,RR}(\Lambda_\chi) &= 1.31 \mathcal{C}_{uusd}^{V8,RR}(m_W) - 0.86 \mathcal{C}_{uusd}^{V1,RR}(m_W) - 0.16 \mathcal{C}_{ddsd}^{V,RR}(m_W), \end{aligned} \quad (3.17)$$

which enter eq. (3.15) and depend on other coefficients. To simplify things further, we can pick $\mathcal{C}_{uusd}^{V1,RR}(m_W) = \mathcal{C}_{ddsd}^{V,RR}(m_W) = 0$ and $\mathcal{C}_{uusd}^{V8,RR}(m_W) = \Lambda_{\text{NP}}^{-3}$, which lead to $\tilde{r}_B^{\text{NP}} = 51$ in eq. (3.16). This exceeds the maximum $r_B^{\text{GN}} = 4.3$ of the GN bound by more than 10 times. For this example, in figure 2 we depict $\mathcal{B}(K \rightarrow \pi S)$ as functions of the S mass m_S with $\Lambda_{\text{NP}} = 1 \text{ TeV}$ (left panel) and 800 GeV (right panel). In the figure, we also exhibit the upper limits on $\mathcal{B}(K_L \rightarrow \pi^0 S)$ and $\mathcal{B}(K^+ \rightarrow \pi^+ S)$ at 90% CL from KOTO 2015 [15]




Figure 2. The branching fractions of $K \rightarrow \pi S$ versus the S mass m_S induced by the dim-7 operators with NP scales $\Lambda_{\text{NP}} = 1 \text{ TeV}$ (left panel) and 800 GeV (right panel), compared to the experimental upper limits from KOTO 2015 [15] and BNL [17] along with the standard GN constraint on $K_L \rightarrow \pi^0 S$ from the BNL result. The blue and black curves are related by $\mathcal{B}(K_L \rightarrow \pi^0 S)_{\text{GN}} = 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ S)_{\text{BNL}}$.

and BNL [17], respectively, along with the empirical GN constraint based on the BNL result: $\mathcal{B}(K_L \rightarrow \pi^0 S)_{\text{GN}} < 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ S)_{\text{BNL}}$, which is reflected by the blue and black curves in each graph. If $\Lambda_{\text{NP}} = 1 \text{ TeV}$, the left panel reveals that the GN inequality is not respected in the $m_S \lesssim 110 \text{ MeV}$ region, while the current experimental limits are satisfied. If Λ_{NP} is smaller, it is possible to break the GN bound with higher m_S values, such as $170 \text{ MeV} \lesssim m_S \lesssim 240 \text{ MeV}$ in the right panel for $\Lambda_{\text{NP}} = 800 \text{ GeV}$. We conclude that it can be violated by dim-7 EFT operators with a NP scale $\Lambda_{\text{NP}} = \mathcal{O}(1 \text{ TeV})$.

3.2.2 $K \rightarrow \pi SS$

For the three-body decays $K \rightarrow \pi SS$, the amplitudes are

$$\begin{aligned} \mathcal{A}_{K^+ \rightarrow \pi^+ SS} &= 2\hat{a}_1 + \hat{a}_2 \frac{m_{K^+}^2 + m_{\pi^+}^2 - \hat{s}}{F_0^2}, \\ \mathcal{A}_{K_L \rightarrow \pi^0 SS} &= -2 \text{Re} \hat{b}_1 - \text{Re} \hat{b}_2 \frac{m_{K^0}^2 + m_{\pi^0}^2 - \hat{s}}{F_0^2}, \end{aligned} \quad (3.18)$$

where \hat{s} designates the invariant mass squared of the SS pair. These bring about another modified definition of r_B ,

$$r_B = \frac{\mathcal{B}(K_L \rightarrow \pi^0 + E_{\text{miss}})}{\mathcal{B}(K^+ \rightarrow \pi^+ + E_{\text{miss}})} = \frac{\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{SM}} + \mathcal{B}(K_L \rightarrow \pi^0 SS)}{\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} + \mathcal{B}(K^+ \rightarrow \pi^+ SS)}. \quad (3.19)$$

As before, we have many options regarding the parameters which can yield a violation of the GN bound. This time we entertain the possibility that the operators with purely left-handed or right-handed quarks do not contribute, implying that $\hat{a}_2 = \hat{b}_2 = 0$ and so




Figure 3. The branching fractions of $K \rightarrow \pi SS$ induced by the dim-8 operators as functions of the NP scale Λ_{NP} for $m_S = 0$, as described in the text. Also displayed are the corresponding SM predictions for $K \rightarrow \pi \nu \bar{\nu}$ (red and blue horizontal bands) and upper limits from KOTO [15] and NA62 [19] (blue and red horizontal thin lines). The light-blue region is excluded by the KOTO bound. The blue dot corresponds to KOTO’s three events.

the branching fractions become, for $m_S = 0$,

$$\mathcal{B}(K^+ \rightarrow \pi^+ SS) = \frac{\tau_{K^+}}{2^9 \pi^3 m_{K^+}^3} \int \Pi_3 |\mathcal{M}_{K^+ \rightarrow \pi^+ 2S}|^2 = \frac{\tau_{K^+} |\hat{a}_1|^2}{2^7 \pi^3 m_{K^+}^3} \int \Pi_3 = 7 \times 10^{11} |\hat{a}_1|^2,$$

$$\mathcal{B}(K_L \rightarrow \pi^0 SS) = \frac{\tau_{K_L}}{2^9 \pi^3 m_{K^0}^3} \int \Pi_3 |\mathcal{M}_{K_L \rightarrow \pi^0 2S}|^2 = \frac{\tau_{K_L} (\text{Re } \hat{b}_1)^2}{2^7 \pi^3 m_{K^0}^3} \int \Pi_3 = 3 \times 10^{12} (\text{Re } \hat{b}_1)^2. \tag{3.20}$$

Supposing additionally that \hat{a}_1 and \hat{b}_1 come only from the $g_{8 \times 8}$ terms in eqs. (3.6) and (3.7) and setting $\mathbf{C}_{sddd}^{V1,LR}/2 = \mathbf{C}_{dds d}^{V1,LR} = \mathbf{C}_{uusd}^{V1,LR} = \mathbf{C}_{sdvu}^{V1,LR} = \Lambda_{\text{NP}}^{-4}$, we derive

$$\hat{a}_1 = \frac{F_0^2}{4} \frac{g_{8 \times 8}}{\Lambda_{\text{NP}}^4}, \quad \hat{b}_1 = \frac{F_0^2}{4} \frac{3g_{8 \times 8}}{\Lambda_{\text{NP}}^4}. \tag{3.21}$$

In figure 3 we plot the resulting branching fractions of $K \rightarrow \pi SS$ as functions of Λ_{NP} . Evidently, in this particular instance, to boost $\mathcal{B}(K_L \rightarrow \pi^0 SS)$ to a level within KOTO’s current sensitivity reach would need Λ_{NP} to be no more than roughly 200 GeV. As can be inferred from this graph in conjunction with eq. (3.19), for Λ_{NP} above this value the GN bound is no longer violated.

4 Summary and conclusions

Motivated by the recent preliminary observation of three anomalous events of $K_L \rightarrow \pi^0 \nu \bar{\nu}$ by the KOTO collaboration, we study in detail the possibility of having new physics responsible for enhancing the $K \rightarrow \pi + E_{\text{miss}}$ modes over their SM expectations. We explore two types of scenarios:

- NP above the EW scale represented by quark-neutrino interactions which do not preserve lepton flavor/number.
- NP above the EW scale with new scalar particles that are sufficiently light to be produced in $K \rightarrow \pi + E_{\text{miss}}$ decays.

The NP is described with an effective Lagrangian above the EW scale that respects the gauge symmetries of the SM. In all the cases considered, we specifically look for true violations of the Grossman-Nir bound through four-quark $\Delta I = 3/2$ interactions.

The NP effects are classified according to the mass dimensionality of the necessary operators. To this end, we catalogue all the operators that can give rise to the reaction $K \rightarrow \pi \mathcal{X}$ with \mathcal{X} standing for one or more particles carrying away the missing energy. As itemized above, we allow \mathcal{X} to comprise: a neutrino-antineutrino pair ($\nu\bar{\nu}$), a pair of neutrinos ($\nu\nu$) or antineutrinos ($\bar{\nu}\bar{\nu}$), an invisible light new scalar boson (S), and a pair of these scalars (SS). These cases require a minimal dimensionality of ten, nine, seven, and eight, respectively. On general grounds, we argue that the scenarios with new scalars (dim-7 or -8 operators) are consistent with sizable boosts in the rate of $K_L \rightarrow \pi^0 + E_{\text{miss}}$ for NP scales above the EW scale.

We construct the effective Lagrangian for each of the cases and, after identifying the $\Delta I = 3/2$ components of the operators, we discuss the renormalization group running of the couplings down to a hadronic scale followed by the matching of the operators onto chiral perturbation theory. We present numerical results illustrating the scale of the NP needed to amplify the $K_L \rightarrow \pi^0 + E_{\text{miss}}$ rate above the GN bound obtained from the measurements of $K^+ \rightarrow \pi^+ + E_{\text{miss}}$.

We find that the production of a single light new scalar via $\Delta I = 3/2$ interactions permits enlargements in the $K_L \rightarrow \pi^0 + E_{\text{miss}}$ rate that are big enough to appear in the KOTO experiment, and we clarify this with figure 2. This is achievable with some degree of tuning among the coefficients of the operators. Our results are attained for stable new scalars, but long-lived ones would also work as they have weaker constraints [17].

The production of a pair of the new light scalars could have substantial rate gains over the SM but not above the GN bound. We depict this in figure 4,⁶ where the blue area illustrates that increases over the SM by factors of a few are possible while keeping $\Lambda_{\text{NP}} \geq v$ [in contrast, to exceed the bound (explain the KOTO events) would require $\Lambda_{\text{NP}} < 200 \text{ GeV}$ ($\Lambda_{\text{NP}} \sim 160 \text{ GeV}$), as indicated in figure 3]. With a different choice of parameters, the charged mode could also be amplified by a similar amount.

The production of two neutrinos, on the other hand, suffers from relatively much greater Λ_{NP} suppression. The restriction $\Lambda_{\text{NP}} \geq v$ results in very small rises over the SM, completely within the uncertainty of the SM predictions and thus unobservable.

⁶To draw the blue region, we use again the example in eq. (3.20), with $\hat{a}_1 = F_0^2 g_{8 \times 8} / (4\Lambda_{\text{NP}}^4)$ as in eq. (3.21), but now let $\text{Re} \hat{b}_1$ vary under the condition $0 \leq |\text{Re} \hat{b}_1| \leq 3|\hat{a}_1|$ and demand $\Lambda_{\text{NP}} \geq v$. With regard to the operator coefficients in eqs. (3.6) and (3.7), one way to accomplish this is to arrange $2\mathbf{C}_{sduu}^{V1,LR} - \mathbf{C}_{sddd}^{V1,LR} + 2\mathbf{C}_{uusd}^{V1,LR} - \mathbf{C}_{ddsd}^{V1,LR} = \Lambda_{\text{NP}}^{-4}$ and $0 \leq \mathbf{C}_{sddd}^{V1,LR} + \mathbf{C}_{ddsd}^{V1,LR} \leq 3$, having taken the others to vanish.




Figure 4. The branching fractions of $K \rightarrow \pi + E_{\text{miss}}$: in red the 90%-CL SM predictions; in green the 1σ BNL E787/E949 result; in brown the 90% NA62 exclusion; in grey the GN bound; and in blue a region accessible with $K \rightarrow \pi SS$ for parameters chosen to enhance mostly the neutral mode with a NP scale $\Lambda_{\text{NP}} \geq v$.

We conclude that continued improvement of the KOTO upper bound on $K_L \rightarrow \pi^0 + E_{\text{miss}}$, even at current levels which are much above the GN bound, provides relevant constraints on possible new physics scenarios.

Acknowledgments

This work was supported in part by the MOST (Grant No. MOST 106-2112-M-002-003-MY3). This work was also supported in part by the Australian Government through the Australian Research Council.

A Isospin decomposition of quark parts of dim-9 operators

In this appendix, for completeness we write down the decomposition of the quark portion of each of the dim-9 operators in eq. (2.2) into its $\Delta I = 1/2, 3/2$ components. This will allow us to see clearly the difference between them. Since additionally each operator also causes a definite change ΔI_3 in the third isospin component, we can first group them according to their ΔI_3 values and then express them as linear combinations of their ΔI terms. Inspecting the operators, we find that 0_1^{usdu} , $\tilde{0}_1^{usdu}$, 0_3^{ddds} , and $\tilde{0}_3^{ddds}$ have $\Delta I_3 = 1/2$, whereas 0_1^{udsu} , $\tilde{0}_1^{udsu}$, 0_3^{ddsd} , and $\tilde{0}_3^{ddsd}$ have $\Delta I_3 = -1/2$. Employing the Clebsch-Gordan decomposition rule, we then get the following results:

- The $\Delta I_3 = 1/2$ operators:

$$0_1^{usdu} = -\frac{1}{3}0_{1,\Delta I=1/2}^{usdu} + \frac{1}{3}0_{1,\Delta I=3/2}^{usdu}, \quad 0_3^{ddds} = \frac{1}{3}0_{3,\Delta I=1/2}^{ddds} - \frac{1}{3}0_{3,\Delta I=3/2}^{ddds}, \quad (\text{A.1})$$

with their components of definite ΔI being given by

$$\begin{aligned}
 \mathcal{O}_{1,\Delta I=1/2}^{usdu} &= \left[(\bar{d}_L \gamma_\mu s_L) (\bar{u}_R \gamma^\mu u_R) - 2(\bar{u}_L \gamma_\mu s_L) (\bar{d}_R \gamma^\mu u_R) - (\bar{d}_L \gamma_\mu s_L) (\bar{d}_R \gamma^\mu d_R) \right] J, \\
 \mathcal{O}_{1,\Delta I=3/2}^{usdu} &= \left[(\bar{d}_L \gamma_\mu s_L) (\bar{u}_R \gamma^\mu u_R) + (\bar{u}_L \gamma_\mu s_L) (\bar{d}_R \gamma^\mu u_R) - (\bar{d}_L \gamma_\mu s_L) (\bar{d}_R \gamma^\mu d_R) \right] J, \\
 \mathcal{O}_{3,\Delta I=1/2}^{dds} &= \left[(\bar{u}_R u_L) (\bar{d}_R s_L) + (\bar{d}_R u_L) (\bar{u}_R s_L) + 2(\bar{d}_R d_L) (\bar{d}_R s_L) \right] J, \\
 \mathcal{O}_{3,\Delta I=3/2}^{dds} &= \left[(\bar{u}_R u_L) (\bar{d}_R s_L) + (\bar{d}_R u_L) (\bar{u}_R s_L) - (\bar{d}_R d_L) (\bar{d}_R s_L) \right] J,
 \end{aligned} \tag{A.2}$$

and similarly $\tilde{\mathcal{O}}_1^{usdu}$ and $\tilde{\mathcal{O}}_3^{dds}$.

- The $\Delta I_3 = -1/2$ operators:

$$\mathcal{O}_1^{usdu} = -\frac{1}{3} \mathcal{O}_{1,\Delta I=1/2}^{usdu} + \frac{1}{3} \mathcal{O}_{1,\Delta I=3/2}^{usdu}, \quad \mathcal{O}_3^{dds} = \frac{1}{3} \mathcal{O}_{3,\Delta I=1/2}^{dds} - \frac{1}{3} \mathcal{O}_{3,\Delta I=3/2}^{dds}, \tag{A.3}$$

with their components of definite ΔI being given by

$$\begin{aligned}
 \mathcal{O}_{1,\Delta I=1/2}^{usdu} &= \left[(\bar{u}_L \gamma_\mu u_L) (\bar{s}_R \gamma^\mu d_R) - 2(\bar{u}_L \gamma_\mu d_L) (\bar{s}_R \gamma^\mu u_R) - (\bar{d}_L \gamma_\mu d_L) (\bar{s}_R \gamma^\mu d_R) \right] J, \\
 \mathcal{O}_{1,\Delta I=3/2}^{usdu} &= \left[(\bar{u}_L \gamma_\mu u_L) (\bar{s}_R \gamma^\mu d_R) + (\bar{u}_L \gamma_\mu d_L) (\bar{s}_R \gamma^\mu u_R) - (\bar{d}_L \gamma_\mu d_L) (\bar{s}_R \gamma^\mu d_R) \right] J, \\
 \mathcal{O}_{3,\Delta I=1/2}^{dds} &= \left[(\bar{u}_R u_L) (\bar{s}_R d_L) + (\bar{u}_R d_L) (\bar{s}_R u_L) + 2(\bar{d}_R d_L) (\bar{s}_R d_L) \right] J, \\
 \mathcal{O}_{3,\Delta I=3/2}^{dds} &= \left[(\bar{u}_R u_L) (\bar{s}_R d_L) + (\bar{u}_R d_L) (\bar{s}_R u_L) - (\bar{d}_R d_L) (\bar{s}_R d_L) \right] J,
 \end{aligned} \tag{A.4}$$

and similarly $\tilde{\mathcal{O}}_1^{usdu}$ and $\tilde{\mathcal{O}}_3^{dds}$.

B RG running of dim-6 four-quark operators for $K \rightarrow \pi S(S)$

The 1-loop QCD RG equations of the Wilson coefficients of the LEFT dim-6 quark operators relevant to the $K \rightarrow \pi S(S)$ transitions are given by [48]

$$\mu \frac{d}{d\mu} \begin{pmatrix} \mathcal{C}_{dds}^{V,LL} \\ \mathcal{C}_{uus}^{V1,LL} \\ \mathcal{C}_{uus}^{V8,LL} \\ \mathcal{C}_{sdu}^{V1,LR} \\ \mathcal{C}_{sdu}^{V8,LR} \\ \mathcal{C}_{sdd}^{V1,LR} \\ \mathcal{C}_{sdd}^{V8,LR} \end{pmatrix} = -\frac{\alpha_s}{2\pi} \begin{pmatrix} -\frac{20}{9} & 0 & -\frac{1}{18} & 0 & -\frac{1}{18} & 0 & -\frac{1}{18} \\ 0 & 0 & -\frac{4}{3} & 0 & 0 & 0 & 0 \\ -\frac{4}{3} & -6 & \frac{5}{3} & 0 & -\frac{1}{3} & 0 & -\frac{1}{3} \\ 0 & 0 & 0 & 0 & \frac{4}{3} & 0 & 0 \\ -\frac{4}{3} & 0 & -\frac{1}{3} & 6 & -\frac{22}{3} & 0 & -\frac{1}{3} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{4}{3} \\ -\frac{4}{3} & 0 & -\frac{1}{3} & 0 & -\frac{1}{3} & 6 & -\frac{22}{3} \end{pmatrix} \begin{pmatrix} \mathcal{C}_{dds}^{V,LL} \\ \mathcal{C}_{uus}^{V1,LL} \\ \mathcal{C}_{uus}^{V8,LL} \\ \mathcal{C}_{sdu}^{V1,LR} \\ \mathcal{C}_{sdu}^{V8,LR} \\ \mathcal{C}_{sdd}^{V1,LR} \\ \mathcal{C}_{sdd}^{V8,LR} \end{pmatrix}, \tag{B.1}$$

$$\mu \frac{d}{d\mu} \begin{pmatrix} \mathcal{C}_{uus}^{S1,LL} \\ \mathcal{C}_{uus}^{S8,LL} \\ \mathcal{C}_{uds}^{S1,LL} \\ \mathcal{C}_{uds}^{S8,LL} \end{pmatrix} = -\frac{\alpha_s}{2\pi} \begin{pmatrix} 8 & -\frac{8}{9} & -\frac{32}{9} & -\frac{56}{27} \\ -4 & -\frac{8}{3} & \frac{8}{3} & -\frac{22}{9} \\ -\frac{32}{9} & -\frac{56}{27} & 8 & -\frac{8}{9} \\ \frac{8}{3} & -\frac{22}{9} & -4 & -\frac{8}{3} \end{pmatrix} \begin{pmatrix} \mathcal{C}_{uus}^{S1,LL} \\ \mathcal{C}_{uus}^{S8,LL} \\ \mathcal{C}_{uds}^{S1,LL} \\ \mathcal{C}_{uds}^{S8,LL} \end{pmatrix}. \tag{B.2}$$

The solutions to these equations between the electroweak scale, which we take to be $\mu = m_W$, and the chiral symmetry breaking $\mu = \Lambda_\chi = 4\pi F_\pi \simeq 1.2 \text{ GeV}$ are

$$\begin{pmatrix} \mathcal{C}_{ddsd}^{V,LL} \\ \mathcal{C}_{uusd}^{V1,LL} \\ \mathcal{C}_{uusd}^{V8,LL} \\ \mathcal{C}_{sduu}^{V1,LR} \\ \mathcal{C}_{sduu}^{V8,LR} \\ \mathcal{C}_{sddd}^{V1,LR} \\ \mathcal{C}_{sddd}^{V8,LR} \end{pmatrix}_{\mu=\Lambda_\chi} = \begin{pmatrix} 0.76 & 0.00 & -0.01 & -0.00 & -0.00 & -0.00 & -0.00 \\ 0.01 & 1.07 & -0.19 & 0.00 & 0.00 & 0.00 & 0.00 \\ -0.16 & -0.86 & 1.31 & -0.01 & -0.03 & -0.01 & -0.03 \\ -0.01 & 0.00 & -0.00 & 1.05 & 0.11 & -0.00 & -0.00 \\ -0.09 & 0.01 & -0.03 & 0.51 & 0.43 & -0.01 & -0.02 \\ -0.01 & 0.00 & -0.00 & -0.00 & -0.00 & 1.05 & 0.11 \\ -0.09 & 0.01 & -0.03 & -0.01 & -0.02 & 0.51 & 0.43 \end{pmatrix} \begin{pmatrix} \mathcal{C}_{ddsd}^{V,LL} \\ \mathcal{C}_{uusd}^{V1,LL} \\ \mathcal{C}_{uusd}^{V8,LL} \\ \mathcal{C}_{sduu}^{V1,LR} \\ \mathcal{C}_{sduu}^{V8,LR} \\ \mathcal{C}_{sddd}^{V1,LR} \\ \mathcal{C}_{sddd}^{V8,LR} \end{pmatrix}_{\mu=m_W}, \quad (\text{B.3})$$

$$\begin{pmatrix} \mathcal{C}_{uusd}^{S1,LL} \\ \mathcal{C}_{uusd}^{S8,LL} \\ \mathcal{C}_{udsu}^{S1,LL} \\ \mathcal{C}_{udsu}^{S8,LL} \end{pmatrix}_{\mu=\Lambda_\chi} = \begin{pmatrix} 2.97 & -0.03 & -1.17 & -0.36 \\ -1.01 & 0.71 & 0.84 & -0.16 \\ -1.17 & -0.36 & 2.97 & -0.03 \\ 0.84 & -0.16 & -1.01 & 0.71 \end{pmatrix} \begin{pmatrix} \mathcal{C}_{uusd}^{S1,LL} \\ \mathcal{C}_{uusd}^{S8,LL} \\ \mathcal{C}_{udsu}^{S1,LL} \\ \mathcal{C}_{udsu}^{S8,LL} \end{pmatrix}_{\mu=m_W}. \quad (\text{B.4})$$

All of these formulas are also valid for the chirality-flipped counterparts of the operators.

C Chiral structure and hadronization of quark operators for $K \rightarrow \pi S(S)$

Here we collect the $SU(3)_L \times SU(3)_R$ irreducible representations of the dim-6 four-quark operators examined in section 3 and the corresponding mesonic operators decomposed into their $\Delta I = 1/2, 3/2$ components. Adopting the normalization convention of ref. [42] for the chiral realization of each operator,⁷ we have

$$\begin{aligned} \mathcal{O}_{ddsd}^{V,LL}|_{27 \times 1} &= \frac{1}{5} [(4\bar{d}_L \gamma^\mu d_L - \bar{s}_L \gamma^\mu s_L)(\bar{s}_L \gamma_\mu d_L) - (\bar{u}_L \gamma^\mu d_L)(\bar{s}_L \gamma_\mu u_L)] - \frac{1}{5} (\bar{q}_L \gamma^\mu q_L)(\bar{s}_L \gamma_\mu d_L) \\ &\Rightarrow \frac{1}{12} g_{27 \times 1} F_0^4 [4\tilde{L}_{\mu 22} \tilde{L}_{23}^\mu - \tilde{L}_{\mu 33} \tilde{L}_{23}^\mu - \tilde{L}_{\mu 21} \tilde{L}_{13}^\mu] \supset \frac{1}{18} g_{27 \times 1} (2\mathcal{Q}_{1/2}^V - 5\mathcal{Q}_{3/2}^V), \end{aligned} \quad (\text{C.1})$$

$$\begin{aligned} \mathcal{O}_{ddsd}^{V,LL}|_{8 \times 1} &= \frac{1}{5} [(\bar{u}_L \gamma^\mu d_L)(\bar{s}_L \gamma_\mu u_L) - (\bar{u}_L \gamma^\mu u_L)(\bar{s}_L \gamma_\mu d_L)] + \frac{2}{5} (\bar{q}_L \gamma^\mu q_L)(\bar{s}_L \gamma_\mu d_L) \\ &\Rightarrow \frac{1}{12} g_{8 \times 1} F_0^4 [\tilde{L}_{\mu 21} \tilde{L}_{13}^\mu - \tilde{L}_{\mu 11} \tilde{L}_{23}^\mu] \supset \frac{1}{6} g_{8 \times 1} \mathcal{Q}_{1/2}^V, \end{aligned} \quad (\text{C.2})$$

$$\begin{aligned} \mathcal{O}_{uusd}^{V1,LL}|_{27 \times 1} &= \frac{1}{5} [3(\bar{u}_L \gamma^\mu u_L)(\bar{s}_L \gamma_\mu d_L) + 2(\bar{u}_L \gamma^\mu d_L)(\bar{s}_L \gamma_\mu u_L)] - \frac{1}{5} (\bar{q}_L \gamma^\mu q_L)(\bar{s}_L \gamma_\mu d_L) \\ &\Rightarrow \frac{1}{12} g_{8 \times 1} F_0^4 [3\tilde{L}_{\mu 11} \tilde{L}_{23}^\mu + 2\tilde{L}_{\mu 21} \tilde{L}_{13}^\mu] \supset \frac{1}{18} g_{27 \times 1} (\mathcal{Q}_{1/2}^V + 5\mathcal{Q}_{3/2}^V), \end{aligned} \quad (\text{C.3})$$

$$\begin{aligned} \mathcal{O}_{uusd}^{V1,LL}|_{8 \times 1} &= \frac{2}{5} [(\bar{u}_L \gamma^\mu u_L)(\bar{s}_L \gamma_\mu d_L) - (\bar{u}_L \gamma^\mu d_L)(\bar{s}_L \gamma_\mu u_L)] + \frac{1}{5} (\bar{q}_L \gamma^\mu q_L)(\bar{s}_L \gamma_\mu d_L) \\ &\Rightarrow \frac{1}{6} g_{8 \times 1} F_0^4 [\tilde{L}_{\mu 11} \tilde{L}_{23}^\mu - \tilde{L}_{\mu 21} \tilde{L}_{13}^\mu] \supset -\frac{1}{3} g_{8 \times 1} \mathcal{Q}_{1/2}^V, \end{aligned} \quad (\text{C.4})$$

⁷Particularly $(\bar{s}_L \gamma_\mu d_L)(\bar{s}_L \gamma^\mu d_L) \Rightarrow \frac{5}{12} g_{27 \times 1} F_0^4 \tilde{L}_{\mu 23} \tilde{L}_{23}^\mu$ and $(\bar{s}_L \gamma_\mu d_L)(\bar{s}_R \gamma^\mu d_R) \Rightarrow \frac{1}{4} g_{8 \times 8} F_0^4 \Sigma_{23} \Sigma_{23}^\dagger$ among the operators with purely left-handed quarks and quarks of mixed chirality, respectively.

$$O_{usd}^{V8,LL}|_{27\times 1} = \frac{1}{3} O_{usd}^{V1,LL}|_{27\times 1}, \quad (C.5)$$

$$\begin{aligned} O_{usd}^{V8,LL}|_{8\times 1} &= \frac{11}{30} [(\bar{u}_L\gamma^\mu d_L)(\bar{s}_L\gamma_\mu u_L) - (\bar{u}_L\gamma^\mu u_L)(\bar{s}_L\gamma_\mu d_L)] + \frac{1}{15} (\bar{q}_L\gamma^\mu q_L)(\bar{s}_L\gamma_\mu d_L) \\ &\Rightarrow \frac{11}{72} g_{8\times 1} F_0^4 [\tilde{L}_{\mu 21}\tilde{L}_{13}^\mu - \tilde{L}_{\mu 11}\tilde{L}_{23}^\mu] \supset \frac{11}{36} g_{8\times 1} \mathcal{Q}_{1/2}^V, \end{aligned} \quad (C.6)$$

$$\begin{aligned} O_{sduu}^{V1,LR}|_{8\times 8} &= (\bar{s}_L\gamma_\mu d_L) \left[(\bar{u}_R\gamma^\mu u_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right] \\ &\Rightarrow \frac{F_0^4}{4} g_{8\times 8} \Sigma_{21}\Sigma_{13}^\dagger \supset \frac{1}{6} g_{8\times 8} (2\mathcal{Q}_{1/2}^S + \mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.7)$$

$$\begin{aligned} O_{sduu}^{V8,LR}|_{8\times 8} &= -\frac{1}{6} (\bar{s}_L\gamma_\mu d_L) \left[(\bar{u}_R\gamma^\mu u_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right] + \frac{1}{2} (\bar{s}_L\gamma_\mu d_L) \left\{ (\bar{u}_R\gamma^\mu u_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right\} \\ &\Rightarrow -\frac{1}{6} \frac{F_0^4}{4} (g_{8\times 8} - 3\tilde{g}_{8\times 8}) \Sigma_{21}\Sigma_{13}^\dagger \supset \frac{1}{36} (3\tilde{g}_{8\times 8} - g_{8\times 8}) (2\mathcal{Q}_{1/2}^S + \mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.8)$$

$$\begin{aligned} O_{sddd}^{V1,LR}|_{8\times 8} &= (\bar{s}_L\gamma_\mu d_L) \left[(\bar{d}_R\gamma^\mu d_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right] \\ &\Rightarrow \frac{F_0^4}{4} g_{8\times 8} \Sigma_{22}\Sigma_{23}^\dagger \supset -\frac{1}{12} g_{8\times 8} (\mathcal{Q}_{1/2}^S + 2\mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.9)$$

$$\begin{aligned} O_{sddd}^{V8,LR}|_{8\times 8} &= -\frac{1}{6} (\bar{s}_L\gamma_\mu d_L) \left[(\bar{d}_R\gamma^\mu d_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right] + \frac{1}{2} (\bar{s}_L\gamma_\mu d_L) \left\{ (\bar{d}_R\gamma^\mu d_R) - \frac{1}{3} (\bar{q}_R\gamma^\mu q_R) \right\} \\ &\Rightarrow -\frac{1}{6} \frac{F_0^4}{4} (g_{8\times 8} - 3\tilde{g}_{8\times 8}) \Sigma_{22}\Sigma_{23}^\dagger \supset \frac{1}{72} (g_{8\times 8} - 3\tilde{g}_{8\times 8}) (\mathcal{Q}_{1/2}^S + 2\mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.10)$$

$$\begin{aligned} O_{usd}^{S1,LL}|_{6\times \bar{6}} &= \frac{1}{2} [(\bar{u}_R u_L)(\bar{s}_R d_L) + (\bar{u}_R d_L)(\bar{s}_R u_L)] \\ &\Rightarrow \frac{1}{2} \frac{F_0^4}{4} g_{6\times \bar{6}} (\Sigma_{23}\Sigma_{11} + \Sigma_{13}\Sigma_{21}) \supset -\frac{1}{24} g_{6\times \bar{6}} (5\mathcal{Q}_{1/2}^S + 4\mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.11)$$

$$\begin{aligned} O_{usd}^{S1,LL}|_{\bar{3}\times 3} &= \frac{1}{2} [(\bar{u}_R u_L)(\bar{s}_R d_L) - (\bar{u}_R d_L)(\bar{s}_R u_L)] \\ &\Rightarrow \frac{1}{2} \frac{F_0^4}{4} g_{\bar{3}\times 3} (\Sigma_{23}\Sigma_{11} - \Sigma_{13}\Sigma_{21}) \supset \frac{1}{8} g_{\bar{3}\times 3} \mathcal{Q}_{1/2}^S, \end{aligned} \quad (C.12)$$

$$\begin{aligned} O_{usd}^{S8,LL}|_{6\times \bar{6}} &= -\frac{1}{12} [(\bar{u}_R u_L)(\bar{s}_R d_L) + (\bar{u}_R d_L)(\bar{s}_R u_L)] + \frac{1}{4} \{(\bar{u}_R u_L)[\bar{s}_R d_L] + (\bar{u}_R d_L)[\bar{s}_R u_L]\} \\ &\Rightarrow \frac{F_0^4}{84} (g_{6\times \bar{6}} - 3\tilde{g}_{6\times \bar{6}}) (\Sigma_{23}\Sigma_{11} + \Sigma_{13}\Sigma_{21}) \supset \frac{g_{6\times \bar{6}} - 3\tilde{g}_{6\times \bar{6}}}{144} (5\mathcal{Q}_{1/2}^S + 4\mathcal{Q}_{3/2}^S), \end{aligned} \quad (C.13)$$

$$\begin{aligned} O_{usd}^{S8,LL}|_{\bar{3}\times 3} &= -\frac{1}{12} [(\bar{u}_R u_L)(\bar{s}_R d_L) - (\bar{u}_R d_L)(\bar{s}_R u_L)] + \frac{1}{4} \{(\bar{u}_R u_L)[\bar{s}_R d_L] - (\bar{u}_R d_L)[\bar{s}_R u_L]\} \\ &\Rightarrow -\frac{F_0^4}{48} (g_{\bar{3}\times 3} - 3\tilde{g}_{\bar{3}\times 3}) (\Sigma_{23}\Sigma_{11} - \Sigma_{13}\Sigma_{21}) \supset \frac{3\tilde{g}_{\bar{3}\times 3} - g_{\bar{3}\times 3}}{48} \mathcal{Q}_{1/2}^S, \end{aligned} \quad (C.14)$$

$$\begin{aligned} O_{udsu}^{S1,LL}|_{6\times \bar{6}} &= O_{usd}^{S1,LL}|_{6\times \bar{6}}, & O_{udsu}^{S1,LL}|_{\bar{3}\times 3} &= -O_{usd}^{S1,LL}|_{\bar{3}\times 3}, \\ O_{udsu}^{S8,LL}|_{6\times \bar{6}} &= O_{usd}^{S8,LL}|_{6\times \bar{6}}, & O_{udsu}^{S8,LL}|_{\bar{3}\times 3} &= -O_{usd}^{S8,LL}|_{\bar{3}\times 3}, \end{aligned} \quad (C.15)$$

where $q^T = (u, d, s)$ and $\tilde{L}_{ij}^\mu = (\Sigma \partial^\mu \Sigma^\dagger)_{ij}$. For the chirality-flipped counterparts of these operators, the irreducible components and chiral realizations can be obtained from the above results by making the exchanges $L \leftrightarrow R$ and $\Sigma \leftrightarrow \Sigma^\dagger$. We observe that among these operators $\mathcal{O}_{ddsd}^{V,LL}|_{8 \times 1}$, $\mathcal{O}_{uusd}^{V1,LL}|_{8 \times 1}$, $\mathcal{O}_{uusd}^{V8,LL}|_{8 \times 1}$, $\mathcal{O}_{uusd,udsu}^{S1,LL}|_{\bar{3} \times 3}$, and $\mathcal{O}_{uusd,udsu}^{S8,LL}|_{\bar{3} \times 3}$ generate exclusively $\Delta I = 1/2$ interactions.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.S. Hagelin and L.S. Littenberg, *Rare Kaon Decays*, *Prog. Part. Nucl. Phys.* **23** (1989) 1 [[INSPIRE](#)].
- [2] L.S. Littenberg, *The CP-violating Decay $K_L \rightarrow \pi^0 \nu \bar{\nu}$* , *Phys. Rev. D* **39** (1989) 3322 [[INSPIRE](#)].
- [3] F. Mescia and C. Smith, *Improved estimates of rare K decay matrix-elements from K_{l3} decays*, *Phys. Rev. D* **76** (2007) 034017 [[arXiv:0705.2025](#)] [[INSPIRE](#)].
- [4] T. Inami and C.S. Lim, *Effects of Superheavy Quarks and Leptons in Low-Energy Weak Processes $K_L \rightarrow \mu \bar{\mu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K^0 \leftrightarrow \bar{K}^0$* , *Prog. Theor. Phys.* **65** (1981) 297 [*Erratum ibid.* **65** (1981) 1772] [[INSPIRE](#)].
- [5] G. Buchalla and A.J. Buras, *The rare decays $K \rightarrow \pi \nu \bar{\nu}$, $B \rightarrow X \nu \bar{\nu}$ and $B \rightarrow l^+ l^-$: An Update*, *Nucl. Phys. B* **548** (1999) 309 [[hep-ph/9901288](#)] [[INSPIRE](#)].
- [6] J. Brod, M. Gorbahn and E. Stamou, *Two-Loop Electroweak Corrections for the $K \rightarrow \pi \nu \bar{\nu}$ Decays*, *Phys. Rev. D* **83** (2011) 034030 [[arXiv:1009.0947](#)] [[INSPIRE](#)].
- [7] A.J. Buras, M. Gorbahn, U. Haisch and U. Nierste, *Charm quark contribution to $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ at next-to-next-to-leading order*, *JHEP* **11** (2006) 002 [*Erratum JHEP* **11** (2012) 167] [[hep-ph/0603079](#)] [[INSPIRE](#)].
- [8] J. Brod and M. Gorbahn, *Electroweak Corrections to the Charm Quark Contribution to $K^+ \rightarrow \pi^+ \nu \bar{\nu}$* , *Phys. Rev. D* **78** (2008) 034006 [[arXiv:0805.4119](#)] [[INSPIRE](#)].
- [9] G. Isidori, F. Mescia and C. Smith, *Light-quark loops in $K \rightarrow \pi \nu \bar{\nu}$* , *Nucl. Phys. B* **718** (2005) 319 [[hep-ph/0503107](#)] [[INSPIRE](#)].
- [10] A.F. Falk, A. Lewandowski and A.A. Petrov, *Effects from the charm scale in $K^+ \rightarrow \pi^+ \nu \bar{\nu}$* , *Phys. Lett. B* **505** (2001) 107 [[hep-ph/0012099](#)] [[INSPIRE](#)].
- [11] CKMfitter Group, *CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories*, *Eur. Phys. J. C* **41** (2005) 1 [[hep-ph/0406184](#)] [[INSPIRE](#)].
- [12] CKMfitter Group, (2020) <http://ckmfitter.in2p3.fr/>.
- [13] PARTICLE DATA Group, *Review of Particle Physics*, *Phys. Rev. D* **98** (2018) 030001 [[INSPIRE](#)].
- [14] Y. Grossman and Y. Nir, *$K_L \rightarrow \pi^0 \nu \bar{\nu}$ beyond the standard model*, *Phys. Lett. B* **398** (1997) 163 [[hep-ph/9701313](#)] [[INSPIRE](#)].
- [15] KOTO collaboration, *Search for the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 X^0$ decays at the J-PARC KOTO experiment*, *Phys. Rev. Lett.* **122** (2019) 021802 [[arXiv:1810.09655](#)] [[INSPIRE](#)].

- [16] E949 collaboration, *New measurement of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ branching ratio*, *Phys. Rev. Lett.* **101** (2008) 191802 [[arXiv:0808.2459](#)] [[INSPIRE](#)].
- [17] BNL-E949 collaboration, *Study of the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ in the momentum region $140 < P_\pi < 199 \text{ MeV}/c$* , *Phys. Rev. D* **79** (2009) 092004 [[arXiv:0903.0030](#)] [[INSPIRE](#)].
- [18] NA62 collaboration, *First search for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ using the decay-in-flight technique*, *Phys. Lett. B* **791** (2019) 156 [[arXiv:1811.08508](#)] [[INSPIRE](#)].
- [19] G. Ruggiero, *New Result on $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ from the NA62 Experiment*, in proceedings of the *International Conference on Kaon Physics 2019 (KAON2019)*, Perugia, Italy, 10–13 September 2019.
- [20] S. Shinohara, *Search for the rare decay $K_L \rightarrow \pi^0 \nu \bar{\nu}$* , in proceedings of the *International Conference on Kaon Physics 2019 (KAON2019)*, Perugia, Italy, 10–13 September 2019.
- [21] K. Fuyuto, W.-S. Hou and M. Kohda, *Loophole in $K \rightarrow \pi \nu \bar{\nu}$ Search and New Weak Leptonic Forces*, *Phys. Rev. Lett.* **114** (2015) 171802 [[arXiv:1412.4397](#)] [[INSPIRE](#)].
- [22] T. Kitahara, T. Okui, G. Perez, Y. Soreq and K. Tobioka, *New physics implications of recent search for $K_L \rightarrow \pi^0 \nu \bar{\nu}$ at KOTO*, *Phys. Rev. Lett.* **124** (2020) 071801 [[arXiv:1909.11111](#)] [[INSPIRE](#)].
- [23] D. Egana-Ugrinovic, S. Homiller and P. Meade, *Light Scalars and the KOTO Anomaly*, *Phys. Rev. Lett.* **124** (2020) 191801 [[arXiv:1911.10203](#)] [[INSPIRE](#)].
- [24] P.S.B. Dev, R.N. Mohapatra and Y. Zhang, *Constraints on long-lived light scalars with flavor-changing couplings and the KOTO anomaly*, *Phys. Rev. D* **101** (2020) 075014 [[arXiv:1911.12334](#)] [[INSPIRE](#)].
- [25] J. Liu, N. McGinnis, C.E.M. Wagner and X.-P. Wang, *A light scalar explanation of $(g-2)_\mu$ and the KOTO anomaly*, *JHEP* **04** (2020) 197 [[arXiv:2001.06522](#)] [[INSPIRE](#)].
- [26] Y. Liao, H.-L. Wang, C.-Y. Yao and J. Zhang, *An imprint of a new light particle at KOTO?*, [arXiv:2005.00753](#) [[INSPIRE](#)].
- [27] Y. Jho, S.M. Lee, S.C. Park, Y. Park and P.-Y. Tseng, *Light gauge boson interpretation for $(g-2)_\mu$ and the $K_L \rightarrow \pi^0 + (\text{invisible})$ anomaly at the J-PARC KOTO experiment*, *JHEP* **04** (2020) 086 [[arXiv:2001.06572](#)] [[INSPIRE](#)].
- [28] J.M. Cline, M. Puel and T. Toma, *A little theory of everything, with heavy neutral leptons*, *JHEP* **05** (2020) 039 [[arXiv:2001.11505](#)] [[INSPIRE](#)].
- [29] M. Fabbrichesi and E. Gabrielli, *Dark-sector physics in the search for the rare decays $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$* , *Eur. Phys. J. C* **80** (2020) 532 [[arXiv:1911.03755](#)] [[INSPIRE](#)].
- [30] T. Li, X.-D. Ma and M.A. Schmidt, *Implication of $K \rightarrow \pi \nu \bar{\nu}$ for generic neutrino interactions in effective field theories*, *Phys. Rev. D* **101** (2020) 055019 [[arXiv:1912.10433](#)] [[INSPIRE](#)].
- [31] X.-G. He, X.-D. Ma, J. Tandean and G. Valencia, *Breaking the Grossman-Nir Bound in Kaon Decays*, *JHEP* **04** (2020) 057 [[arXiv:2002.05467](#)] [[INSPIRE](#)].
- [32] R. Ziegler, J. Zupan and R. Zwicky, *Three Exceptions to the Grossman-Nir Bound*, [arXiv:2005.00451](#) [[INSPIRE](#)].
- [33] X.-G. He, G. Valencia and K. Wong, *Constraints on new physics from $K \rightarrow \pi \nu \bar{\nu}$* , *Eur. Phys. J. C* **78** (2018) 472 [[arXiv:1804.07449](#)] [[INSPIRE](#)].

- [34] D. Rein and L.M. Sehgal, *Long Distance Contributions to the Decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$* , *Phys. Rev. D* **39** (1989) 3325 [INSPIRE].
- [35] G. Buchalla and G. Isidori, *The CP conserving contribution to $K_L \rightarrow \pi^0 \nu \bar{\nu}$ in the standard model*, *Phys. Lett. B* **440** (1998) 170 [hep-ph/9806501] [INSPIRE].
- [36] M. Lu and M.B. Wise, *Long distance contributions to $K^+ \rightarrow \pi^+ \nu \bar{\nu}$* , *Phys. Lett. B* **324** (1994) 461 [hep-ph/9401204] [INSPIRE].
- [37] A. Kobach, *Baryon Number, Lepton Number, and Operator Dimension in the Standard Model*, *Phys. Lett. B* **758** (2016) 455 [arXiv:1604.05726] [INSPIRE].
- [38] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142 [INSPIRE].
- [39] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark*, *Nucl. Phys. B* **250** (1985) 465 [INSPIRE].
- [40] J. Kambor, J.H. Missimer and D. Wyler, *The Chiral Loop Expansion of the Nonleptonic Weak Interactions of Mesons*, *Nucl. Phys. B* **346** (1990) 17 [INSPIRE].
- [41] M.L. Graesser, *An electroweak basis for neutrinoless double β decay*, *JHEP* **08** (2017) 099 [arXiv:1606.04549] [INSPIRE].
- [42] V. Cirigliano, W. Dekens, M. Graesser and E. Mereghetti, *Neutrinoless double beta decay and chiral SU(3)*, *Phys. Lett. B* **769** (2017) 460 [arXiv:1701.01443] [INSPIRE].
- [43] Y. Liao, X.-D. Ma and H.-L. Wang, *Effective field theory approach to lepton number violating decays $K^\pm \rightarrow \pi^\mp l^\pm l^\pm$: short-distance contribution*, *JHEP* **01** (2020) 127 [arXiv:1909.06272] [INSPIRE].
- [44] W. Buchmüller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys. B* **268** (1986) 621 [INSPIRE].
- [45] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085 [arXiv:1008.4884] [INSPIRE].
- [46] E.E. Jenkins, A.V. Manohar and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Operators and Matching*, *JHEP* **03** (2018) 016 [arXiv:1709.04486] [INSPIRE].
- [47] V. Cirigliano, G. Ecker, H. Neufeld and A. Pich, *Isospin breaking in $K \rightarrow \pi\pi$ decays*, *Eur. Phys. J. C* **33** (2004) 369 [hep-ph/0310351] [INSPIRE].
- [48] E.E. Jenkins, A.V. Manohar and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Anomalous Dimensions*, *JHEP* **01** (2018) 084 [arXiv:1711.05270] [INSPIRE].