



Evaluation of OMI operational standard NO₂ column retrievals using in situ and surface-based NO₂ observations

L. N. Lamsal^{1,2}, N. A. Krotkov², E. A. Celarier^{1,2}, W. H. Swartz^{3,2}, K. E. Pickering², E. J. Bucsela⁴, J. F. Gleason², R. V. Martin^{5,6}, S. Philip⁵, H. Irie⁷, A. Cede^{8,2}, J. Herman^{9,2}, A. Weinheimer¹⁰, J. J. Szykman¹¹, and T. N. Knepp^{12,13}

¹Goddard Earth Sciences Technology and Research, Universities Space Research Association, Columbia, MD 21046, USA

²NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

³Johns Hopkins University Applied Physics Laboratory, Laurel, MD 20723, USA

⁴SRI International, Menlo Park, CA 94025, USA

⁵Department of Physics and Atmospheric Science, Dalhousie University, Halifax, NS, B3H 3J5, Canada

⁶Also at Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

⁷Center for Environmental Remote Sensing, Chiba University, 1–33, Yayoicho, Inage-ku, Chiba 263-8522, Japan

⁸LuftBlick, Kreith, Austria

⁹University of Maryland, Joint Center for Earth Systems Technology, Baltimore, MD 21228, USA

¹⁰National Center for Atmospheric Research, Boulder, CO 80307, USA

¹¹United States Environmental Protection Agency Office of Research and Development, Hampton, VA 23681, USA

¹²Science Systems and Applications, Inc., Hampton, VA 23681, USA

¹³NASA Langley Research Center, Hampton, VA 23681, USA

Correspondence to: L. N. Lamsal (lok.lamsal@nasa.gov)

Received: 31 March 2014 – Published in Atmos. Chem. Phys. Discuss.: 5 June 2014

Revised: 12 September 2014 – Accepted: 26 September 2014 – Published: 5 November 2014

Abstract. We assess the standard operational nitrogen dioxide (NO₂) data product (OMNO2, version 2.1) retrieved from the Ozone Monitoring Instrument (OMI) onboard NASA's Aura satellite using a combination of aircraft and surface in situ measurements as well as ground-based column measurements at several locations and a bottom-up NO_x emission inventory over the continental US. Despite considerable sampling differences, NO₂ vertical column densities from OMI are modestly correlated ($r = 0.3\text{--}0.8$) with in situ measurements of tropospheric NO₂ from aircraft, ground-based observations of NO₂ columns from MAX-DOAS and Pandora instruments, in situ surface NO₂ measurements from photolytic converter instruments, and a bottom-up NO_x emission inventory. Overall, OMI retrievals tend to be lower in urban regions and higher in remote areas, but generally agree with other measurements to within $\pm 20\%$. No consistent seasonal bias is evident. Contrasting results between different data sets reveal complexities behind NO₂ validation. Since validation data sets are scarce and are limited in space and time, validation of the global product is still limited in

scope by spatial and temporal coverage and retrieval conditions. Monthly mean vertical NO₂ profile shapes from the Global Modeling Initiative (GMI) chemistry-transport model (CTM) used in the OMI retrievals are highly consistent with in situ aircraft measurements, but these measured profiles exhibit considerable day-to-day variation, affecting the retrieved daily NO₂ columns by up to 40%. This assessment of OMI tropospheric NO₂ columns, together with the comparison of OMI-retrieved and model-simulated NO₂ columns, could offer diagnostic evaluation of the model.

1 Introduction

Nitrogen oxides (NO_x = NO + NO₂) play a key role in atmospheric chemistry by controlling the production of tropospheric ozone, forming aerosol nitrates, and affecting the abundance of the hydroxyl radical (OH) and the lifetimes of greenhouse gases (Solomon et al., 1999; Intergovernmental Panel on Climate Change, 2007). Nitrogen dioxide (NO₂) is

one of the pollutants regulated by the Environmental Protection Agency (EPA), as it is detrimental to human health and ecosystems (EPA, 2009). Major sources of NO_x include combustion, soil emissions, and lighting. Tropospheric NO₂ concentrations are highly variable in space and time due to spatial heterogeneity of NO_x sources and the relatively short lifetime of NO_x in the lower troposphere.

NO₂ is measured locally by in situ monitors and detected remotely in an atmospheric column by ground-based and satellite instruments. NO₂ observations from satellite offer a globally consistent data set, albeit at coarse resolutions of 10 s to 100 s of kilometers, enabling a wide range of applications including many not feasible from in situ observations. Several studies have used satellite observations of NO₂ to evaluate chemical transport models (Martin et al., 2002; van Noije et al., 2006; Lamsal et al., 2008; Kim et al., 2009; Herron-Thorpe et al., 2010; Huijnen et al., 2010), examine spatial and temporal patterns of NO_x emissions (Beirle et al., 2003; Richter et al., 2005; Kim et al., 2006; van der A et al., 2006; Zhang et al., 2007; Boersma et al., 2008a; Lu and Streets, 2012; Wang et al., 2012; Hilboll et al., 2013; Russell et al., 2010, 2012; Duncan et al., 2013), examine NO_x sources (Jaeglé et al., 2005; van der A et al., 2008; Bucseła et al., 2010; de Wildt et al., 2012; Lin, 2012; Ghude et al., 2010, 2013a; Mebust et al., 2011; Mebust and Cohen, 2013), provide top-down constraints on surface NO_x emissions (Martin et al., 2003; Konovalov et al., 2006; Zhao and Wang, 2009; Lin et al., 2010; Lamsal et al., 2011; Ghude et al., 2013b; Vinken et al., 2014), infer NO_x lifetimes (Schaub et al., 2007; Lamsal et al., 2010; Beirle et al., 2011), and estimate surface NO₂ concentrations (Lamsal et al., 2008, 2013; Novotny et al., 2011; Bechle et al., 2013). The quality of the satellite data directly affects every one of these applications and estimates. Careful assessments of the accuracy of retrievals with credible, coincident, independent measurements help ensure reliable analyses.

Tropospheric NO₂ column retrievals from satellites have been evaluated with in situ NO₂ profile measurements from aircraft (Heland et al., 2002; Martin et al., 2006; Boersma et al., 2008a; Bucseła et al., 2008, 2013; Celarier et al., 2008; Hains et al., 2010), NO₂ column measurements from ground-based and airborne instruments (Ionov et al., 2008; Celarier et al., 2008; Brinksmā et al., 2008; Kramer et al., 2008; Irie et al., 2008, 2012; Wenig et al., 2008; Oetjen et al., 2013), in situ surface measurements (Schaub et al., 2006; Blond et al., 2007; Boersma et al., 2009; Lamsal et al., 2008, 2010), and a bottom-up NO_x emission inventory (Lamsal et al., 2010). Aircraft offer precise in situ measurements within vertical spirals covering a spatial domain over a satellite field of view, but these are generally campaign-based experiments spanning only a few days to weeks and are limited by the need to extrapolate below the lowest measurement altitude (e.g. Bucseła et al., 2008). Ground-based NO₂ column observations from the multi-axis differential optical absorption spectroscopy (MAX-DOAS) and direct-sun DOAS are ma-

tured, but assessments with these measurements are still restricted by a limited number of sites. Validation with in situ surface NO₂ measurements from dense networks of commercial molybdenum converter analyzers are complicated by instrument interference (e.g. Steinbacher et al., 2007; Lamsal et al., 2008), and is more appropriate in rural areas (Lamsal et al., 2010). Observations of NO₂ from photolytic converter analyzers (Ryerson et al., 2000) are sparse, but offer useful opportunities to evaluate satellite retrievals. In the United States, the confidence in the estimates of local and regional emissions are at medium to high levels, suggesting low uncertainty in total continental NO_x emissions (NARSTO, 2005). Validation using the US emission data benefits from a large domain coincident with satellite observations and a variety of observational conditions. This study takes advantage of state-of-the-art NO₂ measurement techniques, and exploits the strength of various measurements to assess the quality of the new standard tropospheric NO₂ retrievals (OMNO2, version 2.1) from the Ozone Monitoring Instrument (OMI) under various atmospheric conditions.

Well-validated daily global observations from satellite provide a rich resource to evaluate results from regional air quality (AQ) models and global chemical transport models (CTMs), thereby helping to increase model accuracy. To facilitate satellite-model comparison, the OMNO2 product provides information on vertical NO₂ measurement sensitivity (scattering weights). Combining scattering weights with model-derived vertical NO₂ profile shape allows for the calculation of new air mass factors (AMFs) needed to convert satellite-retrieved slant column densities (SCDs) to vertical column densities (VCDs). Since the assumed vertical distribution of NO₂ in the retrieval is taken from the model subject to evaluation, this approach allows consistent comparison of satellite-observed columns with model-simulated columns. Applying a similar approach for in situ NO₂ measurements from aircraft provides insights into the retrieval uncertainty, as using measured profiles and resulting AMFs indicate how much the satellite retrieval would change when climatological assumptions about profile shape are replaced with specific, observed profile information.

Our main goals here are to assess the operational OMI NO₂ standard product, elucidate errors in retrieved columns due to a priori NO₂ vertical profiles through the use of nearly-coincident NO₂ profiles measured from aircraft, and devise objective methods to compare model-simulated NO₂ columns with satellite retrievals. Section 2 describes the OMI retrievals and various concurrent data sources used in this study. We present validation results in Sect. 3. The impacts of the a priori NO₂ profiles used in the satellite retrievals are discussed in Sect. 4. We discuss the comparison of modeled and OMI NO₂ in Sect. 5. Section 6 summarizes the conclusions of this study.

2 Observations

2.1 OMI Retrieval

The Dutch–Finnish OMI instrument aboard the NASA EOS-Aura satellite provides continuous monitoring of atmospheric NO₂ columns through measurement of hyperspectral solar backscatter in the UV-visible range from 264 to 504 nm (Levelt et al., 2006). The satellite was launched on 15 July 2004, into a polar, sun-synchronous orbit with an equator-crossing time of 13:45 LT (ascending node). OMI observes the atmosphere in 60 cross-track ground pixels measuring 13–26 km along track and 24–128 km across track, achieving daily global coverage.

We use the tropospheric NO₂ columns from OMI standard product (Bucsela et al., 2013) publicly available from the NASA archive: http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2_v003.shtml. The algorithm uses the differential optical absorption spectroscopy (DOAS) technique (Platt, 1994) to determine NO₂ SCD by nonlinear least squares fitting of reference spectra for NO₂, ozone, H₂O and the Ring filling-in effect to the OMI-measured reflectance in the 405–465 nm spectral window (Bucsela et al., 2006; Boersma et al., 2007). The slant column represents the integrated NO₂ abundance along the average light path from the sun, through the atmosphere, to the satellite. The measured SCDs are corrected for instrumental artifacts (stripes Dobber et al., 2008; Bucsela et al., 2013) accounting for cross-track variation of the stratospheric AMF. The AMF, defined as the ratio of the SCD to the VCD, is calculated using a look-up table of vertically resolved NO₂ sensitivities (scattering weights) and various input parameters including viewing geometry, surface reflectivity, effective cloud pressure, cloud radiance fraction, and a priori NO₂ vertical profile shapes (Palmer et al., 2001). The a priori NO₂ profiles are early afternoon (at the OMI overpass time) monthly mean values derived from the Global Modeling Initiative (GMI at 2° × 2.5° CTM (Appendix A) (Strahan et al., 2007). To separate stratospheric and tropospheric columns, the algorithm first applies stratospheric (close to geometric) AMFs to the de-striped measured SCDs to yield initial VCDs. Cloud-free areas of tropospheric contamination in the stratospheric NO₂ field are identified using the a priori GMI monthly mean tropospheric NO₂ columns and OMI cloud measurements. Those regions are then masked and filled in with the stratospheric VCDs measured outside the masked regions, primarily from unpolluted or cloudy areas. The stratospheric field is further smoothed by using a boxcar averaging.

The OMNO2 retrievals used here, version 2.1 (Bucsela et al., 2013), represent a significant advance over previous version 1.0 (Bucsela et al., 2006; Celarier et al., 2008). The main changes include the use of monthly, rather than annual, mean a priori NO₂ profiles, and improvements in the estimates of stratospheric NO₂ columns, correction of calibra-

tion artifacts (de-stripping), and the calculation of scattering weights.

The uncertainties in the retrieval of tropospheric NO₂ columns arise from the uncertainties in the retrieval of slant column densities, the calculation of AMFs, and the separation of stratospheric and troposphere components. The uncertainty in the individual OMI NO₂ slant column is $\sim 0.75 \times 10^{15}$ molec. cm⁻² (Boersma et al., 2004, 2011; Bucsela et al., 2013) and dominates the overall retrieval error over the oceans and remote areas. AMF uncertainties are $\sim 20\%$ in clear-sky and 30–80% under cloudy conditions and dominate overall retrieval errors over continental polluted regions. In this study, we include the data for scenes with cloud radiance fractions less than 0.5 and those unaffected by the OMI row anomaly (Dobber et al., 2008). We use data from all cross-track positions.


2.2 In situ NO₂ measurements from aircraft

In situ NO₂ concentrations were measured from the NASA P-3B aircraft in the Baltimore-Washington, D.C. metropolitan region on 14 flight days in July 2011, as part of the NASA Earth Venture-1 DISCOVER-AQ (Deriving Information on Surface Conditions from Column and Vertically Resolved Observations Relevant to Air Quality, <http://www-air.larc.nasa.gov/missions/discover-aq/discover-aq.html>) field program. Measurements usually began between 7:00 and 10:00 local time and continued for about 8 h. Flights occurred over a range of weather conditions including clean days, pollution episodes, and weekdays and weekends. The P-3B aircraft housed two well-characterized in situ NO₂ measuring instruments: The University of California, Berkeley thermal dissociation laser induced fluorescence (TD-LIF, (Thornton et al., 2000; Wagner et al., 2011) and the National Center for Atmospheric Research (NCAR) 4-channel chemiluminescence instrument (P-CL). The P-CL measures NO₂ by photolysis of NO₂ and chemiluminescence detection of the product NO (Ridley and Grahek, 1990; Ridley et al., 2004). The TD-LIF instrument had a low NO₂ sampling frequency due to an alternating measurement cycle for other species such as peroxy nitrates, alkyl nitrates, and nitric acid, so we use measurements from the NCAR P-CL. The instrument has an NO₂ measurement uncertainty of 10% and a 1 s, 2σ detection limit of 50 ppt, making it useful to measure NO₂ in the free troposphere.

Figure 1 shows a typical in situ NO₂ measurement pattern during DISCOVER-AQ. Flight tracks for this campaign targeted urban air pollution spatially along the Interstate 95 (I-95) corridor in the Baltimore-Washington, D.C. region and vertically over the Chesapeake Bay and six surface air quality monitoring sites (see Table 1). Typically, during each sortie, three vertical spirals were flown over each location, covering altitudes from ~ 300 m, in the boundary layer to ~ 3.3 km, in the free troposphere. Table 1 provides the details on the number of spirals and observations and the measured altitude

Table 1. Aircraft flight parameters.

Site	Location type	Spirals	Minimum and maximum altitude (km)	Number of observations
Beltsville	suburban	43	0.24–2.03	19 867
Padonia	suburban	38	0.35–3.47	27 106
Fair Hill	rural	41	0.26–4.78	41 550
Aldino	rural/suburban	36	0.27–4.82	30 407
Edgewood	coastal/suburban	43	0.25–4.82	35 050
Essex	coastal/urban	38	0.24–3.26	30 269

**Figure 1.** Distribution of NO₂ on 21 July 2011 obtained from P-3B aircraft measurements during the DISCOVER-AQ field campaign in Maryland. More than 190 000 1 s NO₂ measurements were taken during 254 spirals over the entire campaign period.

range. There were a total of 13–19 P-3B spirals over each surface site with 5356–15 827 1 s observations made near the time of the OMI overpass. We found that the limited vertical extent of the aircraft pass over I-95 and the Chesapeake rendered those measurements less useful. We binned the measurements to the pressure grid of the GMI model to directly compare the model profiles with observed profiles, and to estimate the retrieval error due to the difference.

Figures 1 and 2 show the early afternoon (12:00–15:00) NO₂ vertical profiles measured during DISCOVER-AQ. NO₂ mixing ratios over land range over 0.02–28 ppb below 950 hPa, decrease sharply to 0.01–2 ppb at ~ 800 hPa, and are 10–200 ppt above 700 hPa. Over the Chesapeake Bay, NO₂ mixing ratios are generally less than 1 ppb, and the vertical gradient in the profile is less pronounced due to limited surface sources and transported NO₂ downwind. Large spatial and temporal variability in near-surface NO₂ reflect the large spatiotemporal variation in NO_x emissions and differences in local dynamics. NO₂ enhancement and variability over Beltsville and Essex are largely due to local emissions, mostly from traffic. Most sites experienced more than a factor of 2 greater NO₂ concentrations on highly polluted days with a shallow mixed layer on 5, 10, 21, and 28 July.

The aircraft measurements show that NO₂ concentrations within the mixed layer make the largest contributions to tropospheric NO₂ columns. The lowest 1 km of sampled aircraft data contain 64–84 % of the NO₂ column below 5 km. The same altitude range in the GMI profile represents 72–83 %, providing confidence in the GMI simulation. In the free troposphere (2–5 km), NO₂ concentrations from the a priori GMI climatology and aircraft measurements generally agree to within 0.03 ppb. GMI simulations suggest that the NO₂ partial column within first few hundred meters from the ground to the lowest aircraft altitude comprise 30–40 % of the total column. The upper tropospheric column above 5 km is rather small, consisting of 10–15 % of the total column. We inferred the complete P-3B tropospheric NO₂ column by combining the measured values with GMI climatology above the highest aircraft level and extrapolating below the lowest aircraft level. The extrapolation scheme applies the vertical gradient of the NO₂ concentrations between the lowest aircraft altitude (C_M^j) and underneath (C_M^{j-1}) in the GMI profiles to the measured concentration (C^j) to estimate concentration (C^{j-1}):

$$C^{j-1} = \frac{C_M^{j-1}}{C_M^j} \times C^j, \quad (1)$$

where the subscript “M” represents model. In this approach, we assume that the GMI model captures the vertical distribution of NO₂ well.

We first evaluated the extrapolation scheme by comparing the estimated surface NO₂ mixing ratios with NO₂ measurements from a photolytic converter instrument at Padonia. Since NO₂ measurements at the lowest aircraft altitude are on average 45 % lower than the measurements at the ground, extrapolation of aircraft profiles by assuming a constant mixing ratio from the value at the lowest aircraft level will substantially underestimate the true NO₂ near the surface. In Fig. 3, we show a comparison of our estimates using Eq. (1) with surface measurements at Padonia. The extrapolated and measured values are well correlated ($r = 0.64$, $N = 14$), and generally compare well (mean bias = 23 %), although extrapolation could at times overestimate observations when the aircraft encountered elevated plumes with high NO₂ concentrations. Errors in the calculated gradient propagate into the




Figure 2. Early afternoon (12:00 to 15:00) vertical mean profile of NO₂ mixing ratio over Fair Hill, Aldino, Padonia, Beltsville, Edgewood, and Essex in Maryland. The open circles (in orange) represent NO₂ mixing ratios averaged over the GMI pressure grid from each spiral. Error bars represent the 10th to 90th percentiles. Solid black lines connect the mean mixing ratios determined from in situ measurements during the entire campaign. The number of measurements within each GMI pressure grid is shown in the right of each panel. The dotted lines show the surface pressure levels. The bottom-right panel shows the GMI a priori monthly (July) mean NO₂ mixing ratio profile over the DISCOVER-AQ domain.

extrapolated value, degrading the quality of integrated P-3B tropospheric NO₂ columns. Allowing factor of 2 extrapolation errors, the errors in the integrated P-3B tropospheric NO₂ columns are generally less than 20 %.

2.3 Ground-based MAX-DOAS

Tropospheric NO₂ columns were measured by the ground-based MAX-DOAS instruments for several months during 2006–2011 at a remote site in Hedo and a suburban site in Tsukuba, Japan. NO₂ observations at these sites allow us to assess the OMI retrievals for contrasting environments (rural vs. urban).

The MAX-DOAS instrument measures scattered sunlight observations in the UV/visible wavelengths at several elevation angles between the horizon and zenith (e.g. Hönninger

et al., 2004; Irie et al., 2012). Spectral fitting of the MAX-DOAS measured differential structure with absorption cross-section of NO₂ from Vandaele et al. (1998) at 294 K and other interfering species including O₂–O₂, O₃, H₂O, and the Ring and undersampling effects over the 460–490 nm window yields the differential slant column density, i.e., the difference in integrated columns along the average light path between measurements made at low elevation angles and that at an elevation angle of 90°. The accuracy of the retrieved NO₂ slant columns is ~ 10 %, as confirmed by a formal semi-blind intercomparison experiment involving MAX-DOAS observations from different research groups (Roscoe et al., 2010). The NO₂ slant column densities are converted to tropospheric vertical column density by using the AMF calculated with measured aerosol information and the vertical




Figure 3. Comparison of aircraft-measured NO₂ profiles extrapolated to the surface (using Eq. 1) with surface NO₂ measurements with photolytic converter instrument at Padonia during the DISCOVER-AQ field campaign. The dotted line represents the 1 : 1 relationship.

profile of NO₂ and a non-linear iterative inversion scheme (Irie et al., 2012). Additional details on the MAX-DOAS measurements, calibration, and retrieval procedures can be found in Irie et al. (2008) and references therein. Overall errors in the tropospheric NO₂ vertical columns are < 14 %.

The MAX-DOAS instrument observes air masses representative of horizontal distance of about 10 km (Irie et al., 2012), comparable to the OMI spatial resolution. The temporal resolution corresponds to a complete sequence of elevation angles lasting for 30 min. We use the MAX-DOAS measurements taken within 30 min of OMI overpasses to compare with the OMI retrievals.

2.4 Ground-based Pandora

The direct sun total NO₂ column measurements were carried out at 12 DISCOVER-AQ sites (including six aircraft-spiral locations) in Maryland and at the Chemistry and Physics Atmospheric Boundary Layer Experiment (CAPABLE) site at NASA's Langley Research Center in Hampton, Virginia. The CAPABLE site is located in a coastal suburban area, which could experience sporadic local and transported NO_x emissions. Additional details on the CAPABLE site can be found in Knepp et al. (2013). These measurements are useful to examine spatial and temporal variation in the OMI retrievals.

Pandora is a ground-based spectrometer that measures direct solar irradiance over the range 280–525 nm at the spectral resolution of 0.6 nm, allowing the retrieval of the total column abundance of various species, such as O₃, NO₂, HCHO, H₂O, and SO₂ (Herman et al., 2009). An algorithm for the retrieval of NO₂ from Pandora is similar to the direct-sun NO₂ inversion method from a Brewer spectrometer (Cede et al., 2006). The direct-sun DOAS technique is

equally sensitive to stratospheric and tropospheric NO₂, and is not affected by the ring effect. The algorithm initially retrieves the relative NO₂ slant columns by least-square fitting of the difference between the logarithm of a reference irradiance spectrum and the logarithm of measured irradiance spectra with the absorption spectra of NO₂ (Vandaele et al., 1998) at 254.5 K and other atmospheric absorbers, a low-order polynomial, and wavelength shift and squeeze functions in the spectral range 370–500 nm. The temperature dependence of the NO₂ cross section is not accounted for in the fitting process. The differential NO₂ slant columns represent the difference between the absolute slant columns in the measured and the reference spectrum used for normalization. The reference spectrum is an average spectrum measured on clear clean days. The absolute slant column in the reference spectrum is determined by the minimum-amount Langley-extrapolation method, as described in Herman et al. (2009). The direct sun AMF can be approximated as the secant of solar zenith angle and therefore does not require radiative transfer calculations or prior knowledge of the ground reflectivity or NO₂ profile shape.

The Pandora spectrometer provides NO₂ vertical column observations with a clear-sky precision of about 2.7×10^{14} molec. cm⁻² and a absolute accuracy of 2.7×10^{15} molec. cm⁻². NO₂ column retrievals from Pandora have been previously validated against direct-sun multifunction DOAS (MFDOAS) and Fourier transform ultraviolet spectrometer (UVFTS) data and have been found to agree to within 12 % (Piters et al., 2012; Wang et al., 2010; Herman et al., 2009). Here, we compute 30 min Pandora column averages close to the OMI overpass time to compare with the nearest OMI NO₂ columns representing individual field of view (FOV). The maximum allowed collocation radius (distance between the center of the OMI FOV and the Pandora site) is 10 km.

2.5 In situ surface measurements

In situ measurements of surface NO₂ were made at the Southeastern Aerosol Research and Characterization (SEARCH) network, consisting of 7 sites in the Southeastern United States (Edgerton et al., 2006). We use data from two regionally representative sites: Centreville, in Alabama, and Yorkville, in Georgia measured during 2006–2009. Other sites that are either urban/suburban or located in close proximity to urban facilities were not found to be suitable for validation of satellite retrievals. NO₂ measurements are made using photolytic converter analyzers, a measurement method that employs photolysis of ambient NO₂ followed by chemiluminescence detection of the product NO. This method offers highly accurate NO₂ measurements, with an uncertainty < 10 %.




Figure 4. Relative vertical distribution (profile shape) of NO₂ over the six spiral locations during DISCOVER-AQ. The colored lines show the mean profile shapes determined from the in situ measurements. The shape factors are calculated as the ratio of partial columns to total tropospheric column. The dashed line with filled circles shows the profile shape calculated from the GMI model.

3 Evaluation of the OMI Retrieval

3.1 NO₂ profile shapes

We initially evaluate the a priori monthly mean relative vertical distribution (shape factor) of NO₂ used in the OMI NO₂ retrievals with aircraft measurements during the DISCOVER-AQ field campaign. Figure 4 compares average NO₂ shape factors over various locations from aircraft with those calculated with the GMI model. Although the aircraft measurements are qualitatively similar to the model results, differences up to 30% were observed near the surface and in the free-troposphere. The GMI model suggests that 20–30% of the tropospheric NO₂ column is located near the surface (first model layer, ~1000 hPa), while only 5–10% is in the mixed layers between 900–1000 hPa, and less than 3% is in the free-troposphere (<900 hPa). Aircraft measurements indicate the horizontal spatial gradient in the free-tropospheric shape factors, primarily due to the dominant lower tropospheric contributions to the total tropospheric NO₂ columns in urban source regions. These measurements also reveal considerable day-to-day variation in NO₂ profile shapes within a given month, suggesting that the use of a monthly mean profile in the operational algorithm is potentially a significant source of error in individual retrieved tropospheric NO₂ columns.

3.2 Tropospheric NO₂ columns

3.2.1 Comparison with in situ aircraft measurements

In this section, we compare OMI tropospheric NO₂ columns with integrated columns from aircraft spirals at six locations in Maryland during the DISCOVER-AQ field campaign in July 2011. We select only the spirals made within 1 h of the OMI overpass. Not all data from the 14 flight days could be used due to adverse instrumental (row anomaly) or cloudy conditions affecting the OMI data.

Figure 5 shows tropospheric NO₂ columns from OMI and vertically integrated in situ aircraft measurements for several individual flight days. Individual measurements agree to within 20% in 60% of cases at Fair Hill, Aldino, Padonia, and Beltsville. A more substantial difference was observed at Edgewood and Essex, where aircraft measurements were systematically higher than OMI retrievals. These two coastal towns were often impacted by a bay breeze, yielding complex vertical and horizontal distributions of NO₂. Figure 6 shows a summary of comparisons at all DISCOVER-AQ sites. Although OMI and the in situ tropospheric NO₂ columns are highly correlated at some sites, the overall correlation at all sites is rather poor ($r = 0.2$, $N = 59$). The observed discrepancy between the two measurements is primarily due to the difference in spatial sampling, but it could also be due to other reasons, such as errors in OMI tropospheric NO₂ due to inaccurate removal of stratospheric NO₂ on 2 July and partly cloudy conditions obstructing the scene on 20 July.

Figure 7 shows the campaign average tropospheric NO₂ columns observed by the OMI and aircraft instruments. Measurements from both instruments exhibit a distinct spatial variation, with low columns at the rural site Fair Hill and high columns in urban sites such as Beltsville and Essex. NO₂ retrievals from OMI are lower than aircraft measurements by 5.8–22.1%, with the exception of Edgewood and Essex, where aircraft measurements are often up to a factor of 2 higher than OMI retrievals. We quantify the impact of the a priori NO₂ profiles in the OMI retrievals for the observed discrepancy between OMI and in situ measurements in Sect. 4.

3.2.2 Comparison with pandora measurements

We compare OMI total NO₂ columns (sum of tropospheric and stratospheric columns) with Pandora direct sun NO₂ column retrievals at six sites in Maryland during the first DISCOVER-AQ field campaign in July 2011 and at the CAPABLE site at NASA Langley in Hampton, Virginia for 2010–2012. Although analysis of Pandora measurements allows inference of the stratospheric portion of the total NO₂ column (Herman et al., 2009), the separate stratospheric and tropospheric components are not currently available from Pandora. Subtraction of OMI-derived stratospheric




Figure 5. Comparison of tropospheric NO₂ columns retrieved from OMI (squares) with those determined from in situ measurements (filled circles). The figure shows OMI retrievals performed using GMI NO₂ a priori vertical profiles (open squares) and in situ NO₂ measurements (filled squares). Error bars represent errors in the aircraft measurements, extrapolated aircraft profiles, and OMI retrievals.




Figure 6. A summary plot showing comparison of collocated tropospheric (black) and total (orange) NO₂ columns derived from in situ aircraft measurements and retrieved from Pandora with OMI retrievals performed using GMI NO₂ a priori vertical profiles (filled circles) and in situ NO₂ measurements (open circles). The dotted line represents the 1 : 1 relationship.

NO₂ columns from Pandora total column measurements could as well introduce errors in Pandora-derived tropospheric NO₂ columns. Therefore the use of total columns allows us to reduce these errors, and allows more direct comparison between the two measurements.




Figure 7. Comparison of average tropospheric (orange bars) and total (blue squares) NO₂ columns retrieved from OMI with tropospheric NO₂ columns determined from in situ aircraft (black bars) measurements and total columns retrieved from Pandora (filled blue circles) at the six locations in Maryland during the DISCOVER-AQ field campaign. Open bars and squares represent the operational retrievals and filled bars and squares represent the retrievals performed using collocated aircraft-measured NO₂ vertical profiles. The vertical lines represent the standard deviation of the average.

Figures 6 and 8 present a comparison of coincident total NO₂ column retrievals from the OMI and Pandora instruments. The variations of OMI NO₂ are broadly consistent with the Pandora measurements. Although the OMI and Pandora NO₂ columns are fairly correlated ($r = 0.25$, $N = 52$), they generally agree to within 18% at Aldino and Beltsville and within 30% at the other DISCOVER-AQ sites. Occasional large discrepancies are evident, reflecting a




Figure 8. Comparison of total NO₂ columns retrieved from OMI (rectangles) with those retrieved from Pandora measurements (filled circles). Open squares represent the operational retrievals (using GMI profiles), and filled squares represent OMI retrievals performed using collocated aircraft-measured NO₂ vertical profiles. Error bars represent errors in Pandora and OMI retrievals.

combination of enhanced spatial variation and placement of the Pandora spectrometers.

Figure 7 shows campaign average total NO₂ columns measured by Pandora and OMI at six DISCOVER-AQ sites in Maryland. The measurements are in good agreement. NO₂ columns measured with the Pandora are on average < 6% higher at Aldino, Beltsville, and Edgewood, and 9–13% lower than OMI at Padonia and Essex. Inconsistent results at Fair Hill, with a high bias in the OMI retrievals (44%) vs. Pandora and a low bias (6.7%) vs. aircraft measurements, suggest differences in sampling area by the three independent measurement systems.

We also compare long-term observations of the total NO₂ columns by the OMI and Pandora instruments at the CAPABLE site. Figure 9 shows the multi-year monthly mean variation of OMI and Pandora NO₂ columns. NO₂ retrievals from the two instruments are moderately correlated ($r = 0.5$, $N = 163$), with the largest correlation ($r = 0.71$, $N = 40$) in winter and smallest correlation ($r = 0.25$, $N = 33$) in spring. However, the magnitude of the seasonal cycle differs for the two measurements, and they are not in phase. The seasonal variation in Pandora NO₂ columns exhibits a summer maximum and fall minimum, in contrast to the winter maximum and summer minimum in OMI total columns. The monthly mean biases range from -2.8% in January to -28.4% in June (Pandora being higher). The seasonal cycle in tropospheric and stratospheric NO₂ columns retrieved from OMI and simulated from GMI are highly consistent (not shown), providing confidence in the seasonal variation in the OMI retrievals. Several factors could contribute to the observed seasonal biases between the OMI and Pandora retrievals. Due to




Figure 9. Monthly variation of total NO₂ columns at Hampton, VA for 2009–2012, as calculated from Pandora measurements (line with open circles) and OMI measurements (bars). OMI total NO₂ columns are separated into stratospheric (green bars) and tropospheric (orange bars) components. The bars represent the standard deviation of the average.

the close proximity to local traffic at Langley Air Force Base, and the Yorktown power plant, Pandora measurements are influenced by local NO_x emission sources and could exhibit a dampened seasonal tropospheric NO₂ cycle. Also, unlike the OMI retrievals, the Pandora retrievals are based on the NO₂ cross-section at a constant temperature of 255 K (representing the stratosphere and troposphere), which could affect seasonal variation in the retrieved NO₂ columns. However,

the effect of the temperature error in Pandora data is small ($\sim 3.3\%$ per 10° change in NO₂ temperature) and is unlikely to explain the observed seasonal differences. Errors in absolute calibration could lead to $\pm 2.7 \times 10^{15}$ molec. cm⁻² uncertainty in Pandora NO₂ slant columns, with a similar uncertainty in vertical columns in summer, but only half of that in winter, favoring wintertime data, which are in excellent agreement with the OMI retrievals.

3.2.3 Comparison with MAX-DOAS measurements

Tropospheric NO₂ column retrievals from OMI are compared with long-term MAX-DOAS measurements at two Japanese sites, Tsukuba and Hedo, for the period 2006–2011. Figure 10 (left) compares tropospheric NO₂ columns retrieved from OMI and MAX-DOAS instruments. Tropospheric NO₂ columns over Hedo range over $0.2\text{--}3.2 \times 10^{15}$ molec. cm⁻² for MAX-DOAS and $-0.5\text{--}2.8 \times 10^{15}$ molec. cm⁻² for OMI. The stratosphere–troposphere separation scheme in the OMI retrievals could yield slightly negative tropospheric columns in remote areas when measured slant columns are lower than estimated stratospheric slant columns. NO₂ columns over Tsukuba are much higher, reaching 40×10^{15} molec. cm⁻² in both the MAX-DOAS and OMI data. Measurements from the two techniques exhibit a significant spatiotemporal correlation ($r = 0.86$, $N = 626$). The mean relative difference between OMI and MAX-DOAS measurements is -16.3% in Tsukuba and 7.1% in Hedo.

Figure 10 (right) presents the seasonal mean tropospheric NO₂ column from MAX-DOAS measurements and those retrieved from OMI. The seasonal variation of the OMI-retrieved NO₂ columns is consistent with the MAX-DOAS measurements. The seasonal mean NO₂ columns for the MAX-DOAS measurements decrease by a factor of 1.6–1.9 from winter to summer, compared with a factor of 1.4–1.5 for OMI. The relative difference between OMI and MAX-DOAS seasonal mean tropospheric NO₂ columns range from 0.5% in fall to -20.8% in winter at Tsukuba and from -21.3% in winter to 24.8% in spring at Hedo. These results are generally consistent with the comparisons made with aircraft and Pandora observations.

3.2.4 Comparison with in situ surface measurements

We conduct an indirect validation of cloud-free (cloud radiance fraction < 0.5) OMI tropospheric NO₂ columns by comparison with coincident hourly in situ surface NO₂ measurements. This approach requires estimating ground-level NO₂ concentrations from OMI. We follow the method of Lamsal et al. (2008) with improvements as described in Lamsal et al. (2013) that combines coincidentally sampled NO₂ vertical profile taken from a GEOS-Chem nested simulation (see Appendix B) with the OMI observations containing information about the spatial variation of the tropospheric NO₂ columns in the boundary layer. The OMI-derived surface

NO₂ represents the mean mixing ratio in the lowest vertical layer (~ 50 m) of the model.

We compare the OMI-derived surface NO₂ mixing ratios with the in situ measurements at the two rural surface sites, in Yorkville and Centerville for 2006–2010. Figure 11 displays the seasonal average surface NO₂ mixing ratios from the in situ measurements and those derived from the OMI retrievals. The OMI-derived surface NO₂ concentrations are well correlated with the photolytic converter measurements ($r = 0.61$, $N = 700$ for Yorkville and $r = 0.69$, $N = 676$ for Centerville) and exhibit similar seasonal variation with summertime minima. The OMI-derived surface NO₂ are lower than the in situ measurements at Centerville by 11.8% in fall but higher by 4.1% in spring. Discrepancies are larger at Yorkville, where the OMI-derived surface NO₂ overestimates in situ measurements by 8.2% in spring and underestimates by $25\text{--}31\%$ in other seasons.

3.2.5 Comparison with bottom-up emissions

We use an inventory of US NO_x emissions to indirectly validate OMI tropospheric NO₂ columns. We employ the emissions for 2005 as implemented in the GEOS-Chem model (Appendix B). Emissions employed by the GMI (Appendix A) simulation used in the operational retrieval included outdated North American NO_x emissions not suitable for validation. In GEOS-Chem, the bottom-up emissions over the US comprise over 75% of NO_x emissions from anthropogenic activities; the remainder comes from soil, lightning, and biomass burning emissions. In contrast to inventories in developing countries, the US national emission inventory is more complete, accurate, and transparent (NARSTO, 2005), and is expected to be less uncertain ($< 25\%$, Christian Hogrefe, personal communication, 2008) at least in national totals. The largest contributors to the US NO_x emissions include on- and off-road vehicles ($\sim 62\%$) and electricity and industrial power generation ($\sim 27\%$), which exhibit little seasonal variation (EPA, 2009; Lamsal et al., 2010), a characteristic that is useful to assess seasonal variation in OMI retrievals. Difficulty could arise for comparisons focused on county or sectoral levels, where uncertainty in bottom-up emissions could be significant, and in spring and summer, when emissions from soils and biomass burning are at peak levels. To compare the OMI retrievals with NO_x emissions, we follow a simple mass balance approach (Martin et al., 2003; Lamsal et al., 2010), which directly relates OMI tropospheric NO₂ columns (Ω) to surface NO_x emissions (E):

$$E = \frac{E_M}{\Omega_M} \times \Omega. \quad (2)$$

Here, Ω_M is the tropospheric NO₂ column from a GEOS-Chem nested simulation based on the a priori surface NO_x emissions E_M , both sampled at the OMI overpass time. To account for the impact of spatial smearing (Palmer et al., 2003), we considered an approach that accounts for the




Figure 10. Comparison of tropospheric NO₂ columns retrieved from OMI and MAX-DOAS instruments. Observations at Tsukuba and Hedo, Japan, are shown in red and blue, respectively. (left) Scatter plot of OMI tropospheric NO₂ and MAX-DOAS measurements. The regression analysis parameters are given in the legend. The slope was calculated with reduced major-axis linear regression (Hirsch and Gilroy, 1984). The dotted line represents the 1 : 1 relationship and the solid lines represent the 30 % deviation range. (right) Seasonal mean tropospheric NO₂ columns for December–February (DJF), March–May (MAM), June–August (JJA), and September–November (SON) for 2006–2011 from MAX-DOAS (open bars) and OMI (filled bars). The vertical lines are the standard deviation of the seasonal average.




Figure 11. Seasonal variation of NO₂ mixing ratios at rural SEARCH sites for 2006–2010. Open bars represent seasonal mean NO₂ mixing ratios from in situ measurements, and solid bars represent those derived from the OMI tropospheric NO₂ columns. Error bars in in situ measurements represent 10 % errors in the photolytic converter measurements. Error bars in the OMI-derived surface NO₂ represent errors in retrievals including errors in the GEOS-Chem NO₂ profiles.

emissions from eight adjacent model grid cells to estimate surface NO_x emissions ($E_{i,j}$) at grid cell (i, j) from OMI (Toenges-Schüller et al., 2006; Boersma et al., 2008a; Lamsal et al., 2010) with improvements as discussed in Tang et al. (2013):

$$E_{i,j} = \frac{E'_{M_{i,j}}}{\sum_{n=-1}^1 \sum_{m=-1}^1 K_{i,j} E'_{M_{i+m,j+n}}} \times \frac{E_{M_{i,j}}}{\Omega_{M_{i,j}}} \times \Omega_{i,j}. \quad (3)$$

The smoothing kernel (K) is defined as $\frac{1}{p+8} \begin{bmatrix} 1 & 1 & 1 \\ 1 & p & 1 \\ 1 & 1 & 1 \end{bmatrix}$, where p is the smoothing parameter. To determine the value

of p , we applied K to each grid cell in the bottom-up NO_x emission inventory with different p values, and computed the correlation between smoothed 24 h averaged bottom-up NO_x emissions ($E'_{i,j}$) and corresponding modeled tropospheric NO₂ columns. The maximum correlation coefficient corresponding to the optimal value of p was achieved at $p = 12$ as in Boersma et al. (2008a), which we adopt to infer monthly top-down surface NO_x emissions from OMI.

Figure 12 shows the spatial variation of bottom-up and OMI-based top-down NO_x inventories of land surface emissions. Both top-down and bottom-up inventories exhibit similarity in their spatial patterns, with large emissions in major urban centers, reflecting industrialization, dense traffic and population. The top-down and bottom-up annual surface NO_x emissions are strongly correlated ($r = 0.95$, $N = 2706$). The difference between the OMI-derived and bottom-up annual surface NO_x emissions integrated over the continental US is 8.8 %, much lower than the uncertainty in the bottom-up inventory and in the daily OMI retrievals. Excluding the smoothing parameter in the inversion, the difference decreases to 3.5 %. Despite excellent agreement in the total surface NO_x emissions, we observe a pronounced difference of up to a factor of 2 in the magnitude of local and regional NO_x emissions. These differences could arise from errors in the bottom-up emissions, in the OMI retrievals, and from the simple inversion scheme.

Figure 12 (bottom right) shows the ratio of the seasonal area-integrated OMI-derived and bottom-up NO_x emissions over the US. The ratio ranges from 0.91 in July to 1.35 in April. These results suggest consistency between bottom-up emissions and OMI retrievals within the range of their uncertainties.




Figure 12. (top) Annual mean surface NO_x emissions over the United States for 2005. The left panel contains bottom-up emissions from fossil fuels, bio-fuels, biomass burning, and soils. The right panel shows top-down emissions estimated using OMI tropospheric NO₂ columns. The bottom left panel shows the difference between top-down and bottom-up surface NO_x emissions. (bottom right) Monthly mean ratio of area-averaged top-down surface NO_x emissions to bottom-up emissions over the United States.

Table 2. Summary of validation results.

Location	Data sources	Measurement period	Mean difference	Sample size
Beltsville, MD	P-3B	Jul 2011	−6.0 %	8
	Pandora		−5.9 %	
Padonia, MD	P-3B	Jul 2011	−8.0 %	9
	Pandora		9.1 %	
Fair Hill, MD	P-3B	Jul 2011	−22.1 %	8
	Pandora		43.9 %	
Aldino, MD	P-3B	Jul 2011	−19.5 %	8
	Pandora		−5.4 %	
Edgewood, MD	P-3B	Jul 2011	−41.3 %	10
	Pandora		−5.8 %	
Essex, MD	P-3B	Jul 2011	−40.1 %	13
	Pandora		13.1 %	
Hampton, VA	Pandora	2009–2011	−16.8 %	163
Tsukuba, Japan	MAX-DOAS	2006–2007, 2010–2011	−16.3 %	191
Hedo, Japan	MAX-DOAS	2007–2011	7.1 %	514
Yorkville, GA	In situ surface	2006–2009	−1.9 %	700
Centerville, AL	In situ surface	2006–2009	−17.8 %	676
Continental USA	NO _x emission inventory	2005	8.8 %	2706

3.2.6 Synthesis of validation results

Direct validation results of OMI NO₂ retrievals vs. in situ aircraft, MAX-DOAS, and ground direct sun Pandora measurements, and indirect validation results of OMI-derived surface NO₂ vs. in situ surface measurements and top-down

vs. bottom-up emission inventories suggest the scientifically useful quality of the archived tropospheric NO₂ product from the standard OMI operational algorithm. Table 2 contains a summary of these validation results. OMI tropospheric NO₂ data generally correlate well ($r > 0.5$), agree to within $\pm 20\%$ with biases tending to be more negative than positive,




Figure 13. Seasonal mean tropospheric NO₂ columns binned at $0.5^\circ \times 0.667^\circ$ latitude \times longitude over North America for 2005 from GEOS-Chem (first row), OMI standard product (OMI_GMI, second row), and OMI retrievals using the GEOS-Chem NO₂ a priori vertical profiles (OMI_GC, third row). White areas represent regions with insufficient data. The bottom two rows show the difference between (fourth row) GEOS-Chem and OMI_GC, and (fifth row) OMI_GMI and OMI_GC.

and exhibit similarity in monthly/seasonal variation with the independent data sets. These results are impressive considering the inherent limitations associated with the uncertainties in OMI retrievals and currently available validation data sets. Both temporal and spatial incoherence causes complications in comparing satellite observations with ground-based and aircraft measurements and can often result in misleading conclusions. NO₂ in the lower troposphere is short-lived and is concentrated close to emission sources. Ground-based and in situ instruments offer local measurements, in contrast to satellite observations averaged over a large field of view covering several hundred square kilometers. Therefore, differences between the two measurements ought to be expected simply due to NO₂ spatial inhomogeneity. The sampling differences can be reduced by acquiring long time series of NO₂ measurements, preferably in background locations with more homogeneous distributions.

Although OMI tropospheric NO₂ retrievals show promise and generally compare well with ground truth, occasional large differences could be due to errors in OMI tropospheric

NO₂ columns. Principal sources of error in OMI tropospheric column density are radiometric errors, slant column density calculation, the air mass factor, the retrieved cloud parameters, and the stratosphere–troposphere separation procedure. The tropospheric air mass factor is highly sensitive to errors in surface reflectivity in polluted areas with low surface reflectivity (e.g. Boersma et al., 2004). Further, the tropospheric air mass factor is calculated assuming the NO₂ retrieval implicitly accounts for aerosols through OMI-retrieved cloud fraction and surface reflectivity. However, algorithmic bias due to the presence of actual aerosols has not been studied. We quantify the impact of the a priori NO₂ profiles in tropospheric NO₂ retrievals in Sect. 4.

4 Effect of NO₂ profiles in NO₂ retrievals

In this section, we use aircraft in situ NO₂ measurements coincident with OMI observations during the DISCOVER-AQ campaign in Maryland to explore the sensitivity of the retrieved tropospheric columns to the a priori profiles.

Conversion of the slant column (Ω_s) retrieved from the satellite-measured reflectance spectrum y to vertical NO₂ column (Ω_v) requires an AMF (A):

$$\Omega_v = \frac{\Omega_s(y)}{A(x_a, b)}. \quad (4)$$

The AMF, a measure of the sensitivity of $\ln(y)$ to NO₂, depends on both the a priori NO₂ profile x_a and the forward model parameters b , which include the optical geometry and atmospheric and surface properties (surface albedo, cloud fraction, and cloud height). NO₂ is optically thin in the visible; this allows the calculation of AMF with a profile of altitude-dependent scattering weights (w) computed from a radiative transfer model and the a priori NO₂ profile shape:

$$A_{\text{trop}} = \frac{\sum_{\text{surface}}^{\text{tropopause}} w \times x_a}{\sum_{\text{surface}}^{\text{tropopause}} x_a}, \quad (5)$$

where x_a is the partial NO₂ column. The scattering weights include layer-specific correction factors to account for the temperature dependence of the NO₂ absorption cross-section. The summation extending from the surface to the tropopause provides tropospheric AMF (A_{trop}).

We use Eq. (5) to re-compute tropospheric AMF ($A_{\text{trop_ac}}$) using the measured NO₂ vertical profiles from aircraft and re-calculate tropospheric NO₂ column $\Omega_{v, \text{trop_ac}}$ from OMI tropospheric slant columns ($\Omega_{s, \text{trop}}$):

$$\Omega_{v, \text{trop_ac}} = \frac{\Omega_{s, \text{trop}}}{A_{\text{trop_ac}}} = \frac{\Omega_s - \Omega_{s, \text{strat}}}{A_{\text{trop_ac}}}. \quad (6)$$

Here, Ω_s is the de-stripped NO₂ slant column density (measured NO₂ slant column corrected for instrumental artifacts). The stratospheric slant columns ($\Omega_{s, \text{strat}}$) are calculated from the stratospheric NO₂ vertical columns and the stratospheric AMF, both available in the operational data product.

Figures 5–8 contain tropospheric NO₂ columns re-calculated with aircraft-measured NO₂ vertical profiles. The OMI NO₂ retrievals calculated using the aircraft measured profiles differ from the operational retrievals calculated with model-simulated profiles by up to -43% , in line with other estimates using high resolution a priori profiles (Heckel et al., 2011; Russell et al., 2011). Compared to the operational retrievals, the new retrievals are systematically lower by 16–19% in rural locations and higher by 15–21% in urban locations. These results are consistent with the previous study by Hains et al. (2010), who evaluated the impact of a priori profiles in the Dutch NO₂ (DOMINO) retrievals using observations from the Dutch Aerosol and Nitrogen Dioxide Ex-

periments for validation of OMI and SCIAMACHY (DANDELIONS) and Intercontinental Chemical Transport Experiment Phase B (INTEX-B) campaigns. Our use of measured profiles improved the correlation between OMI and aircraft measurements ($r = 0.5$, $N = 59$). Overall, the agreement between OMI and aircraft measurements improved in urban locations by 12–14% and worsened at Aldino and Fair Hill by a similar magnitude. Comparison of the OMI NO₂ retrievals calculated using the aircraft measured profiles with Pandora observations are presented in Figs. 6–8. Except for Fair Hill, the correlation of OMI with Pandora improved with the new retrievals ($r = 0.4$, $N = 52$). The bias of the OMI retrievals against Pandora reduced at Fair Hill, Beltsville, and Edgewood, but increased at Aldino, Padonia, and Essex.

5 Use of scattering weights in applications of OMI to evaluate AQ models

Several studies (e.g. van Noije et al., 2006; Lamsal et al., 2010) have compared model-simulated NO₂ columns with satellite retrievals. Such comparisons require coincident sampling of model output with observations, because inconsistent sampling could lead to significant differences and incorrect interpretation of the data. The most common approach to comparison involves examining and interpreting the difference between satellite observations and model results. This approach of direct comparison is expected to have difficulty when interpreting differences unless the a priori NO₂ vertical profile shapes used in the retrievals are from the model in question. In this section, we offer an example of the use of scattering weights and OMI retrievals to evaluate AQ models.

The operational NO₂ retrieval algorithm uses NO₂ shape factors generated from GMI simulation results, available at the resolution of $2^\circ \times 2.5^\circ$. The coarse-resolution model profiles may not sufficiently capture the actual vertical distribution of NO₂, especially where the horizontal gradient is large. Moreover, over the last decade, anthropogenic emissions of NO_x have undergone rapid changes that may change the local NO₂ shape factor and subsequently affect the retrieval of tropospheric NO₂. Use of profiles obtained from a model simulation performed with updated emissions at high resolution not only lead to more accurate retrievals through improved spatial representation of NO₂ shape factors in the AMF calculation, but it also ensures self-consistency when the OMI retrievals are compared with modeled NO₂ columns (Eskes and Boersma, 2003; Boersma et al., 2004).

Here, we show an example by comparing OMI tropospheric NO₂ retrievals with a model simulation. We consider the GEOS-Chem nested model (Appendix B) for North America that includes updated emissions and performs simulation at high resolution $0.5^\circ \times 0.667^\circ$. As compared to the coarse model simulation, the fine model simulation can provide better representation of the vertical distributions of NO₂

in OMI pixels by considering changes in the NO₂ shape factors related to the changes in NO_x emissions. We use Eq. (5) to re-compute the tropospheric AMF ($A_{\text{trop_GC}}$) using the new profile and use Eq. (6) to re-calculate the tropospheric NO₂ column (OMI_GC) from OMI. For comparison, we use OMI pixels with cloud radiance fraction < 0.5 and surface reflectivity < 0.3 and calculate area-weighted average columns (Level 3) on a $0.5^\circ \times 0.667^\circ$ grid.

Figure 13 shows seasonal mean tropospheric NO₂ columns from OMI and GEOS-Chem for 2005. Both show large NO₂ columns in dense urban areas in eastern North America and major metropolitan areas such as Los Angeles, San Francisco, Denver, and Houston. They exhibit a similar seasonal pattern, with a winter maximum, reflecting longer NO_x lifetime and shallower mixing layer depth in winter. The correlation between the GEOS-Chem model and OMI seasonal NO₂ columns is remarkable ($r = 0.85\text{--}0.92$). The seasonal average GEOS-Chem column is lower than the OMI column by 7% in spring and higher by 24% in summer yet within the estimated uncertainty of OMI retrievals and GEOS-Chem simulation.

OMI tropospheric NO₂ columns exhibit a number of differences with the modeled NO₂ columns (Fig. 13, fourth row). The modeled NO₂ columns are generally larger in some urban areas of the west coast and northeastern US and over Alberta. Simulation from GEOS-Chem also indicates about factor of 2 higher columns in summer in the midwestern US, a major region of soil NO_x emissions. Retrieved columns are higher over the eastern US in spring, eastern Canada in winter, and cleaner background areas in all seasons. Some of these differences could point to certain emission sources that are not well represented in the model or remaining retrieval biases such as due to the treatment of snow (O'Byrne et al., 2010; McLinden et al., 2014). Other sources of model bias include the errors in simulating OH concentrations, N₂O₅ hydrolysis rates, and vertical mixing that affect simulation of NO_x chemistry (van Noije et al., 2006; Valin et al., 2011).

Possible errors in OMI retrievals causing the observed difference cannot be ruled out. Due to several error sources in the AMF calculation, systematic biases in the spatial variation of OMI retrievals are expected. The spatial resolution of surface reflectivity and a priori NO₂ shape factor are coarser than the ground resolution of OMI, yielding errors in AMF. A change in surface reflectivity from 0.01 to 0.1 could alter the AMF by up to 90% (Leitão et al., 2010), which suggests the importance of accurate knowledge of surface properties (McLinden et al., 2014) and potential impact of residual cloud contamination in the climatology of surface reflectivity. Some previous retrieval studies have used high-resolution MODIS albedo data in an attempt to reduce uncertainty in the tropospheric AMF (Russell et al., 2011; Zhou et al., 2009). Lack of explicit treatment of aerosols in the AMF calculation could have a significant impact in the retrieval of tropospheric NO₂, although the effect could be moderate (7%)

over highly polluted areas (Leitão et al., 2010). Errors in retrievals could be quite large in cases of elevated aerosols in downwind areas, if those elevated aerosols are not accounted for.

Figure 13 (fifth row) shows the seasonal mean difference resulting from the use of GMI profile shapes in the AMF calculation. Since the GMI model and GEOS-Chem both use GEOS-5 meteorological fields and have similar tropospheric chemical mechanisms, the difference between the two retrievals is primarily due to differences in emissions. The anthropogenic emissions in the GMI simulation are appropriate for 1999, which is considerably higher than 2005 emissions over nearly all of North America, with the notable exception of Alberta, where it is considerably lower. Resulting changes in local NO₂ profile shape impact tropospheric AMFs and, therefore, change individual retrievals by up to 40% and seasonal averages by 1% in winter and 12% in fall.

6 Conclusions

We compared the OMI tropospheric NO₂ product (OMNO2, version 2.1) to ground-based measurements to assess the data quality, and to aircraft-based measurements, both to compare the retrieved column amounts and to assess the sensitivity of OMI NO₂ to the a priori profiles used in the retrieval. Model profiles were used to estimate tropospheric column amounts from in situ measurements of NO₂ at ground-level. Finally, we investigated the potential improvement of the retrievals that could be realized using a higher-resolution model, with updated emissions inputs, as a source of a priori profiles. Table 2 summarizes the results of these investigations.

We examined NO₂ profiles measured in situ by the NCAR chemiluminescence instrument flown in the P-3B aircraft during the DISCOVER-AQ field campaign in the Baltimore-Washington, D.C. metropolitan region in July 2011. The P-3B aircraft provided NO₂ measurements from ~ 300 m in the boundary layer to ~ 3.3 km in the free-troposphere, allowing evaluation of tropospheric NO₂ from OMI and estimation of retrieval errors due to a priori NO₂ profiles from the Global Modeling Initiative model. The mean relative vertical distribution of NO₂ from aircraft and that calculated with the GMI model are in agreement to within 30%, but observations also reveal a significant day-to-day variability in NO₂ profile shape. Using aircraft NO₂ profiles altered tropospheric AMFs by up to 43% on some days and yielded improved daily NO₂ column retrievals. Coincident OMI and aircraft measurements agree to within 20% for a majority of cases, with low biases in OMI retrievals by 5.8–22.1% at rural and urban locations and by 50% in the coastal towns of Essex and Edgewood. Comparison of total NO₂ column measurements from OMI and Pandora instruments at those locations presented inconsistent results, suggesting low biases in OMI retrievals of < 6% at Aldino, Beltsville, and Edgewood, and high biases of 9–13% at Padonia and

Essex. Spatial inhomogeneity within a satellite ground pixel, stratosphere–troposphere separation in OMI data, and differences in the sampling domain among the three measurements make short-term comparisons difficult.

We used ground-based data for an extended period of time to examine the seasonal variation of tropospheric NO₂ retrievals. Comparison with the MAX-DOAS measurements at a remote location in Hedo and an urban site in Tsukuba in Japan during 2006–2011 suggests that OMI and MAX-DOAS data are highly consistent ($r = 0.86$), with seasonal biases $< 25\%$ and a mean bias of -16.3% at Tsukuba and 7.1% at Hedo in the OMI retrievals. The inconsistent seasonal variation in total NO₂ columns from OMI and Pandora at Hampton, VA, likely arises from the influence of local NO_x emission sources in the Pandora measurements.

As an indirect validation, we derived the ground-level NO₂ from OMI using coincident GEOS-Chem NO₂ profiles and compared them with surface NO₂ measurements at two rural sites (Centerville, AL and Yorkville, GA) of the SEARCH network. The mean seasonal difference between the OMI-derived surface NO₂ and surface measurements ranges from -11.8% (fall) to 4.1% (spring) in Centerville and from -31% (winter) to 8.2% (spring) in Yorkville. Use of well-established seasonal bottom-up surface NO_x emissions inventories over the United States suggested that the monthly mean differences in OMI-derived top-down surface NO_x emissions range from -9% in July to 35% in April.

Overall, despite the typical complexities associated with the validation of satellite retrievals, OMI tropospheric NO₂ columns are consistent with and agree within the uncertainty of the validation data sets. Therefore, the OMI data offer the means to evaluate the fidelity of CTM model results. The comparison of model-simulated NO₂ columns with satellite retrievals should utilize scattering weights (or averaging kernels) that are made available with the OMI data files, to correct for the effect of climatological monthly a priori NO₂ profiles used in the retrievals. OMI retrieval algorithms could benefit from high-resolution surface reflectivity information and a priori NO₂ profiles and from the explicit treatment of aerosols.

The spatial and temporal coverage of the comparisons we have examined in this paper are limited; they may not be representative of other locations and seasons. A coordinated effort in generating validation data sets by including remotely sensed and in situ observations at the ground, with balloon sondes, and from aircraft over a wide geographic region for a long time period will be valuable for assessing satellite retrievals.

Appendix A: GMI model description

Retrieval of tropospheric NO₂ columns from a satellite instrument requires an assumed vertical distribution of NO₂. Because NO₂ in situ profile measurements are very few, and because the spatial variability in NO₂ profiles is quite large, this is best achieved from a global three-dimensional chemical transport model for atmospheric composition. We use the Global Modeling Initiative (GMI) model (Strahan et al., 2007), consisting of a chemical mechanism that combines the stratospheric mechanism described in Douglass et al. (2004) with a version of the tropospheric mechanism in GEOS-Chem (Bey et al., 2001) with modifications as described in Duncan et al. (2007). The model is driven by assimilated meteorological fields from the Goddard Earth Observing System (GEOS) at the NASA Global Modeling and Assimilation Office (GMAO, <http://gmao.gsfc.nasa.gov/>). The GEOS-5 meteorological data are provided every 3–6 h (3 h for surface fields and mixing depths) at 72 pressure levels in the vertical, extending from surface to 0.01 hPa.

The model includes the global anthropogenic emissions from the Global Emission Inventory Activity center [GEIA, (Benkovitz et al., 1996) for the base year of 1985 and scaled to 1995, as described in Bey et al. (2001). The global inventory is replaced by the following regional inventories: the US EPA National Emissions Inventory (NEI) for 1999 over the United States, the Criteria Air Contaminants (CAC) inventory (<http://www.ec.gc.ca/inrp-npri>) for 2000 over Canada, the Big Bend Regional Aerosol and Visibility Observational Study (BRAVO) inventory for 1999 over Mexico (Kuhns et al., 2005), the European Monitoring and Evaluation Programme (EMEP) inventory for 2000 over Europe, and the inventory from Streets et al. (2006) for 2006 over East Asia. The GMI model also includes NO_x emissions from soil, lightning, biomass burning, biofuel, and aircraft sources, as described in Duncan et al. (2007).

In this work, the model simulation was conducted at the resolution of 2° × 2.5° for 3 years (2005–2007). Model outputs were sampled at the local time of OMI overpass. Since monthly mean values capture the seasonal variation, we derived monthly mean values for NO₂ and temperature profiles and tropopause pressures needed for the calculation of the AMF.

Appendix B: GEOS-Chem model description

We use the GEOS-Chem three-dimensional model of tropospheric chemistry (Bey et al., 2001), version 9-01-03 (www.geos-chem.org), to demonstrate the application of scattering weights to re-calculate the OMI tropospheric NO₂ column and to examine the effect of NO₂ profile shape in retrievals of tropospheric NO₂ columns. We employ GEOS-Chem nested simulations (Zhang et al., 2011; Wang et al., 2012; van Donkelaar et al., 2012; Lamsal et al., 2013) with a horizontal grid size of $\frac{1}{2}^\circ \times \frac{2}{3}^\circ$ over North America (10–70° N, 40–140° W). Boundary conditions of the nested region are provided by the global simulation at 2° × 2.5°. The GEOS-Chem simulation is driven by assimilated meteorological data available from the Goddard Earth Observing System GEOS-5 at the NASA GMAO. The model includes a detailed simulation of tropospheric ozone-NO_x-hydrocarbon chemistry as well as of aerosols and their precursors (Bey et al., 2001; Park et al., 2004).

The global anthropogenic emissions in this GEOS-Chem simulation are from EDGAR 3.2FT2000 (Olivier et al., 2001) for 2000, which are scaled to 2005 following van Donkelaar et al. (2008). The global inventory is overwritten by the following regional inventories: The US EPA NEI for 2005 over the United States, the CAC inventory (<http://www.ec.gc.ca/inrp-npri>) for 2005 over Canada, the BRAVO inventory (Kuhns et al., 2005) for 1999 over Mexico, the EMEP inventory for 2005 over Europe, the inventory from Zhang et al. (2007) for 2006 over East Asia. NO_x emissions from soils, lightning, biomass burning, and aircraft are as described in Lamsal et al. (2010, 2011).

The GEOS-Chem simulation of NO_x has been evaluated extensively with in situ and satellite observations and generally agrees to within 30 % of measured NO_x (Martin et al., 2006; Hudman et al., 2007; Boersma et al., 2008b). We conducted a simulation for the year 2005 and sample the model output between 13:00 and 15:00 local time for analysis of the OMI data.

Acknowledgements. The work was supported by NASA's Earth Science Division through an Aura Science team grant. The Dutch–Finnish-built OMI instrument is part of the NASA EOS Aura satellite payload. The OMI instrument is managed by KNMI and the Netherlands Agency for Aero-space Programs (NIVR).

Edited by: J. G. Murphy

References

- Bechle, M. J., Millet, D. B., and Marshall, J. D.: Remote sensing of exposure to NO₂, satellite versus ground-based measurement in a large urban, *Atmos. Environ.*, 69, 345–353, doi:10.1016/j.atmosenv.2012.11.046, 2013.
- Beirle, S., Platt, U., Wenig, M., and Wagner, T.: Weekly cycle of NO₂ by GOME measurements: a signature of anthropogenic sources, *Atmos. Chem. Phys.*, 3, 2225–2232, doi:10.5194/acp-3-2225-2003, 2003.
- Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G., and Wagner, T.: Megacity emissions and lifetimes of nitrogen oxides probed from space, *Science*, 333, 1737–1739, doi:10.1126/science.1207824, 2011.
- Benkovitz, C. M., Scholtz, M. T., Pacyna, J., Terrasón, L., Dignon, J., Voldner, E. C., Spiro, P. A., Logan, J. A., and Graedel, T. E.: Global gridded inventories of anthropogenic emissions of sulfur and nitrogen, *J. Geophys. Res.*, 101, 29239–29253, 1996.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. R., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: model description and evaluation, *J. Geophys. Res.*, 106, 23073–23096, 2001.
- Blond, N., Boersma, K. F., Eskes, H. J., van der A, R. J., Van Roozendaal, M., De Smedt, I., Bergametti, G., and Vautard, R.: Intercomparison of SCIAMACHY nitrogen dioxide observations, in situ measurements and air quality modeling results over Western Europe, *J. Geophys. Res.*, 112, D10311, doi:10.1029/2006JD007277, 2007.
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO₂ retrieval from space, *J. Geophys. Res.*, 109, D04311, doi:10.1029/2003JD003962, 2004.
- Boersma, K. F., Eskes, H. J., Veefkind, J. P., Brinksma, E. J., van der A, R. J., Sneep, M., van den Oord, G. H. J., Levelt, P. F., Stammes, P., Gleason, J. F., and Bucsela, E. J.: Near-real time retrieval of tropospheric NO₂ from OMI, *Atmos. Chem. Phys.*, 7, 2103–2118, doi:10.5194/acp-7-2103-2007, 2007.
- Boersma, K. F., Jacob, D. J., Bucsela, E. J., Perring, A. E., Dirksen, R., van der A, R. J., Yantosca, R. M., Park, R. J., Wenig, M. O., Bertram, T. H., and Cohen, R. C.: Validation of OMI tropospheric NO₂ observations during INTEX-B and application to constrain NO_x emissions over the eastern United States and Mexico, *Atmos. Environ.*, 42, 4480–4497, 2008a.
- Boersma, K. F., Jacob, D. J., Eskes, H. J., Pinder, R. W., Wang, J., and van der A, R. J.: Intercomparison of SCIAMACHY and OMI tropospheric NO₂ columns: observing the diurnal evolution of chemistry and emissions from space, *J. Geophys. Res.*, 113, D16S26, doi:10.1029/2007JD008816, 2008b.
- Boersma, K. F., Jacob, D. J., Trainic, M., Rudich, Y., DeSmedt, I., Dirksen, R., and Eskes, H. J.: Validation of urban NO₂ concentrations and their diurnal and seasonal variations observed from the SCIAMACHY and OMI sensors using in situ surface measurements in Israeli cities, *Atmos. Chem. Phys.*, 9, 3867–3879, doi:10.5194/acp-9-3867-2009, 2009.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905–1928, doi:10.5194/amt-4-1905-2011, 2011.
- Brinksma, E., Pinardi, G., Braak, R., Volten, H., Richter, A., Schonhardt, A., Van Roozendaal, M., Fayt, C., Hermans, C., Dirksen, R., Vlemmix, T., Berkhout, A. J. C., Swart, D. P. J., Oetjen, H., Wittrock, F., Wagner, T., Ibrahim, O. W., de Leeuw, G., Moerman, M., Curier, R. L., Celarier, E. A., Knap, W. H., Veefkind, J. P., Eskes, H. J., Allaart, M., Rothe, R., Pijters, A. J. M., and Levelt, P.: The 2005 and 2006 DANDELIONS NO₂ and aerosol intercomparison campaigns, *J. Geophys. Res.*, 113, D16S46, doi:10.1029/2007JD008808, 2008.
- Bucsela, E. J., Celarier, E. A., Wenig, M. O., Gleason, J. F., Veefkind, J. P., Boersma, K. F., and Brinksma, E. J.: Algorithm for NO₂ vertical column retrieval from the ozone monitoring instrument, *IEEE Trans. Geosci. Remote. Sens.*, 44, 1245–1258, 2006.
- Bucsela, E. J., Perring, A. E., Cohen, R. C., Boersma, K. F., Celarier, E. J., Gleason, J. F., Wenig, M. O., Bertram, T. H., Wooldridge, P. J., Dirksen, R., and Veefkind, J. P.: Comparison of NO₂ in situ aircraft measurements with data from the ozone monitoring instrument, *J. Geophys. Res.*, 113, D16S31, doi:10.1029/2007JD008838, 2008.
- Bucsela, E. J., Pickering, K. E., Huntermann, T. L., Cohen, R. C., Perring, A., Gleason, J. F., Blakeslee, R. J., Albrecht, R. I., Holzworth, R., Cipriani, J. P., Vargas-Navarro, D., Mora-Segura, I., Pacheco-Hernández, A., and Laporte-Molina, A.: Lightning-generated NO_x seen by the Ozone monitoring instrument during NASA'S Tropical Composition, Cloud and Climate Coupling experiment (TC4), *J. Geophys. Res.*, 115, D00J10, doi:10.1029/2009JD013118, 2010.
- Bucsela, E. J., Krotkov, N. A., Celarier, E. A., Lamsal, L. N., Swartz, W. H., Bhartia, P. K., Boersma, K. F., Veefkind, J. P., Gleason, J. F., and Pickering, K. E.: A new stratospheric and tropospheric NO₂ retrieval algorithm for nadir-viewing satellite instruments: applications to OMI, *Atmos. Meas. Tech.*, 6, 2607–2626, doi:10.5194/amt-6-2607-2013, 2013.
- Cede, A., Herman, J., Richter, A., Krotkov, N. A., and Burrows, J. P.: Measurements of nitrogen dioxide total column amounts using a Brewer double spectrophotometer in direct sun mode, *J. Geophys. Res.*, 111, D05304, doi:10.1029/2005JD006585, 2006.
- Celarier, E. A., Brinksma, E. J., Gleason, J. F., Veefkind, J. P., Cede, A., Herman, J. R., Ionov, D., Goutail, F., Pommereau, J. P., Lambert, J. C., van Roozendaal, M., Pinardi, G., Wittrock, F., Schönhardt, A., Richter, A., Ibrahim, O. W., Wagner, T., Borkov, B., Mount, G., Spinei, E., Chen, C. M., Pongetti, T. J., Sander, S. P., Bucsela, E. J., Wenig, M. O., Swart, D. P. J., Volten, H., Kroon, M., and Levelt, P. F.: Validation of ozone monitoring instrument nitrogen dioxide columns, *J. Geophys. Res.*, 113, D15S15, doi:10.1029/2007JD008908, 2008.

- de Wildt, M. R., Eskes, H., and Boersma, K. F.: The global economic cycle and satellite-derived NO₂ trends over shipping lanes, *Geophys. Res. Lett.*, 39, 101802, doi:10.1029/2011GL049541, 2012.
- Dobber, M. R., Kleipool, Q., Dirksen, R., Levelt, P. F., Jaross, G., Taylor, S., Kelly, T., Flynn, L., Leppelmeier, G., and Rozemeijer, N.: Validation of ozone monitoring instrument level 1b data products, *J. Geophys. Res.*, 113, D15S06, doi:10.1029/2007JD008665, 2008.
- Douglass, A. R., Stolarski, R. S., Strahan, S. E., and Connell, P. S.: Radicals and reservoirs in the GMI chemistry and transport model: comparison to measurements, *J. Geophys. Res.*, 109, D16302, doi:10.1029/2004JD004632, 2004.
- Duncan, B. N., Strahan, S. E., Yoshida, Y., Steenrod, S. D., and Livesey, N.: Model study of the cross-tropopause transport of biomass burning pollution, *Atmos. Chem. Phys.*, 7, 3713–3736, doi:10.5194/acp-7-3713-2007, 2007.
- Duncan, B. N., Yoshida, Y., de Foy, B., Lamsal, L. N., Streets, D., Lu, Z., Pickering, K. E., and Krotkov, N. A.: The observed response of the ozone monitoring instrument (OMI) NO₂ column to NO_x emission controls on power plants in the United States: 2005–2011, *Atmos. Environ.*, 81, 102–111, 2013.
- Edgerton, E. S., Hartsell, B. E., Saylor, R. D., Jansen, J. J., Hansen, D. A., and Hidy, G. M.: The southeastern aerosol research and characterization study, part 3: Continuous measurements of fine particulate matter mass and composition, *J. Air Waste Manage. Assoc.*, 56, 1325–1341, 2006.
- EPA: Primary national ambient air quality standard for nitrogen dioxide, Tech. rep., Environmental Protection Agency, Federal Register, Washington DC 20460, Vol. 74, No. 134, 2009.
- Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285–1291, doi:10.5194/acp-3-1285-2003, 2003.
- Flynn, C. M., Pickering, K. E., Crawford, J. H., Lamsal, L. N., Krotkov, N. A., Herman, J., Weinheimer, A., Chen, G., Liu, X., Szykman, J., Tsay, S. C., Laughner, C. P., Hains, J., Lee, P., Dickerson, R. R., Stehr, J. W., and Brent, L.: The relationship between column-density and surface mixing ratio: statistical analysis of O₃ and NO₂ data from the July 2011 Maryland DISCOVER-AQ mission, *Atmos. Environ.*, 92, 429–441, 2014.
- Ghude, S. D., Lal, D. M., Beig, G., van der A, R., and Sable, D.: Rain-induced soil NO_x emission from India during the onset of the summer monsoon: a satellite perspective, *J. Geophys. Res.*, 115, D16304, doi:10.1029/2009JD013367, 2010.
- Ghude, S. D., Pfister, G. G., Jena, C., van der A, R., Emmons, L. K., and Kumar, R.: Satellite constraints of nitrogen oxide emissions from India based on OMI observations and WRF-Chem simulations, *Geophys. Res. Lett.*, 40, 1–6, doi:10.1029/2012GL053926, 2013a.
- Ghude, S. D., Kulkarni, S. H., Jena, C., Pfister, G. G., Beig, G., Fadnavis, S., and van der A, R. J.: Application of satellite observations for identifying regions of dominant sources of nitrogen oxides over the Indian Subcontinent, *J. Geophys. Res.*, 118, 1075–1089, doi:10.1029/2012JD017811, 2013b.
- Hains, J., Boersma, K. F., Kroon, M., Dirksen, R. J., Cohen, R. C., Perring, A. E., Bucsele, E., Volten, H., Swart, D. P. J., Richter, A., Wittrock, F., Schoenhardt, A., Wagner, T., Ibrahim, O. W., van Roozendaal, M., Pinardi, G., Gleason, J. F., Veefkind, J. P., and Levelt, P.: Testing and improving OMI DOMINO tropospheric NO₂ using observations from the DANDELIONS and INTEX-B validation campaigns, *J. Geophys. Res.*, 115, D05301, doi:10.1029/2009JD012399, 2010.
- Heckel, A., Kim, S.-W., Frost, G. J., Richter, A., Trainer, M., and Burrows, J. P.: Influence of low spatial resolution a priori data on tropospheric NO₂ satellite retrievals, *Atmos. Meas. Tech.*, 4, 1805–1820, doi:10.5194/amt-4-1805-2011, 2011.
- Heland, J., Schlager, H., Richter, A., and Burrows, J. P.: First comparison of tropospheric NO₂ column densities retrieved from GOME measurements and in situ aircraft profile measurements, *Geophys. Res. Lett.*, 29, 1983, doi:10.1029/2002GL015528, 2002.
- Herman, J. R., Cede, A., Spinei, E., Mount, G., Tzortziou, M., and Abuhassan, N.: NO₂ column amounts from ground-based Pandora and MFDOAS spectrometers using the direct-sun DOAS technique: intercomparisons and application to OMI validation, *J. Geophys. Res.*, 114, D13307, doi:10.1029/2009JD011848, 2009.
- Herron-Thorpe, F. L., Lamb, B. K., Mount, G. H., and Vaughan, J. K.: Evaluation of a regional air quality forecast model for tropospheric NO₂ columns using the OMI/Aura satellite tropospheric NO₂ product, *Atmos. Chem. Phys.*, 10, 8839–8854, doi:10.5194/acp-10-8839-2010, 2010.
- Hilboll, A., Richter, A., and Burrows, J. P.: Long-term changes of tropospheric NO₂ over megacities derived from multiple satellite instruments, *Atmos. Chem. Phys.*, 13, 4145–4169, doi:10.5194/acp-13-4145-2013, 2013.
- Hirsch, R. M. and Gilroy, E. J.: Methods of fitting a straight line to data: examples in water resources, *Water Res. Bull.*, 20, 705–711, 1984.
- Hönninger, G., von Friedeburg, C., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS), *Atmos. Chem. Phys.*, 4, 231–254, doi:10.5194/acp-4-231-2004, 2004.
- Hudman, R. C., Jacob, D. J., Turquety, S., Leibensperger, E. M., Murray, L. T., Wu, S., Gilliland, A. B., Avery, M., Bertram, T. H., Brune, W., Cohen, R. C., Dibb, J. E., Flocke, F. M., Fried, A., Holloway, J., Neuman, J. A., Orville, R., Perring, A., Ren, X., Sachse, G. W., Singh, H. B., Swanson, A., and Wooldridge, P. J.: Surface and lightning sources of nitrogen oxides over the United States: magnitudes, chemical evolution, and outflow, *J. Geophys. Res.*, 112, doi:10.1029/2006JD007912, 2007.
- Huijnen, V., Eskes, H. J., Poupkou, A., Elbern, H., Boersma, K. F., Foret, G., Sofiev, M., Valdebenito, A., Flemming, J., Stein, O., Gross, A., Robertson, L., D’Isidoro, M., Kioutsioukis, I., Friese, E., Amstrup, B., Bergstrom, R., Strunk, A., Vira, J., Zyryanov, D., Maurizi, A., Melas, D., Peuch, V.-H., and Zerefos, C.: Comparison of OMI NO₂ tropospheric columns with an ensemble of global and European regional air quality models, *Atmos. Chem. Phys.*, 10, 3273–3296, doi:10.5194/acp-10-3273-2010, 2010.
- Intergovernmental Panel on Climate Change (IPCC): Contribution of working group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, in: *Climate Change 2007: The Physical Science Basis*, edited by: Solomon, Qin, S., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., 996 pp., Cambridge University Press, Cambridge, UK, 2007.

- Ionov, D. V., Timofeyev, Y. M., Sinyakov, V. P., Semenov, V. K., Goutail, F., Pommereau, J. P., Bucsel, E. J., Celarier, E. A., and Kroon, M.: Ground-based validation of EOS-Aura OMI NO₂ vertical column data in the midlatitude mountain ranges of Tien Shan (Kyrgyzstan) and Alps (France), *J. Geophys. Res.*, 113, 4437, doi:10.1029/2007JD008659, 2008.
- Irie, H., Kanaya, Y., Akimoto, H., Tanimoto, H., Wang, Z., Gleason, J. F., and Bucsel, E. J.: Validation of OMI tropospheric NO₂ column data using MAX-DOAS measurements deep inside the North China Plain in June 2006: Mount Tai Experiment 2006, *Atmos. Chem. Phys.*, 8, 6577–6586, doi:10.5194/acp-8-6577-2008, 2008.
- Irie, H., Boersma, K. F., Kanaya, Y., Takashima, H., Pan, X., and Wang, Z. F.: Quantitative bias estimates for tropospheric NO₂ columns retrieved from SCIAMACHY, OMI, and GOME-2 using a common standard for East Asia, *Atmos. Meas. Tech.*, 5, 2403–2411, doi:10.5194/amt-5-2403-2012, 2012.
- Jaeglé, L., Steinberger, L., Martin, R. V., and Chance, K.: Global partitioning of NO_x sources using satellite observations: Relative roles of fossil fuel combustion, biomass burning and soil emissions, *Faraday Discuss.*, 130, 4537, doi:10.1039/b502128f, 2005.
- Kim, S. W., Heckel, A., McKeen, S. A., Frost, G. J., Hsie, E. Y., Trainer, M. K., Richter, A., Burrows, J. P., Peckham, S. E., and Grell, G. A.: Satellite-observed US power plant NO_x emission reductions and their impact on air quality, *Geophys. Res. Lett.*, 33, L22812, doi:10.1029/2006GL027749, 2006.
- Kim, S. W., Heckel, A., Frost, G. J., Richter, A., Gleason, J., Burrows, J. P., McKeen, S. A., Hsie, E. Y., Granier, C., and Trainer, M. K.: NO₂ columns in the western United States observed from space and simulated by a regional chemistry model and their implications for NO_x emissions, *J. Geophys. Res.*, 114, D11301, doi:10.1029/2008JD011343, 2009.
- Knepp, T., Pippin, M., Crawford, J., Chen, G., Szykman, J., Long, R., Cowen, L., Cede, A., Abuhassan, N., Herman, J., Delgado, R., Compton, J., Berkoff, T., Fishman, J., Martins, D., Stauffer, R., Thompson, A. M., Weinheimer, A., Knapp, D., Montzka, D., Lenschow, D., and Neil, D.: Estimating surface NO₂ and SO₂ mixing ratios from fast-response total column observations and potential application to geostationary missions, *J. Atmos. Chem.*, D15308, doi:10.1007/s10874-013-9257-6, 2013.
- Konovalov, I. B., Beekmann, M., Richter, A., and Burrows, J. P.: Inverse modelling of the spatial distribution of NO_x emissions on a continental scale using satellite data, *Atmos. Chem. Phys.*, 6, 1747–1770, doi:10.5194/acp-6-1747-2006, 2006.
- Kramer, L. J., Leigh, R. J., Remedios, J. J., and Monks, P. S.: Comparison of OMI and ground-based in situ and MAX-DOAS measurements of tropospheric nitrogen dioxide in an urban area, *J. Geophys. Res.*, 113, D16S39, doi:10.1029/2007JD009168, 2008.
- Kuhns, H., Knipping, E. M., and Vokovich, J. M.: Development of a United States-Mexico emissions inventory for the Big Bend Regional Aerosol and Visibility Observational (BRAVO) study, *J. Air Waste Mgt. Assoc.*, 55, 677–692, 2005.
- Lamsal, L. N., Martin, R. V., van Donkelaar, A., Steinbacher, M., Celarier, E. A., Bucsel, E., Dunlea, E. J., and Pinto, J. P.: Ground-level nitrogen dioxide concentrations inferred from the satellite-borne ozone monitoring instrument, *J. Geophys. Res.*, 113, D16308, doi:10.1029/2007JD009235, 2008.
- Lamsal, L. N., Martin, R. V., van Donkelaar, A., Celarier, E. A., Bucsel, E. J., Boersma, K. F., Luo, R. D. C., and Wang, Y.: In-direct validation of tropospheric nitrogen dioxide retrieved from the OMI satellite instrument: insight into the seasonal variation of nitrogen oxides at northern midlatitudes, *J. Geophys. Res.*, 115, D05302, doi:10.1029/2009JD013351, 2010.
- Lamsal, L. N., Martin, R. V., Padmanabhan, A., van Donkelaar, A., Zhang, Q., Sioris, C. E., Chance, K., Kurosu, T. P., and Newchurch, M. J.: Application of satellite observations for timely updates to global anthropogenic NO_x emission inventories, *Geophys. Res. Lett.*, 38, L05810, doi:10.1029/2010GL046476, 2011.
- Lamsal, L. N., Martin, R. V., Parrish, D. D., and Krotkov, N. A.: Scaling relationship for NO₂ pollution and population size: a satellite perspective, *Environ. Sci. Technol.*, 47, 7855–7861, 2013.
- Leitão, J., Richter, A., Vrekoussis, M., Kokhanovsky, A., Zhang, Q. J., Beekmann, M., and Burrows, J. P.: On the improvement of NO₂ satellite retrievals – aerosol impact on the air-mass factors, *Atmos. Meas. Tech.*, 3, 475–493, doi:10.5194/amt-3-475-2010, 2010.
- Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O. V., and Saari, H.: The ozone monitoring instrument, *IEEE Trans. Geosci. Remote.*, 44, 1093–1101, 2006.
- Lin, J.-T.: Satellite constraint for emissions of nitrogen oxides from anthropogenic, lightning and soil sources over East China on a high-resolution grid, *Atmos. Chem. Phys.*, 12, 2881–2898, doi:10.5194/acp-12-2881-2012, 2012.
- Lin, J.-T., McElroy, M. B., and Boersma, K. F.: Constraint of anthropogenic NO_x emissions in China from different sectors: a new methodology using multiple satellite retrievals, *Atmos. Chem. Phys.*, 10, 63–78, doi:10.5194/acp-10-63-2010, 2010.
- Lu, Z. and Streets, D. G.: Increase in NO_x emissions from Indian thermal power plants during 1996–2010: unit-based inventories and multisatellite observations, *Environ. Sci. Technol.*, 46, 7463–7470, doi:10.1021/es300831w, 2012.
- Martin, R. V., Chance, K., Jacob, D. J., Kurosu, T. P., Spurr, R. J. D., Bucsel, E., Gleason, J. F., Palmer, P. I., Bey, I., Fiore, A. M., Li, Q., Yantosca, R. M., and Koelemeijer, R. B. A.: An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.*, 107, 4437, doi:10.1029/2001JD001027, 2002.
- Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Perner, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emission constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, 108, 4537, doi:10.1029/2003JD003453, 2003.
- Martin, R. V., Sioris, C. E., Chance, K., Ryerson, T. B., Bertram, T. H., Wooldridge, P. J., Cohen, R. C., Neuman, J. A., Swanson, A., and Flocke, F. M.: Evaluation of space-based constraints on global nitrogen oxide emissions with regional aircraft measurements over and downwind of eastern North America, *J. Geophys. Res.*, 111, D15308, doi:10.1029/2005JD006680, 2006.
- McLinden, C. A., Fioletov, V., Boersma, K. F., Kharol, S. K., Krotkov, N., Lamsal, L., Makar, P. A., Martin, R. V., Veeffkind, J. P., and Yang, K.: Improved satellite retrievals of NO₂ and SO₂ over the Canadian oil sands and comparisons with surface measurements, *Atmos. Chem. Phys.*, 14, 3637–3656, doi:10.5194/acp-14-3637-2014, 2014.
- Mebust, A. K. and Cohen, R. C.: Observations of a seasonal cycle in NO_x emissions from fires in African woody savannas, *Geophys. Res. Lett.*, 40, 1451–1455, doi:10.1002/grl.50343, 2013.

- Mebust, A. K., Russell, A. R., Hudman, R. C., Valin, L. C., and Cohen, R. C.: Characterization of wildfire NO_x emissions using MODIS fire radiative power and OMI tropospheric NO₂ columns, *Atmos. Chem. Phys.*, 11, 5839–5851, doi:10.5194/acp-11-5839-2011, 2011.
- NARSTO: Improving emission inventories for effective air quality management across North America, A NARSTO Assessment, Tech. rep., NARSTO, available at: <http://www.narsto.org/> (8 December 2013), 2005.
- Novotny, E. V., Bechle, M. J., Millet, D. B., and Marshall, J. D.: National satellite-based land-use regression: NO₂ in the United States, *Environ. Sci. Technol.*, 45, 4407–4414, doi:10.1021/es103578x, 2011.
- O'Byrne, G., Martin, R. V., van Donkelaar, A., Joiner, J., and Celarier, E. A.: Surface reflectivity from OMI using MODIS to eliminate clouds: effects of snow on UV-Vis trace gas retrievals, *J. Geophys. Res.*, 115, D17305, doi:10.1002/jgrd.50550, 2010.
- Oetjen, H., Baidar, S., Krotkov, N. A., Lamsal, L. N., Lechner, M., and Volkamer, R.: Airborne MAX-DOAS measurements over California: testing the NASA OMI tropospheric NO₂ product, *J. Geophys. Res.*, 118, 7400–7413, doi:10.1002/jgrd.50550, 2013.
- Olivier, J. G. J., Berdowski, J. J. M., Peters, J. A. H. W., Bakker, J., Visschedijk, A. J. H., and Bloos, J. P. J.: Applications of EDGAR. Including a description of EDGAR 3.2: Reference database with trend data for 1970–1995, Tech. rep., RIVM Bilthoven, RIVM report, Bilthoven, The Netherlands, 773301 001/NRP report 410200 051, 2001.
- Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A. M., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: application to formaldehyde retrievals from the global ozone monitoring experiment, *J. Geophys. Res.*, 106, 14539–14550, 2001.
- Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *J. Geophys. Res.*, 108, 4180, doi:10.1029/2002JD002153, 2003.
- Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: implications for policy, *J. Geophys. Res.*, 109, D15204, doi:10.1029/2003JD004473, 2004.
- Peters, A. J. M., Boersma, K. F., Kroon, M., Hains, J. C., Van Roozendaal, M., Wittrock, F., Abuhassan, N., Adams, C., Akrami, M., Allaart, M. A. F., Apituley, A., Beirle, S., Bergwerff, J. B., Berkhout, A. J. C., Brunner, D., Cede, A., Chong, J., Clémer, K., Fayt, C., Frieß, U., Gast, L. F. L., Gil-Ojeda, M., Goutail, F., Graves, R., Griesfeller, A., Großmann, K., Hemerijckx, G., Hendrick, F., Henzing, B., Herman, J., Hermans, C., Hoexum, M., van der Hoff, G. R., Irie, H., Johnston, P. V., Kanaya, Y., Kim, Y. J., Klein Baltink, H., Kreher, K., de Leeuw, G., Leigh, R., Merlaud, A., Moerman, M. M., Monks, P. S., Mount, G. H., Navarro-Comas, M., Oetjen, H., Pazmino, A., Perez-Camacho, M., Peters, E., du Piesanie, A., Pinardi, G., Puentedura, O., Richter, A., Roscoe, H. K., Schönhardt, A., Schwarzenbach, B., Shaiganfar, R., Sluis, W., Spinei, E., Stolk, A. P., Strong, K., Swart, D. P. J., Takashima, H., Vlemmix, T., Vrekoussis, M., Wagner, T., Whyte, C., Wilson, K. M., Yela, M., Yilmaz, S., Zieger, P., and Zhou, Y.: The Cabauw Intercomparison campaign for Nitrogen Dioxide measuring Instruments (CINDI): design, execution, and early results, *Atmos. Meas. Tech.*, 5, 457–485, doi:10.5194/amt-5-457-2012, 2012.
- Platt, U.: Differential Optical Absorption Spectroscopy (DOAS), in: *Air Monitoring by Spectroscopic Techniques*, edited by: Sigrist, M. W., John Wiley, New York, USA, 27–84, 1994.
- Richter, A., Burrows, J. P., Nüß, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide levels over China observed from space, *Nature*, 437, 129–132, 2005.
- Ridley, B. A. and Grahek, F.: A small, low flow, high-sensitivity reaction vessel for NO chemiluminescence detectors, *J. Atmos. Ocean. Tech.*, 7, 307–311, 1990.
- Ridley, B., Ott, L., Pickering, K., Emmons, L., Montzka, D., Weinheimer, A., Knapp, D., Grahek, F., Li, L., Heymsfield, G., McGill, M., Kucera, P., Mahoney, M. J., Schultz, M., and Brasseur, G.: Florida thunderstorms: a faucet of reactive nitrogen to the upper troposphere, *J. Geophys. Res.*, 109, D17305, doi:10.1029/2004JD004769, 2004.
- Roscoe, H. K., Van Roozendaal, M., Fayt, C., du Piesanie, A., Abuhassan, N., Adams, C., Akrami, M., Cede, A., Chong, J., Clémer, K., Friess, U., Gil Ojeda, M., Goutail, F., Graves, R., Griesfeller, A., Grossmann, K., Hemerijckx, G., Hendrick, F., Herman, J., Hermans, C., Irie, H., Johnston, P. V., Kanaya, Y., Kreher, K., Leigh, R., Merlaud, A., Mount, G. H., Navarro, M., Oetjen, H., Pazmino, A., Perez-Camacho, M., Peters, E., Pinardi, G., Puentedura, O., Richter, A., Schönhardt, A., Shaiganfar, R., Spinei, E., Strong, K., Takashima, H., Vlemmix, T., Vrekoussis, M., Wagner, T., Wittrock, F., Yela, M., Yilmaz, S., Boersma, F., Hains, J., Kroon, M., PETERS, A., and Kim, Y. J.: Intercomparison of slant column measurements of NO₂ and O₄ by MAX-DOAS and zenith-sky UV and visible spectrometers, *Atmos. Meas. Tech.*, 3, 1629–1646, doi:10.5194/amt-3-1629-2010, 2010.
- Russell, A. R., Valin, L. C., Bucsele, E. J., Wenig, M. O., and Cohen, R. C.: Space-based constraints on spatial and temporal patterns of NO_x emissions in California, 2005–2008, *Environ. Sci. Technol.*, 44, 3608–3615, doi:10.1021/es903451j, 2010.
- Russell, A. R., Perring, A. E., Valin, L. C., Bucsele, E. J., Browne, E. C., Wooldridge, P. J., and Cohen, R. C.: A high spatial resolution retrieval of NO₂ column densities from OMI: method and evaluation, *Atmos. Chem. Phys.*, 11, 8543–8554, doi:10.5194/acp-11-8543-2011, 2011.
- Russell, A. R., Valin, L. C., and Cohen, R. C.: Trends in OMI NO₂ observations over the United States: effects of emission control technology and the economic recession, *Atmos. Chem. Phys.*, 12, 12197–12209, doi:10.5194/acp-12-12197-2012, 2012.
- Ryerson, T. B., Williams, E. J., and Fehsenfeld, F. C.: An efficient photolysis system for fast-response NO₂ measurements, *J. Geophys. Res.*, 105, 26447–26461, 2000.
- Schaub, D., Boersma, K. F., Kaiser, J. W., Weiss, A. K., Folini, D., Eskes, H. J., and Buchmann, B.: Comparison of GOME tropospheric NO₂ columns with NO₂ profiles deduced from ground-based in situ measurements, *Atmos. Chem. Phys.*, 6, 3211–3229, doi:10.5194/acp-6-3211-2006, 2006.
- Schaub, D., Brunner, D., Boersma, K. F., Keller, J., Folini, D., Buchmann, B., Berresheim, H., and Staehelin, J.: SCIAMACHY tropospheric NO₂ over Switzerland: estimates of NO_x lifetimes and impact of the complex Alpine topography on the retrieval, *At-*

- mos. Chem. Phys., 7, 5971–5987, doi:10.5194/acp-7-5971-2007, 2007.
- Solomon, S., Portmann, R. W., Sanders, R. W., Daniel, J. S., Madson, W., Bartram, B., and Dutton, E. G.: On the role of nitrogen dioxide in the absorption of solar radiation, *J. Geophys. Res.*, 104, 12047–12058, 1999.
- Steinbacher, M., Zellweger, C., Schwarzenbach, B., Bugmann, S., Buchmann, B., Ordóñez, C., Prévôt, A. S. H., and Hueglin, C.: Nitrogen oxides measurements at rural sites in Switzerland: Bias of conventional measurement techniques, *J. Geophys. Res.*, 112, D11307, doi:10.1029/2006JD007971, 2007.
- Strahan, S. E., Duncan, B. N., and Hoor, P.: Observationally derived transport diagnostics for the lowermost stratosphere and their application to the GMI chemistry and transport model, *Atmos. Chem. Phys.*, 7, 2435–2445, doi:10.5194/acp-7-2435-2007, 2007.
- Streets, D. G., Zhang, Q., Wang, L., He, K., Hao, J., Wu, Y., Tang, Y., and Carmichael, G. R.: Revisiting China's CO emissions after the Transport and Chemical Evolution over the Pacific (TRACE-P) mission: synthesis of inventories, atmospheric modeling, and observations, *J. Geophys. Res.*, 111, D14306, doi:10.1029/2006JD007118, 2006.
- Tang, W., Cohan, D. S., Lamsal, L. N., Xiao, X., and Zhou, W.: Inverse modeling of Texas NO_x emissions using space-based and ground-based NO₂ observations, *Atmos. Chem. Phys.*, 13, 11005–11018, doi:10.5194/acp-13-11005-2013, 2013.
- Thornton, J. A., Wooldridge, P. J., and Cohen, R. C.: Atmospheric NO₂: in situ laser-induced fluorescence detection at parts per trillion mixing ratios, *Anal. Chem.*, 72, 528–539, doi:10.1021/ac9908905, 2000.
- Toenges-Schüller, N., Stein, O., Rohrer, F., Wahner, A., Richter, A., Burrows, J. P., Beirle, S., Wagner, T., Platt, U., and Elvidge, C. D.: Global distribution pattern of anthropogenic nitrogen oxide emissions: Correlation analysis of satellite measurements and model calculations, *J. Geophys. Res.*, 111, D05312, doi:10.1029/2005JD006068, 2006.
- Valin, L. C., Russell, A. R., Hudman, R. C., and Cohen, R. C.: Effects of model resolution on the interpretation of satellite NO₂ observations, *Atmos. Chem. Phys.*, 11, 11647–11655, doi:10.5194/acp-11-11647-2011, 2011.
- van der A, R. J., Peters, D. H. M., Eskes, H., Boersma, K. F., Van Roozendaal, M., De Smedt, I., and Kelder, H. M.: Detection of the trend and seasonal variation in tropospheric NO₂ over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594, 2006.
- van der A, R. J., Eskes, H. J., Boersma, K. F., van Noije, T. P. C., Van Roozendaal, M., De Smedt, I., Peters, D. H. M., Kuenen, J. J. P., and Meijer, E. W.: Identification of NO₂ sources and their trends from space using seasonal variability analyses, *J. Geophys. Res.*, 113, doi:10.1029/2007JD009021, 2008.
- van Donkelaar, A., Martin, R. V., Leaitch, W. R., Macdonald, A. M., Walker, T. W., Streets, D. G., Zhang, Q., Dunlea, E. J., Jimenez, J. L., Dibb, J. E., Huey, L. G., Weber, R., and Andreae, M. O.: Analysis of aircraft and satellite measurements from the Intercontinental Chemical Transport Experiment (INTEX-B) to quantify long-range transport of East Asian sulfur to Canada, *Atmos. Chem. Phys.*, 8, 2999–3014, doi:10.5194/acp-8-2999-2008, 2008.
- van Donkelaar, A., Martin, R. V., Pasch, A. N., Szykman, J. J., Zhang, L., Wang, Y., and Chen, D.: Improving the accuracy of daily satellite-derived ground-level fine aerosol concentration estimates for North America, *Environ. Sci. Technol.*, 46, 11971–11978, doi:10.1021/es3025319, 2012.
- van Noije, T. P. C., Eskes, H. J., Dentener, F. J., Stevenson, D. S., Ellingsen, K., Schultz, M. G., Wild, O., Amann, M., Aher-ton, C. S., Bergmann, D. J., Bey, I., Boersma, K. F., Butler, T., Cofala, J., Drevet, J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Martin, R. V., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Richter, A., Rodriguez, J. M., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., and van Roozendaal, M.: Multi-model ensemble simulations of tropospheric NO₂ compared with GOME retrievals for the year 2000, *Atmos. Chem. Phys.*, 6, 2943–2979, doi:10.5194/acp-6-2943-2006, 2006.
- Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., Merienne, M. F., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294 K, *J. Quant. Spectrosc. Ra.*, 59(3–5), 171–184, 1998.
- Vinken, G. C. M., Boersma, K. F., van Donkelaar, A., and Zhang, L.: Constraints on ship NO_x emissions in Europe using GEOS-Chem and OMI satellite NO₂ observations, *Atmos. Chem. Phys.*, 14, 1353–1369, doi:10.5194/acp-14-1353-2014, 2014.
- Wagner, N. L., Dubé, W. P., Washenfelder, R. A., Young, C. J., Pollack, I. B., Ryerson, T. B., and Brown, S. S.: Diode laser-based cavity ring-down instrument for NO₃, N₂O₅, NO, NO₂ and O₃ from aircraft, *Atmos. Meas. Tech.*, 4, 1227–1240, doi:10.5194/amt-4-1227-2011, 2011.
- Wang, S., Pongetti, T. J., Sander, S. P., Spinei, E., Mount, G. H., Cede, A., and Herman, J.: Direct sun measurements of NO₂ column abundances from Table Mountain, California: intercomparison of low- and high-resolution spectrometers, *J. Geophys. Res.*, 115, D13305, doi:10.1029/2009JD013503, 2010.
- Wang, S. W., Zhang, Q., Streets, D. G., He, K. B., Martin, R. V., Lamsal, L. N., Chen, D., Lei, Y., and Lu, Z.: Growth in NO_x emissions from power plants in China: bottom-up estimates and satellite observations, *Atmos. Chem. Phys.*, 12, 4429–4447, doi:10.5194/acp-12-4429-2012, 2012.
- Wenig, M. O., Cede, A. M., Bucsela, E. J., Celarier, E. A., Boersma, F. K., Veefkind, J. P., Brinkma, E. J., Gleason, J. F., and Herman, J. R.: Validation of OMI tropospheric NO₂ column densities using direct-sun mode Brewer measurements at NASA Goddard Space Flight Center, *J. Geophys. Res.*, 113, D16S45, doi:10.1029/2007JD008988, 2008.
- Zhang, L., Jacob, D. J., Smith-Downey, N. V., Wood, D. A., Blewitt, D., Carouge, C., van Donkelaar, A., Jones, D. B. A., Murray, L. T., and Wang, Y.: Improved estimate of the policy-relevant background ozone in the United States using the GEOS-Chem global model with 1/2° × 2/3° horizontal resolution over North America, *Atmos. Environ.*, 45, 6769–6776, 2011.
- Zhang, Q., Streets, D. G., He, K., Wang, Y., Richter, A., Burrows, J. P., Uno, I., Jang, C. J., Chen, D., Yao, Z., and Lei, Y.: NO_x emission trends for China, 1995–2004: the view from the ground and the view from space, *J. Geophys. Res.*, 112, D22306, doi:10.1029/2007JD008684, 2007.

Zhao, C. and Wang, Y.: Assimilation inversion of NO₂ emissions over east Asia using OMI NO₂ column measurements, *Geophys. Res. Lett.*, 36, L06805, doi:10.1029/2008GL037129, 2009.

Zhou, Y., Brunner, D., Boersma, K. F., Dirksen, R., and Wang, P.: An improved tropospheric NO₂ retrieval for OMI observations in the vicinity of mountainous terrain, *Atmos. Meas. Tech.*, 2, 401–416, 2009.