Tropical Journal of Pharmaceutical Research June 2011; 10 (3): 335-340 © Pharmacotherapy Group, Faculty of Pharmacy, University of Benin, Benin City, 300001 Nigeria.

All rights reserved.

Available online at http://www.tjpr.org DOI: 10.4314/tjpr.v10i3.3

Research Article

Evaluation of the Antibacterial Activity of 14 Medicinal Plants in Côte d'Ivoire

GEK Bolou¹, I Bagré^{1*}, K Ouattara¹ and AJ Djaman^{1,2}

¹Biochemical Pharmacodynamy Laboratory, Biosciences Department, Cocody University, Abidgan; ²Biochemical Laboratory of Pasteur Institute of Cote d'Ivoire, Abidgan.

Abstract

Purpose: To evaluate the antibacterial potentials of fourteen ethnobotanically selected plants traditionally used in different parts of Côte d'Ivoire for the treatment of typhoid fever and gastrointestinal disorders.

Method: The antimicrobial activity of the extracts of the plant was tested against a collection strain of Salmonella typhimurium, a clinical strain of Salmonella typhi and Pseudomonas aeruginosa by macrobroth dilution method. Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) were determined.

Results: All the plants had bactericidal activity against at least one of the test microorganisms with both MIC and MBC ranging from 2.5 to > 80 mg/ml. Among the plant extracts, Terminalia glaucescens. (Combretaceae) Bersama abyssinica subsp. paullinioides (Melianthaceae) and Abrus precatorius (Fabaceae) showed the most promising broad spectrum antibacterial properties, inhibiting all of the strains tested, especially S. Typhi and P. aeruginosa, with MBC ranging from 2.5 to 5 mg/ml.

Conclusion: The findings provide support for the use of these plants in traditional medicine for treatment of typhoid fever and gastrointestinal disorders, and are thus, potential sources of drugs that would need to be subjected to further studies.

Keywords: Antimicrobial activity, Ivorian medicinal plants, MIC, MBC.

Received: 10 July 2010

Revised accepted: 24 April 2011

*Corresponding author: E-mail: bagrefreefr@yahoo.fr; Tél: +225-(0)8008901

INTRODUCTION

Medicinal plants have been known for their healing and/or disease-curing qualities for centuries. Some drugs of plant origin used in conventional medical practice are direct plant extracts or plant materials that have been suitably prepared and standardized [1].

Many readily available plants in Côte d'Ivoire are used in traditional folklore medicine for the treatment of typhoid fever and gastrointestinal disorders such as cholera, diarrheoa and dysentery [2,3]. However, several of them have not been investigated from a pharmacological point of view to demonstrate their antibacterial properties, which could support their use as antityphoid, anticholera or antidiarrhoeal remedies in traditional medicine.

The objective of the present study was to evaluate the antibacterial activity of the extracts of some plants used in Côte d'Ivoire traditional healthcare system, against a group of pathogenic bacteria. These plants are used by the indigenous people in different parts of Côte d'Ivoire for the treatment of infectious diseases such as cholera. diarrhoea. dysentery and other gastrointestinal disorders [4,5]. Α total number of fourteen species of plants belonging to eleven different families were tested on S. typhimurium, S. typhi and P. aeruginosa.

EXPERIMENTAL

Plant materials

Fourteen medicinal [Abrus plants precatorius Linn. (Fabaceae), Acanthospermum hispidium Schrank. (Asteraceae), Melanthera scandens Schum. & Thonn. (Asteraceae). Vernonia amygdalina Del. (Poaceae), Bambusa vulgaris Schrad. (Poaceae). Bersama abyssinica subsp. paullinioides Planch. (Melianthaceae). Table 1: Plant species with relevant information

polycarpa Enantia Enal. & Diels Nauclea latifolia SM. (Annonaceae), (Rubiaceae). Phyllanthus amarus Schum. Thonn. (Euphorbiaceae), and Portulaca quadric-folia Linn. (Portulacaceae), Tectona grandis Linn. (Verbenaceae), Terminalia catappa Linn. (Combretaceae), Terminalia glauce-scens Planch. (Combretaceae) and Thoningia sanguinea Vahl. (Balanophoraceae) - were ethnobotanically selected for antimicrobial screening in the present study. The fresh plants were collected from Daloa in Central West Region of Ivory Coast in June 2009. The botanical identification of the plant samples was carried out by Pr Ake Assi, of the Department of Botany, University of Cocody-Abidjan. Voucher specimens of the plant materials were conserved in the herbarium of Centre National de Floristique (CNF) of Abidjan. The plant species, parts used. local name. voucher specimen numbers and the traditional uses of the plants are listed in Table 1.

Bacterial strains

The organisms used were clinical isolates of *Salmonella typhi, Pseudomonas aeruginosa* and a collection strain of *Salmonella typhimurium* ATCC 14028 (provided by Laboratoire de Bactériologie and Virologie of Institut Pasteur de Côte d'Ivoire).

Extraction procedure

The freshly collected parts of the plants were air-dried at room temperature for 7 days and powdered. Briefly, 20 g of the powder of each of the plant materials was separately soaked in 500 ml distilled water for 24 h with constant stirring. The suspension was filtered through Whatman filter paper no. 1. The filtrate was concentrated by vacuum using a rotary evaporator to obtain the dry aqueous extract. In the same way, the ethanol (70 %) extract was also obtained.

| Species | Part of plant used (Local name) | Voucher number | Traditional uses* |
|---|--|-----------------------------------|---|
| Abrus precatorus | Seed (Mlinmlin) | CNF/875 | Dermatose, wound, cough |
| Acanthospermum hispidium | Whole plant (Lukoubassa moni) | CNF/16762 | Paludism, typhoïd fever |
| Melanthera scandens | Whole plant | CNF/17737 | Paludism, typhoïd fever |
| Vernonia amygdalina | Whole plant (Abohoui) | CNF/11694 | Paludism, typhoïd fever |
| Bambusa vulgaris | Leaf (Féfé) | CNF/4715 | Typhoïd fever |
| Bersama abyssinica subsp. Paullinioides | Root (Lêgrêgbégbéhi) | CNF/1451 | Stagnation |
| Enantia polycarpa Nauclea latifolia | Bark (Souin) Leaf (Tôlé) | CNF/17786 CNF/15927 | Paludism, typhoïd fever |
| Phyllanttus amarus | Whole plant (Soundjassi) | CNF/6712 | Paludism, typhoïd fever |
| Portulaca quadrifolia | Whole plant (Sourêh) | CNF/578 | Purgative |
| Tectona grandis | Leaf (Tec) | CNF/842 | Typhoid fever |
| Terminalia catappa Terminalia glaucescens Thoningia sanguinea | Leaf (Oôcôman) Leaf (Wôlô) Flower (Glouglan) | CNF/1564 CNF/1657 CNF/17954 | Dermatose Paludism, typhoïd fever Typhoid fever |

* Sources: refs 19 and 20

Antibacterial tests

Isolates were considered susceptible, less susceptible, or resistant to a particular antimicrobial agent on the basis of the diameters of the inhibitory zones that matched the criteria of the manufacturer's interpretation table, which is based on the guidelines of the National Committee for Clinical Laboratory Standards [6,7].

Minimal inhibitory concentration (MIC) was determined according to Wilkinson and Gentry [8]. Two-fold dilutions of the extract were made in the concentration range of 0.625 to 80.0 mg/ml. The tubes were inoculated with a microorganism suspension at a final density of 10^6 ufc/ml prepared according to the procedure of Société Française de Microbiologie (SFM) [9]. The tubes were incubated at 37 °C for 24 h. The lowest concentration of the tube which did not show any visible growth after macroscopic examination was considered as the MIC.

Minimal bactericidal concentration (MBC) is defined as the concentration producing a 99.9 % reduction in colony forming units (CFU) number in the initial inoculum. It was determined by subculture on nutrient agar as previously described [10]. Briefly, the tubes without growth after 24 h of incubation were subcultured on Mueller Hinton agar in Petri dishes for 24 h. MBC was determined as the lowest concentration that showed no bacterial growth in the subcultures [11,12]. The tests were performed in triplicate. Dilutions of ciprofloxacin served as positive control.

Statistical analysis

The data are presented as mean \pm SEM. All the data were analyzed by one-way ANOVA and differences between the means were assessed with Neuman-Keuls's multiple comparison tests. Differences were considered significant *at p* < 0.05. All analyses were carried out using Graph Pad software, version 5.01 (USA).

RESULTS

The results of MIC and MBC tests are presented in Tables 2 and 3, respectively. Generally, each plant studied exhibited antibacteral activity against at least one tested bacterial strain. The aqueous and ethanol extracts of *B. abyssinica*, *P. amarus*,

T. catappa and *T. glaucescens* presented MIC and MBC values ranging from 2.5 to 80 mg/ml against *S. typhimurium*, *S. typhi* and *P. aeruginosa*. The ethanol extracts of *B. abyssinica and T. catappa* exhibited strong MIC and MBC values of 2.5 mg/ml against *P. aeruginosa*. The ethanol extracts of *T. glaucescens and A. precatorius* showed strong bactericidal activities with MIC and MBC values of 5 mg/ml against *S. typhi* while the aqueous extract of *T. sanguinea* and the ethanol extract of *T. glaucescens* exhibited strong bactericidal activities with MIC and MBC values of 5 and 10 mg/ml, respectively. The ethanol extract of *A. hispidium*, and aqueous extracts of *V. amygdalina* and *B. vulgaris* were inactive against the three strains. Also, the aqueous extract of *A. hispidium*, and ethanol extracts of *V. amygdalina* and *T. grandis* were inactive against *S. typhi* and *P. aeruginosa* strains.

Table 2: Minimum inhibitory concentration (MIC, mg/ml) and minimum bactericidal concentration (MBC, mg/ml) of aqueous plant extracts (mean ± S.E.M., n = 3)

| Extract | <i>Salmonella</i> Typhimurium <i>ATCC 14028</i> | | <i>Salmonella</i> Typhi | | Pseudomonas aeruginosa | |
|--|---|--------------------|-------------------------------|--------------------|----------------------------|--------------------|
| | MIC | MBC | MIC | MBC | MIC | MBC |
| Abrus precatorus | 40.0 ± 5.1 ^b | 80 ^c | 40.0 ± 4.9 ^b | >80 | 10.0± 2.9 ^ª | 20 ^b |
| Acanthospermum hispidium | 20.0 ± 3.2 ^b | 40 ^b | 40.0 ± 4.2 ^b | >80 | 40.0 ± 4.5 ^b | >80 |
| Melanthera scandens | 40.0 ± 4.3 ^b | 80 ^c | 20.0 ± 3.6 ^b | >80 | 20.0 ± 3.4 ^b | 20 ^b |
| Vernonia amygdalina | 40.0 ± 3.8 ^b | >80 | 40.0 ± 5.1 ^b | >80 | 40 ± 4.8^{b} | >80 |
| Bambusa vulgaris | 40.0 ± 4.2^{b} | >80 | 40.0 ± 4.3 ^b | 40 ^b | 40.0 ± 4.8 ^b | 40 ^b |
| Bersama abyssinica subsp. Paullinioides | 40.0 ± 46^{b} | 40 ^b | 40.0 ± | 80 ^c | 5.0 ± 1.1^{a} | 5 ^a |
| Enantia polycarpa | 20.0 ± 4.1 | 40 ^b | 10.0 ± | 10 ^a | >80 | >80 |
| Nauclea latifolia | 40.0 ± 5 4 ^b | 80 ^c | >80 | >80 | 80.0 ± | 80 ^c |
| Phyllanttus amarus | 10.0 ± | 20 ^b | 20.0 ± 3 7 ^b | 40 ^b | 10.0 ± 3 4 ^a | 10 ^a |
| Portulaca quadrifolia | >80 | >80 | >80 | >80 | 40.0 ± 4 7 ^b | 40 ^b |
| Tectona grandis | 10.0 ± 3.2 ^a | 20 ^b | 40.0 ± 5^{b} | >80 | 10.0 ± 2.1^{a} | 40 ^b |
| Terminalia catappa | 20.0 ± 4.3^{b} | 80 ^c | 20.0 ± 4.3 ^b | 80 ^c | 5.0 ± 0.6^{a} | 10 ^a |
| Terminalia glaucescens | 10.0 ± | 20 ^b | 10.0 ± | 20 ^b | 10.0 ± 2.3ª | 10 ^a |
| Thoningia sanguinea | 5.0 ± 1.5^{a} | 10 ^a | 5.0 ± 1.4^{a} | 20 ^b | 20.0 ± | 40 ^b |
| Ciprofloxacin | 0.025 ± 0.004 ^e | 0.025 ^e | 0.025 ± 0.002 ^e | 0.025 ^e | 0.025 ± 0.005° | 0.025 ^e |

Note: Values with the same superscript character in each column are not significantly different (p < 0.05)

| Salmonella Typhimurium ATCC 14028 | | Salmonella Typhi | | Pseudomonas aeruginosa | |
|---|---|---|--|---|--|
| МІС | МВС | МІС | МВС | МІС | МВС |
| 5.0 ± 0.3 ^a | 10 ^a | 5.0 ± 0.7 ^a | 5 ^a | 10.0 ± 3.1 ^a | 20 ^b |
| $80.0 \pm 8.4^{\circ}$ | >80 | 80.0 ± 7.1 ^c | >80 ^c | 80.0 ± 5.9 ^c | >80 |
| 40.0 ± 5.1 ^b | 80 ^c | 20.0 ± 3.2 ^b | 80 ^c | 80.0 ± 6.1 ^c | 80 ^c |
| 80.0 ± 6.5 ^c | 80 ^c | 80.0 ± 8.1 ^c | >80 | 80.0 ± 5.9 ^c | >80 |
| 40.0 ± 4.2^{b} | >80 | 40.0 ± 4.3 ^b | 40 ^b | 40.0 ± 4.8 ^b | 40 ^b |
| 10.0 ± 3.5 ^a | 20 ^b | 10.0 ± 3.2^{a} | 20 ^b | 2.5 ± 0.9^{d} | 2.5 ^d |
| 20.0 ± 4.1 | 40 ^b | 10.0 ± 2.5 ^a | 10 ^a | >80 | >80 |
| 20.0 ± 3.6 | 20 ^b | 20.0 ± 2.9 ^b | 20 ^b | 20.0 ± 2.4 ^b | 20 ^b |
| 10.0 ± 2.3 ^a | 20 ^b | 10.0 ± 1.9 ^ª | 20 ^b | 5.0 ± 1.1 ^a | 10 ^a |
| $40.0 \pm 5.2^{\circ}$ | 80 [°] | $20.0 \pm 2.8^{\circ}$ | 80 [°] | $80.0 \pm 6.7^{\circ}$ | 80° |
| 20.0 ± 3.7 ^b | 40 ^b | 20.0 ± 3.4 ^b | >80 | 20.0 ± 2.4 ^b | >80 |
| $5.0 \pm 1.2^{a} \\ 5.0 \pm 0.9^{a} \\ 20.0 \pm 2.8^{b} \\ 0.025 \pm 0.003^{e}$ | 20 ^b 10 ^a 20 ^b 0.025 ^e | $\begin{array}{l} 10.0 \pm 1.7^{a} \\ 5.0 \pm 0.8^{a} \\ 20.0 \pm 1.7^{b} \\ 0.025 {\pm} 0.007^{e} \end{array}$ | 40 ^b 5 ^a 20 ^b 0.025 ^e | 2.5 ± 0.7^{d} 5 ± 1.5^{a} 10.0 ± 2.6^{a} 0.025 ± 0.005^{e} | 2.5 ^d 10 ^a 20 ^b 0.025 ^e |
| | $\begin{tabular}{ c c c c c } \hline Salmone.\\ \hline Typhimuri & ATCC 140 \\ \hline Typhimuri & ATCC 140 \\ \hline Structure & Structu$ | $\begin{tabular}{ c c c c } \hline Salmonella \\ \hline Typhimurium \\ ATCC 14028 \\ \hline \hline MIC & MBC \\ \hline $5.0 \pm 0.3^a & 10^a \\ $80.0 \pm 8.4^c & >80 \\ \hline $40.0 \pm 5.1^b & 80^c \\ $80.0 \pm 6.5^c & 80^c \\ $40.0 \pm 4.2^b & >80 \\ $10.0 \pm 3.5^a & 20^b \\ \hline $20.0 \pm 4.1 & 40^b \\ $20.0 \pm 3.6 & 20^b \\ $10.0 \pm 2.3^a & 20^b \\ $40.0 \pm 5.2^b & 80^c \\ $20.0 \pm 3.7^b & 40^b \\ $5.0 \pm 1.2^a & 20^b \\ $5.0 \pm 0.9^a & 10^a \\ $20.0 \pm 2.8^b & 20^b \\ $0.025 \pm 0.003^e & 0.025^e \\ \hline \end{tabular}$ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ |

Table 3: Minimum inhibitory concentration (MIC, mg/ml) and minimum bactericidal concentration (MBC, mg/ml) of ethanol plant extracts (mean \pm S.E.M., n = 3)

Note: Values with the same superscript letter in each column are not significantly different (p < 0.05)

DISCUSSION

Phytoconstituents present in plants, namely, flavonoids, alkaloids, tannins and triterpenoids, have demonstrated exciting potentials for the expansion of the range of modern chemotherapies against a wide spectrum of microorganisms [13,14]. The antimicrobial activities of various plants have been reported by several researchers [15,16].

The results of this study show that all the selected plants exhibited varied antimicrobial activities against the tested organisms. The extracts tested were effective antibacterial agents against a group of microorganisms that are implicated in either typhoid fever and/or other gastrointestinal disorders/ infectious diseases such as diarrheoa and dysentery. The ethanol extracts showed the better bactericidal activity. Thus, ethanol is the more suitable extraction solvent for the plants assessed. Previous studies also

indicate that ethanol was the best solvent for extracting antimicrobial substances from some plants [17,18].

Among the fourteen plants evaluated, the extracts of *A. precatorius*, *B. abyssinica* and *T. glaucescens* were the most effective against the selected strains, followed by *T. catappa* and *T. sanguinea* in that order.

This study has not only shown the scientific basis for some of the therapeutic uses of traditional plants, but has also confirmed the ethnomedicinal claims for the selected plants.

CONCLUSION

The results of this study lend some scientific credence to the indigenous uses of the lvorian medicinal plants evaluated for the treatment of infectious diseases and gastrointestinal disorders such as cholera

and diarrhoeal diseases. There is, however, a need for further investigation of the plants that exhibited the highest bactericidal activity with a view to identifying and isolating their active principles.

ACKNOWLEDGEMENT

The authors wish to thank Pr. Aké Assi Laurent of the Department of Botany, University of Cocody- Abidjan, for the botanical identification and collection of the plants used in this study.

REFERENCES

- Donald EU. Medicinal plants research in Nigeria: Retrospect and Prospects. In: Sofowora A, Ed. The State of Medicinal Plants Research in Nigeria. Nigerian Society of Pharmacognosy, Ibadan University Press, Nigeria, 1986; pp 1– 12.
- 2. Maikere-Faniyo R, Van Puyvelde L, Mutwewingabo A, Habiyaremye FX. Study of Rwandese medicinal plants used in the treatment of diarrhoea. J Ethonopharmacol, 1989; 26: 101-109.
- Zirihi GN. Contribution au recensement à l'identification et à laconnaissance de quelques espèces végétales utilisées en médecine traditionelle chez les Bété du département d'Issia, Côte d'Ivoire. Université de Cocody, UFR Biosciences, Thèse de Doctorat de 3ème cycle 1991; p 253
- Zirihi GN, Mambu L, Guédé-Guédé F, Bodo B, Grellier P. In vitro antiplasmodial activity and cytotoxicity of 33 West African plants used for the treatment of malaria. J Ethnopharmacol, 2005; 98: 281-285.
- 5. Grayer RJ, Harborne JB. A survey of antifungal compounds from higher plants. Phytochemistry, 1994; 37: 19–42.
- NCCLŚ. Performance standards for antimicrobial disk susceptibility testing, 12th informational supplement. NCCLS document, 2002; M100-S12. National Committee for Clinical Laboratory Standards, Wayne, Pa.
- Acharyya S, Patra A, Bag PK. Evaluation of the Antimicrobial Activity of Some Medicinal Plants against Enteric Bacteria with Particular Reference to Multi-Drug Resistant Vibrio Cholerae. Trop J Pharm Res, 2009; 8(3): 231-237

- Wilkinson ID, Gentry LO. In vitro comparison of ceftazidime and nine other antimicrobial agents against hospital strains of. Gramnegative bacteria. J Antimicrob Chemother, 1981; 8: 53–56.
- SFM (Société Française de Microbiologie). Comité de l'antibiogramme de la société française de microbiologie. Recommandations : édit. Janvier 2008; p 49
- Riffel A, Medina LF, Stefani V, Santos RC, Bizani D, Brandelli A. In vitro antimicrobial activity of a new series of 1,4-naphthoquinones. Braz J Med Biol Res, 2002; 35(7): 811-818
- 11. Suffredini IB, Bacchi EM, Sakuda TMK, Ohara MT, Younes RN, Varella AD. Antibacterial activity of Apocynaceae extracts and MIC of Tabernaemontana angulata stem organic extract. Rev Bras Ciênc Farm, 2002; 38: 89-94.
- 12. N'guessan JD, Dinzedi MR, Guessennd N, Coulibaly A, Dosso M, Djaman AJ, Guede-Guina F. Antibacterial activity of the aqueous extract of Thonningia sanguinea against extendedspectrum-β- lactamases (ESBL) producing Escherichia coli and Klebsiella pneumoniae strains. Trop J Pharm Res, 2007; 6 (3): 779-783
- 13. Lutterodt GD, Ismail A, Basheer RH, Baharudin HM. Antimicrobial effects of Psidium guajava extracts as one mechanism of its antidiarrhoeal action. Malaysian J Med Sci, 1999; 6 (2): 17-20.
- 14. Marjorie MC. Plant products as antimicrobial agents. Clin Microbiol Rev, 1999; 12(4): 564-582.
- 15. Cowan MM. Plant products as antimicrobial agents. Clin Microbiol Rev, 1999; 22: 564-582.
- Shariff ZU. Modern Herbal Therapy for Common Ailments. Nature Pharmacy Series (Vol 1), Spectrum Books Limited, Ibadan, Nigeria in Association with Safari Books (Export) Limited, United Kingdom, 2001; pp 9-88.
- Jonathan SG, Fasidi IO. Antimicrobial activities of two Nigerian edible macrofungi, Lycoperdon pusilum and L. giganteum. Afr J Biomed Res, 2003; 6: 85-90.
- Dulger G, Aki C. Antimicrobial Activity of the Leaves of Endemic Stachys pseudopinardii in Turkey. Trop J Pharm Res, 2009; 8(4): 371-375
- Aké Assi L. Flore de la Côte d'Ivoire : Catalogue systématique, biogéographique et écologie. Mémoires de botaniques systématiques. Boissiera, Génève, Suisse. 2001, vol. 57, p 315.
- Aké Assi L. Flore de la Côte d'Ivoire : Catalogue systématique, biogéographique et écologie. Mémoires de botaniques systématiques. Boissiera, Génève, Suisse. 2002, vol. 58, p 327.