

Evidence and a conceptual model for the co-occurrence of nitrogen fixation and denitrification in heterotrophic marine sediments

R. W. Fulweiler^{1,*}, S. M. Brown², S. W. Nixon³, B. D. Jenkins^{2,3}

¹Department of Earth and Environment & Department of Biology, Boston University, Boston, Massachusetts 02215, USA

²Department of Cell and Molecular Biology, University of Rhode Island, Kingston, Rhode Island 02881, USA

³Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island 02882, USA

ABSTRACT: Marine waters are often nitrogen (N) limited. Denitrification, the microbial conversion of nitrate to dinitrogen (N₂) gas, is responsible for significant N removal from the coastal ocean. In contrast, nitrogen fixation, the microbial transformation of N₂ to ammonium, is typically regarded as an inconsequential N source. The imbalance between these 2 processes is responsible, at least in part, for N limitation in the coastal ocean. Organic matter quality and quantity has been shown to determine rates of these critical N cycling processes. We hypothesized that the timing of organic matter deposition to the benthos might also be important in determining which process dominates. We tested this hypothesis using a coupled biogeochemical-molecular approach. We report directly measured net sediment denitrification rates and corresponding expression of *nirS*, a gene in the denitrification pathway, with the simultaneous expression of *nifH*, a gene associated with nitrogen fixation. The timing of organic matter deposition determined the magnitude of the net sediment N₂ fluxes. Highest rates of denitrification occurred soon after deposition, and the lowest rates occurred over 200 d after the last deposition event concomitant with increased *nifH* expression. Phylogenetic evidence suggests that sulfur and sulfate reducers are responsible for the nitrogen fixation. Globally, warming water temperatures, changes in light, and reduced nutrient loads through management intervention have been linked to decreases and/or altered phenology of water column productivity. Based on a conceptual model developed here, we suggest that in these systems, heterotrophic sediment nitrogen fixation may become an important component of the nitrogen budget.

KEY WORDS: Denitrification · Heterotrophic nitrogen fixation · Marine sediments · *nifH* expression · Direct sediment N₂ fluxes

— Resale or republication not permitted without written consent of the publisher —

INTRODUCTION

Canonical denitrification is the microbially mediated process that converts nitrate to dinitrogen (N₂) gas, thus removing it from the nitrogen (N) pool that is available to most photosynthetic organisms. The opposing process, nitrogen fixation, converts N₂ gas to biologically available ammonium. The working paradigm is that the imbalance between denitrification and nitrogen fixation helps promote N limitation

in marine systems (Howarth et al. 1988a,b). Various environmental factors, including sediment oxygen concentrations and substrate availability, regulate these 2 processes and often in the same direction (Cornwell et al. 1999). Of particular importance is the link between sediment N₂ fluxes and organic matter, which has been shown to be positively correlated with both denitrification (Seitzinger & Giblin 1996) and nitrogen fixation (Capone & Budin 1982). However, it is not simply the amount of organic matter

*Email: rwf@bu.edu

that regulates these processes but a complicated interplay between organic matter quantity and quality (Fulweiler et al. 2007).

Due to energetic constraints and the typically abundant concentrations of nitrogen in marine systems, it was long thought that heterotrophic nitrogen fixation was likely an unimportant process (Howarth et al. 1988a,b). However, over the last decade this paradigm has begun to shift with observations of direct N_2 uptake in a variety of marine systems. For example, significant rates of directly measured nitrogen fixation have been recorded in temperate and subtropical estuaries (e.g. Fulweiler et al. 2007, Gardner et al. 2006, Gardner & McCarthy 2009). In addition, it has been known for some time that nitrogen fixation is a significant source of nitrogen in some salt marsh (e.g. Jones 1974, Teal et al. 1979, Jones 1982) and seagrass (e.g. McGlathery et al. 1998, Cole & McGlathery 2012) systems.

We previously reported a dramatic seasonal switch in N cycling with high rates of net heterotrophic sediment nitrogen fixation in July and August for a temperate New England estuary (41.3° N, 71.1° W; Narragansett Bay, RI, USA; Fulweiler et al. 2007). Using field observations and a large mesocosm experiment, we linked these rates of nitrogen fixation to a climate induced oligotrophication of the bay brought on by long-term warming (Nixon et al. 2004, Oviatt 2004) and the loss of the winter-spring diatom bloom (Nixon et al. 2009). But as observed for a variety of systems (Farnsworth et al. 1995, Walther et al. 2002, Kahru et al. 2011), climate change in this estuary has also altered the phenology or seasonal sequencing of phytoplankton blooms, leading to more phytoplankton production in the summer and fall and decreased organic deposition on the bottom (Nixon et al. 2009). Thus, we hypothesized that shifts in phytoplankton bloom phenology and the subsequent changes to organic matter deposition to the sediments might also be important in determining whether nitrogen fixation or denitrification dominates.

Using a coupled biogeochemical-molecular approach we tested this hypothesis by altering the timing of phytoplankton deposition in 9 large (4 m²) mesocosm tanks designed to replicate the bottom of the estuary. Our purpose here is to report on the co-occurrence of denitrification and nitrogen fixation in heterotrophic sediments, and on the dynamic response of these sediments in relation to time since organic matter deposition. In addition, we present a conceptual model hypothesizing how and why these 2 opposing processes may take place simultaneously.

MATERIALS AND METHODS

Mesocosm experimental details

Sediments were collected from a site located in mid-Narragansett Bay (41° 35.3' N, 71° 22.3' W) on 29 November 2006 where benthic metabolism and net N_2 fluxes have been measured at various time intervals since the late 1970s (e.g. Nixon et al. 1976, Seitzinger et al. 1980, Fulweiler et al. 2007, Fulweiler & Nixon 2009). This site is completely heterotrophic, approximately 7 m deep (mean low water; mlw), with a 1.1 m tidal range, and receives less than 1–2% of surface light. Sediments were collected using a large box corer (0.25 m²) to a depth of approximately 30 cm and placed into trays; approximately 4 box cores were needed to fill each tray, and we took great care to preserve their vertical structure. The sediments were transported to the Graduate School of Oceanography at the University of Rhode Island, and placed into one of 9 large (4 m²) mesocosm tanks maintained in the dark with 65 cm of overlying water. The overlying water was slowly mixed and exchanged 10% each day with water from the lower West Passage of the bay. The mesocosms mimicked the *in situ* mid-bay site temperatures (~1 to 21°C annual range) and the environmental chambers for N_2 gas flux measurements were maintained at the field temperature at the time of collection. Annual salinity range was ~27 to 32. Surface water column nutrient concentrations for mid Narragansett Bay averaged 4.3 $\mu\text{mol l}^{-1}$ over the 2007 annual cycle (www.gso.uri.edu/phytoplankton/). Nitrate was highest during the winter (12.3 $\mu\text{mol l}^{-1}$) and was undetectable during the spring, summer, and fall. In contrast, ammonium and nitrite were highest in the fall and winter (8.3 and 13.8 $\mu\text{mol l}^{-1}$, respectively) and undetectable in the spring and summer (www.gso.uri.edu/phytoplankton/).

This mesocosm experiment was designed to quantify how the temporal addition of organic matter to the benthos affected sediment N_2 fluxes. The amount of organic matter delivered to the benthos from water column phytoplankton production varies both spatially and temporally. Percentages of organic matter reaching the sediment range from 20–30% in the Gulf of Mexico to approximately 25% off the coast of New England and off the Amazon (Seitzinger & Giblin 1996). In general, a larger proportion of winter-spring blooms is deposited to the bottom than of summer blooms (Keller et al. 2001). The amount of organic matter deposition can also depend on water column depth. Based on our previous experiments

(Fulweiler et al. 2007) and literature values (e.g. Seitzinger & Giblin 1996) we assumed a deposition of 25% (or 50 g C m⁻²) of the winter-spring bloom as observed in Narragansett Bay during the 1970s. The 9 tanks were randomly assigned to a timing treatment and were subsequently amended with a total of 50 g C m⁻² in equal daily aliquots over a 2 wk period in the beginning of either January, April, or June 2007. Carbon was added in the form of commercially available spray-dried phytoplankton with a molar C:N of 10.5. We did not include for comparison mesocosms with no organic matter additions as the infrastructure was not available for this experiment and our goal was to test the impact of timing rather than starvation versus deposition (which we tested previously; Fulweiler et al. 2007, 2008).

Sediment N₂ gas fluxes

In early July (17°C) and again in September (22°C) 2007, sediment cores (10 cm inner diameter, 30.5 cm long) were collected from each of the 9 mesocosms with a pull corer (n = 9, 3 cores from each treatment). The sediment cores were rapidly transported to a water bath in a dark environmental chamber at the Graduate School of Oceanography at the University of Rhode Island. The cores were left uncapped with air bubbling the overlying water for 8 to 12 h. The water overlying each core was carefully replaced with filtered (1 µm) lower Narragansett Bay (salinity ~32) water before each incubation. The cores were then sealed with a gas-tight lid (with no air headspace; mean water headspace was 1.2 l) and replicate water samples for N₂/Ar (Kana et al. 1994) analysis were collected at 5 points over the course of an incubation and preserved with HgCl₂. Incubations lasted between 7 and 8 h allowing for at least a 2 mg O₂ l⁻¹ drop in the overlying water, but the sediment cores were not allowed to go hypoxic. Each water sample was analyzed for dissolved gas concentration using a quadrupole membrane inlet mass spectrometer (MIMS, Bay Instruments) (Kana et al. 1994). The N₂/Ar technique is actually a measure of net N₂ production or consumption (gross denitrification minus gross nitrogen fixation). Over the course of the incubation, the change in N₂ concentration was determined from the change in the measured N₂/Ar multiplied by the Ar concentration at air saturation. N₂ flux across the sediment-water interface was determined for each of the cores using a 5-point linear regression, and rates were then prorated for the volume and area of the core.

Sub-sampling and nucleic acid extractions

After the net N₂ flux incubations were completed, we sub-sampled 2 of 3 cores. From each sediment core, 4 replicate sub-cores (~4 to 8 cm length, 0.25 cm diameter) were collected: 2 for DNA extractions and 2 for RNA extractions. The sub-cores were flash frozen in liquid N₂, and sub-sectioned into 0.5 cm segments from the sediment water interface to 3 cm in depth. Total DNA was extracted using the Powersoil DNA Isolation Kit (MO Bio). For RNA isolation, replicate sub-core samples from each depth were pooled (approximately 0.5 g of wet sediment) to average biological heterogeneity across the larger core and to increase biomass and RNA yield. Total RNA was extracted using the Powersoil RNA Isolation Kit (MO Bio); however, the kit was designed to extract RNA from 2 g of soil. To accommodate this reduction in reaction scale, a quarter of the volumes of the Bead, SR1, SR2, SR3 and SR4 solutions and phenol-chloroform-isoamyl alcohol were used. After the RNA precipitation step, the dried pellet was resuspended with 100 µl of nuclease free water, 10 µl of 10× TURBO DNase buffer and 1 µl of TURBO DNase from the TURBO DNase-free Kit (Ambion) and incubated at 37°C for 30 min. To inactivate the reaction, 10 µl of DNase Inactivation reagent was added and incubated at room temperature for 5 min. The remaining RNA purification steps were carried out using the RNeasy Mini Kit (Qiagen) according to the manufacturer's instructions. cDNA copies of RNA were generated with SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen). For all samples, 8 µl of DNase-treated RNA at a concentration of 4 ng µl⁻¹ was added to the reaction. Each reaction was primed with 1 µl of 2 µM outer reverse primers for both our genes of interest, *nifH3* (Zehr & McReynolds 1989) (5'ATR TTR TTN GCN GCR TA3') and *nirS6R* (Braker et al. 1998) (5'CGT TGA ACT TRC CGG T3'). After the reverse transcriptase was added, the mixture was incubated at 50°C for 50 min. All the other steps followed the instructions of the manufacturer. For every sample, we also included controls that did not contain reverse transcriptase to confirm there was no DNA contamination in the subsequent PCR amplification.

Functional gene sequence analysis

The *nifH* gene from environmental DNA or cDNA was isolated using nested PCR with degenerate outer primers *nifH4* (5'TTY TAY GGN AAR GGN GG3') and

nifH3 (5'ATR TTR TTN GCN GCR TA3'), and inner primers nifH1 (5'TGY GAY CCN AAR GCN GA3') and nifH2 (5'AND GCC ATC ATY TCN CC3') (Zehr & McReynolds 1989, Kirshtein et al. 1991). Both rounds of PCR consisted of an initial denaturation step of 2 min at 94°C; cycling steps that included a denaturation step of 30 s at 94°C; an annealing step of 30 s at 50°C; and an extension step of 1 min at 72°C. All reactions had a final extension step of 7 min at 72°C. First round reactions had 25 cycles and the second round reactions had 30 cycles (Zehr & McReynolds 1989, Kirshtein et al. 1991). *nirS* was amplified using the primer pair nirS1F (5'CCT AYT GGC CGC CRC ART3') and nirS6R (5'CGT TGA ACT TRC CGG T3') (Braker et al. 1998, 2000). After a 2 min initial denaturation step 94°C, a touchdown PCR was performed that consisted of a denaturation step of 30 s at 94°C, an annealing step of 30 s, and an extension step of 1 min at 72°C. During the first 11 cycles, the annealing temperature decreased 0.5°C every cycle starting at 56°C. For the last 25 cycles the annealing temperature was 54°C. A final extension step was performed for 7 min at 72°C (Braker et al. 1998, 2000).

To obtain relative quantities of *nifH* amplification, we determined the intensity of *nifH* amplification bands with the manual band quantification function in GeneTools (SynGene). We normalized the intensity of every detectable band on the gels to that of a 100 ng band on HyperLadder II (Biolone Tauton) for a relative quantification value. Based on a distribution curve of the relative values, samples were then assigned low (<200 ng), medium (200 to 400 ng) or high (>400 ng) expression levels. For every expressed *nifH* and *nirS* amplicon we collected 3 or 4 sequences.

Expressed *nifH* sequences from Narragansett Bay, accession numbers JN645307–JN645499, were combined with the *nifH* sequences (4505) selected as unique sequence representatives by J. P. Zehr (pers. comm.) from the NifH database encompassing all currently available sequences in GenBank (Benson et al. 2009) as of September 2009. Translated NifH protein sequences were aligned using the multiple sequence alignment tool, MUSCLE (Edgar 2004) within the Geneious software package. A *nifH* maximum likelihood tree of aligned protein sequences was constructed in Geneious using PhyML with 1000 bootstrap replicates.

A *nirS* database was created by collecting all the *nirS* sequences from the National Center for Biotechnology Information (NCBI) (Benson et al. 2009) and importing the GenBank files into ARB (Ludwig et

al. 2004). Translated NirS protein sequences were aligned using the multiple sequence alignment tool, MUSCLE (Edgar 2004) within the Geneious software package. A NirS protein maximum likelihood tree including the Narragansett Bay expressed *nirS* sequences, accession numbers JN645500–JN645545, was constructed using the PhyML algorithm in Geneious with 1000 bootstrap replicates.

RESULTS

Sediment N₂ fluxes over the course of this experiment ranged from 27 to 65 μmol N₂-N m⁻² h⁻¹. There was no statistical difference ($t(13) = 2.16$, $p = 0.33$) between the mean net N₂ fluxes measured at 17°C and the mean N₂ fluxes measured at 22°C. However, the relationship between days since last organic matter deposition and the magnitude of the net sediment N₂ fluxes was strongly correlated ($R^2 = 0.87$, $p = 0.0002$). The highest rates of denitrification occurred soon after deposition and the lowest rates occurred over 200 d after the last organic matter deposition (Fig. 1).

In two of the triplicate cores measured for direct N₂ fluxes, we identified and followed the presence and expression of functional genes associated with nitrogen fixation (*nifH*) and denitrification (*nirS*). Both *nifH* and *nirS* genes were detected in all sediment samples analyzed from the sediment-water interface




Fig. 1. Net N₂-N fluxes across the sediment-water interface as a function of number of days since last organic matter deposition. Sediments were collected from the mid-bay historic site (41° 35.3' N, 71° 22.3' W). Triplicate tanks received 50 g C m⁻² at 3 different times in 2007: January (open circles), April (shaded circles), and June (closed circles). Net N₂-N fluxes were measured in July and September 2007 (mean ± SE, n = 3). Both the slope ($F = 27.1159$, $p = 0.0065$) and intercept ($p = 0.0002$) of the regression are significant




Fig. 2. Normalized *nifH* (squares) and *nirS* (circles) expression data from mesocosm sediment samples analyzed by gel electrophoresis. Duplicate cores are shown for each treatment and are aligned in order of days since organic matter deposition. Mean net N₂ flux (N₂-N μmol m⁻² h⁻¹) is shown in the shaded bar. For the January treatment (Day 214), RNA yields were below detection, thus only one profile is shown. Expression levels of *nifH* are designated as low, medium, or high (open, shaded, or closed squares, respectively) from normalized gel image analysis. *nirS* amplification was low in most cores, and samples from which we obtained clones are indicated by open circles. Blank spaces: no gene expression detected; X: no sample available

to a depth of 3.0 cm (data not shown). While *nifH* mRNA transcripts were detected in every treatment, the spatial distribution of *nifH* expression changed with treatment. Specifically, in sediments recently enriched with organic matter, high *nifH* expression was only found at the 3.0 cm depth in one of the replicate cores. In contrast, sediments starved of organic matter had expression throughout the sediment column from the surface to 3.0 cm depth. In fact, the majority of *nifH* expression was seen in the treatment tanks 142 to 214 d after organic matter addition (Fig. 2). This *nifH* expression correlates with our direct N₂ flux measurements which also show the lowest net N₂ fluxes during this time.

We observed no temporal or vertical trend in *nirS* expression signals (Fig. 2). We did detect *nirS* expression in some of the same samples and, importantly, at the same depths in which we also observed *nifH* expression (Fig. 2).

Expressed *nifH* sequences from Narragansett Bay sediments are restricted to 2 phylogenetic groups within *nifH* Clusters I and III, as previously defined (Chien & Zinder 1996), and group with known sulfur and sulfate reducing bacteria (Fig. 3). The majority of




Fig. 3. Maximum likelihood phylogenetic tree of NifH protein sequences from Narragansett Bay sediments; sequences observed from cultured organisms and from those reported in Bertics & Ziebis (2010). Bootstrap values over 50% are indicated. The number inside the group = the number of total sequences within the grouping. *nifH* cluster designations are denoted according to Chien & Zinder (1996). Scale bar = substitutions per site




Fig. 4. Maximum likelihood phylogenetic tree of NirS protein sequences from Narragansett Bay and closely related cultured species with bootstrap values. The number inside (or directly next to) the group = number of total sequences within the grouping. Scale bar = substitutions per site

expressed *nifH* sequences (134) are within group NB3 which is most closely related to *Pelobacter carbinolicus*, a sulfur reducer (Lovley et al. 1995) (Fig. 3). The second largest expressed sequence group, NB7, within Cluster I contains 18 expressed *nifH* sequences and the most closely related bacterium is the sulfate reducer *Desulfovibrio vulgaris* (Fig. 3). Expressed *nirS* sequences were found in 5 main phylogenetic groups, none of which contain any cultivated species (Fig. 4).

DISCUSSION

N₂ fluxes

The experiment we report here was motivated by our observations of a switch in sediment N₂ fluxes from net denitrification to net nitrogen fixation in Narragansett Bay in the summer of 2006 (Fulweiler et al. 2007). In our previous work, we linked the switch from net denitrification to net nitrogen fixation to a decrease in water column productivity over the last 30 yr and the subsequent decline in the amount of

organic matter loading to the benthos. Based on these field observations, we ran a mesocosm experiment which showed that the amount of organic matter deposited to the benthos determined the net sediment N₂ flux (Fulweiler et al. 2007). We found that sediments that received organic matter had a positive sediment N₂ flux or net denitrification rate of 245 $\mu\text{mol N}_2\text{-N m}^{-2} \text{h}^{-1}$. In contrast, sediments that did not receive any organic matter exhibited net nitrogen fixation rates of $-289 \mu\text{mol N}_2\text{-N m}^{-2} \text{h}^{-1}$. In addition, when sediment microcosms exhibiting net nitrogen fixation were given organic matter they switched to net denitrification (Fulweiler et al. 2007, Fulweiler et al. 2008). However, Narragansett Bay and other coastal systems are also experiencing changes in the phenology or seasonal sequencing of phytoplankton blooms. Changes to the timing of phytoplankton blooms may alter both the quantity and quality of organic matter loading to the benthos (Nixon et al. 2009). The purpose of this second large mesocosm experiment was to determine how sediment N₂ fluxes changed over time since the last organic matter deposition. We hypothesized that sediment N₂ production would decrease as number of days since the last organic matter deposition event increased. Further, we anticipated that nitrogen fixation would become a more important process over this time and that we would observe net nitrogen fixation rates and/or an increase in the expression of *nifH*.

As we hypothesized, there was a significant decrease in sediment N₂ fluxes as a function of time since organic matter deposition. Because we measured N₂ fluxes at 2 separate temperatures (17 and 22°C) it is tempting to attribute the observed relationship to temperature. However, there was no significant difference between the mean net N₂ flux of the 9 cores across all treatments at 17°C vs. 22°C. Further, extensive measurements at the same site where these sediments were collected in Narragansett Bay have shown that sediment N₂ fluxes are not correlated with temperature (Seitzinger et al. 1984, Fulweiler & Nixon 2012). In fact, the relationship between temperature and both denitrification and nitrogen fixation appears to be complicated and variable, with no clear pattern across systems. In some cases, rates of denitrification increase with temperature (e.g. Boston Harbor and Massachusetts Bay; Nowicki et al. 1997) while in others they decrease (e.g. Chesapeake Bay; Kemp et al. 1990). Similarly, some have reported an increase in nitrogen fixation with temperature (e.g. Jones 1982) while others have

found the opposite (e.g. Fernandez et al. 2011). While we cannot discount the role of temperature completely, it appears that it is not the main driver of the N_2 flux signal we observed.

We did find a significant relationship between number of days since organic matter deposition and net sediment N_2 fluxes. These experimental results confirm our previous work that found that denitrification responds rapidly and positively to organic matter deposition (Fulweiler et al. 2007, 2008). If we extrapolate the relationship between time since organic matter deposition and sediment N_2 fluxes beyond the data, we can estimate a point when the sediments would switch from a net sink of N (denitrification dominated) to a net source (nitrogen fixation dominated). According to this relationship, the switch should occur if no organic matter were to reach the benthos for approximately 320 d. It seems unlikely that no deposition would reach the sediments for almost a year. Thus, these results suggest that other key factors are important in phytoplankton phenology and subsequent bloom deposition. Changes in the timing of phytoplankton blooms can alter organic matter deposition to the benthos in at least 2 ways. First, organic matter deposition to the benthos is considerably reduced from blooms during warmer water temperatures (e.g. Rudnick & Oviatt 1986, Keller et al. 1999). Second, the C:N ratio of phytoplankton bloom deposition is known to change with temperature. Phytoplankton blooms during cold months deposit organic matter with a low C:N while organic matter deposited during warm months has a higher C:N and thus lower nutritional value (Smetacek 1984, Keller & Riebesell 1989). In Narragansett Bay, both the total amount and timing of production has shifted such that the quality and quantity of organic matter deposited to the benthos has changed. It is this combination that we hypothesize is driving the previously observed nitrogen fixation rates. Perhaps we did not observe net nitrogen fixation in this most recent experiment because we gave all treatments the same amount of organic matter and only focused on days since deposition, not changes in the quality of the organic matter. The quality may be particularly important in terms of nitrogen fixation because we know that nitrogen fixers can use a wide variety of carbon sources (Hansen 1993, Rabus et al. 1993, Annachhatre & Suktrakoolvait 2001, Eckford et al. 2002). Because we only measured positive net N_2 fluxes we cannot say how nitrogen fixation responded to these deposition events. For that, we must turn to our molecular results.

Microbial genetics

It was no surprise that we observed the presence of *nifH* in these sediments as the genetic potential to fix nitrogen is widespread in prokaryotic taxa (Zehr et al. 2003) and in estuarine water columns (Jenkins et al. 2004) and sediments (Burns et al. 2002, Moisander et al. 2007). The expressed *nifH* sequences observed in this study were only found within 2 phylogenetic groups (*nifH* Clusters I and III) that contain sulfur and sulfate reducing bacteria. Included in this group are *Desulfovibrio* spp. and *Desulfobacter* spp., which have been shown to fix nitrogen in culture (Sisler & ZoBell 1951, Widdel 1987). Research in seagrass beds (Welsh et al. 1996) and salt marshes (Gandy & Yoch 1988) has also shown sulfate reducers to be primarily responsible for observed rates of nitrogen fixation. Thus, our phylogenetic data indicating sulfate reducers as the dominant nitrogen fixers are consistent with other findings. In addition, the expressed *nifH* sequences we observed phylogenetically group with DNA sequences recently reported from coastal California sediments: Bertics & Ziebis (2010) measured nitrogen fixation indirectly using the acetylene reduction assay and found that nitrogen fixation significantly decreased when sulfate reduction was inhibited. They proposed that the nitrogen fixation rate data and the phylogenetic placement of the sequences provided evidence that sulfate reducers may largely be responsible for their observed nitrogen fixation. While our phylogenetic data are consistent with their findings, we also observed concurrent *nirS* expression. The overlapping expression of these genes in both space and time indicate that both nitrogen fixation and denitrification co-occur within relatively small spatial scales. However, our directly measured sediment N_2 fluxes highlight that denitrification was still the dominant process during this experiment.

The expression of *nirS* was measurable in fewer samples than *nifH* and more difficult to detect. This may be due to the short half-life of *nirS* mRNA (Hartig & Zumft 1999) or low transcript copy levels in the environment (Nogales et al. 2002, Bulow et al. 2008). The majority of expressed *nirS* sequences in Narragansett Bay were found in groups NB1, NB2, and NB3. These are most closely related to *Azoarcus toluolyticus*, a bacterium notable for its ability to both denitrify and fix nitrogen (Zhou et al. 1995). Group NB1 also contains *nirS* sequence CB3-S-16 (DQ675925), one of only 3 archetypes in which mRNA expression was detected in Chesapeake Bay (Bulow et al. 2008), indicating that these microbes may be an important

group of denitrifiers in estuarine ecosystems. Other groups of expressed *nirS* sequences (NB4 and NB5) are related to the metabolically diverse bacteria *Paracoccus denitrificans*, *Roseobacter denitrificans*, and *Ruegeria pomeroyi*. *P. denitrificans* can grow aerobically or as a chemolithoautotroph, using hydrogen and sulfur compounds as possible energy sources. *R. denitrificans* has genes for inorganic sulfur oxidation, dimethyl sulfoxide reduction, and dimethyl sulfoniopropionate methyltransferase, suggesting this, and related microbes from our samples, might have the potential to metabolize sulfur compounds (Swingley et al. 2007).

A conceptual model

The relationship of expressed genes from nitrogen fixing and denitrifying bacteria in Narragansett Bay sediments with those from organisms known to respire sulfur compounds suggest the nitrogen and

sulfur (S) cycles may be mediated by the same organisms. Further, because the timing and quantity of organic matter deposition appears, at least in part, to control the net sediment N_2 flux, the carbon cycle is intimately linked to both N and S cycling. From the results of this study and numerous others we present the following conceptual model which puts forth our hypothesis explaining the co-occurrence of denitrification and nitrogen fixation in heterotrophic marine sediments (Fig. 5).

Canonical denitrification follows 2 possible pathways in heterotrophic systems: direct or coupled nitrification-denitrification. In direct denitrification, water column nitrate diffuses into anaerobic sediments and is reduced to N_2 gas. In coupled nitrification-denitrification, nitrifying bacteria convert ammonium to nitrate in the aerobic surface sediment layer. This nitrate is then available via diffusion for reduction to N_2 by denitrifying bacteria in deeper anoxic sediments. In most coastal ocean sediments (including these), coupled nitrification-denitrification




Fig. 5. Conceptual model explaining the co-occurrence of nitrogen fixation and denitrification in heterotrophic marine sediments. (A) N cycle under typical organic matter loading: (1) Aerobic bacterial decomposition degrades organic matter releasing ammonium and depleting oxygen, (2) ammonium is nitrified to nitrate, (3) nitrate is then denitrified to N_2 gas via coupled nitrification-denitrification, and released into the overlying water, or (4) via direct denitrification, (5) sulfate reduction occurs in deeper anoxic sediment and may be accompanied by (6) low rates of N-fixation. (B) N cycle under organic matter starvation: (1) With little organic matter to degrade, ammonium concentrations are low, aerobic decomposition rates are repressed, and the oxic sediment layer thickness increases, (2) because reduced ammonium is low and the available carbon is recalcitrant, sulfate reducing bacteria expand in the sediment column and (3) begin to fix N at higher rates, (4) increased sulfate reduction produces toxic hydrogen sulfide which inhibits nitrification and shuts down the pathway for coupled nitrification-denitrification, (5) direct denitrification continues as long as nitrate is available, (6) denitrifiers capable of metabolizing toxic sulfur compounds may begin to grow, perhaps in a relationship with the sulfate reducing bacteria. Black lines show dominance of process (solid: high rates; dashed: medium rates; dotted: low or repressed rates); grey dotted lines show proposed processes

is the dominant denitrification pathway (Seitzinger et al. 1984, Nowicki 1994). This is particularly important because rapid and substantial inhibition of nitrification at hydrogen sulfide concentrations commonly occurs in estuarine sediments (Joye & Hollibaugh 1995) and direct denitrification is inhibited at even lower hydrogen sulfide concentrations (Joye 2002). Joye & Hollibaugh (1995) proposed that this inhibition was the explanation for the routine observation of low sediment nitrification and denitrification accompanied by high rates of ammonium regeneration in summer N cycling studies (Kemp et al. 1990).

It is well established that estuarine sediments exhibit high rates of sulfate reduction because they are rich in organic matter, exposed to high concentrations of sulfate in seawater, and are anaerobic beneath the top few centimeters (Canfield 1989). Despite recent decreases in water column productivity, changes in bloom phenology, and the subsequent decline in the deposition of fresh organic matter to the benthos, Narragansett Bay sediments are still rich in organic matter. But the quality of organic matter available for microbial decomposition in these sediments has changed from a mean C:N ratio of 7.6 in the 1970s to a mean of 10.5 (range 9.5 to 13.9) (Nixon et al. 2009). Thus, a microbial community able to use this recalcitrant carbon could potentially flourish. Sulfate reducers are notoriously flexible in their ability to use a wide variety of carbon sources (Hansen 1993, Rabus et al. 1993, Annachhatre & Suktrakoolvatt 2001).

Building on these studies, we propose that while both nitrogen fixation and denitrification co-occur in marine sediments, they are not necessarily coupled. Instead, nitrogen fixation mediated by sulfate reducers represses coupled nitrification-denitrification because of the toxic by-product of sulfate reduction, hydrogen sulfide (H_2S). Thus, we hypothesize that sulfate reduction provides energy for the fixation of nitrogen while simultaneously producing a toxic by-product that inhibits competition from other microbes for reduced nitrogen. While nitrogen fixation is an energetically intensive process, much of the energy costs go to protecting nitrogenase from oxygen (Großkopf et al. 2012) and thus in anaerobic sediments the energetic costs must be significantly reduced.

Sediment microbial activity is heterogenic both spatially and temporally. All of the sediment N cycling processes described above vary depending on the *in situ* environmental conditions. Many factors, including inputs of organic matter, will help determine the thickness of the aerobic sediment zone

(Brune et al. 2000). As environmental conditions change, so too will the location of the oxic-anoxic sediment boundary layer. While some nitrifying bacteria are motile, those that are not will be exposed to a variety of conditions within the sediment column. In the marine sediments tested here, the lack of fresh organic matter addition to the benthos over time decreased microbial respiration and, in turn, would have increased the depth of the aerobic boundary layer. In 120 d starvation experiments using cores from mid Narragansett Bay and maintained in the dark at a constant 15°C , Kelly & Nixon (1984) found that oxygen uptake declined at about $0.6\% \text{ d}^{-1}$ while NH_4^+ releases declined at twice this rate, thus increasing the C:N of the remaining organic matter. In the mesocosm experiment described here, as the sediment became starved of fresh organic matter inputs, the competition for labile carbon and available N would have increased. Under these conditions, sulfate reducing bacteria would have the advantage as they have the ability to metabolize refractory carbon (using the N they fix) while concurrently repressing competition for available ammonium by inhibiting nitrifying bacteria. The increase in *nifH* expression, from being undetectable in one replicate and observed highly only at 3.0 cm depth in the other on Day 8, to being medium or high throughout the sediment column on Day 214, supports this hypothesis as does our phylogenetic analysis. Furthermore, over this same period we observed a decrease in net N_2 sediment fluxes which indicates that denitrification rates were decreasing or, perhaps more likely given our genetic data, that nitrogen fixation rates were increasing.

Despite the decrease in net denitrification over the course of this experiment, it was still the overall dominant process. The observed N_2 production could have been from coupled nitrification-denitrification taking place within sediment microzones, from direct denitrification, or from anaerobic oxidation of ammonium (anammox). To date, no studies have reported rates of anammox production in Narragansett Bay. However, it is unlikely that anammox is important in this estuary, as anammox is thought to only account for a small portion of the total N_2 production in shallow systems. For example, anammox only accounted for $6\% (\pm 2\%)$ of the total N_2 production in systems 16 m or shallower (Dalsgaard et al. 2005). Autotrophic denitrification could also have occurred where reduced sulfur compounds were used as the electron donor and nitrate was respired (Batchelor & Lawrence 1978). In fact, this seems likely as *nirS* sequences were closely related to bacteria that have

the genetic machinery to metabolize sulfur compounds. These bacteria may function similarly to *Roseobacter denitrificans*, and use reduced sulfur compounds as electron donors and respire nitrate. It is also possible that there is a syntrophic relationship between these denitrifiers and the sulfate reducing bacteria like that seen between sulfate reducers and methanogens (Hoehler et al. 1994).

Conclusions

Decreases in water column primary production in many coastal systems and throughout the global ocean have recently been reported. In some cases these observations are linked to warming water temperatures and changes in light (Nixon et al. 2009, Boyce et al. 2010); in other cases, these decreases are directly linked to management intervention and a reduction in nutrient loading (Carstensen et al. 2006, Greening & Janicki 2006). These ecosystems will continue to have sediments high in recalcitrant carbon despite the reduction in fresh organic matter. We predict that nitrogen fixation mediated by anaerobic bacteria that respire sulfur compounds will be found in these systems and may be a significant component of the nitrogen budget. The question remains whether nitrogen fixation was a dominant process before anthropogenic changes in climate and nutrient loading altered productivity, or whether these systems are bearing the legacy of human impact in this new and unpredicted way.

Acknowledgements. This material is based on work supported by funding from the Rhode Island Sea Grant College Program to B.D.J, R.W.F., and S.W.N.; as well as the National Science Foundation's Chemical Oceanography division to R.W.F. (OCE 0926859) and B.D.J. and S.W.N (OCE 0926875) and to the Rhode Island EPSCoR program (NSF-0554548 & NSF-1004057). We gratefully acknowledge the assistance of B. Buckley and S. Granger in maintaining the mesocosm facility and carrying out this experiment. We thank R. Twilley (University of Louisiana at Lafayette) who supported R.W.F. during her postdoc at Louisiana State University and allowed her to travel to RI to make these measurements. This article is dedicated to C.R.S.

LITERATURE CITED

- Annachhatre AP, Suktrakoolvait S (2001) Biological sulfate reduction using molasses as a carbon source. *Water Environ Res* 73:118–126
- Batchelor B, Lawrence AW (1978) Autotrophic denitrification using elemental sulfur. *J Water Pollut Control Fed* 50:1986–2001
- Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2009) Genbank. *Nucleic Acids Res* 37:D26–D31
- Bertics VJ, Ziebis W (2010) Bioturbation and the role of microniches for sulfate reduction in coastal marine sediments. *Environ Microbiol* 12:3022–3034
- Boyce DG, Lewis MR, Worm B (2010) Global phytoplankton decline over the past century. *Nature* 466:591–596
- Braker G, Fesefeldt A, Witzel KP (1998) Development of PCR primer systems for amplification of nitrite reductase genes (*nirK* and *nirS*) to detect denitrifying bacteria in environmental samples. *Appl Environ Microbiol* 64:3769–3775
- Braker G, Zhou JZ, Wu LY, Devol AH, Tiedje JM (2000) Nitrite reductase genes (*nirK* and *nirS*) as functional markers to investigate diversity of denitrifying bacteria in pacific northwest marine sediment communities. *Appl Environ Microbiol* 66:2096–2104
- Brune A, Frenzel P, Cypionka H (2000) Life at the oxic–anoxic interface: microbial activities and adaptations. *FEMS Microbiol Rev* 24:691–710
- Bulow SE, Francis CA, Jackson GA, Ward BB (2008) Sediment denitrifier community composition and *nirS* gene expression investigated with functional gene microarrays. *Environ Microbiol* 10:3057–3069
- Burns JA, Zehr JP, Capone DG (2002) Nitrogen-fixing phenotypes of Chesapeake Bay and Neuse River estuary sediments. *Microb Ecol* 44:336–343
- Canfield DE (1989) Sulfate reduction and oxic respiration in marine-sediments—implications for organic-carbon preservation in euxinic environments. *Deep-Sea Res A* 36:121–138
- Capone DG, Budin JM (1982) Nitrogen fixation associated with rinsed roots and rhizomes of the eelgrass *Zostera marina*. *Plant Physiol* 70:1601–1604
- Carstensen J, Conley DJ, Andersen JH, Ærtebjerg G (2006) Coastal eutrophication and trend reversal: a Danish case study. *Limnol Oceanogr* 51:398–408
- Chien YT, Zinder SH (1996) Cloning, functional organization, transcript studies, and phylogenetic analysis of the complete nitrogenase structural genes (*nifHDK2*) and associated genes in the archaeon *Methanosarcina barkeri* 227. *J Bacteriol* 178:143–148
- Cole LW, McGlathery KJ (2012) Nitrogen fixation in restored eelgrass meadows. *Mar Ecol Prog Ser* 448:235–246
- Cornwell JC, Kemp WM, Kana TM (1999) Denitrification in coastal ecosystems: methods, environmental controls, and ecosystem level controls, a review. *Aquat Ecol* 33:41–54
- Dalsgaard T, Thamdrup B, Canfield DE (2005) Anaerobic ammonium oxidation (anammox) in the marine environment. *Res Microbiol* 156:457–464
- Eckford R, Cook FD, Saul D, Aislabie J, Foght J (2002) Free-living heterotrophic nitrogen-fixing bacteria isolated from fuel-contaminated Antarctic soils. *Appl Environ Microbiol* 68:5181–5185
- Edgar RC (2004) MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797
- Farnsworth EJ, Núñez-Farfán J, Careaga SA, Bazzaz FA (1995) Phenology and growth of three temperate forest life forms in response to artificial soil warming. *J Ecol* 83:967–977
- Fernandez C, Farías L, Ulloa O (2011) Nitrogen fixation in denitrified marine waters. *PLoS ONE* 6:e20539

- Fulweiler RW, Nixon SW (2009) Responses of benthic–pelagic coupling to climate change in a temperate estuary. *Hydrobiologia* 629:147–156
- Fulweiler RW, Nixon SW (2012) Net sediment N₂ fluxes in a southern New England estuary: variations in space and time. *Biogeochemistry* 111:111–124
- Fulweiler RW, Nixon SW, Buckley BA, Granger SL (2007) Reversal of the net dinitrogen gas flux in coastal marine sediments. *Nature* 448:180–182
- Fulweiler RW, Nixon SW, Buckley BA, Granger SL (2008) Net sediment N₂ fluxes in a coastal marine system—experimental manipulations and a conceptual model. *Ecosystems* 11:1168–1180
- Gandy EL, Yoch DC (1988) Relationship between nitrogen-fixing sulfate reducers and fermenters in salt-marsh sediments and roots of *Spartina alterniflora*. *Appl Environ Microbiol* 54:2031–2036
- Gardner WS, McCarthy MJ (2009) Nitrogen dynamics at the sediment–water interface in shallow, sub-tropical Florida Bay: why denitrification efficiency may decrease with increased eutrophication. *Biogeochemistry* 95:185–198
- Gardner WS, McCarthy MJ, An SM, Sobolev D, Sell KS, Brock D (2006) Nitrogen fixation and dissimilatory nitrate reduction to ammonium (DNRA) support nitrogen dynamics in Texas estuaries. *Limnol Oceanogr* 51:558–568
- Greening H, Janicki A (2006) Toward reversal of eutrophic conditions in a subtropical estuary: Water quality and seagrass response to nitrogen loading reductions in Tampa Bay, Florida, USA. *Environ Manage* 38:163–178
- Großkopf T, Mohr W, Baustian T, Schunck H and others (2012) Doubling of marine dinitrogen-fixation rates based on direct measurements. *Nature* 488:361–364
- Hansen TA (1993) Carbon metabolisms of sulphate-reducing bacteria. In: Odum J, Singleton R (eds) *The sulphate reducing bacteria: contemporary perspectives*. Springer, New York, NY, p 79–84
- Härtig E, Zumft WG (1999) Kinetics of nirS expression (cytochrome cd1 nitrite reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to denitrification: evidence for a denitrification-specific nitrate- and nitrite-responsive regulatory system. *J Bacteriol* 181:161–166
- Hoehler TM, Alperin MJ, Albert DB, Martens CS (1994) Field and laboratory studies of methane oxidation in an anoxic marine sediment: evidence for a methanogen-sulfate reducer consortium. *Global Biogeochem Cycles* 8:451–463
- Howarth RW, Marino R, Cole JJ (1988a) Nitrogen-fixation in fresh-water, estuarine, and marine ecosystems. 2. Biogeochemical controls. *Limnol Oceanogr* 33:688–701
- Howarth RW, Marino R, Lane J, Cole JJ (1988b) Nitrogen fixation in freshwater, estuarine, and marine ecosystems. 1. Rates and importance. *Limnol Oceanogr* 33:669–687
- Jenkins BD, Steward GF, Short SM, Ward BB, Zehr JP (2004) Fingerprinting diazotroph communities in the Chesapeake Bay by using a DNA macroarray. *Appl Environ Microbiol* 70:1767–1776
- Jones K (1974) Nitrogen fixation in a salt marsh. *J Ecol* 62:553–565
- Jones K (1982) Nitrogen fixation in the temperate estuarine intertidal sediments of the River Lune. *Limnol Oceanogr* 27:455–460
- Joye SB (2002) Denitrification in the marine environment. In: Collins G (ed) *Encyclopedia of environmental microbiology*. John Wiley & Sons, New York, NY, p 1010–1019
- Joye SB, Hollibaugh JT (1995) Influence of sulfide inhibition of nitrification on nitrogen regeneration in sediments. *Science* 270:623–625
- Kahru M, Brotas V, Manzano-Sarabia M, Mitchell BG (2011) Are phytoplankton blooms occurring earlier in the Arctic? *Glob Change Biol* 17:1733–1739
- Kana TM, Darkangelo C, Hunt MD, Oldham JB, Bennett GE, Cornwell JC (1994) Membrane inlet mass spectrometer for rapid high-precision determination of N₂, O₂, and Ar in environmental water samples. *Anal Chem* 66:4166–4170
- Keller AA, Riebesell U (1989) Phytoplankton carbon dynamics during a winter-spring diatom bloom in an enclosed marine ecosystem: primary production, biomass and loss rates. *Mar Biol* 103:131–142
- Keller AA, Oviatt CA, Walker HA, Hawk JD (1999) Predicted impacts of elevated temperature on the magnitude of the winter-spring phytoplankton bloom in temperate coastal waters: a mesocosm study. *Limnol Oceanogr* 44:344–356
- Keller AA, Taylor C, Oviatt C, Dorrington T, Holcombe G, Reed L (2001) Phytoplankton production patterns in Massachusetts Bay and the absence of the 1998 winter-spring bloom. *Mar Biol* 138:1051–1062
- Kelly JR, Nixon SW (1984) Experimental studies of the effect of organic deposition on the metabolism of a coastal marine bottom community. *Mar Ecol Prog Ser* 17:157–169
- Kemp WM, Sampou P, Caffrey J, Mayer M, Henriksen K, Boynton WR (1990) Ammonium recycling versus denitrification in Chesapeake Bay sediments. *Limnol Oceanogr* 35:1545–1563
- Kirshtein JD, Paerl HW, Zehr J (1991) Amplification, cloning, and sequencing of a *nifH* segment from aquatic microorganisms and natural communities. *Appl Environ Microbiol* 57:2645–2650
- Lovley DR, Phillips EJP, Lonergan DJ, Widman PK (1995) Fe(III) and S⁰ reduction by *Pelobacter carbinolicus*. *Appl Environ Microbiol* 61:2132–2138
- Ludwig W, Strunk O, Westram R, Richter L and others (2004) ARB: a software environment for sequence data. *Nucleic Acids Res* 32:1363–1371
- McGlathery KJ, Risgaard-Petersen N, Christensen PB (1998) Temporal and spatial variation in nitrogen fixation activity in the eelgrass *Zostera marina* rhizosphere. *Mar Ecol Prog Ser* 168:245–258
- Moisander PH, Morrison AE, Ward BB, Jenkins BD, Zehr JP (2007) Spatial-temporal variability in diazotroph assemblages in Chesapeake Bay using an oligonucleotide *nifH* microarray. *Environ Microbiol* 9:1823–1835
- Nixon SW, Oviatta CA, Hale S (1976) Nitrogen regeneration and the metabolism of coastal bottom communities. In: Anderson JM, Macfayden A (eds) *The role of terrestrial and aquatic organisms in decomposition processes*. Blackwell Scientific Publications, Oxford, p 269–283
- Nixon SW, Granger S, Buckley BA, Lamont M, Rowell B (2004) A one hundred and seventeen year coastal water temperature record from Woods Hole, Massachusetts. *Estuaries* 27:397–404
- Nixon SW, Fulweiler RW, Buckley BA, Granger SL, Nowicki BL, Henry KM (2009) The impact of changing climate on phenology, productivity, and benthic–pelagic coupling in Narragansett Bay. *Estuar Coast Shelf Sci* 82:1–18
- Nogales B, Timmis KN, Nedwell DB, Osborn AM (2002) Detection and diversity of expressed denitrification

- genes in estuarine sediments after reverse transcription-PCR amplification from mRNA. *Appl Environ Microbiol* 68:5017–5025
- Nowicki BL (1994) The effect of temperature, oxygen, salinity, and nutrient enrichment on estuarine denitrification rates measured with a modified nitrogen gas flux technique. *Estuar Coast Shelf Sci* 38:137–156
- Nowicki BL, Requentina E, VanKeuren D, Kelly JR (1997) Nitrogen losses through sediment denitrification in Boston Harbor and Massachusetts Bay. *Estuaries* 20: 626–639
- Oviatt CA (2004) The changing ecology of temperate coastal waters during a warming trend. *Estuaries* 27:895–904
- Rabus R, Nordhaus R, Ludwig W, Widdel F (1993) Complete oxidation of toluene under strictly anoxic conditions by a new sulfate-reducing bacterium. *Appl Environ Microbiol* 59:1444–1451
- Rudnick DT, Oviatt CA (1986) Seasonal lags between organic carbon deposition and mineralization in marine sediments. *J Mar Res* 44:815–837
- Seitzinger SP, Giblin AE (1996) Estimating denitrification in North Atlantic continental shelf sediments. *Biogeochemistry* 35:235–260
- Seitzinger S, Nixon S, Pilson MEQ, Burke S (1980) Denitrification and N₂O production in near-shore marine sediments. *Geochim Cosmochim Acta* 44:1853–1860
- Seitzinger SP, Nixon SW, Pilson MEQ (1984) Denitrification and nitrous oxide production in a coastal marine ecosystem. *Limnol Oceanogr* 29:73–83
- Sisler FD, ZoBell CE (1951) Nitrogen fixation by sulfate reducing bacteria indicated by nitrogen/argon ratios. *Science* 113:511–512
- Smetacek V (1984) The supply of food to the benthos. In: Fasham M (ed) *Flows of energy and materials in marine ecosystems*. Plenum Press, NATO Conference Series, p 517–547
- Swingley WD, Sadekar S, Mastrian SD, Matthies HJ and others (2007) The complete genome sequence of *Roseobacter denitrificans* reveals a mixotrophic rather than photosynthetic metabolism. *J Bacteriol* 189:683–690
- Teal JM, Valiela I, Berlo D (1979) Nitrogen fixation by rhizosphere and free-living bacteria in salt marsh sediments. *Limnol Oceanogr* 24:126–132
- Walther GR, Post E, Convey P, Menzel A and others (2002) Ecological responses to recent climate change. *Nature* 416:389–395
- Welsh DT, Bourgues S, deWit R, Herbert RA (1996) Seasonal variation in rates of heterotrophic nitrogen fixation (acetylene reduction) in *Zostera noltii* meadows and uncolonised sediments of the Bassin d'Arcachon, southwest France. *Hydrobiologia* 329:161–174
- Widdel F (1987) New types of acetate-oxidizing, sulfate-reducing *Desulfobacter* species, *D. hydrogenophilus* sp. nov., *D. latus* sp. nov., and *D. curvatus* sp. nov. *Arch Microbiol* 148:286–291
- Zehr JP, McReynolds LA (1989) Use of degenerate oligonucleotides for amplification of the *nifH* gene from the marine cyanobacterium *Trichodesmium thiebautii*. *Appl Environ Microbiol* 55:2522–2526
- Zehr JP, Jenkins BD, Short SM, Steward GF (2003) Nitrogenase gene diversity and microbial community structure: a cross-system comparison. *Environ Microbiol* 5:539–554
- Zhou J, Fries MR, Cheesanford JC, Tiedje JM (1995) Phylogenetic analyses of a new group of denitrifiers capable of anaerobic growth on toluene and description of *Azoarcus toluolyticus* sp. nov. *Int J Syst Bacteriol* 45:500–506

*Editorial responsibility: Ronald Kiene,
Mobile, Alabama, USA*

*Submitted: March 26, 2012; Accepted: December 21, 2012
Proofs received from author(s): April 19, 2013*