

Evidence for brain glial activation in chronic pain patients

Marco L. Loggia,^{1,2,*} Daniel B. Chonde,¹ Oluwaseun Akeju,³ Grae Arabasz,¹ Ciprian Catana,¹ Robert R. Edwards,^{2,4} Elena Hill,⁵ Shirley Hsu,¹ David Izquierdo-Garcia,¹ Ru-Rong Ji,^{2,6} Misha Riley,¹ Ajay D. Wasan,^{2,4,7} Nicole R. Zürcher,¹ Daniel S. Albrecht,¹ Mark G. Vangel,¹ Bruce R. Rosen,^{1,8} Vitaly Napadow^{1,2,9} and Jacob M. Hooker¹

Although substantial evidence has established that microglia and astrocytes play a key role in the establishment and maintenance of persistent pain in animal models, the role of glial cells in human pain disorders remains unknown. Here, using the novel technology of integrated positron emission tomography-magnetic resonance imaging and the recently developed radioligand ¹¹C-PBR28, we show increased brain levels of the translocator protein (TSPO), a marker of glial activation, in patients with chronic low back pain. As the Ala147Thr polymorphism in the *TSPO* gene affects binding affinity for ¹¹C-PBR28, nine patient-control pairs were identified from a larger sample of subjects screened and genotyped, and compared in a matched-pairs design, in which each patient was matched to a *TSPO* polymorphism-, age- and sex-matched control subject (seven Ala/Ala and two Ala/Thr, five males and four females in each group; median age difference: 1 year; age range: 29–63 for patients and 28–65 for controls). Standardized uptake values normalized to whole brain were significantly higher in patients than controls in multiple brain regions, including thalamus and the putative somatosensory representations of the lumbar spine and leg. The thalamic levels of TSPO were negatively correlated with clinical pain and levels of circulating proinflammatory cytokines, suggesting that TSPO expression exerts pain-protective/anti-inflammatory effects in humans, as predicted by animal studies. Given the putative role of activated glia in the establishment and or maintenance of persistent pain, the present findings offer clinical implications that may serve to guide future studies of the pathophysiology and management of a variety of persistent pain conditions.

- 1 MGH/MIT/HMS Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA 02129, USA
- 2 Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02155, USA
- 3 Department of Anesthesia, Critical Care and Pain Medicine, MGH/HMS, Boston, MA 02114, USA
- 4 Department of Psychiatry, Brigham and Women's Hospital, HMS, Boston, MA 02155, USA
- 5 Tufts University School of Medicine, Boston, MA 02111, USA
- 6 Departments of Anesthesiology and Neurobiology, Duke University Medical Center, Durham, NC 27705, USA
- 7 Departments of Anesthesiology and Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, PA 15206, USA
- 8 Division of Health Sciences and Technology, Harvard-Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- 9 Department of Biomedical Engineering, Kyung Hee University, Seoul 130-872, Republic of Korea

Correspondence to: Marco L. Loggia,
Massachusetts General Hospital,
Building 149, suite 2301,
Charlestown,
MA 02129
E-mail: marco@nmr.mgh.harvard.edu

Keywords: chronic pain; ^{11}C -PBR28; translocator protein (18kDa); TSPO; neuroinflammation; glia

Abbreviations: LBP = low back pain; SUV = standardized uptake value

Introduction

Until recently, chronic pain has been thought to arise primarily from neuronal dysfunction within nociceptive pathways of the nervous system (Hains and Waxman, 2006). In the last decade, however, a paradigm shift has occurred in the field of pain neurobiology. Animal studies have clearly demonstrated that microglia and astrocytes in the CNS, as well as neuro-glia interactions, play a key role in the establishment and maintenance of persistent pain (Tsuda *et al.*, 2003; Watkins *et al.*, 2007; Calvo *et al.*, 2012; Ji *et al.*, 2013). Both microglia and astrocytes respond to pathological events in CNS, such as strokes, trauma or neurodegenerative diseases, by undergoing a series of cellular responses collectively known as ‘glial activation’ (Gehrmann *et al.*, 1995; Pekny *et al.*, 2014). This response includes proliferation, morphological changes, increased or *de novo* expression of cell surface markers or receptors, and the production of cytokines and other inflammatory mediators. Normally, glial activation is an adaptive defensive mechanism that can contribute to handling acute stress, limiting tissue damage, and restoring homeostasis. However, when malfunctioning (and, in particular when it does not get resolved during the post-acute or early chronic stage after an injury event) (Rolls *et al.*, 2009) glial activation can have deleterious effects, and turn into the primary pathogenic element (Pekny and Pekna, 2014). Several animal studies have now established that glial activation is a key contributing factor in persistent pain. It is well demonstrated that activated microglia and astrocytes can produce cytokines such as tumour necrosis factor alpha (TNFA) and interleukin 1 beta (IL1B), and these cytokines are thought to play an essential role in the pathogenesis of chronic pain (Watkins *et al.*, 2007; Uceyler and Sommer, 2012). It has also been shown that TNFA and IL1B can directly modulate spinal cord synaptic transmission to induce central sensitization and enhance pain states (Kawasaki *et al.*, 2008). Moreover, the intraspinal injection of activated glia produces tactile allodynia, a hallmark of neuropathic pain, in naive rats (Tsuda *et al.*, 2003). Conversely, the injection of drugs inhibiting glial activation can inhibit, delay or reverse pain (Meller *et al.*, 1994; Watkins *et al.*, 1997; Guo *et al.*, 2007; Okada-Ogawa *et al.*, 2009). These observations, alongside other mounting evidence from laboratory models, suggest that chronic pain may result from gliopathy (Ji *et al.*, 2013).

In humans, some observations suggest that activated glia may contribute to the pathophysiology of chronic pain. For instance, post-mortem immunohistochemical studies in the spinal cord of patients with complex regional pain syndrome (Del Valle *et al.*, 2009) and HIV-related neuropathic pain (Shi *et al.*, 2012), as well as CSF sampling in patients

with fibromyalgia and chronic low back pain (LBP) (Brisby *et al.*, 1999; Kadetoff *et al.*, 2012), support a role for glia in chronic pain. Despite these observations, however, no study has yet demonstrated *in vivo* glial activation in humans suffering from chronic pain. Clinically, acknowledging the role of glial activation in pain disorders holds significant promise for the improved diagnostic accuracy of pain disorders. It may also provide a biological basis for the assessment of existing treatments, and the development of novel ones.

In this study we tested the hypothesis that patients with chronic pain demonstrate *in vivo* activation of brain glia. To assess this hypothesis, we imaged the brain of individuals diagnosed with chronic LBP as well as pain-free healthy volunteers using a Siemens 3 T integrated positron emission tomography/magnetic resonance imaging (PET/MRI) scanner and the recently developed PET radioligand, ^{11}C -PBR28 (Brown *et al.*, 2007; Briard *et al.*, 2008). ^{11}C -PBR28 binds to the translocator protein (18kDa) (TSPO), a protein upregulated in activated microglia and reactive astrocytes in animal models of pain (Hernstadt *et al.*, 2009; Wei *et al.*, 2013), and a putative imaging biomarker of inflammation (Cagnin *et al.*, 2007). In addition, as evidence suggests that proinflammatory cytokines are secreted by glial cells in various animal models of pain (Ji *et al.*, 2013), we assessed blood levels of interleukin 6 (IL6), IL1B and TNFA, and evaluated their association with our imaging findings.

Materials and methods

Study design

The study was conducted at the Athinoula A. Martinos Center for Biomedical Imaging at Massachusetts General Hospital. The protocol was approved by the Institutional Review Board and the Radioactive Drug Research Committee.

Subjects

Nineteen patients diagnosed with chronic LBP for at least 2 years (either with or without radicular pain complaints) and 25 healthy controls with no history of chronic pain were initially screened to participate in the study. Individuals were excluded if they had any PET/MRI contraindications (including pregnancy, metallic implants, claustrophobia), had a history of major medical disorders, or were on benzodiazepines or blood thinners (see Supplementary Table 1 for pain history and demographic information).

The Ala147Thr polymorphism in the *TSPO* gene predicts binding affinity for ^{11}C -PBR28, with the Ala/Ala, Ala/Thr and Thr/Thr genotypes being associated with high, mixed and low affinity binding, respectively (Owen *et al.*, 2012; Kreisl *et al.*, 2013; Yoder *et al.*, 2013). Thus, all participants

were tested for this polymorphism and individuals with predicted low-binding affinity (Thr/Thr) were excluded. Given the effect of the TSPO polymorphism on binding affinity for ^{11}C -PBR28, and the fact that the effects of sex and age on ligand binding are unknown, a matched-pairs design was adopted, in which each patient was matched to a TSPO polymorphism-, age- and sex-matched control. Of the 44 subjects initially screened, nine chronic LBP/control matching pairs (with two patients matching the same control) were identified among the initial pool, consisting of seven Ala/Ala and two Ala/Thr pairs, and five male and four female pairs, with a median age difference of 1 year (Supplementary Table 1).

Screening visit

All participants considered potentially eligible after an initial phone screening were recruited to participate in a 2-h characterization and training visit. In this visit, venous blood was drawn in order to have all participants genotyped for the Ala147Thr TSPO polymorphism, and a urine test was performed to ensure that none of the subjects were on benzodiazepines, or taking illegal drugs. At the end of the visit, all participants completed the Beck Depression Inventory (Turner and Romano, 1984; Geisser *et al.*, 1997) and the Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983). In addition, the patients completed the McGill Pain Questionnaire (MPQ), short form (Melzack, 1987).

Imaging visit

On a separate date, eligible participants were invited to participate in an imaging visit. At the beginning of the visit, venous blood was drawn to assess levels of circulating proinflammatory cytokines: IL6, IL1B and TNFA (Supplementary Table 2). These specific proinflammatory cytokines were assessed because they are secreted by glial cells in various animal models of pain (Ji *et al.*, 2013).

Brain imaging was performed with a Siemens PET/MRI scanner (Catana *et al.*, 2008) consisting of a dedicated brain avalanche photodiode-based PET scanner operating in the bore of a 3 T whole-body magnetic resonance scanner equipped with an 8-channel head coil. The use of integrated PET/MRI allowed us to collect structural MRI simultaneously with the PET data. Patients were scanned with ^{11}C -PBR28, a recently developed TSPO radioligand that displays an 80-fold higher *in vivo* specific binding than the earlier generation TSPO radioligand ^{11}C -(R)-PK11195 (Kreisl *et al.*, 2010). ^{11}C -PBR28 was produced in-house using a procedure modified from the literature (Imaizumi *et al.*, 2007).

At the beginning of the imaging visit, we performed a series of magnetic resonance scans, including a multi-echo MPRAGE volume (repetition time/echo time 1/ echo time 2/ echo time 3/ echo time 4 = 2530/1.64/3.5/5.36/7.22 ms, flip angle = 7°, voxel size = 1 mm isotropic) for the purpose of anatomical localization, spatial normalization of the imaging data, as well as generation of attenuation correction maps (Izquierdo-Garcia *et al.*, 2014). The radioligand was then injected as an intravenous bolus, with a median administered dose (interquartile range) of 11.2 (0.3) mCi for patients with chronic LBP and 11.2 (0.6) mCi for controls, and a median specific activity at time of injection of 2.7 (0.8) mCi/nmol for chronic LBP and 2.0 (1.2) mCi/nmol for controls (both values not significantly different across groups). PET data were acquired over the course of 90 min for all but two subjects (see below) and

stored in list-mode format. During the imaging visit, all subjects except one rated their level of pain using a verbal 0–100 numerical ratings scale (pain: 0 = ‘no pain’, 100 = ‘most intense pain tolerable’). The remaining subject (a healthy volunteer) was scanned under a separate protocol, which did not call for ratings of pain before and after the functional runs (but was otherwise identical in the procedures and parameters of the imaging visit).

Data analysis

Using in-house software, standardized uptake values (SUV; i.e. mean radioactivity/injected dose/weight) were computed for each subject. SUVs have been previously used as a measure of TSPO expression in both humans (Hirvonen *et al.*, 2012; Fujita *et al.*, 2013) and rodents (Shao *et al.*, 2013). SUVs were computed voxel-wise from the 60–90 min post-injection PET data, except for two subjects (one patient and one control) for whom SUVs were computed from 72–89 min and 60–86 min, respectively, because of unavailability of the full 60–90 min frame. The two subjects matching these subjects with incomplete data were reconstructed with identical frame onset and duration for the group analyses, to ensure that differences in the PET framing scheme would not affect our results. To maximize accuracy in the reconstruction, SUVs were generated in a two-step iterative process. First, a preliminary SUV image was created for each subject using an attenuation correction map (mu-map) computed from the MPRAGE in its native space (Izquierdo-Garcia *et al.*, 2014). To account for motion that may have occurred between the time the MPRAGE and the 60–90 min PET data were acquired, a new mu-map was created from the MPRAGE registered to this temporary SUV map. This registration was performed using spmregister, a tool from the FreeSurfer suite (<http://surfer.nmr.mgh.harvard.edu>) (Dale *et al.*, 1999) that performs a rigid-body registration of a functional volume to its relative FreeSurfer anatomical volume with Normalised Mutual Information using spm_coreg from the SPM suite (<http://www.fil.ion.ucl.ac.uk/spm/>) (Friston, 2003). A final SUV map was created using this new mu-map, now well in register with the 60–90 min PET data. SUV maps were then normalized to MNI space using non-linear registration (FNIRT, from the FSL suite; FMRIB's Software Library, version 4.1.9, www.fmrib.ox.ac.uk/fsl/) (Smith *et al.*, 2004). Spatially-normalized SUV images were then spatially smoothed (full-width at half-maximum = 8 mm) to improve signal-to-noise ratio, and intensity-normalized to a mean of 1 (SUVs) in order to account for global signal differences across subjects, such as those introduced by differences in the Ala147Thr polymorphism, which affects global binding affinity.

Although the subjects scanned were carefully selected from a larger sample size to match for genetic and demographic factors, the size of our final sample was relatively small. For this reason, imaging and behavioural group comparisons were performed using non-parametric testing. As glial activation has been postulated to spread transynaptically from the site of an injury, following the lesioned neural pathways (Banati, 2003), we hypothesized that in patients with chronic LBP glial activation would be detected in early pain processing regions, as they are most proximal to the pathological sites in the back (e.g. spinal cord/nerve roots). Thus, a thalamus-targeted region of interest analysis was first performed to test this regionally-specific

hypothesis. In this analysis, a matched-pairs test (sign test) was performed on the mean SUVRs extracted from the voxels within the right and left thalamus labels of the Harvard-Oxford Subcortical Structural Atlas (Centre for Morphometric Analyses, http://www.cma.mgh.harvard.edu/fsl_atlas.html), thresholded at the arbitrary value of 30.

We then performed a whole-brain, voxel-wise, matched-pairs analysis with the purpose of identifying (i) which thalamic subregion would be driving the effect in the region of interest analyses (if any was detected); and (ii) additional regions of glial activation within the entire brain. This analysis was conducted using the non-parametric randomize tool from the FSL suite (Nichols and Holmes, 2002), with 10 000 permutations and 5 mm variance smoothing (which increases power with smaller sample sizes: <http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Randomise/UserGuide>). Finally, as exposure to opioids may also lead to a glial reaction *per se* (Watkins *et al.*, 2009), we performed the same SUVR group analysis after the exclusion of the data from the two patients on opioids (and their matching controls; seven versus seven), and limited our search to significant clusters from the original analysis. In all cases, whole-brain voxel-wise group comparisons maps were thresholded using threshold-free cluster enhancement (Smith and Nichols, 2009), using a corrected threshold of $P = 0.05$. For both region of interest and whole-brain voxel-wise analyses, the comparison of nine patient-control pairs was repeated (for consistency sake) in two separate matched-pairs analyses, each using one of the two patients matched to the same control. In the voxel-wise analyses, we first performed a whole-brain search using the data from the patient that best matched the control in terms of age; subsequently the voxel-wise analyses were repeated using the data from the other patient, limiting our search to significant clusters from the first analysis.

Group comparison of behavioural and blood data was performed using the non-parametric sign test or, where applicable, the Fisher's exact test. These analyses were performed using Statistica v.10 (StatSoft). For the exploratory assessment of the relation between imaging and behavioural/blood measures, it was necessary to assess the association between imaging and clinical parameters, adjusting for the TSPO polymorphism. We did this by using multiple regression analysis. To be consistent in our use of non-parametric methods, we used a non-parametric regression method, Generalized Additive Models. To display the relationship between variables in scatter plots, we computed residuals adjusting for the effect of genotype. We performed these analyses using the *gam* library in the R statistics package (<http://www.R-project.org>). Statistical significance was determined by thresholding at $P = 0.05$.

Results

Higher brain TSPO levels in patients with chronic LBP

In the thalamic region of interest analysis, ^{11}C -PBR28 SUVs, normalized to whole brain, (SUVRs) were significantly higher in patients with chronic LBP than controls (left thalamus: $P \leq 0.01$; right thalamus: $P \leq 0.05$; Fig. 1A). The voxel-wise distribution of thalamic SUVRs (Fig. 1B) revealed that in control subjects non-zero

median voxel counts were observed only below values of 1.4, whereas in patients with chronic LBP a substantial number of voxels demonstrated values higher than 1.4 in both hemispheres (medians: 57.5 and 64 in the left and right thalamus, respectively).

The examination of individual thalamic SUVRs (Fig. 1C) shows that, strikingly, each patient exhibited higher SUVRs than his/her age- sex- and *TSPO* genotype-matched control in the thalamus. In all patients, the areas of maximal *TSPO* levels were consistently localized in dorsomedial subregions of the thalamus, as illustrated by the 3D rendering (Fig. 1C).

In the whole-brain voxel-wise analyses, SUVRs were significantly higher in thalamus, pre- and postcentral gyri and paracentral lobule (Fig. 2 and Supplementary Table 3). The peak group difference was observed in left thalamus, consistent with the mediodorsal nucleus. There were no brain regions for which the healthy controls showed statistically higher SUVRs than the patients with chronic LBP. When examined at the exploratory threshold of $P < 0.01$ uncorrected for multiple comparisons (Supplementary Fig. 1), additional regions demonstrated higher SUVRs in chronic LBP than controls, including insulae, middle cingulate cortex, ventromedial prefrontal cortex, posterior cingulate cortex, supplementary motor area and basal ganglia. Moreover, excluding the two patients on opioids (and their matching controls), yielded similar results (Supplementary Table 4). Thus, while exposure to opioids may also lead to a glial reaction (Watkins *et al.*, 2009), our results do not seem to be confounded by opioid intake.

TSPO expression as a protective mechanism?

In the patients, ^{11}C -PBR28 imaging metrics, corrected for genotype, were negatively associated with pain outcome measures (Fig. 3) and circulating levels of proinflammatory cytokines (Fig. 3). First, the number of thalamic voxels with a SUVR value > 1.4 was negatively associated with pain levels at the time of scan, the total score on the McGill Pain Questionnaire ($P < 0.05$), and the levels of IL1B ($P < 0.05$), but not with IL6 or TNFA levels (P -values > 0.3). Second, the average SUVRs extracted from the thalamic cluster statistically significant in the voxel-wise analysis were negatively associated with levels of IL1B ($P < 0.001$) and IL6 ($P < 0.05$), but not with TNFA levels ($P = 0.38$). The association of thalamic [^{11}C]PBR28 SUVR with the McGill Pain Questionnaire total scores was negative, but did not reach statistical significance ($P = 0.11$). No association was observed between this imaging metric and pain levels during the scan ($P = 0.25$).

Discussion

Our study demonstrates the occurrence of glial activation, as measured by an increase in ^{11}C -PBR28 binding, in the brain of patients with chronic pain. Increased tracer




Figure I Evidence for glial activation in the thalamus of chronic LBP patients. (A) Boxplots are presented for the mean ^{11}C -PBR28 SUVRs extracted for all 10 patients with chronic LBP and nine control subjects from the thalamic regions of interest

binding was observed most prominently in the thalamus, and with remarkable across-subject consistency (Fig. 1C).

Within the primary somatosensory and motor cortices (S1/M1) the SUVRs were higher in the putative sensorimotor representations of the lumbar spine [in the postcentral gyrus; Boendermaker *et al.* (2014), and leg (in the paracentral lobule); Loggia *et al.* (2012)]. This spatial pattern is consistent with the majority of patients suffering from pain in the lower back and leg(s), and is suggestive of somatotopically organized glial activation in S1/M1 (which in turn is also consistent with the observation that spinal glial activation generally follows somatotopic boundaries after unilateral spared nerve injury in rats) (Beggs and Salter, 2007).

In the last decade, animal research has led to the increased recognition of the importance of glial cells (such as microglia and astrocytes), and their interaction with neuronal cells, in the pathogenesis of pain conditions (Tsuda *et al.*, 2003; Watkins *et al.*, 2007; Calvo *et al.*, 2012; Ji *et al.*, 2013). For instance, nerve injury induces a profound activation and proliferation of spinal microglia (Liu *et al.*, 2000; Beggs and Salter, 2007; Echeverry *et al.*, 2008; Beggs *et al.*, 2012; Calvo and Bennett, 2012) and the upregulation of a variety of receptors in these cells, such as the adenosine triphosphate (ATP) receptor P2RX4 (also known as P2X₄) (Tsuda *et al.*, 2003) and the chemokine receptor CX3CR1 (Verge *et al.*, 2004), which induce hyperalgesia. Activated microglial cells produce inflammatory mediators, including proinflammatory cytokines and brain-derived neurotrophic factor (BDNF; Coull *et al.*, 2005; Ji and Suter, 2007) that activate or sensitize nociceptive neurons. The intrathecal injection of activated microglia produces tactile allodynia in naive rats, suggesting that microglial activation is sufficient to induce pain sensitization (Tsuda *et al.*, 2003). Finally, pharmacological inhibition of microglial activation prevents or delays neuropathic pain (Raghavendra *et al.*, 2003; Ledebner *et al.*, 2005). Similar to microglia, astrocytes have been shown to have a role in the induction and maintenance of pain sensitization (Ji *et al.*, 2006; Ren and Dubner, 2010; Ji *et al.*, 2013). For instance, in trigeminal models of inflammatory hyperalgesia (Guo *et al.*, 2007) or following trigeminal nerve injury (Okada-Ogawa *et al.*, 2009), astrocytes exhibit hypertrophy, and express enzymes, such as nitric oxide

Figure I Continued

(insert). The *P*-values refer to matched-pairs analyses (sign test) performed using nine chronic LBP-control matching pairs. The analyses were repeated twice, each time using one of the two patients matching the same control, with statistically significant results in both analyses. **P* < 0.05, ***P* < 0.01. (B) Voxel-wise distribution of thalamic SUVRs, showing that patients with chronic LBP have a substantial number of voxels at values ≥ 1.4 (green arrows), whereas controls have a median voxel count of 0. (C) Individual thalamic SUVRs are presented as axial sections (left), and 3D rendering of values higher than the threshold of 1.4 (right). Each row displays SUVRs for each patient-control matched pair. TSPO polymorphism (Ala/Ala or Ala/Thr) is indicated.




Figure 2 Whole-brain voxel-wise analyses. (A) Median SUVR map from healthy controls ($n = 9$) and patients with chronic LBP ($n = 10$) are presented. Matched-pairs tests (nine versus nine) revealed significantly higher TSPO levels in patients, in thalamus, pre- and postcentral gyri, and paracentral lobule ($P < 0.05$ corrected for multiple comparisons; permutation testing, 10 000 permutations). As two patients were matching the same controls, the analyses were performed first using the patient best matching the control. A second analysis was performed using the other patient, limiting our search to significant clusters from the first analysis, with identical results. (B) Boxplots for each of the four regions demonstrating statistically higher SUVRs in patients are shown for illustrative purposes. postc. = postcentral; g. = gyrus; parac. lob. = postcentral lobule; prec. = precentral.




Figure 3 Anti-inflammatory and anti-nociceptive role of TSPO. SUVRs were negatively associated with pain outcomes (left) and blood levels of proinflammatory cytokines (right). The scatterplots show the residuals adjusting for the effect of genotype. MPQ = McGill Pain Questionnaire.

synthase (Meller *et al.*, 1994), as well as inflammatory mediators, such as the proinflammatory cytokine IL1B (Guo *et al.*, 2007) and chemokines such as CXCL2 (Chen *et al.*, 2014), that contribute to hyperalgesia and allodynia. Thermal and mechanical hyperalgesia are inhibited or attenuated by the injection of agents that disrupt either astroglial function (such as fluorocitrate, a glial metabolic inhibitor) (Meller *et al.*, 1994; Watkins *et al.*, 1997; Guo *et al.*, 2007; Okada-Ogawa *et al.*, 2009), or the action of glial products (such as IL1 receptor antagonists) (Watkins

et al., 1997). Taken together, these studies demonstrate that microglia and astrocytes play an important role in the pathogenesis of persistent pain in animals.

Although a plethora of animal studies has demonstrated that glial cells are involved in the establishment and maintenance of persistent pain, no study has previously demonstrated *in vivo* glial activation in humans suffering from chronic pain. Our observations have several potential implications. Firstly, they provide a rationale for exploring the role of glia as therapeutic target for chronic pain. In

animals, drugs that reduce glial activation (e.g. propentofylline and minocycline) have been found to potently inhibit proinflammatory cytokines, thereby suppressing the development of neuropathic pain (Mika, 2008; Leblanc *et al.*, 2011). Importantly, some of these molecules are already FDA approved to treat human conditions of different aetiologies and testing for new, chronic pain-related indications would therefore be immediately possible. Two recent clinical trials (8- and 12-weeks long, respectively) have indicated that low-dose naltrexone (LDN) may have a clinically beneficial impact on fibromyalgia pain (Younger and Mackey, 2009; Younger *et al.*, 2013). As LDN is thought to produce anti-inflammatory effects mainly by antagonizing the activity of glial cells (Mattioli *et al.*, 2010), these studies suggest that glial modulators may be beneficial for fibromyalgia and perhaps other subgroups of chronic pain patients. On the other hand, it should be noted that some clinical trials have had negative results. A recent clinical trial assessing the efficacy of propentophylline to reduce pain in post-herpetic neuralgia was negative (Landry *et al.*, 2012). Another study suggested that perioperative minocycline administration did not improve persistent pain after lumbar discectomy (Martinez *et al.*, 2013). However, methodological concerns with these studies (Watkins *et al.*, 2012) limit the significance of these negative study outcomes. In particular, the duration of trial design was unusually short in both studies. In the former study, propentophylline was administered for 4 weeks, and in the latter minocycline was administered peri-operatively for only 8 days, in both cases a significantly shorter duration than 12 weeks, as is more typically adopted in clinical trials. Moreover, as studies have shown that it is easier to prevent than reverse neuropathic pain using glial modulators (Raghavendra *et al.*, 2003), longer trial durations may be needed to achieve therapeutic efficacy (Watkins *et al.*, 2012). Other factors may also contribute to explain the negative results in these trials, including the dosage and potential interaction with other drugs or food intake (which may not have allowed for the drug to reach CNS sites at meaningful levels), or the choice of patient population (particularly for the propentofylline trial, as the pre-clinical evidence in support of a role for glia in post-herpetic neuralgia is more tenuous than in other chronic pain conditions) (Watkins *et al.*, 2012).

The possibility to image pain-related glial activation *in vivo*, which we document in the present study, may help to identify patients most likely to benefit from this therapeutic approach, and to identify optimal treatment duration or dosage. Given the putative role of activated glia in many challenging issues associated with pain management, such as the induction of opioid-induced hyperalgesia and tolerance (Eidson and Murphy, 2013; Ferrini *et al.*, 2013), the present findings offer clinical implications that may serve to guide future studies of the pathophysiology and management of a variety of persistent pain conditions.

In this study, glial activation was assessed using brain levels of TSPO, formerly called peripheral benzodiazepine

receptor (Banati *et al.*, 1997; Papadopoulos *et al.*, 2006). As experimental animal models and human post-mortem studies of CNS disorders have reliably shown concomitant and co-localized increases in TSPO expression and markers for activated astrocytes and/or microglia, TSPO expression is widely acknowledged as a marker of glial activation in CNS injury and disease. For instance, co-localization of glial activation and TSPO upregulation was observed in rodent models of experimental autoimmune encephalomyelitis, in rodent models of multiple sclerosis, as well as in human multiple sclerosis lesions (Vowinckel *et al.*, 1997; Banati *et al.*, 2000; Chen *et al.*, 2004; Chen and Guilarte, 2006; Ji *et al.*, 2008; Cosenza-Nashat *et al.*, 2009; Abourbeh *et al.*, 2012), in non-human primate models of HIV encephalitis (Venneti *et al.*, 2007; Cosenza-Nashat *et al.*, 2009), as well as in human HIV encephalitis (Cosenza-Nashat *et al.*, 2009), in both human post-mortem and experimental rodent models of ischaemia (Rojas *et al.*, 2007; Cosenza-Nashat *et al.*, 2009; Martin *et al.*, 2010) and Alzheimer's disease (Ji *et al.*, 2008; Cosenza-Nashat *et al.*, 2009; Gulyas *et al.*, 2009), and in rodent models of ethanol and trimethyl neurotoxicity (Kuhlmann and Guilarte, 2000; Maeda *et al.*, 2007). More pertinent to our study, TSPO was upregulated in spinal astrocytes and microglia in Complete Freund's Adjuvant (CFA)-induced monoarthritis of the tibio-tarsal joint (Hernstadt *et al.*, 2009) and following L5 spinal nerve ligation pain (Wei *et al.*, 2013). Given these observations, the increased ¹¹C-PBR28 levels we observed in patients with chronic pain can be interpreted as evidence of glial activation. Interestingly, the involvement of glial subtypes in neuroinflammatory responses seems to depend on the time course of disease. In several animal models, initial TSPO upregulation following acute CNS insult is accompanied by a predominantly microglial response that typically peaks and begins dissipating several days to weeks after injury. This rapid microglial response is paralleled by a delayed but steadily increasing astrocytic component (Kuhlmann and Guilarte, 2000; Chen *et al.*, 2004; Chen and Guilarte, 2006; Martin *et al.*, 2010; Liu *et al.*, 2014). Human post-mortem data seem to corroborate this phase-dependent glial contribution: in acute multiple sclerosis lesions, microglia and macrophages represent most TSPO+ cells, whereas astrocytes are the dominant TSPO+ cells in chronic, silent lesions (Cosenza-Nashat *et al.*, 2009). Because our patients suffered from years of pain, it is plausible that astrocytes provided a significant contribution to the increased PET signal observed in our data.

Although TSPO is a marker of activated glia, a phenomenon thought to be responsible for the amplification of pain signals within the CNS (Ji *et al.*, 2013), TSPO expression itself has been shown in animals to exert inhibitory effects on neuroinflammation (Wei *et al.*, 2013; Bae *et al.*, 2014; Wang *et al.*, 2014), and to promote recovery from neuropathic pain (Wei *et al.*, 2013), likely through the stimulation of steroidogenesis (Batarseh and Papadopoulos, 2010; Wei *et al.*, 2013). In fact, studies suggest that one of the

functions of TSPO in activated glia is to limit the magnitude of inflammatory responses after their initiation (Wang *et al.*, 2014). Our observation that ^{11}C -PBR28 SUVRs negatively correlate with levels of circulating proinflammatory cytokines and pain further corroborate the hypothesis that TSPO expression has anti-inflammatory and pain-protective effects. The negative correlations found in our study therefore support the contention that TSPO ligands may be a novel therapeutic target for the treatment of pathological pain (Wei *et al.*, 2013), as previously suggested for a variety of conditions (Rupprecht *et al.*, 2009, 2010). As TSPO expression is upregulated in activated glia and was found to negatively correlate with peripheral cytokines and pain in our study, low levels of TSPO might be interpreted differently between groups. For example, in healthy volunteers, low levels could simply reflect low levels of glial activation. On the other hand, patients with chronic pain exhibiting lower TSPO levels may have impairments in TSPO expression in activated glia, and therefore in their ability to limit the glial responses after their initiation. Clearly, additional studies are required to elucidate the relationship between ^{11}C -PBR28 and peripheral markers of inflammation, particularly as peripheral cytokine levels are often found to be decoupled from those within the CNS (Bromander *et al.*, 2012).

Future studies, including some currently already underway in our laboratory, will need to determine whether different pain populations present differences in the spatial distribution of glial activation. The discovery of 'glial signatures' of chronic pain states might lead to the identification of objective imaging markers that could (i) complement the patient's subjective assessment and other measures (e.g. quantitative sensory testing) to guide clinical practice; and (ii) reduce the patient heterogeneity which has traditionally led to poor signal-to-noise ratios in most clinical drug evaluation studies (Gomez-Mancilla *et al.*, 2005). Finally, as glial cells respond to very subtle changes in their microenvironment that even precede pathological changes that are detectable histologically (de Vries *et al.*, 2006; Cagnin *et al.*, 2007), glial activation might be an early marker of the alterations that have been shown to occur in the brains of chronic pain patients (Tracey and Bushnell, 2009). This might allow early identification of individuals at risk of transitioning from acute to chronic pain, thus optimizing treatment strategies.

In sum, our findings demonstrate a role of glia in human pain disorders, support the role of the assessment of glial activation and TSPO expression in selective brain areas as an imaging marker and potential treatment target for chronic pain disorders in humans.

Acknowledgements

We would like to thank Patricia McCarthy, Nancy Shearer and the Massachusetts General Hospital Clinical Research Centre for providing nursing support, Drs Douglas Greve,

Marc Normandin, Nicolas Guehl and Jean Logan for help with data processing, Dr Julie Price for helpful comments, Dr Nazem Atassi for sharing imaging resources, Drs Randy Gollub and Alexander Guimaraes for their help with obtaining informed consent, Ms Jiaxuan (Jessie) Wang for her support with the study logistics and Ms Ekaterina Protsenko for editorial assistance.

Funding

This project was supported by Harvard Catalyst Advanced Imaging Pilot Grant (J.M.H.), 1R21NS087472-01A1 (M.L.L.) 1UL1TR001102-01, 8UL1TR000170-05, from the National Centre for Advancing Translational Science, and 1UL1RR025758-04 from the National Centre for Research Resources, Harvard Clinical and Translational Science Centre, and financial contributions from Harvard University and its affiliated academic healthcare centres. The content is solely the responsibility of the authors and does not necessarily represent the official views of Harvard Catalyst, Harvard University and its affiliated academic healthcare centres, or the National Institutes of Health. The authors have no conflicts of interest or competing financial interests to declare.

Supplementary material

Supplementary material is available at *Brain* online.

References

- Abourbeh G, Theze B, Maroy R, Dubois A, Brulon V, Fontyn Y, et al. Imaging microglial/macrophage activation in spinal cords of experimental autoimmune encephalomyelitis rats by positron emission tomography using the mitochondrial 18 kDa translocator protein radioligand [(1)(8)F]DPA-714. *J Neurosci* 2012; 25; 32: 5728–36.
- Bae KR, Shim HJ, Balu D, Kim SR, Yu SW. Translocator protein 18 kDa negatively regulates inflammation in microglia. *J Neuroimmune Pharmacol* 2014; 9: 424–37.
- Banati RB, Myers R, Kreutzberg GW. PK ('peripheral benzodiazepine')—binding sites in the CNS indicate early and discrete brain lesions: microautoradiographic detection of [3H]PK11195 binding to activated microglia. *J Neurocytol* 1997; 26: 77–82.
- Banati RB, Newcombe J, Gunn RN, Cagnin A, Turkheimer F, Heppner F, et al. The peripheral benzodiazepine binding site in the brain in multiple sclerosis: quantitative *in vivo* imaging of microglia as a measure of disease activity. *Brain* 2000; 123(Pt 11): 2321–37.
- Banati RB. Neuropathological imaging: *in vivo* detection of glial activation as a measure of disease and adaptive change in the brain. *Br Med Bull* 2003; 65: 121–31.
- Batarseh A, Papadopoulos V. Regulation of translocator protein 18 kDa (TSPO) expression in health and disease states. *Mol Cell Endocrinol* 2010; 327: 1–12.
- Beggs S, Salter MW. Stereological and somatotopic analysis of the spinal microglial response to peripheral nerve injury. *Brain Behav Immun* 2007; 21: 624–33.
- Beggs S, Trang T, Salter MW. P2X4R + microglia drive neuropathic pain. *Nat Neurosci* 2012; 15: 1068–73.

- Boendermaker B, Meier ML, Luechinger R, Humphreys BK, Hotz-Boendermaker S. The cortical and cerebellar representation of the lumbar spine. *Hum Brain Mapp* 2014; 35: 3962–71.
- Brisby H, Olmarker K, Rosengren L, Cederlund CG, Rydevik B. Markers of nerve tissue injury in the cerebrospinal fluid in patients with lumbar disc herniation and sciatica. *Spine (Phila Pa 1976)* 1999; 24: 742–6.
- Briard E, Zoghbi SS, Imaizumi M, Gourley JP, Shetty HU, Hong J, et al. Synthesis and evaluation in monkey of two sensitive ¹¹C-labeled aryloxyanilide ligands for imaging brain peripheral benzodiazepine receptors *in vivo*. *J Med Chem* 2008; 51: 17–30.
- Bromander S, Anckarsater R, Kristiansson M, Blennow K, Zetterberg H, Anckarsater H, et al. Changes in serum and cerebrospinal fluid cytokines in response to non-neurological surgery: an observational study. *J Neuroinflammation* 2012; 9: 242.
- Brown AK, Fujita M, Fujimura Y, Liow JS, Stabin M, Ryu YH, et al. Radiation dosimetry and biodistribution in monkey and man of ¹¹C-PBR28: a PET radioligand to image inflammation. *J Nucl Med* 2007; 48: 2072–9.
- Cagnin A, Kassiou M, Meikle SR, Banati RB. Positron emission tomography imaging of neuroinflammation. *Neurotherapeutics* 2007; 4: 443–52.
- Calvo M, Bennett DL. The mechanisms of microgliosis and pain following peripheral nerve injury. *Exp Neurol* 2012; 234: 271–82.
- Calvo M, Dawes JM, Bennett DL. The role of the immune system in the generation of neuropathic pain. *Lancet Neurol* 2012; 11: 629–42.
- Catana C, Procissi D, Wu Y, Judenhofer MS, Qi J, Pichler BJ, et al. Simultaneous *in vivo* positron emission tomography and magnetic resonance imaging. *Proc Natl Acad Sci USA* 2008; 105: 3705–10.
- Cosenza-Nashat M, Zhao ML, Suh HS, Morgan J, Natividad R, Morgello S, et al. Expression of the translocator protein of 18 kDa by microglia, macrophages and astrocytes based on immunohistochemical localization in abnormal human brain. *Neuropathol Appl Neurobiol* 2009; 35: 306–28.
- Chen MK, Baidoo K, Verina T, Guilarte TR. Peripheral benzodiazepine receptor imaging in CNS demyelination: functional implications of anatomical and cellular localization. *Brain* 2004; 127(Pt 6): 1379–92.
- Chen G, Park CK, Xie RG, Berta T, Nedergaard M, Ji RR. Connexin-43 induces chemokine release from spinal cord astrocytes to maintain late-phase neuropathic pain in mice. *Brain* 2014; 137(Pt 8): 2193–209.
- Chen MK, Guilarte TR. Imaging the peripheral benzodiazepine receptor response in central nervous system demyelination and remyelination. *Toxicol Sci* 2006; 91: 532–9.
- Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, et al. BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature* 2005; 438: 1017–21.
- Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage* 1999; 9: 179–94.
- Del Valle L, Schwartzman RJ, Alexander G. Spinal cord histopathological alterations in a patient with longstanding complex regional pain syndrome. *Brain Behav Immun* 2009; 23: 85–91.
- Echeverry S, Shi XQ, Zhang J. Characterization of cell proliferation in rat spinal cord following peripheral nerve injury and the relationship with neuropathic pain. *Pain* 2008; 135: 37–47.
- Eidson LN, Murphy AZ. Blockade of Toll-like receptor 4 attenuates morphine tolerance and facilitates the pain relieving properties of morphine. *J Neurosci* 2013; 33: 15952–63.
- Ferrini F, Trang T, Mattioli TA, Laffray S, Del'Guidice T, Lorenzo LE, et al. Morphine hyperalgesia gated through microglia-mediated disruption of neuronal Cl⁻ homeostasis. *Nat Neurosci* 2013; 16: 183–92.
- Friston K. Introduction: experimental design and statistical parametric mapping. In: Frackowiak RSJ, Ashburner JT, Penny WD, Zeki S, Friston K, Frith CD, et al. editors. *Human brain function*. London, UK: Academic Press; 2003.
- Fujita M, Mahanty S, Zoghbi SS, Ferraris Araneta MD, Hong J, Pike VW, et al. PET reveals inflammation around calcified Taenia solium granulomas with perilesional edema. *PLoS One* 2013; 8: e74052.
- Gehrmann J, Matsumoto Y, Kreutzberg GW. Microglia: intrinsic immunoeffector cell of the brain. *Brain Res Brain Res Rev* 1995; 20: 269–87.
- Geisser ME, Roth RS, Robinson ME. Assessing depression among persons with chronic pain using the Center for Epidemiological Studies-Depression Scale and the Beck Depression Inventory: a comparative analysis. *Clin J Pain* 1997; 13: 163–70.
- Gomez-Mancilla B, Marrer E, Kehren J, Kinnunen A, Imbert G, Hillebrand R, et al. Central nervous system drug development: an integrative biomarker approach toward individualized medicine. *NeuroRx* 2005; 2: 683–95.
- Gulyas B, Makkai B, Kasa P, Gulya K, Bakota L, Varszegi S, et al. A comparative autoradiography study in post mortem whole hemisphere human brain slices taken from Alzheimer patients and age-matched controls using two radiolabelled DAA1106 analogues with high affinity to the peripheral benzodiazepine receptor (PBR) system. *Neurochem Int* 2009; 54: 28–36.
- Guo W, Wang H, Watanabe M, Shimizu K, Zou S, LaGraize SC, et al. Glial-cytokine-neuronal interactions underlying the mechanisms of persistent pain. *J Neurosci* 2007; 27: 6006–18.
- Hains BC, Waxman SG. Activated microglia contribute to the maintenance of chronic pain after spinal cord injury. *J Neurosci* 2006; 26: 4308–17.
- Hernstadt H, Wang S, Lim G, Mao J. Spinal translocator protein (TSPO) modulates pain behavior in rats with CFA-induced monoarthritis. *Brain Res* 2009; 1286: 42–52.
- Hirvonen J, Kreisl WC, Fujita M, Dustin I, Khan O, Appel S, et al. Increased *in vivo* expression of an inflammatory marker in temporal lobe epilepsy. *J Nuclear Med* 2012; 53: 234–40.
- Imaizumi M, Kim HJ, Zoghbi SS, Briard E, Hong J, Musachio JL, et al. PET imaging with ¹¹C-PBR28 can localize and quantify upregulated peripheral benzodiazepine receptors associated with cerebral ischemia in rat. *Neurosci Lett* 2007; 411: 200–5.
- Izquierdo-Garcia D, Hansen AE, Förster S, Benoit D, Schachoff S, Fürst S, et al. An SPM8-based approach for attenuation correction combining segmentation and non-rigid template formation: application to simultaneous PET/MR brain imaging. *J Nucl Med* 2014; 55: 1825–30.
- Ji RR, Berta T, Nedergaard M. Glia and pain: Is chronic pain a gliopathy? *Pain* 2013; 154(Suppl 1): S10–28.
- Ji RR, Kawasaki Y, Zhuang ZY, Wen YR, Decosterd I. Possible role of spinal astrocytes in maintaining chronic pain sensitization: review of current evidence with focus on bFGF/JNK pathway. *Neuron Glia Biol* 2006; 2: 259–69.
- Ji B, Maeda J, Sawada M, Ono M, Okauchi T, Inaji M, et al. Imaging of peripheral benzodiazepine receptor expression as biomarkers of detrimental versus beneficial glial responses in mouse models of Alzheimer's and other CNS pathologies. *J Neurosci* 2008; 28: 12255–67.
- Ji RR, Suter MR. p38 MAPK, microglial signaling, and neuropathic pain. *Mol Pain* 2007; 3: 33.
- Kadetoff D, Lampa J, Westman M, Andersson M, Kosek E. Evidence of central inflammation in fibromyalgia-increased cerebrospinal fluid interleukin-8 levels. *J Neuroimmunol* 2012; 242: 33–8.
- Kawasaki Y, Zhang L, Cheng JK, Ji RR. Cytokine mechanisms of central sensitization: distinct and overlapping role of interleukin-1beta, interleukin-6, and tumor necrosis factor-alpha in regulating synaptic and neuronal activity in the superficial spinal cord. *J Neurosci* 2008; 28: 5189–94.
- Kreisl WC, Fujita M, Fujimura Y, Kimura N, Jenko KJ, Kannan P, et al. Comparison of [(11)C]-(R)-PK 11195 and [(11)C]PBR28, two radioligands for translocator protein (18 kDa) in human

- and monkey: Implications for positron emission tomographic imaging of this inflammation biomarker. *Neuroimage* 2010; 49: 2924–32.
- Kreisl WC, Jenko KJ, Hines CS, Lyoo CH, Corona W, Morse CL, et al. A genetic polymorphism for translocator protein 18 kDa affects both *in vitro* and *in vivo* radioligand binding in human brain to this putative biomarker of neuroinflammation. *J Cereb Blood Flow Metab* 2013; 33: 53–8.
- Kuhlmann AC, Guilarte TR. Cellular and subcellular localization of peripheral benzodiazepine receptors after trimethyltin neurotoxicity. *J Neurochem* 2000; 74: 1694–704.
- Landry RP, Jacobs VL, Romero-Sandoval EA, DeLeo JA. Propentofylline, a CNS glial modulator does not decrease pain in post-herpetic neuralgia patients: *in vitro* evidence for differential responses in human and rodent microglia and macrophages. *Exp Neurol* 2012; 234: 340–50.
- Leblanc BW, Zerah ML, Kadasi LM, Chai N, Saab CY. Minocycline injection in the ventral posterolateral thalamus reverses microglial reactivity and thermal hyperalgesia secondary to sciatic neuropathy. *Neurosci Lett* 2011; 498: 138–42.
- Ledeboer A, Sloane EM, Milligan ED, Frank MG, Mahony JH, Maier SF, et al. Minocycline attenuates mechanical allodynia and proinflammatory cytokine expression in rat models of pain facilitation. *Pain* 2005; 115: 71–83.
- Liu L, Rudin M, Kozlova EN. Glial cell proliferation in the spinal cord after dorsal rhizotomy or sciatic nerve transection in the adult rat. *Exp Brain Res* 2000; 131: 64–73.
- Liu X, Li W, Dai L, Zhang T, Xia W, Liu H, et al. Early repeated administration of progesterone improves the recovery of neuropathic pain and modulates spinal 18kDa-translocator protein (TSPO) expression. *J Steroid Biochem Mol Biol* 2014; 143: 130–40.
- Loggia ML, Edwards RR, Kim J, Vangel MG, Wasan AD, Gollub RL, et al. Disentangling linear and nonlinear brain responses to evoked deep tissue pain. *Pain* 2012; 153: 2140–51.
- Maeda J, Higuchi M, Inaji M, Ji B, Haneda E, Okauchi T, et al. Phase-dependent roles of reactive microglia and astrocytes in nervous system injury as delineated by imaging of peripheral benzodiazepine receptor. *Brain Res* 2007; 1157: 100–11.
- Martin A, Boisgard R, Theze B, Van Camp N, Kuhnast B, Damont A, et al. Evaluation of the PBR/TSPO radioligand [(18)F]DPA-714 in a rat model of focal cerebral ischemia. *J Cereb Blood Flow Metab* 2010; 30: 230–41.
- Martinez V, Szekely B, Lemarie J, Martin F, Gentili M, Ben Ammar S, et al. The efficacy of a glial inhibitor, minocycline, for preventing persistent pain after lumbar discectomy: a randomized, double-blind, controlled study. *Pain* 2013; 154: 1197–203.
- Mattioli TA, Milne B, Cahill CM. Ultra-low dose naltrexone attenuates chronic morphine-induced gliosis in rats. *Mol Pain* 2010; 6: 22.
- Meller ST, Dykstra C, Grzybycki D, Murphy S, Gebhart GF. The possible role of glia in nociceptive processing and hyperalgesia in the spinal cord of the rat. *Neuropharmacology* 1994; 33: 1471–8.
- Melzack R. The short-form McGill Pain questionnaire. *Pain* 1987; 30: 191–7.
- Mika J. Modulation of microglia can attenuate neuropathic pain symptoms and enhance morphine effectiveness. *Pharmacol Rep* 2008; 60: 297–307.
- Nichols TE, Holmes AP. Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Hum Brain Mapp* 2002; 15: 1–25.
- Okada-Ogawa A, Suzuki I, Sessle BJ, Chiang CY, Salter MW, Dostrovsky JO, et al. Astroglia in medullary dorsal horn (trigeminal spinal subnucleus caudalis) are involved in trigeminal neuropathic pain mechanisms. *J Neurosci* 2009; 29: 11161–71.
- Owen DR, Yeo AJ, Gunn RN, Song K, Wadsworth G, Lewis A, et al. An 18-kDa translocator protein (TSPO) polymorphism explains differences in binding affinity of the PET radioligand PBR28. *J Cereb Blood Flow Metab* 2012; 32: 1–5.
- Papadopoulos V, Baraldi M, Guilarte TR, Knudsen TB, Lacapere JJ, Lindemann P, et al. Translocator protein (18kDa): new nomenclature for the peripheral-type benzodiazepine receptor based on its structure and molecular function. *Trends Pharmacol Sci* 2006; 27: 402–9.
- Pekny M, Pekna M. Astrocyte reactivity and reactive astrogliosis: costs and benefits. *Physiol Rev* 2014; 94: 1077–98.
- Pekny M, Wilhelmsson U, Pekna M. The dual role of astrocyte activation and reactive gliosis. *Neurosci Lett* 2014; 565: 30–8.
- Raghavendra V, Tanga F, DeLeo JA. Inhibition of microglial activation attenuates the development but not existing hypersensitivity in a rat model of neuropathy. *J Pharmacol Exp Ther* 2003; 306: 624–30.
- Ren K, Dubner R. Interactions between the immune and nervous systems in pain. *Nat Med* 2010; 16: 1267–76.
- Rojas S, Martin A, Arranz MJ, Pareto D, Purroy J, Verdaguer E, et al. Imaging brain inflammation with [(11)C]PK11195 by PET and induction of the peripheral-type benzodiazepine receptor after transient focal ischemia in rats. *J Cereb Blood Flow Metab* 2007; 27: 1975–86.
- Rolls A, Shechter R, Schwartz M. The bright side of the glial scar in CNS repair. *Nat Rev Neurosci* 2009; 10: 235–41.
- Rupprecht R, Papadopoulos V, Rammes G, Baghai TC, Fan J, Akula N, et al. Translocator protein (18 kDa) (TSPO) as a therapeutic target for neurological and psychiatric disorders. *Nat Rev Drug Discov* 2010; 9: 971–88.
- Rupprecht R, Rammes G, Eser D, Baghai TC, Schule C, Nothdurfter C, et al. Translocator protein (18 kD) as target for anxiolytics without benzodiazepine-like side effects. *Science* 2009; 325: 490–3.
- Shao X, Wang X, English SJ, Desmond T, Sherman PS, Quesada CA, et al. Imaging of carrageenan-induced local inflammation and adjuvant-induced systemic arthritis with [(11)C]PBR28 PET. *Nucleic Med Biol* 2013; 40: 906–11.
- Shi Y, Gelman BB, Lisinicchia JG, Tang SJ. Chronic-pain-associated astrocytic reaction in the spinal cord dorsal horn of human immunodeficiency virus-infected patients. *J Neurosci* 2012; 32: 10833–40.
- Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 2004; 23(Suppl 1): S208–19.
- Smith SM, Nichols TE. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* 2009; 44: 83–98.
- Vowinckel E, Reutens D, Becher B, Verge G, Evans A, Owens T, et al. PK11195 binding to the peripheral benzodiazepine receptor as a marker of microglia activation in multiple sclerosis and experimental autoimmune encephalomyelitis. *J Neurosci Res* 1997; 50: 345–53.
- de Vries EF, Dierckx RA, Klein HC. Nuclear imaging of inflammation in neurologic and psychiatric disorders. *Curr Clin Pharmacol* 2006; 1: 229–42.
- Tracey I, Bushnell MC. How neuroimaging studies have challenged us to rethink: is chronic pain a disease? *J Pain* 2009; 10: 1113–20.
- Tsuda M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, et al. P2X4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* 2003; 424: 778–83.
- Turner JA, Romano JM. Self-report screening measures for depression in chronic pain patients. *J Clin Psychol* 1984; 40: 909–13.
- Uceyler N, Sommer C. Cytokine-related and histological biomarkers for neuropathic pain assessment. *Pain Manag* 2012; 2: 391–8.
- Venneti S, Wang G, Wiley CA. Activated macrophages in HIV encephalitis and a macaque model show increased [3H](R)-PK11195 binding in a PI3-kinase-dependent manner. *Neurosci Lett* 2007; 426: 117–22.
- Verge GM, Milligan ED, Maier SF, Watkins LR, Naeve GS, Foster AC. Fractalkine (CX3CL1) and fractalkine receptor (CX3CR1) distribution in spinal cord and dorsal root ganglia

- under basal and neuropathic pain conditions. *Eur J Neurosci* 2004; 20: 1150–60.
- Wang M, Wang X, Zhao L, Ma W, Rodriguez IR, Fariss RN, et al. Macrogliamicroglia interactions via TSPO signaling regulates microglial activation in the mouse retina. *J Neurosci* 2014; 34: 3793–806.
- Watkins LR, Hutchinson MR, Johnson KW. Commentary on Landry et al.: “Propentofylline, a CNS glial modulator, does not decrease pain in post-herpetic neuralgia patients: *in vitro* evidence for differential responses in human and rodent microglia and macrophages”. *Exp Neurol* 2012; 234: 351–3.
- Watkins LR, Hutchinson MR, Ledebner A, Wieseler-Frank J, Milligan ED, Maier SF. Norman Cousins Lecture. Glia as the “bad guys”: implications for improving clinical pain control and the clinical utility of opioids. *Brain Behav Immun* 2007; 21: 131–46.
- Watkins LR, Hutchinson MR, Rice KC, Maier SF. The “toll” of opioid-induced glial activation: improving the clinical efficacy of opioids by targeting glia. *Trends Pharmacol Sci* 2009; 30: 581–91.
- Watkins LR, Martin D, Ulrich P, Tracey KJ, Maier SF. Evidence for the involvement of spinal cord glia in subcutaneous formalin induced hyperalgesia in the rat. *Pain* 1997; 71: 225–35.
- Wei XH, Wei X, Chen FY, Zang Y, Xin WJ, Pang RP, et al. The upregulation of translocator protein (18 kDa) promotes recovery from neuropathic pain in rats. *J Neurosci* 2013; 33: 1540–51.
- Younger J, Mackey S. Fibromyalgia symptoms are reduced by low-dose naltrexone: a pilot study. *Pain Med* 2009; 10: 663–72.
- Younger J, Noor N, McCue R, Mackey S. Low-dose naltrexone for the treatment of fibromyalgia: findings of a small, randomized, double-blind, placebo-controlled, counterbalanced, crossover trial assessing daily pain levels. *Arthritis Rheum* 2013; 65: 529–38.
- Yoder KK, Nho K, Risacher SL, Kim S, Shen L, Saykin AJ. Influence of TSPO genotype on ¹¹C-PBR28 standardized uptake values. *J Nuclear Med* 2013; 54: 1320–2.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983; 67: 361–70.