

RESEARCH ARTICLE

Open Access

Evidence for positive selection of taurine genes within a QTL region on chromosome X associated with testicular size in Australian Brahman cattle

Russell E Lyons¹, Nguyen To Loan², Leanne Dierens¹, Marina R S Fortes^{3*}, Matthew Kelly³, Sean S McWilliam¹, Yutao Li¹, Rowan J Bunch¹, Blair E Harrison¹, William Barendse¹, Sigrid A Lehnert¹ and Stephen S Moore^{3*}

Abstract

Background: Previous genome-wide association studies have identified significant regions of the X chromosome associated with reproductive traits in two *Bos indicus*-influenced breeds: Brahman cattle and Tropical Composites. Two QTL regions on this chromosome were identified in both breeds as strongly associated with scrotal circumference measurements, a reproductive trait previously shown to be useful for selection of young bulls. Scrotal circumference is genetically correlated with early age at puberty in both male and female offspring. These QTL were located at positions 69–77 and 81–92 Mb respectively, large areas each to which a significant number of potential candidate genes were mapped.

Results: To further characterise these regions, a bioinformatic approach was undertaken to identify novel non-synonymous SNP within the QTL regions of interest in Brahman cattle. After SNP discovery, we used conventional molecular assay technologies to perform studies of two candidate genes in both breeds. Non-synonymous SNP mapped to Testis-expressed gene 11 (*Tex11*) were associated ($P < 0.001$) with scrotal circumference in both breeds, and associations with percentage of normal sperm cells were also observed ($P < 0.05$). Evidence for recent selection was found as *Tex11* SNP form a haplotype segment of *Bos taurus* origin that is retained within Brahman and Tropical Composite cattle with greatest reproductive potential.

Conclusions: Association of non-synonymous SNP presented here are a first step to functional genetic studies. Bovine species may serve as a model for studying the role of *Tex11* in male fertility, warranting further in-depth molecular characterisation.

Keywords: Bovine, Reproductive trait, Candidate gene, Haplotype analysis, Association study

Background

Indicine cattle (*Bos indicus*) are socially- and economically-important breeds globally and dominate beef production systems in tropical and sub-tropical regions, and are an important resource in developing nations for food security in vulnerable communities [1]. Presence of indicine cattle is largely due to these breeds being better able to tolerate high temperature and humid environments common in these regions. *Bos indicus* also have improved tolerance to ticks and other tropical challenges compared to taurine

breeds (*Bos taurus*) commonly found in temperate regions [2,3]. However, it is well documented that indicine breeds are inferior in terms of reproductive efficiency in comparison to taurine cattle [4-6]. For example, Brahman bulls typically reach puberty at a later age compared to taurine breeds, with downstream impacts on a number of economically-important indicators of herd productivity and profitability including generation interval and calving rates [7-9]. Tropically-adapted composite herds have also been developed with the aim of retaining the beneficial adaptations of indicine breeds for environmental stressors, while integrating taurine genes for production traits including reproductive performance [10-12]. Improving reproductive performance of indicine cattle will provide

* Correspondence: m.fortes@uq.edu.au; s.moore3@uq.edu.au

³Queensland Alliance for Agriculture and Food Innovation, Centre for Animal Science, The University of Queensland, Brisbane, Qld 4072, Australia
Full list of author information is available at the end of the article

positive impacts in terms of both food security and environmental impacts through increased efficiency in livestock production in areas where hunger is an issue and demand for animal protein is increasing.

Fertility of individual sires is an important production and economic trait affecting reproductive performance of the whole herd. However, fertility is not an easily defined trait, comprising a variety of important heritable traits for selection of early maturing animals [13]. Whereas estimates of heritability for female fertility traits are low, heritability estimates for testicular traits are moderate to high and therefore can be used for effective selection [13-15]. For example, scrotal circumference (SC) is an easy to measure and consistent indicator trait for puberty in bulls with heritability across breeds ranging from medium to high [16]. Also, SC is favourably correlated to other traits affecting both male and female reproductive performance, including sperm quality, percentage of normal sperm, semen concentration, body weight, age at first pregnancy, progesterone levels and age at puberty in heifers [15-24]. Genetic selection that uses SC measurements to improve male fertility can impact positively on whole herd reproductive performance. Likewise, percentage of normal sperm (PNS) impacts directly on bull fertility, as it is a meaningful predictor of calf output per bull [25]. Therefore, measurements of SC and PNS are used as indicators in breeding programs for selecting fertile bulls [18,20].

A limitation in the use of SC and PNS as selection tools is that these reproductive traits cannot be measured before bulls reach 12–24 months of age. Time constraint can limit the attractiveness of these traits as selection tools. Identifying genetic markers associated with these traits would potentially overcome this limitation as they can be measured in DNA samples extracted from blood at birth.

Recent genome-wide association studies (GWAS) in Brahman and Tropical Composite cattle has resulted in the identification of a number of genomic regions associated with SC, PNS and other reproductive traits [26-30]. Fortes et al. [29] reported a 30 Mb region from approximately 62 to 92 Mb positions on chromosome X associated with SC in Brahman. This region was also associated with the age at which a bull achieves a scrotal circumference of 26 cm (AGE26), regarded as puberty [27]. This finding was confirmed in a subsequent study on a Tropical Composite population [30], with significant polymorphisms grouped in two discrete regions at 69–77 Mb and 81–92 Mb positions on the X chromosome.

These confirmed QTL regions warrant further investigation and in the current study, two candidate genes, androgen receptor (*AR*) and testis expressed 11 (*Tex11*), were investigated for their association with SC and PNS measurements in Brahman and Tropical Composite

populations. Androgen Receptor was chosen for its proximity to the QTL peak and its well-studied roles during embryogenesis and puberty for male phenotype development and sexual maturation [31]. Androgens, principally 5 α -dihydrotestosterone (DHT) and testosterone, are essential for the maintenance of behaviour and function of male reproduction and exert their effects by interacting with AR. Both native ligands of this receptor, testosterone and DHT bind to AR to activate target gene expression at the transcriptional level. Androgen-AR activity results in the promotion of maintenance and development of male reproduction and male phenotype [31]. The *Tex11* gene is also in the same QTL region in chromosome X [29,30], and has previously been reported to be essential for male meiosis and fertility in humans and pigs [32,33].

The aim of this study was to characterise the variations in these genes including 5'UTR, test their association with male reproductive traits and investigate the origin of favourable haplotypes. Precedence for the importance of polymorphisms in the 5'UTR gene region include SNP within this region of the bovine growth hormone receptor gene, which has a marked effect on beef production traits [34], and the association of a SNP within the 5'UTR of the bovine lactoferrin (*LF*) gene with reproductive parameters and uterine infection in dairy cattle [35]. Identification of novel single nucleotide polymorphisms (SNP) markers was possible through a combination of bioinformatic analyses of genome sequence data of representative Brahmans and target sequence analysis within the study population. Discovered markers were used to test these candidate genes for their association with observed phenotypic variation of SC and PNS across both Brahman and Tropical Composite populations. Haplotypes across the X chromosome were determined to be of *Bos taurus* or *Bos indicus* origin. Haplotypes from across the entire X chromosome were contrasted with *Tex11* haplotypes to verify the origin of haplotypes and search for evidence of recent selection.

Results

SNP discovery using Brahman genomes

Genome sequences from 16 Brahman bulls, sires of the studied population, were initially compared to the published *Bos taurus* UMD3.1 Bovine Genome Assembly on chromosome X between positions 60,168,532 and 93,999,806 bp. This region has previously been shown to contain 2 QTL associated with SC, PNS and other reproductive traits in both Brahman and Tropical Composite bulls [26-30]. Using Samtools [36], 139,409 SNP were identified, including 247 non-synonymous SNP (nsSNP) affecting 95 genes of which 192 SNP (78%) have not been previously reported in the dbSNP

short genetic variations database (NCBI). All new SNP have been entered into dbSNP.

Bioinformatic analysis initially identified 555 SNP across the ~210 kb region of the *AR* genomic region equaling an average of 1 SNP per 380 bp. No SNP were identified within the coding region of the gene, although many SNP or short indels are present surrounding the intron-exon boundaries. For *Tex11* 530 SNP were identified across ~187 kb equaling an average of 1 SNP per 352 bp. Three previously unreported nsSNP were identified within exons 1, 10 and 23. It has previously been estimated that 20–30% of nsSNP affect protein function [37,38]. The virtues of using these SNP as targets for the current study included the ease of assay development and potential functionality of SNP. Additionally, these SNP were located across a large genomic area (135.7 kb) spanning multiple exons of the candidate gene.

AR SNP and indel validation by direct sequencing of animals with extreme SC

Initial screening of selected SNP by PCR and sequencing within 5'UTR, intron-exon boundary regions and 3' UTR of the *AR* gene was performed using small panels of phenotypically extreme SC animals from within the Brahman bull population. The sequencing confirmed that 13 markers (12 SNP and 1 indel) previously identified bioinformatically were represented within the Brahman population. Furthermore, the sequencing

exercise identified 3 markers (2 SNP and 1 indel) that were not observed in bioinformatic analyses (Table 1). Of particular interest were a cluster of 4 polymorphisms (2 SNP and 2 indel) within the 5'UTR region (X:88,621,605-88,621,617 bp). These markers formed three haplotypes subsequently designated as 5AR_{CC}, 5AR_{TTAA} and 5AR_{TTAALONG} (Figure 1).

AR 5'UTR association study

Due to the difficulty in developing a high-throughput assay for this region, only a small number (n = 259) of Brahman bulls were characterised by direct sequencing with haplotype frequencies of 5AR_{CC}, 5AR_{TTAA} and 5AR_{TTAALONG} being 59.1%, 36.3% and 4.6% respectively. Despite the relatively small sample size limiting the power of the subsequent association analysis, statistical analysis based upon the G > T SNP at position 88,621,617 across the 259 animals reveals that the SNP was associated with SC12, SC18 and SC24 with *P* values of 0.011, 0.0128 and 0.0144, respectively. This G > T SNP served as a tag SNP to the haplotypes due to perfect LD between these SNP.

A possible mechanism for the effect of the *AR* 5'UTR polymorphism may be changes to transcription factor binding sites across this region, as demonstrated through *in silico* studies (Table 2). Of particular interest is the presence of 4 putative SRY binding sites in 5AR_{TTAA} compared to only 2 comparable sites in each of 5AR_{CC} and 5AR_{TTAALONG}. SRY is a testis-expressed protein encoded

Table 1 Novel SNP or indels identified in the *AR* gene within the Brahman population using a small sample of high and low SC Brahman bulls

SNP	Position on chr. X ^a	Gene Region	Base change B1-B2	Primer pair [#]
1	88,621,617	5'UTR	C - A	For 5UTR + Rev exon1
2	88,621,614	5'UTR	C - A	"
3	88,621,609*	5'UTR	: - T	"
4	88,621,605	5'UTR	: - T	"
5	88,619,468	Intron 1	G - A	For exon 1 + Rev intron 1
6	88,491,830*	Intron 2	C - T	For Intron 1 + Rev Intron 2
7	88,449,481	Intron 2	T - G	For Intron 2 + Rev Intron 3
8	88,418,830	Intron 4	G - A	For Intron 4 + Rev intron 5
9	88,418,772	Intron 4	C - T	"
10	88,418,724	Intron 4	C - T	"
11	88,418,702	Intron 4	G - A	"
12	88,411,385	Intron 6	A - G	For Intron 5 + Rev intron 6
13	88,411,823	Intron 6	A - G	For Intron 6 + Rev Intron 7
14	88,411,595	Intron 7	C - A	"
15	88,411,276*	Intron 7	T - A	"
16	88,411,702	3 UTR	C - T	For Intron 7 + Rev 3 UTR

^aBased up UMD3.1 Bovine Genome Assembly.

*Not identified in bioinformatic analyses.

[#]In naming of primers, "For" indicated a forward primer and "Rev" indicated a reverse primer.

Table 3 AR and Tex11 SNP association analysis in the Brahman cattle population

Trait	SNP	P	Allele	Effect	SE	%Va
PNS18	AR1_In4	0.1159	T	-3.793	2.408	0.425
PNS18	Tex11_r38k	0.4506	G	2.387	3.153	0.097
PNS18	Tex11_g297d	0.0155	A	6.821	2.809	1.099
PNS18	Tex11_r696h	0.4933	G	2.168	3.168	0.077
PNS24	AR1_In4	0.7291	T	0.5082	1.467	0.014
PNS24	Tex11_r38k	0.0796	G	-3.456	1.968	0.374
PNS24	Tex11_g297d	0.0081	A	4.578	1.723	0.912
PNS24	Tex11_r696h	0.0906	G	-3.43	2.023	0.355
SC12	AR1_In4	0.0001	T	-0.5531	0.1405	3.249
SC12	Tex11_r38k	4.38 x 10 ⁻⁸	G	1.038	0.1881	6.579
SC12	Tex11_g297d	0.4797	A	-0.119	0.1679	0.120
SC12	Tex11_r696h	1.30 x 10 ⁻⁷	G	1.012	0.1902	6.037
SC18	AR1_In4	5.04 x 10 ⁻⁸	T	-0.8811	0.1604	4.820
SC18	Tex11_r38k	8.25 x 10 ⁻¹⁴	G	1.625	0.2142	9.427
SC18	Tex11_g297d	0.0862	A	-0.3299	0.192	0.540
SC18	Tex11_r696h	5.79 x 10 ⁻¹³	G	1.582	0.2165	8.625
SC24	AR1_In4	4.64 x 10 ⁻¹⁰	T	-0.9485	0.1506	5.506
SC24	Tex11_r38k	6.24 x 10 ⁻¹⁴	G	1.544	0.2026	8.390
SC24	Tex11_g297d	0.0617	A	-0.3413	0.1825	0.570
SC24	Tex11_r696h	4.46 x 10 ⁻¹³	G	1.507	0.2052	7.716

Significance (P), allele substitution effect (Effect) for the named allele of each SNP, its standard error (SE) and percentage of additive variance (%Va) explained by 3 SNP genotypes in *Tex11* and 1 SNP in *AR* on Scrotal Circumference (SC) at 12, 18 and 24 months of age and on the percentage of normal sperm (PNS) at 18 and 24 months of age.

these chromosomal segments can have significant effects on body weight or other production traits [44]. Upon examination across the entire genome or entire chromosome X, cattle carrying the favourable *Tex11_r38k* allele for SC were found to have similar *Bos taurus* component when compared to bulls carrying the less favourable allele (2.6% and 2.3% respectively). However, when animals were grouped on the basis of allelic variants at the four SNP (*AR_In4* [G > A], *Tex11_g297d* [A > G], *Tex11_r38k* [A > G], *Tex11_r696h* [A > G]), striking differences were found in the average frequency of *Bos taurus* haplotypes between these groups (Figure 2). The difference between groups was most pronounced around the *Tex11_r38k* and *Tex11_r696h* loci. For a 5 haplotype region (each haplotype containing 10 SNP), animals carrying the G allele, favourable for higher SC, had higher levels of *Bos taurus* haplotypes (67% on average) compared to those bulls carrying the A allele at these loci (only 1.33% on average). A similar but less pronounced effect was observed for *AR_In4* (18% compared to 3%) and *Tex11_g297d* (16% compared to 3%). The increase in the proportion of *Bos taurus* haplotypes at approximately 70–100 Mb on chromosome X encompasses the candidate

Table 4 AR and Tex11 SNP association analysis in the tropical composite population

Trait	SNP	P	Allele	Effect	SE	%Va
PNS18	AR1_In4	0.0127	T	-6.384	2.551	5.646
PNS18	Tex11_r38k	0.0072	G	-6.181	2.29	6.911
PNS18	Tex11_g297d	0.5780	A	-1.655	2.992	0.299
PNS18	Tex11_r696h	0.0063	G	-6.266	2.282	7.169
PNS24	AR1_In4	0.0784	T	-4.137	2.347	2.368
PNS24	Tex11_r38k	0.0002	G	-7.838	2.101	11.102
PNS24	Tex11_g297d	0.0594	A	-5.127	2.715	2.863
PNS24	Tex11_r696h	0.0006	G	-7.31	2.121	9.746
SC12	AR1_In4	5.02 x 10 ⁻⁶	T	-1.427	0.3097	7.976
SC12	Tex11_r38k	4.00 x 10 ⁻¹⁵	G	-2.155	0.2669	23.754
SC12	Tex11_g297d	3.35 x 10 ⁻⁵	A	-1.517	0.3628	7.096
SC12	Tex11_r696h	2.22 x 10 ⁻¹⁶	G	-2.28	0.2691	26.837
SC18	AR1_In4	4.98 x 10 ⁻⁷	T	-1.565	0.3077	10.584
SC18	Tex11_r38k	4.88 x 10 ⁻¹⁴	G	-2.066	0.2674	24.086
SC18	Tex11_g297d	7.45 x 10 ⁻⁴	A	-1.222	0.3604	5.079
SC18	Tex11_r696h	1.44 x 10 ⁻¹⁴	G	-2.137	0.2705	26.009
SC24	AR1_In4	1.66 x 10 ⁻⁷	T	-1.548	0.2922	10.772
SC24	Tex11_r38k	3.03 x 10 ⁻¹²	G	-1.839	0.2579	19.853
SC24	Tex11_g297d	0.0066	A	-0.9469	0.3471	3.173
SC24	Tex11_r696h	2.18 x 10 ⁻¹²	G	-1.865	0.2597	20.607

Significance (P), allele substitution effect (Effect) for the named allele of each SNP, its standard error (SE) and percentage of additive variance (%Va) explained by 3 SNP genotypes in *Tex11* and 1 SNP in *AR* on Scrotal Circumference (SC) at 12, 18 and 24 months of age and on the percentage of normal sperm (PNS) at 18 and 24 months of age.

genes and the previously characterised QTL region associated with SC, PNS and other reproductive traits in both Brahman and Tropical Composite bulls. This suggests that this region of *Bos taurus* origin may have been retained in Brahmans and Tropical Composites through positive selection due to its favourable phenotypic outcomes including larger SC related to younger age at puberty and higher fertility.

Conclusions

In this study, two putative candidate genes (*AR* and *Tex11*) were investigated for their association with SC and PNS measurements, recognised fertility traits in Brahman and Tropical Composite bulls. These genes were selected for this study because of their position, which mapped to reported QTL [29,30], and because of their known functions. Our results provide strong evidence for the biological role played by *AR* and in particular *Tex11* in male reproduction in cattle. These two genes emerge as strong candidate genes for explaining SC and PNS variation in tropically adapted cattle breeds such as Brahman and Tropical Composite, with varying degrees of taurine and indicine content. Evidence for




Figure 2 Distribution of the proportion of *B. taurus* alleles across the X chromosome in the Brahman bull population based upon grouping based upon candidate marker alleles.

recent positive selection of a favourable *Bos taurus* haplotype was shown and potential causative mutations were described, such as the nsSNP in *Tex11* and the 5'UTR SNP in *AR* that might affect SRY transcription factor binding sites. Knowledge about causative mutations merits confirmation and functional studies. These causative mutations can be used to improve animal breeding and selection as well as being potential models for male fertility issues in mammals.

Methods

The flow of procedures that formed this study is summarized in the illustration provided (Figure 3) and detailed below.

Animals


Animal Care and Use Committee approval was not required for this study because the data were obtained from existing phenotypic databases and DNA storage banks as described below.

Data from 1,118 Brahman bulls born between 2004 and 2008 were available for the current study. These bulls were the progeny of 55 sires from a larger Cooperative Research Centre for Beef Genetic Technologies (Beef CRC) breeding experiment to investigate genetics associated with

whole herd profitability [12,45-48]. Data from 1,085 Tropical Composite bulls born between 2007 and 2009, developed with combinations of Belmont Red, Charbray, Santa Gertrudis and Senepol breeds that represent a genotype with 50% tropically adapted and 50% unadapted genetics were available to use in the current study [47]. These bulls were the progeny of 56 sires also from the Beef CRC. Experimental design, trait measurement details and quantitative genetic analyses of these Brahman and Tropical Composite populations have been described previously [12,47].

Phenotypic data

Traits utilized in this study are components of male fertility: scrotal circumference (SC) at 12 (SC12), 18 (SC18) and 24 (SC24) months and percentage of normal sperm (PSN) at 18 (PNS18) and 24 (PNS24) months. These traits were measured at the bull breeding soundness evaluations (BBSE). Scrotal circumference was measured in centimetres using a standard metal tape [49]. As part of BBSE, bulls having a SC of 20 cm or more were subjected to electro-ejaculation. The percentage of normal sperm (PNS) was based on sperm morphology assessment which was carried out by a laboratory technician accredited by the Australian Cattle Veterinarians [29,30,49].



Bioinformatic analyses of genome sequences

As part of the larger Beef CRC project, the genomes of sixteen Brahman bulls were sequenced at CSIRO Animal, Food and Health Sciences in St Lucia (unpublished data). These bulls were from the cohort of 55 sires used to generate the population used in GWAS analyses [26-28,30,42]. In brief, DNA from all bulls were size selected to the range 400–600 bp, DNA libraries for sequencing by reversible terminator chemistry were prepared for each animal separately using the standard Illumina kit, and 100 bp paired-end reads were collected using an Illumina HiScan sequencer. Raw data reads were analysed using the CASAVA 1.8 software package (www.illumina.com/software/genome_analyzer_software.ilmn), where reads were matched to the UMD3.1 Bovine Genome Assembly. A single BAM format file was created for each sire.

Using Samtools [36], raw genomic data covering the 60–94 Mb region of chromosome X were extracted for each of the 16 animals as individual bam files, and all were aligned to the *Bos taurus* UMD3.1 Bovine Genome Assembly. Variant calls were outputted in VCF format. Variant Effect Predictor (VEP) was used to predict the functional consequences of detected variants [50]. *In silico* prediction of DNA transcription factor binding sites was performed using TFSEARCH [51].

Androgen receptor gene characterisation

The AR gene is highly conserved across populations, with no nsSNP detected by bioinformatic analyses. Classic PCR and sequencing methodologies across exon-intron boundaries were used to confirm novel non-coding SNP that may be experimentally valuable for further characterisation of this region in studied populations. For sequencing of the AR gene, DNA samples were selected from 17 bulls representing the extreme phenotypes for SC. Sequencing was performed for 10 animals with the highest SC measurements, high SC, and 7 animals with the lowest SC measurements, low SC. Primer design used the *Bos taurus* reference gene sequence (ENSBTAG00000022255). Primer pairs were designed to amplify a number of intronic regions immediately up- or downstream of known exons, with a full list of primer pairs in Additional file 3: Table S3.

PCR amplification was performed in an Eppendorf Master Cycler Gradient EP-S thermal cycler. PCR reactions (25 µl) contained 50–100 ng DNA, 1x GoTaq® PCR master mix (Promega, Australia) and 0.5 µM of each primer. The reaction profile used was 94 °C for 2 mins, 35 cycles of 94°C for 10 s, primer-dependent annealing temperature (Additional file 3: Table S3) for 20s and 72°C for 2 mins, followed by a final extension at 72°C for 10 mins. PCR products were separated on 1% agarose gels with 1xTAE (Tris-acetate-EDTA) and visualized on UV transilluminator. PCR products with the expected size range (Additional file 3: Table S3) were sequenced with the forward and reverse primers using BIG DYE 3.1 terminator mix on an ABI 3130xl Genetic Analyser (Applied Biosystems, CA, USA). DNA sequence data were analyzed using Sequencher™ 4.1 (GeneCodes, MI, USA).

For a number of PCR products, deletions predicated the need for cloning of products prior to sequencing. When necessary, PCR products from individual animals were excised from the gel and purified using the QIAquick gel extraction kit (QIAGEN, Doncaster, VIC, Australia). PCR products were then cloned into pGem-Teasy vector (Promega, Australia). Putative positive clones were sequenced with M13 forward and reverse primers.

Genotyping of selected SNP

Custom TaqMan® assays were developed for a novel non-coding SNP in the AR gene (AR_In4) located in the

fourth intron, as well as 3 novel nsSNP spanning the *Tex11* gene (Tex11_r38k, Tex11_g297d, Tex11_r696h). Primers and probes were developed using the Custom TaqMan® Array Design Tool, and are listed in Additional file 1 Table S1.

Genotyping was performed by allelic discrimination using custom TaqMan® SNP genotyping assays, following the manufacturer's instructions. Briefly, 5 µl PCR reactions were carried out containing 2.5 µl of TaqMan® Universal PCR Master Mix (Applied Biosystems, New Jersey, USA), 10 ng of DNA template and 0.25 µl of TaqMan® assay primers and FAM/VIC labelled probes by Applied Biosystems as Assays-by-Design™ (Applied Biosystems, Foster City, CA, USA). All thermal cycling experiments were performed in 384 well plates on a Gene Amp 9700 (Applied Biosystems). Amplification conditions consisted of 50°C for 2 min, 95°C for 10 min followed by 40 cycles of 95°C for 30 s and 60°C for 1 min, and finally 25°C until removed from the thermal cycler. End-point reads were then performed on the Applied Biosystems ViiA™ 7 Real-Time PCR System, and allelic discrimination analysis was performed using ViiA™ 7RUO software (Life Technologies, CA, USA).

Statistical analyses

The association of each SNP with SC12, SC18, SC24, PNS18 and PNS24 was examined for genotyped Brahman bulls using a mixed model analysis of variance with ASREML software [52]. The analysis performed was similar to those performed in previous GWAS and quantitative genetic analyses [12,29,30]. Briefly, the mixed model can be written as follows:

$$y_i = X\beta + Z\mu + S_j a_j + e_i$$

Where y_i represents the phenotypic measurement for the i th animal, X is the incidence matrix relating fixed effects in β with observations in y , Z is the incidence matrix relating to random additive polygenic effects of animal in μ with observations in y and S_j is the observed animal genotype for the j^{th} SNP (coded as 0, 1 or 2 to represent the number of copies of the B allele), a_j is the estimated SNP effect, lastly e_i is the random residual effect. The same fixed effects were used for each trait. These fixed effects included contemporary group (animals born in the same year and raised together) and a second term that was the interaction of year and month of birth.

The percentage of the genetic variance accounted by the j^{th} SNP was estimated according to the formula $\%V_j = 100 \cdot \frac{2p_j q_j a_j^2}{\sigma_g^2}$ where p_j and q_j are the allele frequencies for the j^{th}

SNP estimated across the entire population, a_j is the estimated additive effect of the j^{th} SNP on the trait under analysis, and σ_g^2 is the REML estimate of the (poly-) genetic variance for the trait.

The origin of haplotypes on the X chromosome was determined by using the method described by Bolormaa et al. [44]. This method classifies haplotypes as either *Bos taurus* or *Bos indicus* in origin depending on their relative frequency in a reference population. In brief, haplotypes of 10 consecutive SNP were built and their frequencies calculated in 1,105 Brahman bulls (reference population for *Bos indicus*) and 3,666 Angus, Hereford, Murray Grey and Shorthorn bulls (reference population for *Bos taurus*) of the Beef CRC. The probability that each segment was of *Bos taurus* origin (b_i) was calculated using the following formula:

$$b_i = \frac{p_{Bti}}{p_{Bti} + p_{Bii}}$$

Where p_{Bti} is the frequency of haplotype segment i in *Bos taurus* animals and p_{Bii} is the frequency of haplotype segment i in *Bos indicus* (Brahman) animals. Segments were classified as *Bos taurus* if $b_i > 0.6$ and *Bos indicus* if $b_i < 0.4$. Haplotype segments with b_i between 0.4 and 0.6 were classified as undetermined origin.

Genomic examination of frequency of taurine and indicine specific alleles

To determine whether there was a general increase in the proportion of *Bos taurus* alleles across the genome or X chromosome in animals carrying the G allele of the Tex11_r38k SNP, the percentage of *Bos taurus* in each animal was estimated using either all 50 k autosomal SNP or all X chromosome SNP from the 50 k chip, using Admixture [53]. The data set used to estimate taurine content was 3,666 *Bos taurus* animals (Angus, Shorthorn Murray Grey and Hereford) and 1,032 Brahman bulls, all extracted from The Beef CRC database, with 50 k genotyping detailed before [29,30,54]. The mean and standard deviation of *Bos taurus* proportion was then calculated and used to compare animals with the A or G alleles of Tex11_r38k.

Availability of supporting data

New SNP were submitted to NCBI and the list of SS reference numbers is provided in the additional material section (Additional file 4). Sequence data was submitted to the 1000 bull genomes project, which is a collection of sequence data, intended as a resource for the bovine research community (<http://www.1000bullgenomes.com/>).

Additional files

Additional file 1: Table S1. SNP genotyped and nucleotide sequences of primers and probes used in TaqMan® Assays.

Additional file 2: Table S2. Mean and Standard deviation of estimated *B. taurus* content of animals carrying the A and G allele of Tex11_r36k calculated using different marker sets from full autosomes (40 k) through subset of X chromosome (628).

Additional file 3: Table S3. Forward and reverse primers used to amplify the intron-exon boundaries, 5'UTR and 3'UTR regions of the AR gene.

Additional file 4: Supplementary list of SS reference numbers for new polymorphisms as provided by NCBI.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

REL, SAL, NTL, SSM, LD and MRSF participated in study design. NTL, LD, REL performed experiments and analysed data. SSMcW provided support in the bioinformatic analysis. MK and YL performed statistical analyses. WB, RJB and BEH contributed genomic data for bioinformatic analyses. REL drafted the manuscript. All authors aided in the manuscript preparation.

Acknowledgements

The authors acknowledge that this research uses resources generated by the Cooperative Research Centre for Beef Genetic Technologies (Beef CRC). In particular, we wish to acknowledge Dr. Richard Holroyd for leading the experiments that created the Beef CRC male fertility dataset and Ms Bronwyn Venus for the sperm morphology assessment. Additional financial support for genotyping Brahman and Tropical Composite bulls was provided by Meat and Livestock Australia (projects: B.NBP.0723 and B.NBP.0604).

Author details

¹CSIRO, Animal, Food and Health Sciences, Queensland Bioscience Precinct, Brisbane, Qld 4067, Australia. ²School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Qld 4067, Australia. ³Queensland Alliance for Agriculture and Food Innovation, Centre for Animal Science, The University of Queensland, Brisbane, Qld 4072, Australia.

Received: 11 October 2013 Accepted: 18 December 2013

Published: 10 January 2014

References

1. Thornton PK: **Livestock production: recent trends, future prospects.** *Philos Trans R Soc B Biol Sci* 2010, **365**:2853–2867.
2. Burrow HM: **Importance of adaptation and genotype x environment interactions in tropical beef breeding systems.** *Animal* 2012, **6**:729–740.
3. Sartori R, Bastos MR, Baruselli PS, Gimenes LU, Ereno RL, Barros CM: **Physiological differences and implications to reproductive management of *Bos taurus* and *Bos indicus* cattle in a tropical environment.** *Soc Reprod Fertil Suppl* 2010, **67**:357–375.
4. Warnick AC: **Reproductive performance.** *Crossbreeding Beef Cattle.* Gainesville: University of Florida Press; 1963.
5. Wythe LD: **Genetic and Environmental Effects on Characters Related to Productive Ability of the American Brahman.** Texas A&M University; 1970.
6. Reynolds WL, DeRouen TM, Meyerhoeffer DC, Bellows RA: **Effect of Percentage Zebu Breeding, Inbreeding and Weight at Different Periods on Calving Percent of Brangus and Africander-Angus Heifers.** *J Anim Sci* 1971, **32**:500–506.
7. Lunstra DD, Ford JJ, Echtenkamp SE: **Puberty in beef bulls: hormone concentrations, growth, testicular development, sperm production and sexual aggressiveness in bulls of different breeds.** *J Anim Sci* 1978, **46**:1054–1062.
8. Lunstra DD, Cundiff LV: **Growth and pubertal development in Brahman-, Boran-, Tuli-, Belgian Blue-, Hereford- and Angus-sired F1 bulls.** *J Anim Sci* 2003, **81**:1414–1426.
9. Abeygunawardena H, Dematawewa CM: **Pre-pubertal and postpartum anestrus in tropical Zebu cattle.** *Anim Reprod Sci* 2004, **82–83**:373–387.
10. Casas E, Thallman R, Cundiff L: **Birth and weaning traits in crossbred cattle from Hereford, Angus, Brahman, Boran, Tuli, and Belgian Blue sires.** *J Anim Sci* 2011, **89**:979–987.
11. Burns BM, Fordyce G, Holroyd RG: **A review of factors that impact on the capacity of beef cattle females to conceive, maintain a pregnancy and wean a calf—Implications for reproductive efficiency in northern Australia.** *Anim Reprod Sci* 2010, **122**:1–22.
12. Corbet NJ, Burns BM, Johnston DJ, Wolcott ML, Corbet DH, Venus BK, Li Y, McGowan MR, Holroyd RG: **Male traits and herd reproductive capability in tropical beef cattle. 2. Genetic parameters of bull traits.** *Anim Prod Sci* 2013, **53**:101–113.
13. Cammack KM, Thomas MG, Enns RM: **Review: Reproductive Traits and Their Heritabilities in Beef Cattle.** *Prof Anim Sci* 2009, **25**:517–528.
14. Morris CA, Wilson JA, Bennett GL, Cullen NG, Hickey SM, Hunter JC: **Genetic parameters for growth, puberty, and beef cow reproductive traits in a puberty selection experiment.** *N Z J Agric Res* 2000, **43**:83–91.
15. Evans JL, Golden BL, Bourdon RM, Long KL: **Additive genetic relationships between heifer pregnancy and scrotal circumference in Hereford cattle.** *J Anim Sci* 1999, **77**:2621–2628.
16. Vargas CA, Elzo MA, Chase CC Jr, Chenoweth PJ, Olson TA: **Estimation of genetic parameters for scrotal circumference, age at puberty in heifers, and hip height in Brahman cattle.** *J Anim Sci* 1998, **76**:2536–2541.
17. Smith BA, Brinks JS, Richardson GV: **Relationships of sire scrotal circumference to offspring reproduction and growth.** *J Anim Sci* 1989, **67**:2881–2885.
18. Boligon AA, Silva JAV, Sesana RC, Sesana JC, Junqueira JB, Albuquerque LG: **Estimation of genetic parameters for body weights, scrotal circumference, and testicular volume measured at different ages in Nellore cattle.** *J Anim Sci* 2010, **88**:1215–1219.
19. Martin LC, Brinks JS, Bourdon RM, Cundiff LV: **Genetic effects on beef heifer puberty and subsequent reproduction.** *J Anim Sci* 1992, **70**:4006–4017.
20. Corbet NJ, Burns BM, Corbet DH, Crisp JM, Johnson DJ, McGowan MR, Venus BK, Holroyd RG: **Bull traits measured early in life as indicators of herd fertility.** In *19th Conference of the Association for the Advancement of Animal Breeding and Genetics.* Perth, Australia; 2011. **19**:55–58.
21. Eler JP, Silva J, Evans JL, Ferraz JBS, Dias F, Golden BL: **Additive genetic relationships between heifer pregnancy and scrotal circumference in Nellore cattle.** *J Anim Sci* 2004, **82**:2519–2527.
22. Martinez-Velazquez G, Gregory KE, Bennett GL, Van Vleck LD: **Genetic relationships between scrotal circumference and female reproductive traits.** *J Anim Sci* 2003, **81**:395–401.
23. Moser DW, Bertrand JK, Benyshek LL, McCann MA, Kiser TE: **Effects of selection for scrotal circumference in Limousin bulls on reproductive and growth traits of progeny.** *J Anim Sci* 1996, **74**:2052–2057.
24. Siddiqui MA, Bhattacharjee J, Das ZC, Islam MM, Islam MA, Haque MA, Parrish JJ, Shamsuddin M: **Crossbred bull selection for bigger scrotum and shorter age at puberty with potentials for better quality semen.** *Reprod Domest Anim* 2008, **43**:74–79.
25. Holroyd RG, Doogan W, De Faveri J, Fordyce G, McGowan MR, Bertram JD, Vankan DM, Fitzpatrick LA, Jayawardhana GA, Miller RG: **Bull selection and use in northern Australia. 4. Calf output and predictors of fertility of bulls in multiple-sire herds.** *Anim Reprod Sci* 2002, **71**:67–79.
26. Hawken RJ, Zhang YD, Fortes MRS, Collis E, Barris WC, Corbet NJ, Williams PJ, Fordyce G, Holroyd RG, Walkley JRW, et al: **Genome-wide association studies of female reproduction in tropically adapted beef cattle.** *J Anim Sci* 2012, **90**:1398–1410.
27. Fortes MRS, Lehnert SA, Bolormaa S, Reich C, Fordyce G, Corbet NJ, Whan V, Hawken RJ, Reverter A: **Finding genes for economically important traits: Brahman cattle puberty.** *Anim Prod Sci* 2012, **52**:143–150.
28. Henshall J, Hawken R, Dominik S, Barendse W: **Estimating the effect of SNP genotype on quantitative traits from pooled DNA samples.** *Genet Sel Evol* 2012, **44**:12.
29. Fortes MR, Reverter A, Hawken RJ, Bolormaa S, Lehnert SA: **Candidate genes associated with testicular development, sperm quality, and hormone levels of inhibin, luteinizing hormone, and insulin-like growth factor 1 in Brahman bulls.** *Biol Reprod* 2012, **87**:58.
30. Fortes MRS, Reverter A, Kelly M, McCulloch R, Lehnert SA: **Genome-wide association study for inhibin, luteinizing hormone, insulin-like growth**

- factor 1, testicular size and semen traits in bovine species. *Andrology* 2013, **1**:644–650.
31. Li J, Al-Azzawi F: Mechanism of androgen receptor action. *Maturitas* 2009, **63**:142–148.
 32. Tang L, Zeng W, Clark RK, Dobrinski I: Characterization of the porcine testis-expressed gene 11 (Tex11). *Spermatogenesis* 2011, **1**:147–151.
 33. Yang F, Gell K, van der Heijden GW, Eckardt S, Leu NA, Page DC, Benavente R, Her C, Hoog C, McLaughlin KJ, Wang PJ: Meiotic failure in male mice lacking an X-linked factor. *Genes Dev* 2008, **22**:682–691.
 34. Maj A, Oprzadek J, Oprzadek A, Dymnicki E, Zwierzchowski L: Polymorphism in the 5'-noncoding region of the bovine growth hormone receptor gene and its association with meat production traits in cattle. *Anim Res* 2004, **53**:503–514.
 35. Abolfazl H, Hassan S, Abdolah M, Abbas Rowshan G: Characterization of single nucleotide polymorphism in the 5'-untranslated region (5'-UTR) of Lactoferrin gene and its association with reproductive parameters and uterine infection in dairy cattle. *Vet Res Forum* 2012, **3**:37–43.
 36. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup: The Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics* 2009, **25**:2078–2079.
 37. Sunyaev S, Ramensky V, Koch I, Lathe W 3rd, Kondrashov AS, Bork P: Prediction of deleterious human alleles. *Hum Mol Genet* 2001, **10**:591–597.
 38. Chasman D, Adams RM: Predicting the functional consequences of non-synonymous single nucleotide polymorphisms: structure-based assessment of amino acid variation. *J Mol Biol* 2001, **307**:683–706.
 39. Koopman P, Munsterberg A, Capel B, Vivian N, Lovellbadge R: Expression of a Candidate Sex-Determining Gene during Mouse Testis Differentiation. *Nature* 1990, **348**:450–452.
 40. Gubbay J, Collignon J, Koopman P, Capel B, Economou A, Munsterberg A, Vivian N, Goodfellow P, Lovellbadge R: A Gene-Mapping to the Sex-Determining Region of the Mouse Y-Chromosome Is a Member of a Novel Family of Embryonically Expressed Genes. *Nature* 1990, **346**:245–250.
 41. Yuan X, Lu ML, Li T, Balk SP: SRY interacts with and negatively regulates androgen receptor transcriptional activity. *J Biol Chem* 2001, **276**:46647–46654.
 42. Siadkowska E, Zwierzchowski L, Oprzadek J, Strzałkowska N, Bagnicka E, Krzyżewski J: Effect of polymorphism in IGF-1 gene on production traits in Polish Holstein-Friesian cattle. *Anim Sci Paper Rep* 2006, **24**:225–237.
 43. Fortes MR, Li Y, Collis E, Zhang Y, Hawken RJ: The IGF1 pathway genes and their association with age of puberty in cattle. *Anim Genet* 2013, **44**:91–95.
 44. Bolormaa S, Hayes BJ, Hawken RJ, Zhang Y, Reverter A, Goddard ME: Detection of chromosome segments of zebu and taurine origin and their effect on beef production and growth. *J Anim Sci* 2011, **89**:2050–2060.
 45. Corbet NJ, Burns BM, Corbet DH, Johnston DJ, Crisp JM, McGowan MR, Prayaga KC, Venus BK, Holroyd RG: Genetic variation in growth, hormonal and seminal traits of young tropically adapted bulls. In *18th Conference of the Association for the Advancement of Animal Breeding and Genetics*. Barossa Valley, Australia; 2009. **18**:121–124.
 46. Johnston DJ, Barwick SA, Corbet NJ, Fordyce G, Holroyd RG, Williams PJ, Burrow HM: Genetics of heifer puberty in two tropical beef genotypes in northern Australia and associations with heifer- and steer-production traits. *Anim Prod Sci* 2009, **49**:399–412.
 47. Burns BM, Corbet NJ, Corbet DH, Crisp JM, Venus BK, Johnston DJ, Li Y, McGowan MR, Holroyd RG: Male traits and herd reproductive capability in tropical beef cattle. 1. Experimental design and animal measures. *Anim Prod Sci* 2013, **53**:87–100.
 48. Barwick SA, Wolcott ML, Johnston DJ, Burrow HM, Sullivan MT: Genetics of steer daily and residual feed intake in two tropical beef genotypes, and relationships among intake, body composition, growth and other post-weaning measures. *Anim Prod Sci* 2009, **49**:351–366.
 49. Fordyce G, Entwistle K, Norman S, Perry V, Gardiner B, Fordyce P: Standardising bull breeding soundness evaluations and reporting in Australia. *Theriogenology* 2006, **66**:1140–1148.
 50. McLaren W, Pritchard B, Rios D, Chen YA, Flicek P, Cunningham F: Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. *Bioinformatics* 2010, **26**:2069–2070.
 51. Heinemeyer T, Wingender E, Reuter I, Hermjakob H, Kel AE, Kel OV, Ignatieva EV, Ananko EA, Podkolodnaya OA, Kolpakov FA, *et al*: Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids Res* 1998, **26**:362–367.
 52. Gilmour AR, Cullis BR, Gogel BJ, Welham SJ, Thompson R: *ASReml, user guide. Release 2.0*. Hemel Hempstead: VSN International Ltd; 2006.
 53. Alexander DH, Novembre J, Lange K: Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* 2009, **19**:1655–1664.
 54. Fortes MR, Kemper K, Sasazaki S, Reverter A, Pryce JE, Barendse W, Bunch R, McCulloch R, Harrison B, Bolormaa S, *et al*: Evidence for pleiotropism and recent selection in the PLAG1 region in Australian Beef cattle. *Anim Genet* 2013, **44**:636–647.

doi:10.1186/1471-2156-15-6

Cite this article as: Lyons *et al*: Evidence for positive selection of taurine genes within a QTL region on chromosome X associated with testicular size in Australian Brahman cattle. *BMC Genetics* 2014 **15**:6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

