

ARTICLE

Received 7 Mar 2011 | Accepted 7 Dec 2011 | Published 17 Jan 2012

DOI: 10.1038/ncomms1635

Evidence for the late formation of hydrous asteroids from young meteoritic carbonates

Wataru Fujiya¹, Naoji Sugiura¹, Hideyuki Hotta¹, Koji Ichimura¹ & Yuji Sano²

The accretion of small bodies in the Solar System is a fundamental process that was followed by planet formation. Chronological information of meteorites can constrain when asteroids formed. Secondary carbonates show extremely old ^{53}Mn - ^{53}Cr radiometric ages, indicating that some hydrous asteroids accreted rapidly. However, previous studies have failed to define accurate Mn/Cr ratios; hence, these old ages could be artefacts. Here we develop a new method for accurate Mn/Cr determination, and report a reliable age of $4,563.4 \pm 0.4/-0.5$ million years ago for carbonates in carbonaceous chondrites. We find that these carbonates have identical ages, which are younger than those previously estimated. This result suggests the late onset of aqueous activities in the Solar System. The young carbonate age cannot be explained if the parent asteroid accreted within 3 million years after the birth of the Solar System. Thus, we conclude that hydrous asteroids accreted later than differentiated and metamorphosed asteroids.

¹ Department of Earth and Planetary Science, Graduate School of Science, the University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

² Atmosphere and Ocean Research Institute, the University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa-shi, Chiba 277-8564, Japan. Correspondence and requests for materials should be addressed to W.F. (email: fujiya@eps.s.u-tokyo.ac.jp).

An understanding of the evolution of the asteroid belt can constrain the origin and early history of the Solar System. It is suggested that the accretion time of asteroids determines the degree of metamorphism (and/or melting) they experienced¹, because a short-lived radionuclide of ²⁶Al (half-life: 0.73×10^6 years) is the dominant heat source of metamorphism, and hence, the abundance of the heat source is a function of accretion time. Several lines of evidence from the dating of meteorites have shown that the parent bodies of differentiated meteorites accreted earlier than chondrites^{2,3}. These observations indicate that accretion time is indeed an important controlling factor of asteroidal thermal history, whereas other factors, such as water concentrations in asteroids, could have minor roles. However, the accretion time of hydrous asteroids has been controversial. CM chondrites, which are carbonaceous chondrites named for their type specimen of the Mighei meteorite, experienced aqueous alteration at low temperatures⁴, indicating late accretion. However, the ⁵³Mn–⁵³Cr ages (⁵³Mn decays to ⁵³Cr with a half-life of 3.7×10^6 years) of the carbonates in CM chondrites revealed by secondary ion mass spectrometry (SIMS) are very old (ref. 5 and references therein), with ages comparable to the oldest solids in the Solar System (that is, calcium–aluminium–rich inclusions (CAIs); 4,568.2 million years ago (Myr ago)⁶). Although the ages of the secondary minerals in CI chondrites (named for their type specimen of the Ivuna meteorite) may not be as old (4,567 to 4,554 Myr ago^{7–12}), one must conclude that at least some hydrous asteroids, such as the CM chondrite parent body, accreted very rapidly because the accretion time should be constrained by the oldest alteration product.

It should be noted, however, that the accuracies of Mn–Cr ages depend on accurate determinations of ⁵⁵Mn/⁵²Cr ratios. Accurate ⁵⁵Mn/⁵²Cr measurements of meteoritic carbonates require a carbonate standard with a known chemical composition (that is, ⁵⁵Mn/⁵²Cr atomic ratio) because the sensitivities of Mn and Cr in SIMS strongly depend on analyte chemical compositions (the so-called ‘matrix effect’), and hence, the data obtained for unknown samples should be calibrated using the standard. However, a suitable carbonate standard is not available in natural terrestrial environments, which is why previous studies have used other materials, such as San Carlos olivine, as alternative standards. This use has led to systematic errors on the ⁵⁵Mn/⁵²Cr of meteoritic samples and their ages. To overcome this problem, we synthesized a carbonate (calcite: CaCO₃) standard doped with Mn and Cr, and we demonstrated that the Mn/Cr relative sensitivity of carbonates is significantly different from that of San Carlos olivine¹³, that is, systematic errors of ~2.2 Myr indeed exist in previous studies for carbonate dating. This result is crucial for a high-resolution chronology of the early Solar System.

In the present study, we determined the accurate Mn–Cr ages of carbonates in four CM chondrites using the carbonate standard. We found that the CM carbonates have an identical age of $4,563.4^{+0.4}_{-0.5}$ Myr ago, which is younger than previous estimates. The young carbonate age suggests the late onset of aqueous activities in the Solar System. Based on this result, we conclude that the parent asteroid of CM chondrites accreted at ~3.5 Myr after the birth of the Solar System.

Results

Initial ⁵³Mn/⁵⁵Mn ratios and absolute ages of CM carbonates. We performed Mn–Cr isotope measurements of the standard and carbonates in the CM chondrites with different alteration indexes¹⁴ (calcites in Murchison CM 2.5 and Y791198 CM 2.4 and dolomites in ALH83100 CM 2.1 and Sayama CM 2.1) using a NanoSIMS 50. The Mn/Cr relative sensitivity data are presented in Supplementary Figure S1 and Supplementary Table S1. For all samples, the ⁵³Cr excesses correlate well with the ⁵⁵Mn/⁵²Cr, which indicates the *in-situ* decay of ⁵³Mn (Fig. 1; Table 1). The analyses were performed for several carbonate grains in each meteorite. Hence, it is likely that

the carbonates dispersed in each meteorite formed at the same time within the analytical uncertainties. The initial ⁵³Mn/⁵⁵Mn ratios, (⁵³Mn/⁵⁵Mn)₀, were estimated from the isochron slopes, and the uncertainty in the Mn/Cr relative sensitivity was propagated to the errors on (⁵³Mn/⁵⁵Mn)₀. The (⁵³Mn/⁵⁵Mn)₀ values were found to be $(2.59 \pm 0.74) \times 10^{-6}$, $(3.31 \pm 0.68) \times 10^{-6}$, $(2.71 \pm 0.40) \times 10^{-6}$ and $(3.28 \pm 0.30) \times 10^{-6}$ for the carbonates in Murchison, Y791198, ALH83100 and Sayama, respectively (errors are 2σ). In calculating the absolute ages of carbonates, the LEW 86010 angrite with a known Pb–Pb age¹⁵ and (⁵³Mn/⁵⁵Mn)₀ (ref. 16) was referenced as a time anchor. We assume a homogeneous distribution of ⁵³Mn in the early Solar System, which is likely to be correct at least among samples from chondrites, the eucrite parent body, Earth and Mars¹⁷. The initial ⁵³Mn/⁵⁵Mn ratios of the carbonates correspond to absolute ages of $4,562.6^{+1.4}_{-1.9}$ Myr ago for Murchison, $4,563.9^{+1.1}_{-1.3}$ Myr ago for Y791198, $4,562.8^{+0.8}_{-1.0}$ Myr ago for ALH83100 and $4,563.9^{+0.6}_{-0.7}$ Myr ago for Sayama.

Accuracies of the carbonate ages. It should be stressed that we calculated a Cr isotopic ratio by summing all ⁵³Cr counts, then dividing by the sum of ⁵²Cr counts over the entire measurement, rather than averaging isotopic ratios (Methods). This is an important difference from earlier studies, that is, an isotope ratio has been traditionally calculated by averaging a number of ratios collected over the course of a measurement, which leads to a positive bias especially in the case of low counting rates¹⁸.

The meteoritic carbonates studied in this study have experienced only low temperatures and it is unlikely that thermally driven volume diffusion has disturbed the Mn–Cr system. Also, other mechanisms of redistribution of the parent and daughter isotopes, such as grain boundary diffusion, can be ruled out because the cathodoluminescence observations of the carbonates show well-preserved zoning of trace elements including Mn. Therefore, we are sure that these ages are indigenous and there are no possibilities of age biases caused by the disturbance of the Mn–Cr system.

In summary, we successfully determined the carbonate ages of the CM chondrites, which are free from the potential biases resulting from the matrix-mismatched standard, the calculation of isotopic ratios and/or the disturbance of the Mn–Cr system.

Discussion

It is worth noting that the carbonate ages are younger than those reported previously for the same meteorites. These ages are identical within the errors, irrespective of the degrees of aqueous alteration (Fig. 2). Therefore, we can define a formation age of $4,563.4^{+0.4}_{-0.5}$ Myr ago (that is, ~4.8 Myr after CAI formation) for carbonates in CM chondrites using the weighted average. Our data indicate a later accretion of the CM chondrite parent asteroid, and therefore, later onsets of aqueous alteration in the Solar System than previously inferred from the oldest carbonates in the CM chondrites. The problem of the very early formation of carbonates contemporaneous with CAIs is solved using the proper Mn/Cr relative sensitivity. It is also suggested that the carbonates in the CM chondrites precipitated in a relatively short time. This suggestion is contrary to earlier studies suggesting a rough correlation between alteration indexes and carbonate ages of CM chondrites⁵.

Here we consider the accretion time of the CM chondrite parent body. Our new carbonate ages determine the younger side of the accretion time (~4.8 Myr after CAI formation). The older side could be constrained by the ⁵³Mn–⁵³Cr or ²⁶Al–²⁶Mg ages of chondrules. Unfortunately, chondrules in CM chondrites have not been dated yet, and their dating may be complicated by extensive aqueous alteration. Therefore, we rely on the Al–Mg ages of chondrules in CO chondrites (named for their type specimen of the Ornans meteorite), because they are related to CM chondrites in chemical and oxygen isotopic compositions^{19,20}. Previous studies suggested that the




Figure 1 | Mn–Cr isochron diagrams for carbonates in four CM chondrites. (a) Calcite in the Murchison CM 2.5 chondrite. (b) Calcite in the Y791198 CM 2.4 chondrite. (c) Dolomite in the ALH83100 CM 2.1 chondrite. (d) Dolomite in the Sayama CM 2.1 chondrite. The vertical axis shows $^{53}\text{Cr}/^{52}\text{Cr}$ described as the permil deviation ($\delta^{53}\text{Cr}$) from the value of the synthetic calcite standard. Error bars ($\pm 2\sigma$) on both $^{53}\text{Cr}/^{52}\text{Cr}$ and $^{55}\text{Mn}/^{52}\text{Cr}$ represent the external reproducibility or the internal precision, whichever is larger. MSWD, mean square weighted deviation.

forming events of CO chondrules took place 1.7 to 3.2 Myr after CAI formation^{21,22}. Thus, the accretion time of the CM chondrite parent body would be 3.2 to 4.8 Myr after the birth of the Solar System.

On the basis of the chronological information discussed above, we simulate the thermal history of the CM chondrite parent body and estimate the accretion time. We consider the static water model^{20,23} here, although the fluid flow model has been investigated in some previous works^{24–27}. The thermal history is modelled using a one-dimensional heat conduction equation in radial form. See Supplementary Methods for details. We assume that an asteroid of 60 km in diameter accreted instantaneously at 3.5 Myr (case A) or 3.0 Myr (case B) after CAI formation, although the results are essentially similar for different asteroid sizes, if the asteroid is larger than 60 km in diameter (Supplementary Fig. S2). The parent body consists of a homogeneous mixture of rock, void space and water ice. The heat source is the decay energy of ^{26}Al , and the reaction heat of hydration is also included. The parameters and variables used in the model are presented in Supplementary Table S2 and Supplementary Methods. According to the oxygen isotopic compositions²³ and a simulation of chemistry during aqueous alteration²⁸, the calcite in Murchison is likely to have precipitated at $<25^\circ\text{C}$ in the earliest stage of aqueous alteration (when less than a few percent of rock reacted with water). The clumped-isotope thermometry also suggests that the calcite in Murchison formed at a low temperature ($20\text{--}33^\circ\text{C}$)²⁹. On the other hand, the dolomite in Sayama seems to have formed at a higher temperature ($>120^\circ\text{C}$), as inferred from the small isotopic fractionation of oxygen between the dolomite and the matrix³⁰.

In the model set-up described above, the calculated thermal history of case A well reproduces the temperatures of carbonate formation at $4,563.4^{+0.4}_{-0.5}$ Myr ago (Fig. 3a). In the intermediate region of the asteroid, the peak temperature is less than $\sim 50^\circ\text{C}$, and the

progress of aqueous alteration is mild. In the inner part, on the other hand, the peak temperature is $\sim 150^\circ\text{C}$, and alteration occurs rapidly. While less altered CM chondrites, such as Murchison, could be derived from the outer part of the asteroid, more altered ones, such as Sayama, come from the inner part. In contrast, case B gives a rapid increase in temperature caused by more abundant ^{26}Al , even if the hydration heat is not considered, and as a consequence, the carbonates are likely to have precipitated at $\sim 4,564.5$ Myr ago (Fig. 3b). Other heat sources, such as impact heating, inductive heating or decay energies of other radioactive nuclides, if any, would assist the ^{26}Al heating of the parent body, resulting in a more rapid temperature increase. Therefore, such early accretion as case B is ruled out considering our high-precision Mn–Cr data. Although parameters such as water fraction and the size of the asteroid are not well constrained, it is important to stress that there is no parameter space where carbonates could form at 4.8 Myr after CAI formation, if the parent body accreted within 3.0 Myr after CAI formation, because the timing at which the aqueous alteration began was determined predominantly by the accretion time of the asteroid.

An accretion time of >3 Myr after CAI formation is later than the formation of chondrules in the least metamorphosed chondrites (typically ~ 2 Myr after CAI formation³). On the other hand, the W isotopic compositions of iron meteorites suggest their formation within 1.5 Myr after CAI formation². If the carbonate ages of CI chondrites reported so far also have systematic errors, like those of the CM chondrites, they would become comparable with or younger than the carbonate ages of the CM chondrites^{8,10–12}. In any case, the early formation of hydrous asteroids suggested by old carbonates in CM chondrites is no longer well founded, and we conclude that hydrous asteroids accreted later than differentiated and metamorphosed asteroids. This conclusion is consistent with the composi-

Table 1 | Mn–Cr isotope data of carbonates in four CM chondrites.

Grain	⁵⁵ Mn (counts)	⁵² Cr (counts)	⁵³ Cr (counts)	⁵² Cr (cps/nA)	⁵⁵ Mn/ ⁵² Cr	Error	δ ⁵³ Cr (%)	Error
<i>Murchison</i>								
Calcite_A	5.45.E+06	386	158	0.835	22,228	2,263	486	281
Calcite_A	4.20.E+06	1,094	368	2.50	5,684	374	218	147
Calcite_A	7.61.E+06	416	181	0.807	27,437	2,689	578	281
Calcite_A	7.71.E+05	20,237	5,669	112	54	4	16	31
Calcite_B	1.37.E+06	217	79	1.14	10,026	1,361	326	348
Calcite_B	3.29.E+06	379	136	0.783	12,686	1,303	304	260
Calcite_C	2.96.E+06	937	294	3.40	4,659	306	139	152
Calcite_D	2.33.E+06	1,606,380	443,868	5,501	3	0.2	2	3
Calcite_E	9.12.E+05	189	60	0.242	7,747	1,128	148	341
<i>Y791198</i>								
Calcite_A	2.12.E+06	143	62	0.478	20,750	3,465	574	478
Calcite_A	5.44.E+06	147	94	0.492	51,556	8,491	1,319	612
Calcite_B	1.78.E+06	300	109	2.68	8,057	931	317	295
Calcite_C	4.06.E+06	143	93	0.478	40,119	6,699	1,358	627
Calcite_C	7.32.E+05	13,389	3,815	44.6	79	5	34	38
Calcite_D	1.57.E+06	215	80	2.11	9,859	1,345	352	354
Calcite_D	2.11.E+06	57	41	0.382	50,585	13,372	1,575	1,057
Calcite_E	1.83.E+06	326	99	2.12	7,724	855	107	254
Calcite_E	4.65.E+05	192	63	0.822	3,369	486	192	346
Calcite_F	4.79.E+06	101	70	0.338	66,967	13,295	1,512	780
Calcite_F	6.76.E+06	148	156	0.495	65,268	10,712	2,819	875
Calcite_G	1.56.E+06	296	118	1.10	7,389	859	449	315
<i>ALH83100</i>								
Dolomite_A	9.23.E+06	1,064	374	5.85	13,372	879	273	153
Dolomite_B	6.13.E+06	578	241	4.73	15,987	1,331	512	232
Dolomite_C	2.32.E+07	9,698	3,033	40.7	4,074	268	134	47
Dolomite_C	1.19.E+07	1,151	469	3.84	17,468	1,148	477	162
Dolomite_C	5.93.E+06	1,223,345	341,521	4,078	25	2	12	4
Dolomite_D	5.42.E+06	12,393	3,688	88.5	672	44	79	40
Dolomite_E	8.93.E+06	15,632	4,417	52.1	963	63	25	35
Dolomite_F	1.36.E+07	683	330	2.28	33,109	2,534	751	235
Dolomite_F	1.20.E+07	1,543	571	5.14	12,641	831	341	131
Dolomite_F	8.15.E+06	740	312	2.57	18,788	1,381	530	207
Dolomite_F	6.08.E+06	364	146	5.68	23,753	2,491	453	285
Dolomite_F	4.75.E+06	1,656	491	5.52	4,570	300	75	110
<i>Sayama</i>								
Dolomite_A	1.22.E+07	186	183	1.82	88,891	13,043	2,575	745
Dolomite_A	2.02.E+07	562	351	1.87	51,629	4,354	1,266	308
Dolomite_A	9.03.E+06	99	123	0.865	124,314	25,030	3,513	1,221
Dolomite_B	6.71.E+06	65	89	0.578	140,802	35,012	3,978	1,628
Dolomite_B	6.02.E+06	97	99	0.896	84,327	17,148	2,709	1,061
Dolomite_C	1.46.E+07	604	339	2.01	34,879	2,837	1,037	276
Dolomite_C	1.34.E+07	423	295	1.50	45,455	4,422	1,531	384
Dolomite_D	9.32.E+06	190	142	1.14	67,826	9,840	1,715	602
Dolomite_D	6.12.E+06	141	97	1.23	59,134	9,972	1,494	659
Dolomite_D	9.89.E+06	235	174	1.16	59,160	7,724	1,693	539
Dolomite_D	6.33.E+06	2,578,365	713,592	18,157	4	0.2	5	7
Dolomite_E	2.19.E+07	881	530	2.94	35,729	2,407	1,183	240
Dolomite_E	8.60.E+06	221	189	0.830	55,344	7,448	2,109	616
Dolomite_F	6.19.E+06	141	111	1.12	60,012	10,125	1,875	730
Dolomite_F	8.84.E+06	197	180	0.973	62,850	8,964	2,324	686

⁵⁵Mn, ⁵²Cr and ⁵³Cr show the total counts during analyses after the correction for the dynamic background. ⁵⁵Mn/⁵²Cr is calculated considering the time dependence of the RSF (see Methods for details). Errors on both ⁵⁵Mn/⁵²Cr and δ⁵³Cr represent the external reproducibility or the internal precision, whichever is larger.

tional structure of the asteroid belt, which shows that asteroids with spectra indicative of high-temperature silicate minerals are concentrated between 1.8 and 3 AU, whereas dark C-rich types seem to be located farther away from the Sun³¹.

Methods

NanoSIMS measurement condition. We performed the Mn–Cr isotope measurements using a NanoSIMS 50 installed at the Atmosphere and Ocean Research Institute at the University of Tokyo. Both the meteorites and the synthetic calcite

standard were embedded in epoxy and polished in the same way. A focused O[−] primary ion beam of 1 nA was accelerated to 16 keV and irradiated on the surfaces of the samples. Before the measurements, an intense primary beam of ~3 nA was rastered over 20×20 to 50×50 μm² on the meteoritic carbonates to remove surface contaminations and to obtain secondary ion images of carbonates to choose measurement spots. Secondary ions were extracted from the sample surfaces by an accelerating voltage of 8 kV, and ⁴³Ca⁺, ⁵²Cr⁺, ⁵³Cr⁺ and ⁵⁵Mn⁺ ions were detected in a combined peak-jumping/multi-detection mode. The magnet was cycled through two field settings and the waiting times were 2 seconds each. In the first setting, ⁴³Ca⁺, ⁵²Cr⁺ and ⁵⁵Mn⁺ were simultaneously detected with three secondary electron multipliers (SEMs). The measurement time was 2 seconds.




Figure 2 | Summary and comparison of Mn-Cr ages of carbonates.

The filled symbols represent Mn-Cr ages in this study using the calcite standard, and the open symbols represent previously reported data (ref. 5 and references therein). Also shown are the Pb-Pb age of a type-B CAI from the NWA2364 CV3 chondrite⁶ (named for their type specimen of the Vigarano meteorite) and the Pb-Pb age of the LEW86010 angrite¹⁵ used as a time anchor. The carbonates in the four CM chondrites analysed here show an identical formation age ($4,563.4^{+0.4}_{-0.5}$; shown by the vertical grey band). Error bars ($\pm 2\sigma$) are based on the uncertainties of the Mn/Cr RSF and the isochron slopes (see Methods for details).

Subsequently, $^{53}\text{Cr}^+$ was detected with the same SEM as that for $^{52}\text{Cr}^+$. The measurement time was 5 seconds. The measurements consisted of 150 cycles of the two magnetic field settings. The difference between sensitivities of two SEMs for $^{52}\text{Cr}^+$ and $^{55}\text{Mn}^+$ was corrected when evaluating the Mn/Cr relative sensitivity of carbonates, although this difference was cancelled when we calculated the $^{55}\text{Mn}/^{52}\text{Cr}$ of the meteoritic samples and normalized it to the standard value. The dynamic background of the SEM for $^{52}\text{Cr}^+$ was measured by applying a -10 V offset relative to the peak centre on the deflection plates in front of the SEMs. The background ranged from 0.01 to 0.02 cps.

Mn/Cr relative sensitivity of carbonates. We synthesised a calcite standard doped with Mn and Cr to evaluate the relative sensitivity factor (RSF; defined as $(^{55}\text{Mn}^+ / ^{52}\text{Cr}^+)_{\text{ion}} / (^{55}\text{Mn} / ^{52}\text{Cr})_{\text{atomic}}$) of carbonates. See ref. 13 for details of the production. The synthetic calcite has concentration gradients of Mn and Cr in a single grain. The concentrations decrease from the centre to periphery of a grain due to the fractional crystallization of Mn and Cr. Therefore, we analysed the grains for small regions within $100 \times 100 \mu\text{m}^2$ of the grain centre. In these limited areas, the heterogeneities of Mn and Cr are up to 20%.

The $^{55}\text{Mn}/^{52}\text{Cr}$ atomic ratios were determined with an electron probe micro analyser (EPMA) around measurement spots with a NanoSIMS. In the EPMA analyses, the Mg, Ca, Cr, Mn and Fe concentrations were determined with a JEOL JXA-8900L. A sample current of 3 nA was accelerated to 15 keV and rastered over $\sim 10 \times 10 \mu\text{m}^2$ on the sample surface to avoid damage by electron bombardment. The measurement times of the Mg, Ca, Cr, Mn and Fe concentrations and the background were 10 seconds and 5 seconds, respectively.

In the NanoSIMS analyses, we found that the $^{43}\text{Ca}^+$, $^{52}\text{Cr}^+$ and $^{55}\text{Mn}^+$ intensities and the $^{55}\text{Mn}^+ / ^{52}\text{Cr}^+$ decrease with time for the standard and meteoritic carbonates during a single measurement (Supplementary Fig. S1). This effect is too large to be attributable to the heterogeneities of Mn and Cr in a grain, but rather is due to the charge-up of the calcite. Hence, we used the ‘time-dependent RSF’ (hereafter described as ‘RSF(*t*)’) to calibrate the $^{55}\text{Mn}/^{52}\text{Cr}$ atomic ratios of the meteoritic samples. The RSF(*t*) was defined for each measurement cycle (the measurement consisted of 150 cycles of two magnetic field settings). We measured the RSF approximately ten times in each measurement session (the overall measurement consists of three sessions, see Supplementary Fig. S1 and Supplementary Table S1), and the average of approximately ten RSF values at a given time (measurement cycle) was used as the RSF(*t*) at the time. Then, we calculated the $^{55}\text{Mn}^+_{\text{convert.}}$ of the meteoritic carbonates at a given time as $^{55}\text{Mn}^+_{\text{convert.}} = ^{55}\text{Mn}^+ / \text{RSF}(t)$. Finally, we determined the mean $^{55}\text{Mn}/^{52}\text{Cr}$ atomic ratios of the meteoritic carbonates (Table 1), as $^{55}\text{Mn}/^{52}\text{Cr} = \Sigma(^{55}\text{Mn}^+_{\text{convert.}}) / \Sigma(^{52}\text{Cr}^+)$. We did not use the average of the $^{55}\text{Mn}^+_{\text{convert.}} / ^{52}\text{Cr}^+$ values in the 150 cycles as the mean $^{55}\text{Mn}/^{52}\text{Cr}$ atomic ratio, because the intensity of $^{52}\text{Cr}^+$ was usually quite low, and $^{55}\text{Mn}^+_{\text{convert.}} / ^{52}\text{Cr}^+$ was often infinite (no $^{52}\text{Cr}^+$ ion was detected in the




Figure 3 | Thermal history of CM chondrite parent body without fluid flow.

Temperature (solid lines) and the extent of alteration (dotted lines) are given at three depths (*r*: distance from the centre) in the asteroid with a radius of 30 km. The vertical grey band shows the formation age of carbonates ($4,563.4^{+0.4}_{-0.5}$). (a) Accretion at 3.5 Myr after CAI formation (case A). The extent of alteration is defined by $(1 - \text{olivine}_{\text{res.}} / \text{olivine}_{\text{ini.}})$, where $\text{olivine}_{\text{res.}}$ and $\text{olivine}_{\text{ini.}}$ are the moles of the residual and initial olivine, respectively. (b) Accretion at 3.0 Myr after CAI formation (case B). Note that the hydration reaction is not included in case B.

cycle). The internal errors on $^{55}\text{Mn}/^{52}\text{Cr}$ were calculated as

$$\sigma_{^{55}\text{Mn}/^{52}\text{Cr}} = ^{55}\text{Mn} / ^{52}\text{Cr} \times \sqrt{1 / (\Sigma ^{55}\text{Mn}^+) + 1 / (\Sigma ^{52}\text{Cr}^+)}$$

Note that $\Sigma(^{55}\text{Mn}^+)$ instead of $\Sigma(^{55}\text{Mn}^+_{\text{convert.}})$ was used in calculating the internal error.

The uncertainty in the RSF (σ_{RSF}) is difficult to estimate. In the three measurement sessions, the average RSF(*t*) values are 0.636, 0.675 and 0.690, respectively (Supplementary Table S1). These values vary within 8.1%, although the error in each session is less than 2.4%. This variation seems to be caused by the behaviours of the $^{55}\text{Mn}^+ / ^{52}\text{Cr}^+$ changes, that is, degrees of charge-up. The degrees of charge-up are likely to be closely related to crater depth produced by primary ion bombardment, which is controlled by the intensity of the primary ion beam in each measurement session. In each measurement session, we found that the change was reproducible, as shown in Supplementary Figure S1, because we kept the measurement conditions exactly the same including the primary ion beam intensity, which is why the error of the RSF in each session is smaller than the variation among the three sessions. Thus, we are quite sure that the $^{55}\text{Mn}/^{52}\text{Cr}$ of the meteoritic carbonates is properly corrected for the RSF in each measurement session. In this study, we conservatively adopted a σ_{RSF} of 8.1%. We consider that this value can fully describe the uncertainty in the estimated $^{55}\text{Mn}/^{52}\text{Cr}$ ratios, or rather, might be overestimated. This uncertainty was propagated to the final errors on the initial $^{53}\text{Mn}/^{55}\text{Mn}$ ratios, $(^{53}\text{Mn}/^{55}\text{Mn})_0$, which were calculated as

$$\sigma_{(^{53}\text{Mn}/^{55}\text{Mn})_0} = \left(^{53}\text{Mn} / ^{55}\text{Mn} \right)_0 \times \sqrt{\sigma_{\text{slope}}^2 + \sigma_{\text{RSF}}^2}$$

where σ_{slope} is the 95% confident interval of the isochron slope based on $\sigma_{^{55}\text{Mn}/^{52}\text{Cr}}$ and $\sigma_{^{53}\text{Cr}/^{52}\text{Cr}}$ using the York fit program ‘Isoplot 3.41’ (ref. 32). $\sigma_{^{53}\text{Cr}/^{52}\text{Cr}}$ will be defined in the next subsection.

Chromium isotope measurements. In contrast to the RSF, the Cr isotopic composition is stable for the synthetic calcite standard during a single measurement. The mean values of $^{53}\text{Cr}/^{52}\text{Cr}$ are 0.1103 (1), 0.1102 (3) and 0.1099 (2) in the three measurement sessions, respectively, where the numbers in parentheses represent the uncertainties of the last digits (Supplementary Table S1). We calculated the $^{53}\text{Cr}/^{52}\text{Cr}$ of the meteoritic carbonates as $^{53}\text{Cr}/^{52}\text{Cr} = \Sigma(^{53}\text{Cr}^+) / \Sigma(^{52}\text{Cr}^+)$. We did not use the average of the $^{53}\text{Cr}^+ / ^{52}\text{Cr}^+$ values in the 150 cycles as the mean $^{53}\text{Cr}/^{52}\text{Cr}$ ratio for the same reason as that for $^{55}\text{Mn}/^{52}\text{Cr}$. The $^{53}\text{Cr}/^{52}\text{Cr}$ of the meteoritic carbonates is described as the permil deviation ($\delta^{53}\text{Cr}$) from the standard value (Table 1). The internal errors on $^{53}\text{Cr}/^{52}\text{Cr}$ ($\sigma_{^{53}\text{Cr}/^{52}\text{Cr}}$) were calculated as

$$\sigma_{^{53}\text{Cr}/^{52}\text{Cr}} = \sqrt{\frac{^{53}\text{Cr}}{^{52}\text{Cr}} \times \frac{1}{\left(\frac{\Sigma(^{52}\text{Cr}^+)}{^{52}\text{Cr}}\right)^2} + \frac{1}{\left(\frac{\Sigma(^{53}\text{Cr}^+)}{^{53}\text{Cr}}\right)^2}}$$

References

- Grimm, R. E. & McSween, H. Y. Jr. Heliocentric zoning of the asteroid belt by aluminum-26 heating. *Science* **259**, 653–655 (1993).
- Qin, L. *et al.* Rapid accretion and differentiation of iron meteorite parent bodies inferred from ^{182}Hf - ^{182}W chronometry and thermal modeling. *Earth Planet. Sci. Lett.* **273**, 94–104 (2008).
- Krot, A. N. *et al.* Origin and chronology of chondritic components: a review. *Geochim. Cosmochim. Acta* **73**, 4963–4997 (2009).
- Brearley, A. J. in *Meteorites and the Early Solar System II* (eds Lauretta, D. S. & McSween, H. Y. Jr) 587–624 (Univ. Arizona Press, 2006).
- de Leuw, S., Rubin, A. E., Schmitt, A. K. & Wasson, J. T. ^{53}Mn - ^{53}Cr systematics of carbonates in CM chondrites: implications for the timing and duration of aqueous alteration. *Geochim. Cosmochim. Acta* **73**, 7433–7442 (2009).
- Bouvier, A. & Wadhwa, M. The age of the Solar System redefined by the oldest Pb-Pb age of a meteoritic inclusion. *Nat. Geosci.* **3**, 637–641 (2010).
- Macdougall, J. D., Lugmair, G. W. & Kerridge, J. F. Early Solar System aqueous activity: Sr isotope evidence from the Orgueil CI meteorite. *Nature* **307**, 249–251 (1984).
- Endress, M., Zinner, E. & Bischoff, A. Early aqueous activity on primitive meteorite parent bodies. *Nature* **379**, 701–703 (1996).
- Hohenberg, C. M., Pravdivtseva, O. & Meshik, A. Reexamination of anomalous I-Xe ages: Orgueil and Murchison magnetites and Allegan feldspar. *Geochim. Cosmochim. Acta* **64**, 4257–4262 (2000).
- Hoppe, P., Macdougall, J. D. & Lugmair, G. W. High spatial resolution ion microprobe measurements refine chronology of carbonate formation in Orgueil. *Meteorit. Planet. Sci.* **42**, 1309–1320 (2007).
- Petit, M. *et al.* Duration and sequence of carbonate crystallization on the Orgueil protolith: ^{53}Mn - ^{53}Cr systematics of their evolution in O and C isotopic composition. *Lunar Planet. Sci. Conf. XL A1657* (2009).
- Petit, M. *et al.* ^{53}Mn - ^{53}Cr ages of Kaidun carbonates. *Meteorit. Planet. Sci.* **46**, 275–283 (2011).
- Sugiura, N., Ichimura, K., Fujiya, W. & Takahata, N. Mn/Cr relative sensitivity factors for synthetic calcium carbonate measured with a NanoSIMS ion microprobe. *Geochem. J.* **44**, e11–e16 (2010).
- Rubin, A. E., Trigo-Rodríguez, J. M., Huber, H. & Wasson, J. T. Progressive aqueous alteration of CM carbonaceous chondrites. *Geochim. Cosmochim. Acta* **71**, 2361–2382 (2007).
- Amelin, Y. U-Pb ages of angrites. *Geochim. Cosmochim. Acta* **72**, 221–232 (2008).
- Lugmair, G. W. & Shukolyukov, A. Early solar system timescales according to ^{53}Mn - ^{53}Cr systematics. *Geochim. Cosmochim. Acta* **62**, 2863–2886 (1998).
- Trinquier, A. *et al.* ^{53}Mn - ^{53}Cr systematics of the early Solar System revisited. *Geochim. Cosmochim. Acta* **72**, 5146–5163 (2008).
- Ogliore, R. C., Huss, G. R. & Nagashima, K. in *Nuclear Instruments and Methods in Physics Research B* **269**, 1910–1918 (2011).
- Kallemeyn, G. W. & Wasson, J. T. The compositional classification of chondrites I The carbonaceous chondrite groups. *Geochim. Cosmochim. Acta* **45**, 1217–1230 (1981).
- Clayton, R. N. & Mayeda, T. K. Oxygen isotope studies of carbonaceous chondrites. *Geochim. Cosmochim. Acta* **63**, 2089–2104 (1999).
- Kunihiro, T., Rubin, A. E., McKeegan, K. D. & Wasson, J. T. Initial $^{26}\text{Al}/^{27}\text{Al}$ ratio in carbonaceous-chondrite chondrules: too little ^{26}Al to melt asteroids. *Geochim. Cosmochim. Acta* **68**, 2947–2957 (2004).
- Kurahashi, E., Kita, N. T., Nagahara, H. & Morishita, Y. ^{26}Al - ^{26}Mg systematics of chondrules in a primitive CO chondrite. *Geochim. Cosmochim. Acta* **72**, 3865–3882 (2008).
- Clayton, R. N. & Mayeda, T. K. The oxygen isotope record in Murchison and other carbonaceous chondrites. *Earth Planet. Sci. Lett.* **67**, 151–161 (1984).
- Grimm, R. E. & McSween, H. Y. Jr. Water and the thermal evolution of carbonaceous chondrite parent bodies. *Icarus* **82**, 244–280 (1989).
- Cohen, B. A. & Coker, R. F. Modeling of Liquid water on CM meteorite parent bodies and implications for amino acid racemization. *Icarus* **145**, 369–381 (2000).
- Young, E. D., Ash, R. D., England, P. & Rumble, D. Fluid flow in chondritic parent bodies: deciphering the compositions of planetesimals. *Science* **286**, 1331–1335 (1999).
- Palguta, J., Schubert, G. & Travis, B. Fluid flow and chemical alteration in carbonaceous chondrite parent bodies. *Earth Planet. Sci. Lett.* **296**, 235–243 (2010).
- Zolensky, M. E., Bourcier, W. L. & Gooding, J. L. Aqueous alteration on the hydrous asteroids: results of EQ3/6 computer simulations. *Icarus* **78**, 411–425 (1989).
- Guo, W. & Eiler, J. M. Temperatures of aqueous alteration and evidence for methane generation on the parent bodies of the CM chondrites. *Geochim. Cosmochim. Acta* **71**, 5565–5575 (2007).
- Nakamura, T., Noguchi, T. & Yoneda, S. The conditions of aqueous alteration reactions recorded in the Sayama CM2 chondrite. *Meteorit. Planet. Sci.* **38**, A5080 (2003).
- Ghosh, A., Weidenschilling, S. J., McSween, H. Y. Jr. & Rubin, A. in *Meteorites and the Early Solar System II* (eds Lauretta, D. S. & McSween, H. Y. Jr.) 555–565 (Univ. Arizona Press, 2006).
- Ludwig, K. R. in *Berkeley Geochronology Center Special Publication No. 4*, 1–72 (2003).

Acknowledgements

We thank H. Hiyagon for critical discussions, N. Takahata for technical help with the NanoSIMS analyses, H. Yoshida for technical help with the EPMA analyses, and S. Yoneda and T. Nakamura for providing the Sayama meteorite.

Author contributions

W.F. and N.S. prepared the polished thin sections of the meteoritic samples and the standard. W.F. performed the NanoSIMS analyses, and collected, processed and refined the data. Y.S. provided technical support for the NanoSIMS analyses. N.S. and K.I. developed the method to synthesise the calcite standard doped with Mn and Cr. W.F. performed the EPMA analyses. H.H. produced the numerical code to calculate the thermal history of the asteroid. W.F. designed the study and prepared the manuscript. All authors discussed the results and commented on the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Fujiya, W. *et al.* Evidence for the late formation of hydrous asteroids from young meteoritic carbonates. *Nat. Commun.* **3**:627 doi: 10.1038/ncomms1635 (2012).