

# UC Office of the President

## Recent Work

### Title

Evidence-informed practical recommendations for increasing physical activity among persons living with HIV.

### Permalink

<https://escholarship.org/uc/item/7qp3q2p7>

### Journal

AIDS (London, England), 33(6)

### ISSN

0269-9370

### Authors

Montoya, Jessica L  
Jankowski, Catherine M  
O'Brien, Kelly K  
et al.

### Publication Date

2019-01-22

### DOI

10.1097/qad.0000000000002137

Peer reviewed

# Evidence-informed practical recommendations for increasing physical activity among persons living with HIV

Jessica L. Montoya<sup>a</sup>, Catherine M. Jankowski<sup>b</sup>, Kelly K. O'Brien<sup>c</sup>,  
Allison R. Webel<sup>d</sup>, Krisann K. Oursler<sup>e,f</sup>, Brook L. Henry<sup>a</sup>,  
David J. Moore<sup>a</sup> and Kristine M. Erlandson<sup>g</sup>

*AIDS* 2019, **33**:931–939

**Keywords:** exercise, guidelines, HIV, physical activity, resistance training

## Introduction

With the advent of effective antiretroviral therapy (ART), the care of persons living with HIV (PLWH) is shifting focus to the management of age-related chronic health conditions (e.g. metabolic syndrome and cardiovascular disease), syndromes of ageing (e.g. dementia and frailty) and side effects related to ART. Non-ART polypharmacy is common among PLWH and associated with an increasing risk of hospitalization and mortality [1]; thus, nonpharmacologic management of comorbidities is critical for PLWH who may experience an earlier onset and a greater burden of comorbidities. Routine engagement in health-enhancing behaviours, including physical activity, may help prevent and manage comorbid health conditions and syndromes of ageing common among PLWH [2]. Physical activity refers to any bodily movement produced by muscle contraction that causes energy expenditure, whereas exercise is a subset of physical activity that involves planned, repetitive body movement with the intent to increase well being and energy level to allow for independent participation in

physical activities [3]. The goal of this narrative review is to summarize key literature from the past 10 years examining the benefits of physical activity and to outline recommendations to prescribe and support physical activity engagement among PLWH.

The second edition of Physical Activity Guidelines for Americans issued by the Department of Health and Human Services (HHS) [4] proposes that adults – even those with chronic conditions and disability – engage in at least 150–300 min of moderate-intensity or 75–150 min of vigorous-intensity aerobic physical activity per week, as well as muscle strengthening activities on two or more days a week. Balance training is additionally recommended as part of older adults' weekly physical activity to reduce fall risk. Importantly, HHS emphasizes that moving more and sitting less will benefit nearly everyone, with the most sedentary and least active individuals experiencing the greatest benefit from small physical activity increases [5]. Although some PLWH may have unique physical limitations that must be accommodated in order for them to safely engage in physical activity, the take-home

<sup>a</sup>Department of Psychiatry, University of California San Diego, La Jolla, California, <sup>b</sup>University of Colorado Anschutz Medical Campus, College of Nursing, Aurora, Colorado, USA, <sup>c</sup>Department of Physical Therapy; Rehabilitation Sciences Institute; Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada, <sup>d</sup>Frances Payne Bolton School of Nursing, Case Western Reserve University, Cleveland, Ohio, <sup>e</sup>Geriatric Research and Education, Salem Veterans Medical Center, Salem, <sup>f</sup>Virginia Tech Carilion School of Medicine, Roanoke, Virginia, and <sup>g</sup>Division of Infectious Diseases; Division of Geriatric Medicine, Department of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA.

Correspondence to Kristine M. Erlandson, MD, MS, 12700 E. 19<sup>th</sup> Avenue, Mail Stop B168; Aurora, CO 80045, USA.

Tel: +1 303 724 4941; fax: +1 303 724 4926; e-mail: Kristine.Erlandson@ucdenver.edu

Received: 8 November 2018; revised: 14 December 2018; accepted: 17 December 2018.

recommendation is that physical activity participation is key to maximizing health and function. For additional information regarding physical activity recommendations and safety precautions for patients with combinations of cardiovascular disease risk factors (i.e. obesity, arterial hypertension, diabetes mellitus and dyslipidaemia), we refer the reader to an expert consensus statement on physical activity prescription [6].

Despite the well established health benefits of physical activity (summarized in Table 1) [7–41], rates among PLWH remain low. A meta-analysis of 24 studies involving nearly 4000 PLWH found that only half of the individuals engaged in 150 min of moderate-intensity physical activity [42]. Among middle-aged PLWH, 26% reported no moderate, vigorous, or muscle strengthening physical activity, similar to rates in the U.S. general population [43]. These physical activity trends are consistent across settings: 32% of Vietnamese adults with HIV reported low or no physical activity [44,45], with similar rates in the Swiss HIV Cohort study (41%) [46], Germany (39%) [47] and Brazil (44%) [7]. Despite limited engagement in physical activity, results of a qualitative study found that PLWH understood the health promotion benefits of physical activity and thought physical activity should be a greater priority in their life [48].

### Alternative types of physical activity

In addition to traditional aerobic (e.g. walking, biking and swimming) and resistance exercises (Table 1), lower intensity physical activity such as yoga is associated with positive benefits among PLWH, including an improvement in quality of life [8], reduction of depressive symptoms [9] and reduction of blood pressure [9,49]. Tai Chi was also associated with improved well being and balance in PLWH [50]. The benefits of high-intensity exercise are similarly evident: Erlandson *et al.* [10] recently demonstrated that among older PLWH, high-intensity aerobic and resistance exercise (based on target heart rate and resistance load) improved endurance and strength to a similar, if not greater, extent than moderate-intensity exercise. Oursler *et al.* [51] similarly demonstrated significant gains in endurance among older PLWH exercising at a higher than a more moderate-intensity aerobic programme. These studies of high-intensity exercise have indicated no reason to dissuade PLWH – young or old – from progressing to high-intensity exercise following several weeks of moderate-intensity training. More details about the effectiveness of various types of physical activity among PLWH can be found in a recent meta-analysis [52].

### Barriers to physical activity among persons living with HIV

Understanding barriers to physical activity specific to, or more pronounced among, PLWH is essential to developing effective, patient-centred physical activity recommendations. Across populations, physical activity is

affected by diverse factors, including intrapersonal, interpersonal and environmental barriers.

#### *Intrapersonal barriers*

In a systematic review of 45 studies of physical activity in PLWH [53], lower engagement in physical activity was consistently associated with demographic (i.e. older age and fewer years of formal education), HIV-specific and biologic (i.e. being on ART, lower CD4<sup>+</sup> T-cell counts, lipodystrophy, lower cardiorespiratory fitness and opportunistic infections) and psychological (i.e. lower motivation, depression and worse self-perceived physical function) variables. Moreover, the experience of these physical and mental health challenges that hinder engagement in physical activity may be unpredictable and episodic for PLWH [54]. Many PLWH experience great symptom burden, including neuropathy [55], fatigue [56] and reduced cardiorespiratory fitness [57], with even greater impairments among older PLWH [58]. Reduced cardiorespiratory fitness among PLWH as indicated by impaired peripheral oxygen uptake [49], dysfunction of skeletal muscle mitochondria [59], reduction in mitochondrial DNA content [60] and/or limitations in lung function [61] (e.g. impaired carbon monoxide diffusion capacity [62]) may interfere with adaptation to physical activity. Aerobic and resistance exercise, however, may reduce symptoms such as HIV-related fatigue [11,12]. Adherence to a moderate-intensity, home-based, aerobic exercise [12] or a supervised aerobic and resistance exercise intervention has shown reduced fatigue among PLWH, with increased benefit among those with high adherence [11].

#### *Interpersonal barriers*

Social factors, including worries about HIV disclosure and stigma [63] and a lack of social support [64], can also hinder participation in physical activity. Interviews of PLWH who participated in a community-based exercise programme, however, indicated that participants found the environment less stigmatizing than initially feared, suggesting the malleability of social determinants of physical activity engagement. Healthcare providers' recommendations can also play a key role in perceptions and engagement in physical activity among PLWH [65]. When providers only focus on HIV-related health outcomes and do not recommend routine physical activity, PLWH have less recognition of the health benefits of physical activity [65].

#### *Environmental barriers*

Environmental barriers to physical activity may be more pronounced among PLWH, such as concern about physical safety [66]. Although fitness centres may be ideal venues for engaging in physical activity, access may be limited by physical and financial accessibility [67,68], and/or concerns regarding potential stigma related to body image [54]. Thus, when developing a physical activity plan, environmental barriers may need to be

**Table 1. Summary of key findings from published studies (2009–2018) investigating the effects of physical activity on different body systems among adults living with HIV.**

Physical activity	Brief summary of findings
<i>Intervention training studies</i>	
<b>Aerobic + resistance</b>	
60 min aerobic + resistance 3x/week for 24 weeks [11]	Fatigue: ↑ vitality Anthropometry/body composition: ↓ % body fat, HC, ↑ muscle mass, ↔ BMI, WC, WHR Cardiometabolic profile: ↓ resting HR, glucose, ↑ metabolic equivalent, ↔ haemoglobin, TC, HDL, LDL, TRIG, maximum oxygen consumption Immunological profile: ↑ CD4 <sup>+</sup> T-cell count QOL/Mental Health: ↑ QOL
40–60 min aerobic + resistance 4x/week for 12 weeks [13]	Anthropometry/body composition: ↓ waist circumference, ↔ body weight, BMI Cardiometabolic profile: ↓ DBP, ↔ SBP, TRIG, TC, HDL, LDL, fasting glucose, % w/MetS Inflammation: ↔ hsCRP Physical function/frailty: ↑ physical function
Aerobic + resistance 3x/week for 24 weeks [10]	Inflammation: ↓ IL-8, ↔ IL-4, IL-5, IL-6, IL-10, TNF-α, IFN-γ, GM-CSF
20 min aerobic + resistance 3x/weekly for 16 weeks [14]	
Aerobic and/or resistance	
30 min aerobic exercise or progressive resistance 3x/week for 12 weeks [15]	Pain: ↓ neuropathic pain
60 min brisk walking with or without 30 min circuit-training 3x/week for 12 weeks [16]	Anthropometry/body composition: ↓ body weight, BMI, WC, HC Cardiometabolic profile: ↓ TC, LDL Inflammation: ↓ hsCRP, IL-6, d-dimer, IL-18
<b>Aerobic alone</b>	
High-intensity aerobic interval 3x/week for 12 weeks [17]	Anthropometry/body composition: ↓ Body weight, BMI, body fat %, WC, HC Cardiometabolic profile: ↓ HbA1c, DBP, ↔ HOMA-IR, SBP, TRIG, TC, HDL, LDL Cardiometabolic profile: ↑ capacity of the endothelium to release t-PA
45–60 min aerobic 5–7x/week for 12 weeks [18]	
20–40 min light aerobic (walking or jogging) 3x/week for 12 weeks [19]	Mitochondrial function: ↑ Peripheral blood mononuclear cells mitochondrial respiratory capacity, spare respiratory capacity and nonmitochondrial respiration Fatigue: ↑ time to fatigue (measured by treadmill time), VO <sub>2</sub> max
30 min aerobic treadmill training 2x/week for 6 weeks [20]	
Moderate-to-vigorous PA (≥2690 counts/min for ≥10 min); accelerometry [21]	Anthropometry/body composition: ↑ weight loss, ↔ BMI changes
<b>Resistance alone</b>	
Circuit resistance 3x/week for 8 weeks [22]	Anthropometry/body composition: ↓ fat mass, ↑ lean mass, ↔ body weight Immunological profile: ↑ CD4 <sup>+</sup> T-cell count Physical function/frailty: ↑ strength, muscle mass, upper-arm and forearm circumference Inflammation: ↔ IFN-γ, IL-1β, IL-2, IL-4, IL-6, IL-10, TNF-α, VCAM-1, cortisol
Resistance 3x/week for 6 weeks [23]	Anthropometry/body composition: ↑ lean body mass, ↓ Body fat %, bod fat mass, ↔ body mass, BMI Cardiometabolic profile: ↑ HDL, ↓ TC, LDL, TRIG, CRP Inflammation: ↓ IL-1B, IL-6, IL-8, TNF-α, ↑ IL-10
Nonlinear resistance 3x/week for 12 weeks [24,25]	Physical function/frailty: ↑ muscular strength, ↔ bone mineral content Anthropometry/body composition: ↔ body weight, BMI, lean mass, fat mass Immunological profile: ↑ CD4 <sup>+</sup> T-cell count, CD4 <sup>+</sup> /CD8 <sup>+</sup> ratio
Resistance 2x/week for 52 weeks [26]	
<b>Yoga</b>	
6 days of training followed by daily home-based and 1x/week supervised yoga for 12 weeks [8]	Anthropometry/body composition: ↔ BMI Immunological profile: ↔ CD4 <sup>+</sup> T-cell count QOL/Mental Health: ↑ QOL
60 min yoga 6x/week for 4 weeks [9]	Immunological profile: ↑ CD4 <sup>+</sup> T-cell count QOL/Mental Health: ↓ depression symptoms
60 min yoga 2–3x/week for 20 weeks [27]	Anthropometry/body composition: ↔ weight, fat mass Cardiometabolic profile: ↓ resting SBP, DBP, ↔ lipids, glucose tolerance QOL/Mental Health: ↔ QOL
60 min yoga 2x/week for 8 weeks [28]	QOL/Mental Health: ↑ QOL (perceived stress and recent distress)
<i>Observational studies</i>	
Step counts; accelerometry [29]	Cardiometabolic profile: ↓ HOMA-IR, ↔ CAC score, SBP, DBP, TC Inflammation: ↓ IL-6 Immunological profile: ↔ CD4 <sup>+</sup> T-cell count
Moderate PA; accelerometry [30]	Cardiometabolic profile: ↓ insulin resistance, TRIG Inflammation: ↓ CRP, ↔ IL-6
Moderate-vigorous activity (≥70 min/day); SenseWear arm band [31]	
S-R >600 MET min/week relative to inactive; short-version IPAQ [7]	Anthropometry/body composition: ↓ WHR, ↔ % overweight, WC, body fat percentage Cardiometabolic profile: ↓ % hypertension, % diabetes Physical function/frailty: ↓ odds of frailty
S-R of ≥3 days/week of moderate/high PA; IPAQ [32]	
S-R of ≥3 days/week of moderate/high PA; IPAQ [33]	Disability: ↓ odds of impairment in activities of daily living

Table 1 (continued)

Physical activity	Brief summary of findings
S-R of high PA ( $\geq 3$ days/week of vigorous PA or $\geq 7$ days combination of walking, moderate and vigorous activity); IPAQ [34]	Neurocognition: $\downarrow$ odds of impairment in learning, memory, motor function
S-R of moderate PA; IPAQ [35]	Neurocognition: $\downarrow$ odds of neurocognitive impairment, instrumental activities of daily living dependence
Moderate-vigorous PA (min); NIAID Adult AIDS Clinical Trials Group Physical Activity Assessment [36]	Bone: $\uparrow$ BMD at the total hip and lumbar spine
S-R of PA; NIAID Adult AIDS Clinical Trials Group Physical Activity Assessment [37]	Anthropometry/body composition: $\leftrightarrow$ BMI Cardiometabolic profile: $\downarrow$ leptin, hyperaemic VTI, $\leftrightarrow$ HDL, LDL, TRIG, HOMA-IR, carotid distensibility, pericardial fat, flow mediated dilation, mean-mean IMT <i>Inflammation</i> : $\downarrow$ IL-6, hsCRP, $\leftrightarrow$ sCD163, sCD14, % CD14+ CD16+ monocytes, % CD14dimCD16+ monocytes
S-R of home-based exercise (min); 7-day diary [12]	Fatigue: $\downarrow$ fatigue
S-R of $>4$ h/week of PA, relative to $<4$ h/week or no PA [38]	Bone: $\uparrow$ BMD
S-R moderate-high PA, relative to low PA [39]	Anthropometry/body composition: $\downarrow$ waist circumference, $\leftrightarrow$ waist/hip ratio, BMI Cardiometabolic profile: $\downarrow$ SBP, DBP, Framingham risk score
S-R of PA in last 72 h [40]	Neurocognition: $\uparrow$ neurocognitive functioning (higher baseline and maintenance of neurocognitive function over time)
S-R of PA in last 72 h [41]	Neurocognition: $\downarrow$ rates of global neurocognitive impairment, driven by $\downarrow$ rates of impairment in working memory and speed of information processing

Note:  $\uparrow$  denotes increase/improvement;  $\downarrow$  denotes decrease/decline;  $\leftrightarrow$  denotes no significant change/difference. BMD, body mineral density; BP, blood pressure; CAC, coronary artery calcium; GM-CSF, granulocyte-macrophage colony-stimulating factor; HbA1c, haemoglobin A1c; HC, hip circumference; HDL, high-density lipoprotein cholesterol; HOMA-IR, haemostatic model assessment of insulin resistance; HR, heart rate; hsCRP, high-sensitivity C-reactive protein; IFN- $\gamma$ , interferon gamma; IL, interleukin; IPAQ, International Physical Activity Questionnaire; LDL, low-density lipoprotein cholesterol; MetS, metabolic syndrome; QOL, quality of life; sCD14, soluble cluster of differentiation 14; sCD163, soluble cluster of Differentiation 163; S-R, self-report; TC, total cholesterol; TNF- $\alpha$ , tumour necrosis factor-alpha; t-PA, tissue-type plasminogen activator; TRIG, triglycerides; VCAM-1, vascular cell adhesion protein 1; WC, waist circumference; WHR, waist-to-hip ratio.

considered and solutions for overcoming these barriers may need to be identified (e.g. encouraging outdoor activity, access to indoor facilities such as a shopping mall or short bouts of physical activity throughout the day that do not require gym access or equipment).

Collectively, this body of literature on barriers to physical activity among PLWH indicates an interplay of intrapersonal, interpersonal and environmental factors. Considering potential barriers to physical activity and plausible solutions are essential components of a physical activity action plan for PLWH. PLWH may benefit from an individualized approach to promoting physical activity, such as a gradual increase in activity, a different type of physical activity or physical activity environment, additional social support and/or a physical activity prescription that is tailored to the goals, abilities and interests of the individual in order to restore and achieve levels of cardiorespiratory fitness and physical function similar to HIV-uninfected peers.

### Behaviour change strategies for increasing engagement in physical activity

The American Heart Association recommends the routine assessment and promotion of physical activity as a 'vital sign' with every patient at every visit [69]. Assessment and promotion of physical activity in the healthcare setting has been found to be so effective at

improving patient outcomes that some healthcare systems have begun including physical activity as a vital sign in the electronic medical record [69]. The current extensive primary care guidelines for PLWH include detailed screening and management recommendations for hypertension, hyperlipidaemia and other common health conditions but do not mention physical activity assessment or counselling either routinely or in the management of comorbidities. The primary care guidelines for PLWH do recommend healthcare encounters every 3–6 months [70]: these frequent healthcare encounters provide excellent opportunities for healthcare providers to inquire about and promote physical activity among PLWH. A recent review confirms that assessment and promotion of physical activity by healthcare providers has a small to moderate positive effect on increasing physical activity levels, with larger effects observed when multiple behaviour change strategies are implemented [71].

One simple, effective behaviour change strategy is a physical activity prescription that is collaboratively developed by the patient and provider [72,73]. We adapted the physical activity prescription from the 'Exercise is Medicine' website [74] and included several behaviour change strategies that can be personalized (Fig. 1). The adapted physical activity prescription form allows healthcare providers to encourage patients to progress to higher levels of physical activity and/or work

General Exercise Guidelines for Positive Aging

Why should I exercise? Exercise may help you to:

- Improve mood & physical health
- Sleep better
- Decrease medications
- Improve overall quality of life

Is it safe? Yes! YES!

Studies show that moderate exercise (exercise that raises your heart rate, makes you break a sweat, but not so hard that you cannot talk) is generally safe for people with any chronic condition. Regardless of your starting point, gradually work toward a long-term exercise goal.

What's a good long-term exercise goal?

The basic exercise recommendations for all adults include either:

- At least 150 minutes of moderate aerobic activity per week and muscle-strengthening activities on 2 or more days per week, or
- 75 minutes of vigorous-intensity aerobic activity a week and muscle-strengthening activities on 2 or more days per week

TIP: You can add up exercise time in short periods throughout the day. Walking for 10 minutes morning, noon, and evening equals 30 minutes of walking.

R for Health: Get Up and Get Moving

Congratulations on deciding to increase your physical activity! Here is the plan we discussed to start you on your way.

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Aerobic Activity

Type: Walk Run Swim Bike Other \_\_\_\_\_

Days per Week start with:	1	2	3	4	5	6	7
gradually increase to:	1	2	3	4	5	6	7

Intensity

start with:	Light (a casual walk)	Moderate (a brisk walk)	Vigorous (jogging or running)
gradually increase to:	Light (a casual walk)	Moderate (a brisk walk)	Vigorous (jogging or running)

How should I start exercising?

Try a step-by-step approach:

Step 1: Pick an exercise

- What kind of exercise do I enjoy, or would be willing to do?
- What kind of exercise fits into my day?

Step 2: Set a short-term goal that you can accomplish

- "This week, I will walk for 10 minutes on 3 days."

Step 3: Set a long-term goal

- "Six months from now, I will be able to take a brisk walk for 20 minutes, 5 days of the week."

Step 4: Develop an action plan!

- Set a specific plan for how you will accomplish your short and long-term goals.

Step 5: Monitor your activity

- This could include a paper log, a pedometer, wrist monitor (such as FitBit), a smart phone program, or inexpensive hip-worn pedometer.

Step 6: Schedule time for activity!

- Block the time you plan to exercise on your calendar. Set an alert on your phone to remind you. Pick a time of the day that you are less likely to cancel because of fatigue or other commitments.

Step 7: Recruit a buddy or a group or tell a friend about your plan!

Minutes per Day

Start with:	10	20	30	45	60+
Gradually increase to:	10	20	30	45	60+

Steps per Day:

Start with:	2,500	5,000	7,500	10,000	12,500+
Gradually increase to:	2,500	5,000	7,500	10,000	12,500+

For next week, only change one thing at a time

Strength Training

- There are benefits to muscle strengthening done two days per week.
  - It is best to do exercise to strengthen all major muscle groups: legs, hips, back, chest, abdomen, shoulders, arms.
  - For each exercise, 8-12 repetitions is optimal.
  - Examples include resistance exercises using body weight (e.g., push-ups, lunges) or resistance bands, sit-ups, and heavy gardening.
- We will review this plan at your next visit.

Health Care Provider: \_\_\_\_\_  
Patient's Signature: \_\_\_\_\_

Fig. 1. Example physical activity prescription.

towards meeting the HHS recommendations by gradually increasing the amount of time, intensity or number of times a week they are physically active. For sedentary or physically inactive PLWH, prescribing movement breaks or walking may be a practical initial strategy to increase physical activity, as it does not require special skills or equipment [75].

Effective behaviour change strategies that consider readiness to engage in physical activity in the context of living with a sometimes unpredictable and episodic illness are needed to optimally promote adherence to a physical activity prescription among PLWH [68]. On the basis of HIV-specific literature in combination with existing behaviour change techniques recommended for all persons [76,77], we recommend adoption of behaviour change strategies, such as self-monitoring of physical activity; goal setting and action planning; prompts, cues or scheduling; and social support. Providing these strategies to patients may increase the likelihood of initiation and maintenance or sustained uptake of physical activity, ultimately leading to improved health and wellness in this population. We briefly summarize the benefits of the following approaches.

*Self-monitoring* is a useful physical activity behaviour change technique [77] that can include the use of wearable motion-sensing technologies (e.g. pedometers, Fitbit) [78], smartphone applications and online tracking resources such as Go4Life from the National Institute on Aging [79]. Self-monitoring can also occur using low-technology strategies such as a pen and paper diary. Results from a scoping study indicated that wireless activity monitors are increasingly used among PLWH as an outcome measure of physical activity; however, evidence of their effectiveness to enhance physical activity among PLWH is scarce [80].

*Goal setting and action planning* are effective behavioural strategies for increasing engagement in physical activity [81]. Goal setting encourages specific behavioural resolution (e.g. engaging in more physical activity this week), while action planning involves detailed planning of what the person will do, when they will engage in the specified behaviour and for how long. Combining behaviour change techniques, a person can self-monitor physical activity (e.g. steps per day) and then set a behavioural goal (e.g. increasing the number of steps by 10% per week). Action planning builds upon goal setting by linking behaviours to situational cues or an existing behaviour (i.e. associative learning). For example, physical activity can be linked with the use of transportation (e.g. exiting one stop early on the subway or bus, parking at the outskirts of a parking lot and taking the stairs), to establish a habit.

*Prompts, cues and scheduling* that remind PLWH to engage in physical activity may drive habit formation and

improve long-term physical activity adherence. Pre-scheduled activity is more often adhered to than relying on impromptu self-motivation [82]. A calendar, alarms and/or cell phone reminders to set aside a specific time for physical activity can support physical activity scheduling. An ongoing study is evaluating the effectiveness of a novel Short Message Service intervention (iSTEP) to increase moderate physical activity among PLWH [83,84].

*Social support* is strongly linked to physical activity engagement. Recent data clearly describe the prominent role that healthcare providers have in integrating health promotion into routine HIV care [65]. Thus, providers can leverage their strong patient relationships to emphasize a holistic concept of well being that includes physical activity [65]. Persons are more likely to engage in physical activity if they are linked to a similarly motivated person with whom they are able to engage in physical activity, creating a 'buddy system' (e.g. making a 'contract' with others to achieve specified levels of physical activity or participating in group exercise) [85]. In addition, the social environment of group exercise can enhance motivation and adherence to physical activity among older PLWH [86].

## Conclusion

Similar to the general population, approximately one-third of PLWH do not achieve the minimum physical activity guidelines despite the numerous health benefits across the life span. HIV providers have a responsibility to assess and promote physical activity in the routine healthcare of all PLWH. These evidence-based practice recommendations can be used as a tool to facilitate engagement in physical activity among PLWH.

## Acknowledgements

J.L.M, C.M.J, D.J.M and K.M.E. provided substantial contribution to the conception of this opinion piece and collaboratively wrote the first draft of the manuscript. All authors critically revised and provided final approval of the manuscript.

We acknowledge Roberto Gallardo for his contribution to the design and revision of Fig. 1.

K.K.O'B. is supported by a Canada Research Chair (CRC) in Episodic Disability and Rehabilitation. K.E. is supported by the National Institute of Aging of the National Institutes of Health K23AG050260 and AG054366.

## Conflicts of interest

There are no conflicts of interest.

## References

- Justice AC, Gordon KS, Skanderson M, Edelman EJ, Akgun KM, Gibert CL, et al. **Nonantiretroviral polypharmacy and adverse health outcomes among HIV-infected and uninfected individuals.** *AIDS* 2018; **32**:739–749.
- d'Ettore G, Ceccarelli G, Giustini N, Mastroianni CM, Silvestri G, Vullo V. **Taming HIV-related inflammation with physical activity: a matter of timing.** *AIDS Res Hum Retroviruses* 2014; **30**:936–944.
- Caspersen CJ, Powell KE, Christenson GM. **Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research.** *Public Health Rep* 1985; **100**:126–131.
- U.S. Department of Health and Human Services. *Physical activity guidelines for Americans*. 2nd ed. Washington, DC: U.S. Department of Health & Human Services; 2018.
- Kelly P, Kahlmeier S, Gotschi T, Orsini N, Richards J, Roberts N, et al. **Systematic review and meta-analysis of reduction in all-cause mortality from walking and cycling and shape of dose response relationship.** *Int J Behav Nutr Phys Act* 2014; **11**:132.
- Hansen D, Niebauer J, Cornelissen V, Barna O, Neunhauserer D, Stettler C, et al. **Exercise prescription in patients with different combinations of cardiovascular disease risk factors: a consensus statement from the EXPERT working group.** *Sports Med* 2018; **48**:1781–1797.
- Silveira EA, Santos A, Falco MO, Cardoso RC, Vitorino PVO. **Association of physical inactivity with hypertension and low educational level in people living with HIV/AIDS.** *AIDS Care* 2018; **30**:1004–1009.
- Mawar N, Katendra T, Bagul R, Bembalkar S, Vedamurthachar A, Tripathy S, et al. **Sudarshan Kriya yoga improves quality of life in healthy people living with HIV (PLHIV): results from an open label randomized clinical trial.** *Indian J Med Res* 2015; **141**:90–99.
- Naoroibam R, Metri KG, Bhargav H, Nagaratna R, Nagendra HR. **Effect of Integrated Yoga (IY) on psychological states and CD4 counts of HIV-1 infected patients: a randomized controlled pilot study.** *Int J Yoga* 2016; **9**:57–61.
- Erlandson KM, MaWhinney S, Wilson M, Gross L, McCandless SA, Campbell TB, et al. **Physical function improvements with moderate or high-intensity exercise among older adults with or without HIV infection.** *AIDS* 2018.
- Ogalha C, Luz E, Sampaio E, Souza R, Zarife A, Neto MG, et al. **A randomized, clinical trial to evaluate the impact of regular physical activity on the quality of life, body morphology and metabolic parameters of patients with AIDS in Salvador, Brazil.** *J Acquir Immune Defic Syndr* 2011; **57** (Suppl 3):S179–185.
- Webel AR, Perazzo J, Decker M, Horvat-Davey C, Sattar A, Voss J. **Physical activity is associated with reduced fatigue in adults living with HIV/AIDS.** *J Adv Nurs* 2016; **72**:3104–3112.
- Cutrono SE, Lewis JE, Perry A, Signorile J, Tiozzo E, Jacobs KA. **The effect of a community-based exercise program on inflammation, metabolic risk, and fitness levels among persons living with HIV/AIDS.** *AIDS Behav* 2016; **20**:1123–1131.
- Pedro RE, Candido N, Guariglia DA, Melo BP, Bertolini DA, Peres SB, et al. **Exercise improves cytokine profile in HIV-infected people: a randomized clinical trial.** *Cytokine* 2017; **99**:18–23.
- Maharaj SS, Yakasai AM. **Does a rehabilitation program of aerobic and progressive resisted exercises influence HIV-induced distal neuropathic pain?** *Am J Phys Med Rehabil* 2018; **97**:364–369.
- Bonato M, Galli L, Passeri L, Longo V, Pavei G, Bossolasco S, et al. **A pilot study of brisk walking in sedentary combination antiretroviral treatment (cART)-treated patients: benefit on soluble and cell inflammatory markers.** *BMC Infect Dis* 2017; **17**:61.
- Stoa EM, Meling S, Nyhus LK, Glenn S, Mangerud KM, Helgerud J, et al. **High-intensity aerobic interval training improves aerobic fitness and HbA1c among persons diagnosed with type 2 diabetes.** *Eur J Appl Physiol* 2017; **117**:455–467.
- Connick E, Greiner J, Dow C, Diehl K, Stauffer B, DeSouza C. **Exercise, oxidative stress, and fibrinolytic function in HIV-1 infected adults.** *Conference on Retroviruses and Opportunistic Infections*; Seattle, Washington, USA: 13–15 February; 2017.
- Kocher M, McDermott M, Lindsey R, Shikuma CM, Gerschenon M, Chow DC, et al. **Short communication: HIV patient systemic mitochondrial respiration improves with exercise.** *AIDS Res Hum Retroviruses* 2017; **33**:1035–1037.
- Hand GA, Phillips KD, Dudgeon WD, William Lyerly G, Larry Durstine J, Burgess SE. **Moderate intensity exercise training reverses functional aerobic impairment in HIV-infected individuals.** *AIDS Care* 2008; **20**:1066–1074.
- Webel AR, Moore SM, Longenecker CT, Currie J, Horvat Davey C, Perazzo J, et al. **Randomized controlled trial of the System-CHANGE intervention on behaviors related to cardiovascular risk in HIV+ adults.** *J Acquir Immune Defic Syndr* 2018; **78**:23–33.
- Ghayomzadeh M, SeyedAlinaghi S, Shamsi MM, Rezaei S, Earnest CP, Akbarnejad S, et al. **Effect of 8-week of hospital-based resistance training program on TCD4+ cell count and anthropometric characteristic of HIV patients in Tehran, Iran: a randomized controlled trial.** *J Strength Cond Res* 2017 [Epub ahead of print].
- Vingren JL, Curtis JH, Levitt DE, Duplanty AA, Lee EC, McFarlin BK, et al. **Adding resistance training to the standard of care for inpatient substance abuse treatment in men with human immunodeficiency virus improves skeletal muscle health without altering cytokine concentrations.** *J Strength Cond Res* 2018; **32**:76–82.
- Zanetti HR, da Cruz LG, Lourenco CL, Ribeiro GC, Ferreira de Jesus Leite MA, Neves FF, et al. **Nonlinear resistance training enhances the lipid profile and reduces inflammation marker in people living with HIV: a randomized clinical trial.** *J Phys Act Health* 2016; **13**:765–770.
- Zanetti HR, Cruz LG, Lourenco CL, Neves Fde F, Silva-Vergara ML, Mendes EL. **Nonlinear resistance training reduces inflammatory biomarkers in persons living with HIV: a randomized controlled trial.** *Eur J Sport Sci* 2016; **16**:1232–1239.
- Souza PM, Jacob-Filho W, Santarem JM, Silva AR, Li HY, Burattini MN. **Progressive resistance training in elderly HIV-positive patients: does it work?** *Clinics (Sao Paulo)* 2008; **63**:619–624.
- Cade WT, Reeds DN, Mondy KE, Overton ET, Grassino J, Tucker S, et al. **Yoga lifestyle intervention reduces blood pressure in HIV-infected adults with cardiovascular disease risk factors.** *HIV Med* 2010; **11**:379–388.
- Agarwal RP, Kumar A, Lewis JE. **A pilot feasibility and acceptability study of yoga/meditation on the quality of life and markers of stress in persons living with HIV who also use crack cocaine.** *J Altern Complement Med* 2015; **21**:152–158.
- Webel AR, Perazzo J, Longenecker CT, Jenkins T, Sattar A, Rodriguez M, et al. **The influence of exercise on cardiovascular health in sedentary adults with human immunodeficiency virus.** *J Cardiovasc Nurs* 2018; **33**:239–247.
- Forde C, Loy A, O'Dea S, Mulcahy F, Gormley J, Daly C. **Physical activity is associated with metabolic health in men living with HIV.** *AIDS Behav* 2018; **22**:1965–1971.
- Wirth MD, Jagers JR, Dudgeon WD, Hebert JR, Youngstedt SD, Blair SN, et al. **Association of markers of inflammation with sleep and physical activity among people living with HIV or AIDS.** *AIDS Behav* 2015; **19**:1098–1107.
- Erlandson KM, Wu K, Koletar SL, Kalayjian RC, Ellis RJ, Taiwo B, et al. **Association between frailty and components of the frailty phenotype with modifiable risk factors and antiretroviral therapy.** *J Infect Dis* 2017; **215**:933–937.
- Johs NA, Wu K, Tassiopoulos K, Koletar SL, Kalayjian RC, Ellis RJ, et al. **Disability among middle-aged and older persons with human immunodeficiency virus infection.** *Clin Infect Dis* 2017; **65**:83–91.
- Monroe AK, Zhang L, Jacobson LP, Plankey MW, Brown TT, Miller EN, et al. **The association between physical activity and cognition in men with and without HIV infection.** *HIV Med* 2017; **18**:555–563.
- Fazeli PL, Marquine MJ, Dufour C, Henry BL, Montoya J, Gouaux B, et al. **Physical activity is associated with better neurocognitive and everyday functioning among older adults with HIV disease.** *AIDS Behav* 2015; **19**:1470–1477.
- Perazzo JD, Webel AR, Alam SMK, Sattar A, McComsey GA. **Relationships between physical activity and bone density in people living with HIV: results from the SATURN-HIV study.** *J Assoc Nurses AIDS Care* 2018; **29**:528–537.
- Dirajlal-Fargo S, Webel AR, Longenecker CT, Kinley B, Labbato D, Sattar A, et al. **The effect of physical activity on cardiometabolic health and inflammation in treated HIV infection.** *Antivir Ther* 2016; **21**:237–245.



38. Erlandson KM, Lake JE, Sim M, Falutz J, Prado CM, Domingues da Silva AR, et al. **Bone mineral density declines twice as quickly among HIV-infected women compared with men.** *J Acquir Immune Defic Syndr* 2018; **77**:288–294.
39. Kazooba P, Kasamba I, Mayanja BN, Lutaakome J, Namakoola I, Salome T, et al. **Cardiometabolic risk among HIV-POSITIVE Ugandan adults: prevalence, predictors and effect of long-term antiretroviral therapy.** *Pan Afr Med J* 2017; **27**:40.
40. Dufour CA, Marquine MJ, Fazeli PL, Umlauf A, Henry BL, Zlatar Z, et al. **A longitudinal analysis of the impact of physical activity on neurocognitive functioning among HIV-infected adults.** *AIDS Behav* 2018; **22**:1562–1572.
41. Dufour CA, Marquine MJ, Fazeli PL, Henry BL, Ellis RJ, Grant I, et al. **Physical exercise is associated with less neurocognitive impairment among HIV-infected adults.** *J Neurovirol* 2013; **19**:410–417.
42. Vancampfort D, Mugisha J, De Hert M, Probst M, Firth J, Gorczynski P, et al. **Global physical activity levels among people living with HIV: a systematic review and meta-analysis.** *Disabil Rehabil* 2018; **40**:388–397.
43. Clingerman EM. **Participation in physical activity by persons living with HIV disease.** *J Assoc Nurses AIDS Care* 2003; **14**:59–70.
44. Bui TV, Blizzard CL, Luong KN, Truong Nle V, Tran BQ, Otahal P, et al. **Physical activity in Vietnam: estimates and measurement issues.** *PLoS One* 2015; **10**:e0140941.
45. Dang AK, Nguyen LH, Nguyen AQ, Tran BX, Tran TT, Latkin CA, et al. **Physical activity among HIV-positive patients receiving antiretroviral therapy in Hanoi and Nam Dinh, Vietnam: a cross-sectional study.** *BMJ Open* 2018; **8**:e020688.
46. Marzel A, Kouyos RD, Reinschmidt S, Balzer K, Garon F, Spitaleri M, et al. **Dietary patterns and physical activity correlate with total cholesterol independently of lipid-lowering drugs and antiretroviral therapy in aging people living with human immunodeficiency virus.** *Open Forum Infect Dis* 2018; **5**:ofy067.
47. Stein L, Hechler D, Jessen AB, Neumann K, Jessen H, Beneke R. **Sports behaviour among HIV-infected versus noninfected individuals in a Berlin cohort.** *Int J STD AIDS* 2012; **23**:25–29.
48. Vader K, Simonik A, Ellis D, Kesbian D, Leung P, Jachyra P, et al. **Perceptions of 'physical activity' and 'exercise' among people living with HIV: a qualitative study.** *Int J Ther Rehabil* 2017; **24**:473–482.
49. Cade WT, Fantry LE, Nabar SR, Keyser RE. **Decreased peak arteriovenous oxygen difference during treadmill exercise testing in individuals infected with the human immunodeficiency virus.** *Arch Phys Med Rehabil* 2003; **84**:1595–1603.
50. Galantino ML, Shepard K, Krafft L, Laperriere A, Ducette J, Sorbello A, et al. **The effect of group aerobic exercise and t'ai chi on functional outcomes and quality of life for persons living with acquired immunodeficiency syndrome.** *J Altern Complement Med* 2005; **11**:1085–1092.
51. Oursler KK, Sorkin JD, Ryan AS, Katzel LI. **A pilot randomized aerobic exercise trial in older HIV-infected men: insights into strategies for successful aging with HIV.** *PLoS One* 2018; **13**:e0198855.
52. Voigt N, Cho H, Schnall R. **Supervised physical activity and improved functional capacity among adults living with HIV: a systematic review.** *J Assoc Nurses AIDS Care* 2018; **29**:667–680.
53. Vancampfort D, Mugisha J, Richards J, De Hert M, Probst M, Stubbs B. **Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies.** *Disabil Rehabil* 2018; **40**:1618–1629.
54. Montgomery CA, Henning KJ, Kantarzi SR, Kideckel TB, Yang CF, O'Brien KK. **Experiences participating in a community-based exercise programme from the perspective of people living with HIV: a qualitative study.** *BMJ Open* 2017; **7**:e015861.
55. Kaku M, Simpson DM. **HIV neuropathy.** *Curr Opin HIV AIDS* 2014; **9**:521–526.
56. Barroso J, Leserman J, Harmon JL, Hammill B, Pence BW. **Fatigue in HIV-infected people: a three-year observational study.** *J Pain Symptom Manage* 2015; **50**:69–79.
57. Oursler KK, Katzel LI, Smith BA, Scott WB, Russ DW, Sorkin JD. **Prediction of cardiorespiratory fitness in older men infected with the human immunodeficiency virus: clinical factors and value of the six-minute walk distance.** *J Am Geriatr Soc* 2009; **57**:2055–2061.
58. Oursler KK, Sorkin JD, Smith BA, Katzel LI. **Reduced aerobic capacity and physical functioning in older HIV-infected men.** *AIDS Res Hum Retroviruses* 2006; **22**:1113–1121.
59. Ortmeyer HK, Ryan AS, Hafer-Macko C, Oursler KK. **Skeletal muscle cellular metabolism in older HIV-infected men.** *Physiol Rep* 2016; **4**:pii: e12794.
60. Payne BA, Gardner K, Chinnery PF. **Mitochondrial DNA mutations in ageing and disease: implications for HIV?** *Antivir Ther* 2015; **20**:109–120.
61. Campo M, Oursler KK, Huang L, Goetz MB, Rimland D, Hoo GS, et al. **Association of chronic cough and pulmonary function with 6-min walk test performance in HIV infection.** *J Acquir Immune Defic Syndr* 2014; **65**:557–563.
62. Kristoffersen US, Lebeck AM, Mortensen J, Gerstoft J, Gutte H, Kjaer A. **Changes in lung function of HIV-infected patients: a 4.5-year follow-up study.** *Clin Physiol Funct Imaging* 2012; **32**:288–295.
63. Ley C, Barrio MR, Leach L. **Social-ecological, motivational and volitional factors for initiating and maintaining physical activity in the context of HIV.** *Open AIDS J* 2015; **9**:96–103.
64. Rehm KE, Konkle-Parker D. **Physical activity levels and perceived benefits and barriers to physical activity in HIV-infected women living in the deep south of the United States.** *AIDS Care* 2016; **28**:1205–1210.
65. Webel AR, Perazzo JD, Dawson-Rose C, Smith C, Nicholas PK, Rivero-Mendez M, et al. **A multinational qualitative investigation of the perspectives and drivers of exercise and dietary behaviors in people living with HIV.** *Appl Nurs Res* 2017; **37**:13–18.
66. Roos R, Myezwa H, van Aswegen H. **Not easy at all but I am trying: barriers and facilitators to physical activity in a South African cohort of people living with HIV participating in a home-based pedometer walking programme.** *AIDS Care* 2015; **27**:235–239.
67. Li A, McCabe T, Silverstein E, Dragan S, Salbach NM, Zobeiry M, et al. **Community-based exercise in the context of HIV: factors to consider when developing and implementing community-based exercise programs for people living with HIV.** *J Int Assoc Provid AIDS Care* 2017; **16**:267–275.
68. Simonik A, Vader K, Ellis D, Kesbian D, Leung P, Jachyra P, et al. **Are you ready? Exploring readiness to engage in exercise among people living with HIV and multimorbidity in Toronto, Canada: a qualitative study.** *BMJ Open* 2016; **6**:e010029.
69. Lobelo F, Rohm Young D, Sallis R, Garber MD, Billinger SA, Duperly J, et al. **Routine assessment and promotion of physical activity in healthcare settings: a scientific statement from the American Heart Association.** *Circulation* 2018; **137**:e495–e522.
70. Aberg JA, Gallant JE, Ghanem KG, Emmanuel P, Zingman BS, Horberg MA, et al. **Primary care guidelines for the management of persons infected with HIV: 2013 update by the HIV medicine association of the Infectious Diseases Society of America.** *Clin Infect Dis* 2014; **58**:e1–e34.
71. Sanchez A, Bully P, Martinez C, Grandes G. **Effectiveness of physical activity promotion interventions in primary care: a review of reviews.** *Prev Med* 2015; **Suppl 76**:S56–S67.
72. Cress ME, Buchner DM, Prohaska T, Rimmer J, Brown M, Macera C, et al. **Best practices for physical activity programs and behavior counseling in older adult populations.** *J Aging Phys Act* 2005; **13**:61–74.
73. Neale AV, Singleton SP, Dupuis MH, Hess JW. **The use of behavioral contracting to increase exercise activity.** *Am J Health Promot* 1990; **4**:441–447.
74. American College of Sports Medicine. **Exercise is medicine.** 2018. [https://www.exerciseismedicine.org/assets/page\\_documents/EIM%20Prescription%20pad%201-up.pdf](https://www.exerciseismedicine.org/assets/page_documents/EIM%20Prescription%20pad%201-up.pdf)
75. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. **The physical activity guidelines for Americans.** *JAMA* 2018; **320**:2020–2028.
76. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. **Effective techniques in healthy eating and physical activity interventions: a meta-regression.** *Health Psychol* 2009; **28**:690–701.
77. Michie S, Ashford S, Sniehotta FF, Dombrowski SU, Bishop A, French DP. **A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating behaviours: the CALO-RE taxonomy.** *Psychol Health* 2011; **26**:1479–1498.

78. Goode AP, Hall KS, Batch BC, Huffman KM, Hastings SN, Allen KD, *et al.* **The impact of interventions that integrate accelerometers on physical activity and weight loss: a systematic review.** *Ann Behav Med* 2017; **51**:79–93.
79. National Institute on Aging. Go4Life. 2018. <https://go4life.nia.nih.gov> [Accessed 11 May 2018]
80. Dagenais M, Cheng D, Salbach NM, Brooks D, O'Brien KK. **Wireless physical activity monitor use among adults living with HIV: a scoping review.** *Rehab Oncol* (in press).
81. McEwan D, Harden SM, Zumbo BD, Sylvester BD, Kaulius M, Ruissen GR, *et al.* **The effectiveness of multicomponent goal setting interventions for changing physical activity behaviour: a systematic review and meta-analysis.** *Health Psychol Rev* 2016; **10**:67–88.
82. Gardner B, Lally P, Wardle J. **Making health habitual: the psychology of 'habit-formation' and general practice.** *Br J Gen Pract* 2012; **62**:664–666.
83. Henry BL, Moore DJ. **Preliminary findings describing participant experience with iSTEP, an mHealth intervention to increase physical activity and improve neurocognitive function in people living with HIV.** *J Assoc Nurses AIDS Care* 2016; **27**:495–511.
84. Montoya JL, Wing D, Knight A, Moore DJ, Henry BL. **Development of an mHealth ontervention (iSTEP) to promote physical activity among people living with HIV.** *J Int Assoc Provid AIDS Care* 2015; **14**:471–475.
85. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, *et al.* **The effectiveness of interventions to increase physical activity. A systematic review.** *Am J Prev Med* 2002; **22** (Suppl 4):73–107.
86. Neff HA, Kellar-Guenther Y, Jankowski CM, Worthington C, McCandless SA, Jones J, *et al.* **Turning disability into ability: barriers and facilitators to initiating and maintaining exercise among older men living with HIV.** *AIDS Care* 2018:1–5.