

RESEARCH ARTICLE

Evidence of Drought Stress Memory in the Facultative CAM, *Aptenia cordifolia*: Possible Role of Phytohormones

Eva Fleta-Soriano, Marta Pintó-Marijuan, Sergi Munné-Bosch*

Departament de Biologia Vegetal, Facultat de Biologia, Universitat de Barcelona, Avinguda Diagonal, 643, E-08028, Barcelona, Spain

* smunne@ub.edu



OPEN ACCESS

Citation: Fleta-Soriano E, Pintó-Marijuan M, Munné-Bosch S (2015) Evidence of Drought Stress Memory in the Facultative CAM, *Aptenia cordifolia*: Possible Role of Phytohormones. PLoS ONE 10(8): e0135391. doi:10.1371/journal.pone.0135391

Editor: Belay T. Ayele, University of Manitoba, CANADA

Received: May 6, 2015

Accepted: July 21, 2015

Published: August 14, 2015

Copyright: © 2015 Fleta-Soriano et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The research was fully funded by the Generalitat de Catalunya under the ICREA Academia award given to SMB. The funder had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript. MPM was supported by a postdoctoral fellowship from Generalitat de Catalunya (2013 BP-B 00235).

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Although plant responses to drought stress have been studied in detail in several plant species, including CAM plants, the occurrence of stress memory and possible mechanisms for its regulation are still very poorly understood. In an attempt to better understand the occurrence and possible mechanisms of regulation of stress memory in plants, we measured the concentrations of phytohormones in *Aptenia cordifolia* exposed to reiterated drought, together with various stress indicators, including leaf water contents, photosynthesis and mechanisms of photo- and antioxidant protection. Results showed that plants exposed to drought stress responded differently if previously challenged with a first drought. Gibberellin levels decreased upon exposure to the first drought and remained lower in double-stressed plants compared with those exposed to stress for the first time. In contrast, abscisic acid levels were higher in double- than single-stressed plants. This occurred in parallel with alterations in hydroperoxide levels, but not with malondialdehyde levels, thus suggesting an increased oxidation state that did not result in oxidative damage in double-stressed plants. It is concluded that (i) drought stress memory occurs in double-stressed *A. cordifolia* plants, (ii) both gibberellins and abscisic acid may play a role in plant response to repeated periods of drought, and (iii) changes in abscisic acid levels in double-stressed plants may have a positive effect by modulating changes in the cellular redox state with a role in signalling, rather than cause oxidative damage to the cell.

Introduction

Stress memory in plants, also known as stress imprint or priming, is considered to be an important component of the behavioral ecology of plants and it is becoming an increasingly important part of plant stress physiology textbooks nowadays. Stress memory is defined as the capacity of organisms to respond better to a given stress factor when individuals have already been challenged previously with the same stimulus relative to those that have not been exposed to the stress before [1]. Indeed, there is an increasing interest in stress memory effects, since

this feature has important implications in plant stress physiology [2]. Unfortunately, the occurrence of stress memory, either leading to positive or negative effects in plant stress responses, and the mechanisms underlying stress memory are still very poorly understood [2]. Therefore, a better knowledge on stress memory effects is urgently needed not only to better understand the physiology and ecology of plants, but also to improve crop production and environmental management practices.

Mechanisms of stress memory may largely vary depending on the organizational level to what the studies are carried out, from changes in leaf anatomy to epigenomics, including phenological, biochemical and physiological mechanisms that may operate in an integrated way to fulfill a role in plant stress tolerance. Among these mechanisms, it appears that phytohormones may have a prominent role. Gibberellins (GAs) have long been known to be involved in vernalization, which implies epigenetic changes and long-term memory effects [3]. On the other hand, recent studies suggest that abscisic acid (ABA) may be involved in short-term drought stress acclimation in the model plant, *Arabidopsis thaliana*. It has been shown that *A. thaliana* increase the transcription of several ABA-induced genes in response to reiterated dehydration, while maintaining leaf water contents [4,5]. Guard cells appear to have a dehydration stress memory so that plants produce ABA to keep partially closed stomata in order to reduce water loss under reiterated water deficit conditions [6].

Crassulacean acid metabolism (CAM) is an adaptive mechanism to survive in extreme habitats characterized by severe drought in which the carbon dioxide is assimilated during the night avoiding an excessive water loss [7]. In fact, CAM plants are able to keep a minimal metabolically active state for a long time during severe droughts, while they are able to recover quickly during re-watering [8]. Therefore, drought stress memory in CAM plants, which are specialized in drought stress tolerance, may have a tremendous biological significance. While drought stress responses have been extensively studied in CAM plants [9–11], the occurrence of stress memory in CAM plants and the possible mechanisms for its regulation are still very poorly understood.

Here, we hypothesized that the CAM plant, *Aptenia cordifolia* may show a drought stress memory in plant response to drought stress, in which phytohormones could play a role. Since *A. cordifolia* is an invasive species, this capacity could help displace other species less resistant to drought stress and colonize new habitats. Specifically, in the present study, we examined whether or not *A. cordifolia* show any stress memory to reiterated drought. With this aim, we measured the endogenous levels of phytohormones, together with various markers of physiological stress, in double- compared with single-stressed plants.

Materials and Methods

Plant material, treatments and sampling

Sixty plants of baby sun rose (*Aptenia cordifolia* (L.f.) Schwantes) were purchased in a local garden (Ca L'Agustí, Barcelona, Spain) and were transferred to 0.5 L-pots with peat:perlite:vermiculite (2:1:1, v/v). Plants were grown in a greenhouse at the Faculty of Biology of the University of Barcelona (Barcelona, Spain). Prior to experiments, plants were watered 3 times per week with half-diluted Hoagland nutrient solution. Experiments, which started on 11th June 2014, consisted in developing two water regimes on plants: CS plants were watered for 13 days, exposed to drought by withholding water for 10 days, and then re-watered for 4 days; while SS plants were stressed by withholding water for 9 days, recovered for 4 days and then exposed again to drought for 10 days, followed by a final recovery of 4 days. Therefore, SS were double-stressed, while CS plants were stressed during a single period. Samplings were performed at the beginning of the experiment (day 0) and after 9, 13, 23 and 27 days of treatments, that is at the

points of maximum stress and during recovery. All measurements were performed at midday (between 11 and 13h solar time). At each sampling point, fully-expanded young leaves of 7 individuals were used to estimate the endogenous contents of phytohormones, together with various physiological indicators, including leaf water contents and gas exchange, chlorophyll fluorescence, levels of photosynthetic pigments and antioxidants, and the extent of lipid peroxidation in leaves. Samples for phytohormone and other biochemical analyses were collected, immediately frozen in liquid nitrogen and stored at -80°C until analysis.

Hormonal profiling

The extraction and analysis of GAs, including the bioactive GA_1 and GA_4 , and their precursors GA_9 , GA_{19} , GA_{20} and GA_{24} , the bioactive auxin, indole-3-acetic acid (IAA), the cytokinins, zeatin (Z), zeatin riboside (ZR), 2-isopentenyl adenine (2iP) and isopentenyl adenosine (iPA), and the stress-related phytohormones, ABA, jasmonic acid (JA) and salicylic acid (SA) were carried out by UPLC-MS/MS as described [12]. Deuterium-labelled phytohormones were used as internal standards.

Leaf water contents, gas exchange and chlorophyll fluorescence

Samples were weighed to estimate the fresh matter (FW), immersed in distilled water at 4°C for 24h to estimate the turgid matter (TW) and then oven-dried at 80°C to constant weight to estimate the dry matter (DW). Relative water content (RWC) was then calculated as $100 \times (\text{FW} - \text{DW}) / (\text{TW} - \text{DW})$. Net photosynthesis (A), stomatal conductance (g_s) and the maximum efficiency of photosystem II photochemistry (F_v/F_m) were estimated by using a portable infrared gas analyzer with a leaf chamber fluorometer (LI-COR 6400 system, LI-COR, Lincoln, NE, USA). Light intensity was set at $700 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ with a 10% of blue light; CO_2 concentration at 400 ppm; leaf temperature at $20\text{--}25^{\circ}\text{C}$, and relative humidity ranging 50–60% with a flow of $500 \mu\text{mol s}^{-1}$. The F_v/F_m ratio, which was measured after adapting the leaves to darkness for 2 h, was calculated as $(F_m - F_0) / F_m$, where F_m and F_0 are the maximum and basal fluorescence yields, respectively, of dark-adapted leaves [13].

Photosynthetic pigments and antioxidants

For pigment and tocopherol analysis, leaf samples (50 mg) were ground in liquid nitrogen and extracted with cold methanol (v/v) using ultrasonication. After centrifuging at 8000 rpm for 10 min and 4°C , the supernatant was collected and the pellet was re-extracted with the same solvent until it was colourless. Then, supernatants were pooled and filtered through a $0.5 \mu\text{m}$ syringe filter. Total chlorophylls and carotenoids were estimated spectrophotometrically as described [14]. Levels of neoxanthin and violaxanthin, ABA precursors, were measured by high performance liquid chromatography (HPLC) as described [15]. Tocopherols were measured by HPLC as described [16].

Estimation of lipid peroxidation

The extent of lipid peroxidation was estimated by measuring the levels of lipid hydroperoxides (primary stable products of lipid peroxidation) and malondialdehyde (MDA) equivalents (secondary products of lipid peroxidation) in leaves. Lipid hydroperoxides levels were estimated spectrophotometrically following the ferrous oxidation-xylenol orange assay as described [17]. MDA levels were estimated spectrophotometrically following the thiobarbituric acid-reactive assay considering the effect of potential interfering compounds, as described [18].

Statistical analysis

In the first set of results, which included a time-course evolution of water contents, leaf gas exchange and phytohormone levels, data was analyzed by using two-way factorial analysis of variance (ANOVA) with treatment and time (sampling day) as factors, and by additionally using Duncan posthoc tests. In the second set of results, in which differences in photosynthetic pigments, antioxidants and the extent of lipid peroxidation in a single time point were analyzed, Student's t-tests were used. In all cases, differences were considered significant at a probability level of $P < 0.05$. All statistical tests were carried out using the SPSS 15.0 statistical package (SPSS, Inc., Chicago, IL, USA).

Results and Discussion

Drought-stressed *A. cordifolia* plants induce CAM metabolism

A number of CAM plants are extremely resistant to drought, as they are commonly adapted to deserts and other arid or semi-arid environments [19]. Aside from stomatal closure during the day, which largely increases water use efficiency, CAM plants adopt a series of strategies to resist water deprivation during long periods, including, among others, specific hormonal responses and activation of mechanisms of photo- and antioxidant protection [9]. *A. cordifolia* was first described as a facultative CAM species [20], and later thought to be an obligate CAM [21]. In the present study, however, it is shown that it is a facultative CAM plant that opens stomata at midday when water is available and closing them completely when water is withheld (Fig 1). After 9 days of withholding water, the RWC decreased from 68% to 40% at midday. This was associated with a sharp reduction in CO₂ assimilation and stomatal conductance rates, which reached values of and close to zero, respectively (Fig 1). The stress caused to plants, as indicated by RWC values, was quite severe. When challenged with a new drought, plants reduced again the RWC and gas exchange values to a similar extent. Double-stressed plants did not respond differently as those stressed for the first time in terms of leaf water contents and gas exchange (Fig 1), thus indicating that a second stress did impact neither positively nor negatively on these parameters.

GAs and ABA levels reveal stress memory

Among the analysed phytohormones, the bioactive GA₄ was the one showing the most relevant results (Figs 2 and 3, Table 1). GA₄ levels decreased during drought and did not recover, so that double-stressed plants showed slightly, but consistently lower GA₄ levels throughout the experiment compared with plants challenged by drought for the first time (Fig 2). While GA₉ levels were not affected by drought stress, the endogenous concentrations of GA₂₄ increased during the first drought, to decrease later during the second drought in double-stressed plants (Fig 2). Since GA₂₄ is a precursor of GA₄ [22], it is likely that conversion to bioactive GAs is reduced during the first drought, thus leading to an accumulation of GA precursors. When challenged again with a second stress, however, it seems that this effect disappears, so that GA precursors do not accumulate (despite bioactive GA levels were kept at low levels). Although still to be confirmed using enzymatic assays and molecular tools, these results suggest a memory effect on GA metabolism in plant response to reiterated drought, in analogy to the regulation of GA metabolism by vernalization [23]. It is noteworthy that levels of GAs from the GA₄ pathway were affected by reiterated stress (Figs 2 and 3), and that neither auxin, cytokinins, salicylic acid nor jasmonic acid levels were affected by repeated periods of drought (Table 1).

ABA showed differences in double-stressed plants compared with plants challenged with drought for the first time (Fig 4). ABA levels increased 40-fold in response to the first drought




Fig 1. Relative water content (RWC), CO₂ assimilation (A) and stomatal conductance (g_s) in leaves of *A. cordifolia*. Data represent the mean \pm SE of $n = 7$ individuals. Significant differences between groups were tested by two-way analysis of variance (ANOVA) and Duncan posthoc tests.

doi:10.1371/journal.pone.0135391.g001




Fig 2. Endogenous concentrations of gibberellin 4 (GA₄), and its precursors, gibberellin 9 (GA₉) and gibberellin 24 (GA₂₄) in leaves of *A. cordifolia*. Data represent the mean \pm SE of $n = 7$ individuals. Significant differences between groups were tested by two-way analysis of variance (ANOVA) and Duncan posthoc tests.

doi:10.1371/journal.pone.0135391.g002




Fig 3. Endogenous concentrations of gibberellin 1 (GA₁), and its precursor, gibberellin 20 (GA₂₀) in leaves of *A. cordifolia*. Data represent the mean ± SE of *n* = 7 individuals. Significant differences between groups were tested by two-way analysis of variance (ANOVA) and Duncan posthoc tests.

doi:10.1371/journal.pone.0135391.g003

to recover later to initial pre-drought values. During the second period of stress, however, ABA levels increased more in double-stressed plants compared with plants challenged by drought for the first time (ABA values reaching 4.2 vs. 3.4 μg/g DW, respectively, Fig 4). Despite double-stressed plants did not respond differently in terms of leaf water contents and gas exchange (Fig 1), they did so in terms of ABA accumulation, enhancing the endogenous levels of this phytohormone during the second stress compared with plants challenged by drought for the first time. Increases of ABA levels under reiterated stress may have an effect of growth regulation [24], osmotic adjustment [25] and antioxidant responses [26–28]. Furthermore, previous

Table 1. P values of the analysis of variance (ANOVA) to test the effect of treatment, sampling time and its interaction on the levels of phytohormones in leaves of *A. cordifolia*.

Hormone	Treatment	Time	Interaction
IAA	NS	0.001	0.003
iPA	NS	0.001	NS
2iP	NS	0.001	0.008
Z	NS	0.041	NS
ZR	NS	0.001	0.001
SA	NS	NS	0.001
JA	NS	0.042	0.001

IAA, indole-3-acetic acid; iPA, isopentenyl adenosine; 2iP, isopentenyl adenine; Z, zeatin; ZR, zeatin riboside; SA, salicylic acid; JA, jasmonic acid. NS, not significant ($P > 0.050$).

doi:10.1371/journal.pone.0135391.t001

studies have shown that ABA exerts a protective role under reiterated drought by reprogramming gene expression [4–6]. In the present study, plants did not suffer photo-inhibitory damage to the photosynthetic apparatus, as indicated by constant F_v/F_m values (Table 2), and plants were able to fully recover after 4 days of re-watering (Fig 1), thus indicating that the observed changes in the endogenous levels of phytohormones may indeed have a positive effect on the physiology of double-stressed plants.

ABA could exert a drought memory effect by modulating antioxidant responses. By comparing various photo-oxidative stress and lipid peroxidation markers in double- vs. single-stressed plants, it was found that reiterated stress had a significant impact on the chlorophyll a/




Fig 4. Endogenous concentrations of abscisic acid (ABA) in leaves of *A. cordifolia*. Data represent the mean \pm SE of $n = 7$ individuals. Significant differences between groups were tested by two-way analysis of variance (ANOVA) and Duncan posthoc tests.

doi:10.1371/journal.pone.0135391.g004

Table 2. F_v/F_m ratio, photosynthetic pigment and antioxidant levels, and extent of lipid peroxidation in double-stressed plants (SS) compared with plants exposed to drought for the first time (CS). Data represent the mean \pm SE of $n = 7$.

Parameter	CS	SS
F_v/F_m	0.68 \pm 0.02	0.68 \pm 0.01
Chlorophyll <i>a+b</i> (mg/gDW)	11.4 \pm 1.0	10.6 \pm 0.3
Chlorophyll <i>a/b</i> (g/g)	1.98 \pm 0.01	2.04 \pm 0.01*
Carotenoids/ Chlorophyll <i>a+b</i> (mg/g)	206 \pm 2	207 \pm 2
Neoxanthin/Chlorophyll <i>a+b</i> (mg/g)	12.5 \pm 1.4	12.0 \pm 1.6
Violaxanthin/Chlorophyll <i>a+b</i> (mg/g)	27.3 \pm 0.2	26.3 \pm 0.4*
α -Tocopherol/Chlorophyll <i>a+b</i> (mg/g)	21.9 \pm 3.5	23.8 \pm 2.0
γ -Tocopherol/ Chlorophyll <i>a+b</i> (mg/g)	7.3 \pm 1.1	6.5 \pm 0.5
Ascorbate (μ mol/gDW)	11.41 \pm 0.67	10.3 \pm 0.58
Dehydroascorbate/Total ascorbate (%)	12.43 \pm 1.95	11.3 \pm 1.95
Lipid hydroperoxides (μ mol equiv. H_2O_2 /gDW)	3.9 \pm 0.4	7.4 \pm 0.8*
MDA (nmol equiv./gDW)	9.6 \pm 2.7	13.6 \pm 2.9

* Significant differences between treatments (Student's *t*-test, $P < 0.050$)

doi:10.1371/journal.pone.0135391.t002

b ratio and the extent of lipid peroxidation in leaves (Table 2). The chlorophyll *a/b* ratio increased, however, by 3% only in double- compared with single-stressed plants. In contrast, the extent of lipid peroxidation increased by 90% in double- vs. single-stressed plants, as indicated by lipid hydroperoxide levels, respectively. In contrast, malondialdehyde levels were not affected by reiterated drought (Table 2). Lipid hydroperoxide and malondialdehyde levels are the primary and secondary stable products of lipid peroxidation, respectively [29]. Therefore, results obtained suggest that double-stressed plants experienced an increased oxidative stress compared with plants challenged by drought for the first time. However, neither

Table 3. Correlation coefficients and *P* values (in parentheses) of the Spearman rank's correlations between the endogenous concentrations of all phytohormones analyzed and the levels of α - and γ -tocopherol in *A. cordifolia* leaves. Significant correlations are indicated in bold (Bonferroni corrected, $P < 0.004$).

Hormone	α -Tocopherol	γ -Tocopherol
GA ₁	-0.010 (0.470)	-0.177 (0.107)
GA ₄	-0.286 (0.012)	-0.316 (0.011)
GA ₉	0.001 (0.497)	-0.160 (0.111)
GA ₂₀	0.141 (0.138)	-0.066 (0.32)
GA ₂₄	0.157 (0.112)	-0.128 (0.191)
IAA	0.057 (0.331)	-0.217 (0.045)
Z	0.048 (0.357)	-0.262 (0.020)
ZR	0.046 (0.362)	0.072 (0.290)
2iP	0.279 (0.015)	0.285 (0.012)
iPA	-0.223 (0.042)	-0.272 (0.018)
ABA	0.331 (0.005)	0.552 (<0.001)
JA	-0.125 (0.167)	-0.218 (0.044)
SA	-0.224 (0.040)	-0.374 (0.001)

GAs, gibberellins; IAA, indole-3-acetic acid; Z, zeatin; ZR, zeatin riboside; 2iP, isopentenyl adenine; iPA, isopentenyl adenosine; SA, salicylic acid; JA, jasmonic acid.

doi:10.1371/journal.pone.0135391.t003

malondialdehyde levels nor the F_v/F_m ratio were affected by reiterated drought, thus indicating absence of increased photo-oxidative damage in double- vs. single-stressed plants [30]. Results obtained appear therefore to be consistent with a hormonal and redox-related gene reprogramming, an aspect that warrants further research. In this respect, it has been previously shown that ABA may trigger activation of antioxidant defences, including vitamin E biosynthesis [31,32]. Despite neither α - nor γ -tocopherol increased in double- vs. single-stressed plants (Table 2), a correlative analysis revealed that ABA concentrations positively correlated with γ -tocopherol levels (Table 3). A relationship between ABA and vitamin E biosynthesis is also supported by previous correlative studies in the same species [9] and by studies on model plants, showing that tocopherol-biosynthesis genes have ABA-response elements in their promoter region [33]. Furthermore, analysis of ABA precursors, the carotenoids neoxanthin and violaxanthin, revealed a reduction by 3.7% in violaxanthin levels in double- compared with single-stressed plants (Table 2). Although this reduction appears to be small, it should be considered that it is in the order of 1 mg/g chlorophyll (equivalent to 1.5 mmol/mol chlorophyll), which corresponds to 350 nmol/g DW, while the observed enhanced ABA levels in double- compared with single-stressed plants were in the order of 0.8 μ g/g DW (Fig 4), which corresponds to 3 nmol/g DW. Therefore, a 3.7% reduction in violaxanthin levels was more than sufficiently enough to account for the 24% increase of ABA levels in double- compared with single-stressed plants. Further research is however needed to confirm a memory effect on ABA metabolism by directly measuring gene expression, amounts and/or activity of 9-*cis*-epoxycarotenoid dioxygenases, the committed step in ABA biosynthesis from carotenoids [34]. Moreover, it is still to be demonstrated whether or not ABA catabolism is additionally affected in double- compared with single-stressed plants.

Conclusions

Phytohormones may be involved in plant response to reiterated drought stress in the invasive CAM plant, *A. cordifolia*. GA levels decreased upon exposure to the first drought and remained lower in double- compared with single-stressed plants. In contrast, ABA levels were higher in double- than in single-stressed plants. This occurred in parallel with alterations in primary products of lipid peroxidation, but not with changes in malondialdehyde levels and the F_v/F_m ratio, thus suggesting an increased oxidative stress that did not result in photo-oxidative damage in double-stressed plants. It is concluded that (i) drought stress memory occurs in double-stressed *A. cordifolia* plants, (ii) both GAs and ABA may play a role in plant response to repeated periods of drought, and (iii) changes in ABA levels in double-stressed plants may have a positive effect by modulating changes in the cellular redox state with a role in signalling, rather than cause oxidative damage to the cell. In other words, changes in ABA levels in double-stressed plants may be associated with an increased oxidative stress that did not result in photo-oxidative damage. Further research is needed to better understand the GA-, ABA- and redox-mediated effects in plant stress acclimation to reiterated drought in plants.

Acknowledgments

We are very grateful to the Serveis dels Camps Experimentals and Serveis Científico-tècnics (Universitat de Barcelona) for technical assistance. We also thank Jana Cela, Alba Cotado and Verónica Tijero for their help in samplings and pigment analyses, and Maren Müller for her help in phytohormone analysis.

Author Contributions

Conceived and designed the experiments: EFS MPM SMB. Performed the experiments: EFS MPM. Analyzed the data: EFS MPM SMB. Contributed reagents/materials/analysis tools: SMB. Wrote the paper: EFS SMB.

References

1. Trewavas A. Green plants as intelligent organisms. *Trends Plant Sci.* 2005; 10: 413–419. PMID: [16054860](#)
2. Bruce TJA, Matthes MC, Napier JA, Pickett JA. Stressful “memories” of plants: Evidence and possible mechanisms. *Plant Sci.* 2007; 173: 603–608.
3. Songling B, Takanori S, Daisuke S, Akiko I, Hiroshi F, Takaya M. Transcriptome analysis of Japanese pear (*Pyrus pyrifolia* Nakai.) flower buds transitioning through endodormancy. *Plant Cell Physiol.* 2013; 54: 1132–1151. doi: [10.1093/pcp/pct067](#) PMID: [23624675](#)
4. Ding Y, Fromm M, Avramova Z. Multiple exposures to drought ‘train’ transcriptional responses in *Arabidopsis*. *Nat. Commun.* 2012; 3: 740. doi: [10.1038/ncomms1732](#) PMID: [22415831](#)
5. Virilouvet L, Fromm M. Physiological and transcriptional memory in guard cells during repetitive dehydration stress. *New Phytol.* 2015; 205: 596–607. doi: [10.1111/nph.13080](#) PMID: [25345749](#)
6. Virilouvet L, Ding Y, Fujii H, Avramova Z, Fromm M. ABA signaling is necessary but not sufficient for RD29B transcriptional memory during successive dehydration stresses in *Arabidopsis thaliana*. *Plant J.* 2014; 79: 150–161. doi: [10.1111/tpj.12548](#) PMID: [24805058](#)
7. Herrera A. Crassulacean acid metabolism and fitness under water deficit stress: if not for carbon gain, what is facultative CAM good for? *Ann. Bot.* 2009; 103: 645–653. doi: [10.1093/aob/mcn145](#) PMID: [18708641](#)
8. Rayder L, Ting IP. CAM-idling in *Hoya carnos*a (Asclepiadaceae). *Photosynth Res.* 1983; 4: 203–211.
9. Cela J, Arrom L, Munné-Bosch S. Diurnal changes in photosystem II photochemistry, photoprotective compounds and stress-related phytohormones in the CAM plant, *Aptenia cordifolia*. *Plant Sci.* 2009; 177: 404–410.
10. Mito PT, Mercier H. Abscisic acid and nitric oxide signaling in two different portions of detached leaves of *Guzmania monostachia* with CAM up-regulated by drought. *J. Plant Physiol.* 2013; 170: 996–1002. doi: [10.1016/j.jplph.2013.02.004](#) PMID: [23523467](#)
11. Winter K, Garcia M, Holtum JAM. Drought-stress-induced up-regulation of CAM in seedlings of a tropical cactus, *Opuntia elatior*, operating predominantly in the C₃ mode. *J. Exp. Bot.* 2011; 62: 4037–4042. doi: [10.1093/jxb/err106](#) PMID: [21504876](#)
12. Müller M, Munné-Bosch S. Rapid and sensitive hormonal profiling of complex plant samples by liquid chromatography coupled to electrospray ionization tandem mass spectrometry. *Plant Meth.* 2011; 7: 37.
13. van Kooten O, Snel JFH. The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynth. Res.* 1990; 25: 147–150. doi: [10.1007/BF00033156](#) PMID: [24420345](#)
14. Lichtenthaler HK. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Meth. Enzymol.* 1987; 148: 350–382.
15. Munné-Bosch S, Alegre L. Changes in carotenoids, tocopherols and diterpenes during drought and recovery, and the biological significance of chlorophyll loss in *Rosmarinus officinalis* plants. *Planta* 2000; 210: 925–931. PMID: [10872224](#)
16. Amaral JS, Casal C, Torres D, Seabra RM, Oliveira BPP. Simultaneous determination of tocopherols and tocotrienols in hazelnuts by a normal phase liquid chromatographic method. *Anal. Sci.* 2005; 21: 1545–1548. PMID: [16379404](#)
17. DeLong JM, Prange RK, Hodges DM, Forney CF, Bishop MC, Quilliam M. Using a modified ferrous oxidation-xylene orange (FOX) assay for detection of lipid hydroperoxides in plant tissue. *J. Agric. Food Chem.* 2002; 50: 248–254. PMID: [11782190](#)
18. Hodges DM, DeLong JM, Forney CF, Prange RK. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* 1999; 207: 604–611.
19. Cushman JC. Crassulacean Acid Metabolism. A plastic photosynthetic adaptation to arid environments. *Plant Physiol.* 2001; 127: 1439–1448. PMID: [11743087](#)
20. Treichel S. Crassulaceen säurestoffwechsel bei einem salztoleranten Vertreter der Aizoacea: *Aptenia cordifolia*. *Plant Sci. Lett.* 1975; 4: 141–144.

21. Herppich WB, Peckmann K. Responses of gas exchange, photosynthesis, nocturnal acid accumulation and water relations of *Aptenia cordifolia* to short-term drought and rewatering. *J. Plant Physiol.* 1997; 150: 141–144.
22. Crozier A, Turnbull CGN, Malcolm JM, Graebe JE. Gibberellin metabolism in cell-free preparations from *Phaseolus coccineus*. In: Takahashi BO, Phinney BO, McMillan J, editors. *Gibberellins*. New York, N.Y., USA: Springer; 1991. pp. 83–93.
23. Thomas SG, Rieu I, Stever CM. Gibberellin metabolism and signaling. In: Litwack G, editor. *Plant hormones, vitamins and hormones*. Vol. 72. San Diego, California, USA: Elsevier Academic Press; 1991. pp. 289–338, 1991.
24. Sharp RE, LeNoble ME. ABA, ethylene and the control of shoot and root growth under water stress. *J. Exp. Bot.* 2001; 53: 33–37.
25. Verslues PE, Bray EA. Role of abscisic acid (ABA) and *Arabidopsis thaliana* ABA-insensitive loci in low water potential-induced ABA and proline accumulation. *J. Exp. Bot.* 2006; 57: 201–212. PMID: [16339784](#)
26. Alonso R, Berli FJ, Bottini R, Picoli P. Acclimation mechanisms elicited by sprayed abscisic acid, solar UV-B and water deficit in leaf tissues of field-grown grapevines. *Plant Physiol. Biochem.* 2015; 91: 56–60. doi: [10.1016/j.plaphy.2015.03.011](#) PMID: [25885355](#)
27. Bao G, Zhuo C, Qian C, Xiao T, Guo Z, Lu S. Co-expression of NCED and ALO improves vitamin C level and tolerance to drought and chilling in transgenic tobacco and stylo plants. *Plant Biotechnol. J.* 2015 in press, doi: [10.1111/pbi.12374](#)
28. Morales M, Garcia QS, Munné-Bosch S. Ecophysiological response to seasonal variations in water availability in the arborescent, endemic plant *Vellozia gigantea*. *Tree Physiol.* 2015; 35: 253–265. doi: [10.1093/treephys/tpv012](#) PMID: [25769340](#)
29. Coudray C, Richard MJ, Favier AE. Determination of primary and secondary lipid peroxidation products: Plasma lipid hydroperoxides and thiobarbituric acid reactive substances. In: Favier AE, editor. *Analysis of free radicals in biological systems*. Basel, Switzerland: Birkhauser; 1995. pp. 185–200.
30. Pintó-Marijuan M, Munné-Bosch S. Photo-oxidative stress markers as a measure of abiotic stress-induced leaf senescence in plants: advantages and limitations. *J. Exp. Bot.* 2014; 65: 3845–3857. doi: [10.1093/jxb/eru086](#) PMID: [24683180](#)
31. El Kayal W, Keller G, Debayles C, Kumar R, Weier D, Teulières C, et al. Regulation of tocopherol biosynthesis through transcriptional control of tocopherol cyclase during cold hardening in *Eucalyptus gunnii*. *Physiol. Plant.* 2006; 126: 212–223.
32. Munné-Bosch S, Falara V, Pateraki I, López-Carbonell M, Cela J, Kanellis AK. Physiological and molecular responses of the isoprenoid biosynthetic pathway in a drought-resistant Mediterranean shrub, *Cistus creticus* exposed to water deficit. *J. Plant Physiol.* 2009; 166: 136–145. doi: [10.1016/j.jplph.2008.02.011](#) PMID: [18455260](#)
33. Chaudhary N, Khurana P. Vitamin E biosynthesis genes in rice: molecular characterization, expression profiling and comparative phylogenetic analysis. *Plant Sci.* 2009; 177: 479–491.
34. Xiong L, Zhu J-K. Regulation of abscisic acid biosynthesis. *Plant Physiol.* 2003; 133: 29–36. PMID: [12970472](#)