

## RESEARCH PAPER

Evidence that the plant cannabinoid cannabigerol is a highly potent  $\alpha_2$ -adrenoceptor agonist and moderately potent 5HT<sub>1A</sub> receptor antagonist

MG Cascio\*, LA Gauson\*, LA Stevenson, RA Ross and RG Pertwee

School of Medical Sciences, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen, UK

**Background and purpose:** Cannabis is the source of at least seventy phytocannabinoids. The pharmacology of most of these has been little investigated, three notable exceptions being  $\Delta^9$ -tetrahydrocannabinol, cannabidiol and  $\Delta^9$ -tetrahydrocannabivarin. This investigation addressed the question of whether the little-studied phytocannabinoid, cannabigerol, can activate or block any G protein-coupled receptor.

**Experimental approach:** The [<sup>35</sup>S]GTP $\gamma$ S binding assay, performed with mouse brain membranes, was used to test the ability of cannabigerol to produce G protein-coupled receptor activation or blockade. Its ability to displace [<sup>3</sup>H]CP55940 from mouse CB<sub>1</sub> and human CB<sub>2</sub> cannabinoid receptors and to inhibit electrically evoked contractions of the mouse isolated vas deferens was also investigated.

**Key results:** In the brain membrane experiments, cannabigerol behaved as a potent  $\alpha_2$ -adrenoceptor agonist (EC<sub>50</sub> = 0.2 nM) and antagonized the 5-HT<sub>1A</sub> receptor agonist, *R*-(+)-8-hydroxy-2-(di-*n*-propylamino)tetralin (apparent *K*<sub>B</sub> = 51.9 nM). At 10  $\mu$ M, it also behaved as a CB<sub>1</sub> receptor competitive antagonist. Additionally, cannabigerol inhibited evoked contractions of the vas deferens in a manner that appeared to be  $\alpha_2$ -adrenoceptor-mediated (EC<sub>50</sub> = 72.8 nM) and displayed significant affinity for mouse CB<sub>1</sub> and human CB<sub>2</sub> receptors.

**Conclusions and implications:** This investigation has provided the first evidence that cannabigerol can activate  $\alpha_2$ -adrenoceptors, bind to cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptors and block CB<sub>1</sub> and 5-HT<sub>1A</sub> receptors. It will now be important to investigate why cannabigerol produced signs of agonism more potently in the [<sup>35</sup>S]GTP $\gamma$ S binding assay than in the vas deferens and also whether it can inhibit noradrenaline uptake in this isolated tissue and in the brain.

*British Journal of Pharmacology* (2010) **159**, 129–141; doi:10.1111/j.1476-5381.2009.00515.x; published online 4 December 2009

**Keywords:** cannabigerol; CP55940; mouse vas deferens;  $\alpha_2$ -adrenoceptor; 5-HT<sub>1A</sub> receptor; CB<sub>1</sub> receptor; clonidine; dexmedetomidine; maprotiline; *R*-(+)-8-hydroxy-2-(di-*n*-propylamino)tetralin

**Abbreviations:** 8-OH-DPAT, *R*-(+)-8-hydroxy-2-(di-*n*-propylamino)tetralin; anandamide, arachidonoyl ethanolamide; BSA, bovine serum albumin;  $\beta$ , $\gamma$ -methylene-ATP,  $\beta$ , $\gamma$ -methylene adenosine 5'-triphosphate disodium salt; CHO, Chinese hamster ovary; CP55940, (–)-*cis*-3-[2-hydroxy-4-(1,1-dimethylheptyl)phenyl]-*trans*-4-(3-hydroxypropyl)cyclohexanol; DMSO, dimethyl sulphoxide; GTP $\gamma$ S, guanosine-5'-O-(3-thiotriphosphate); *R*-(+)-WIN55212, *R*-(+)-[2,3-dihydro-5-methyl-3-(4-morpholinylmethyl)pyrrolo-[1,2,3-*de*]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone mesylate; rimonabant, *N*-(piperidin-1-yl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide hydrochloride; WAY100635, *N*-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-*N*-2-pyridinylcyclohexanecarboxamide maleate

## Introduction


*Cannabis sativa* is the natural source of a set of at least seventy C<sub>21</sub> compounds that are known collectively as phytocannab-

inoids (see ElSohly and Slade, 2005). To date, pharmacological research has focused primarily on just three of these compounds. One of these is  $\Delta^9$ -tetrahydrocannabinol, the main psychoactive constituent of cannabis, the others being the non-psychoactive phytocannabinoid, cannabidiol and  $\Delta^9$ -tetrahydrocannabivarin, which at low doses can block cannabinoid receptor-mediated actions of  $\Delta^9$ -tetrahydrocannabinol (see Pertwee, 2008). Both  $\Delta^9$ -tetrahydrocannabinol and cannabidiol are present in a currently licensed medicine, Sativex®, and  $\Delta^9$ -tetrahydrocannabivarin has

Correspondence: Professor RG Pertwee, School of Medical Sciences, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK. E-mail: rgp@abdn.ac.uk

\*These authors contributed equally to this investigation.

Received 2 July 2009; revised 3 August 2009; accepted 17 August 2009



**Figure 1** The chemical structure of cannabigerol.

therapeutic potential for the management of disorders such as obesity and drug dependence (see Pertwee, 2008). It is important that more research is directed at exploring the pharmacology of the many other cannabinoids present in cannabis, not least because such research will help to identify any additional therapeutic applications of these phytocannabinoids, compounds that were described in the title of a recent *BJP* commentary as 'a neglected pharmacological treasure trove' (Mechoulam, 2005).

The present investigation focused on the little-studied phytocannabinoid, cannabigerol (Figure 1) which was first detected in cannabis and synthesized by Gaoni and Mechoulam (1964) and subsequently found not to induce  $\Delta^9$ -THC-like psychopharmacological effects *in vivo* (Grünfeld and Ederly, 1969; Mechoulam *et al.*, 1970). Our main objective was to establish whether this compound can activate or block any G protein-coupled receptor as indicated by stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse whole brain membranes or by blockade of such stimulation when this is induced by another compound. We also investigated the ability of cannabigerol to displace [ $^3$ H]CP55940 both from CB<sub>1</sub> binding sites in these membranes and from CB<sub>2</sub> binding sites in membranes prepared from Chinese hamster ovary (CHO) cells transfected with human CB<sub>2</sub> receptors (receptor nomenclature follows Alexander *et al.*, 2008). Some of our experiments were carried out with the mouse isolated vas deferens, a tissue in which cannabinoid receptor agonists can inhibit electrically evoked contractions (Devane *et al.*, 1992; Pertwee *et al.*, 1995). They are thought to do this by targeting prejunctional neuronal cannabinoid CB<sub>1</sub> receptors in a manner that inhibits neuronal release of the contractile neurotransmitters, ATP and noradrenaline (Trendelenburg *et al.*, 2000; see von Kügelgen and Starke, 1991; Pertwee, 1997; Schlicker and Kathmann, 2001).

The results we obtained extend a recent finding that cannabigerol can activate TRPA1 transient receptor potential channels and block the activation of TRPM8 transient receptor potential channels *in vitro* (De Petrocellis *et al.*, 2008) by providing evidence that it can indeed also target certain G protein-coupled receptors. Some of the results described in this paper have been presented to the International Cannabinoid Research Society (Gauson *et al.*, 2008; 2009; Cascio *et al.*, 2009).

## Methods

### Animals

All animal care and experimental procedures complied with the UK Animals (Scientific Procedures) Act, 1986 and associ-

ated guidelines for the use of experimental animals. MF1 mice aged 6 to 7 weeks and weighing 30 to 35 g were purchased from Harlan UK Ltd. (Blackthorn, UK), whereas C57BL/6 CB<sub>1</sub> receptor knockout mice and their wild-type litter mates were obtained from NIH (Rockville, MD, USA). Mice were maintained on a 12/12 h light/dark cycle with free access to food and water. All experiments were performed with tissues obtained from adult male mice.

### CHO cells

Chinese hamster ovary cells stably transfected with cDNA encoding human cannabinoid CB<sub>2</sub> receptors were maintained at 37°C and 5% CO<sub>2</sub> in Dulbecco's modified Eagle's medium nutrient mixture F-12 HAM supplemented with 2 mM L-glutamine, 10% foetal bovine serum, 0.6% penicillin-streptomycin and G418 (600  $\mu$ g·mL<sup>-1</sup>). These CHO-hCB<sub>2</sub> cells were passaged twice a week using a non-enzymatic cell dissociation solution.

### Membrane preparation

Binding assays with [ $^3$ H]CP55940 and with [ $^{35}$ S]GTP $\gamma$ S were performed with mouse whole brain membranes, prepared as described by Thomas *et al.* (2004) or with CHO-CB<sub>2</sub> cell membranes (Ross *et al.*, 1999a). The hCB<sub>2</sub> transfected cells were removed from flasks by scraping and then frozen as a pellet at -20°C until required. Before use in a radioligand binding assay, cells were defrosted, diluted in Tris-buffer (50 mM Tris-HCl and 50 mM Tris-Base) and homogenized with a 1-mL handheld homogenizer. Protein assays were performed using a Bio-Rad Dc kit (Hercules, CA, USA).

### Radioligand displacement assay

The assays were carried out with [ $^3$ H]CP55940 and Tris-binding buffer [50 mM Tris-HCl, 50 mM Tris-Base, 0.1% bovine serum albumin (BSA); pH 7.4], total assay volume 500  $\mu$ L, using the filtration procedure described previously by Ross *et al.* (1999b). Binding was initiated by the addition of either brain membranes (33  $\mu$ g protein per well) or transfected hCB<sub>2</sub>-cells (25  $\mu$ g protein per well). All assays were performed at 37°C for 60 min before termination by addition of ice-cold Tris-binding buffer and vacuum filtration using a 24-well sampling manifold (Brandel Cell Harvester) and Brandel GF/B filters that had been soaked in wash buffer at 4°C for at least 24 h (Brandel Inc., Gaithersburg, MD, USA). Each reaction well was washed six times with a 1.2 mL aliquot of Tris-binding buffer. The filters were oven-dried for 60 min and then placed in 5 mL of scintillation fluid (Ultima Gold XR, PerkinElmer). Radioactivity was quantified by liquid scintillation spectrometry. Specific binding was defined as the difference between the binding that occurred in the presence and absence of 1  $\mu$ M unlabelled CP55940. The concentration of [ $^3$ H]CP55940 used in our displacement assays was 0.7 nM. Compounds under investigation were stored as stock solutions of 10 mM in dimethyl sulphoxide (DMSO), the vehicle concentration in all assay wells being 0.1% DMSO. The binding parameters for [ $^3$ H]CP55940, determined by fitting data from saturation-binding experiments to a one-site

saturation plot using GraphPad Prism, were 2336 fmol·mg<sup>-1</sup> protein ( $B_{\max}$ ) and 2.31 nM ( $K_d$ ) in mouse brain membranes (Thomas *et al.*, 2004) and 215 pmol·mg<sup>-1</sup> ( $B_{\max}$ ) and 4.3 nM ( $K_d$ ) in hCB<sub>2</sub>-transfected cells.

#### [<sup>35</sup>S]GTPγS binding assay

The method for measuring agonist-stimulated [<sup>35</sup>S]GTPγS binding to cannabinoid CB<sub>1</sub> receptors was adapted from the methods of Kurkinen *et al.* (1997) and Breivogel *et al.* (2001). The assays were carried out with GTPγS binding buffer (50 mM Tris-HCl, 50 mM Tris-Base, 5 mM MgCl<sub>2</sub>, 1 mM EDTA, 100 mM NaCl, 1 mM dithiothreitol, 0.1% BSA) in the presence of [<sup>35</sup>S]GTPγS and GDP, in a final volume of 500 μL. Binding was initiated by the addition of [<sup>35</sup>S]GTPγS to the wells. Nonspecific binding was measured in the presence of 30 μM GTPγS. The drugs were incubated in the assay for 60 min at 30°C. The reaction was terminated by a rapid vacuum filtration method using Tris-binding buffer as described previously, and the radioactivity was quantified by liquid scintillation spectrometry. In all the [<sup>35</sup>S]GTPγS-binding assays, we used 0.1 nM [<sup>35</sup>S]GTPγS, 30 μM GDP and a protein concentration of 5 μg per well. Additionally, mouse brain membranes were preincubated for 30 min at 30°C with 0.5 U·mL<sup>-1</sup> adenosine deaminase (200 U·mL<sup>-1</sup>) to remove endogenous adenosine. Agonists and antagonists were stored at -20°C as 10 mM stock solutions dissolved in distilled water (yohimbine) or DMSO.

#### Vas deferens experiments

Vasa deferentia were obtained from albino MF1 mice weighing 36 to 53 g. The tissues were mounted vertically in 4 mL organ baths. They were then subjected to electrical stimulation of progressively greater intensity followed by an equilibration procedure in which they were exposed to alternate periods of stimulation (2 min) and rest (10 min) until contractions with consistent amplitudes were obtained (Thomas *et al.*, 2004). These contractions were monophasic and isometric and were evoked by 0.5 s trains of pulses of 110% maximal voltage (train frequency 0.1 Hz; pulse frequency 5 Hz; pulse duration 0.5 ms).

Except in our experiments with phenylephrine, all drug additions were made to the organ baths after the equilibration period and there was no washout between these additions. In most experiments there was an initial application of a potential antagonist or its vehicle. This was followed 28 min later by a 2 min period of electrical stimulation at the end of which the lowest of a series of concentrations of the twitch inhibitors, cannabigerol, clonidine, dexmedetomidine or maprotiline was applied. After a period of rest, the tissues were electrically stimulated for 2 min and then subjected to a further addition of twitch inhibitor. This cycle of drug addition, rest and 2 min stimulation was repeated so as to construct cumulative concentration-response curves. Only one concentration-response curve was constructed per tissue (Pertwee *et al.*, 1996). The rest period was 13 min in the experiments with cannabigerol, dexmedetomidine and maprotiline and 3 min in the clonidine experiments.

In experiments with β,γ-methylene-ATP, no electrical stimuli were applied after the equilibration procedure. Log

concentration-response curves of β,γ-methylene-ATP were constructed cumulatively without washout. Cannabigerol was added 30 min before the first addition of β,γ-methylene-ATP, each subsequent addition of which was made immediately after the effect of the previous dose had reached a plateau (dose cycles of 1 to 2 min). Only one addition of phenylephrine was made to each tissue and this was carried out 30 min after the addition of cannabigerol or its vehicle.

#### Analysis of data

Values have been expressed as means and variability as SEM or as 95% confidence limits. The concentration of the compounds under investigation that produced a 50% displacement of radioligand from specific binding sites ( $IC_{50}$  value) was calculated using GraphPad Prism and the corresponding  $K_i$  values were calculated using the equation of Cheng and Prusoff (1973). Net agonist stimulated [<sup>35</sup>S]GTPγS binding values were calculated by subtracting basal binding values (obtained in the absence of agonist) from agonist-stimulated values (obtained in the presence of agonist) as detailed elsewhere (Ross *et al.*, 1999a). Inhibition of the electrically evoked twitch response of the vas deferens has been expressed in percentage terms, and this has been calculated by comparing the amplitude of the twitch response after each addition of a twitch inhibitor with its amplitude immediately before the first addition of the inhibitor. Contractile responses to phenylephrine and β,γ-methylene-ATP have been expressed as increases in tension (g). Values for  $EC_{50}$ , maximal effect ( $E_{\max}$ ) and SEM or 95% confidence limits of these values have been calculated by nonlinear regression analysis using the equation for a sigmoid concentration-response curve (GraphPad Prism).

Unless stated otherwise, apparent dissociation constant ( $K_B$ ) values for antagonism of agonists by yohimbine in the vas deferens or by yohimbine or cannabigerol in the [<sup>35</sup>S]GTPγS binding assay have been calculated by Schild analysis (GraphPad Prism). These  $K_B$  values were calculated only from data obtained in experiments in which yohimbine or cannabigerol produced a right-ward shift in the log concentration response curve of an agonist that was indicated by (2 + 2) dose parallel line analysis to be statistically significant and not to deviate significantly from parallelism (Pertwee *et al.*, 2002). In one set of experiments, the effect of one or other of five concentrations of cannabigerol on the log concentration response curve of 8-OH-DPAT was determined. For these experiments, the  $K_B$  of cannabigerol was calculated from the intercept on the x-axis ( $-\log K_B$ ) of the best-fit straight line of a plot of  $\log(x - 1)$  against  $\log B$  constructed by linear regression analysis (GraphPad Prism). The equation for this Schild plot is  $\log(x - 1) = \log B - \log K_B$ , where x (the 'concentration ratio') is the concentration of 8-OH-DPAT that elicits a particular response in the presence of cannabigerol at a concentration, B, divided by the concentration of 8-OH-DPAT that elicits a response of the same size in the absence of cannabigerol. This equation predicts a slope of unity for all receptor-mediated interactions between agonists and antagonists that are competitive and reversible (Tallarida *et al.*, 1979).  $\log(x - 1)$  values were determined by (2 + 2) dose parallel line analysis as described previously (Pertwee *et al.*, 2002). Mean values obtained *in vitro*

have been compared with zero using the one-sample *t*-test and with each other using Student's two-tailed *t*-test for unpaired data or one-way analysis of variance (ANOVA) followed by Dunnett's test (GraphPad Prism). A *P* value of 0.05 or less was considered to be significant.

### Materials

Cannabigerol was supplied by GW Pharmaceuticals (Porton Down, Wiltshire, UK) and rimonabant (SR141716A) was obtained from Sanofi-Aventis (Montpellier, France). Phenylephrine hydrochloride,  $\beta$ ,  $\gamma$ -methyleneadenosine 5'-triphosphate disodium salt ( $\beta$ ,  $\gamma$ -methylene-ATP), arachidonoyl ethanolamide (anandamide), clonidine hydrochloride and *N*-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-*N*-2-pyridinylcyclohexanecarboxamide maleate (WAY100635) were purchased from Sigma-Aldrich (Poole, Dorset, UK) and *R*-(+)-[2,3-dihydro-5-methyl-3-(4-morpholinylmethyl)pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone (*R*-(+)-WIN55212), (-)-*cis*-3-[2-hydroxy-4-(1,1-dimethylheptyl)phenyl]-*trans*-4-(3-hydroxypropyl)cyclohexanol (CP55940), dexmedetomidine hydrochloride, yohimbine hydrochloride, maprotiline hydrochloride and *R*-(+)-8-hydroxy-2-(di-*n*-propylamino)tetralin (8-OH-DPAT) from Tocris (Bristol, UK). For the binding experiments, [<sup>3</sup>H]CP55940 (160 Ci·mmol<sup>-1</sup>) and [<sup>35</sup>S]GTP $\gamma$ S (1250 Ci·mmol<sup>-1</sup>) were obtained from PerkinElmer Life Sciences Inc. (Boston, MA, USA), GTP $\gamma$ S and adenosine deaminase from Roche Diagnostic (Indianapolis, IN, USA) and GDP from Sigma-Aldrich. Phenylephrine hydrochloride,  $\beta$ ,  $\gamma$ -methylene-ATP and maprotiline were dissolved in a 0.9% aqueous solution of NaCl (saline) and yohimbine and clonidine in distilled water. All other compounds were dissolved in pure DMSO. In the vas deferens experiments, all compounds were added to organ baths in a volume of 10  $\mu$ L.

## Results

### *Cannabigerol is a potent stimulator of [<sup>35</sup>S]GTP $\gamma$ S binding to brain membranes*


In our initial experiments, we investigated the effect of cannabigerol on [<sup>35</sup>S]GTP $\gamma$ S binding to MF1 mouse brain membranes. We found that at concentrations in the picomolar and low nanomolar range, this cannabinoid produced a concentration-related stimulation of [<sup>35</sup>S]GTP $\gamma$ S binding to MF1 mouse brain membranes (Figure 2). Experiments performed with brain membranes obtained from either CB<sub>1</sub><sup>+/-</sup> or CB<sub>1</sub><sup>-/-</sup> C57BL/6J mice yielded similar results (Table 1) and also showed that, as in MF1 mouse brain membranes (Figure 2),

cannabigerol at 1  $\mu$ M had no significant effect on [<sup>35</sup>S]GTP $\gamma$ S binding and, at 10  $\mu$ M, produced a marked inhibitory effect.

The mean percentage inhibition of [<sup>35</sup>S]GTP $\gamma$ S binding was found to be 30.1%  $\pm$  3.8 (*n* = 17) in MF1 mouse brain membranes (Figure 2), 14.2%  $\pm$  3.4 (*n* = 6) in CB<sub>1</sub><sup>+/-</sup> C57BL/6J mouse brain membranes and 21.8%  $\pm$  4.6 (*n* = 8) in CB<sub>1</sub><sup>-/-</sup> C57BL/6J mouse brain membranes and each of these mean values is significantly less than zero (*P* < 0.01; one-sample *t*-test). These results suggest that neither the stimulatory effect nor the inhibitory effect of cannabigerol on [<sup>35</sup>S]GTP $\gamma$ S binding to mouse brain membranes is CB<sub>1</sub> receptor-mediated.

### *Cannabigerol behaves as an $\alpha_2$ -adrenoceptor agonist in the mouse isolated vas deferens*

Further evidence that cannabigerol is not a CB<sub>1</sub> receptor agonist was obtained from experiments performed with the mouse isolated vas deferens. More specifically, although cannabigerol shared the ability of established cannabinoid receptor agonists such as CP55940 and  $\Delta^9$ -tetrahydrocannabinol (Pertwee *et al.*, 1995) to produce a concentration-related inhibition of electrically evoked contractions (Figure 3A), no right-ward shift in the log concentration response curve of cannabigerol was produced by rimonabant at 100 nM (data not shown), a concentration that equals or exceeds




**Figure 2** The effect of cannabigerol on [<sup>35</sup>S]GTP $\gamma$ S binding to whole brain membranes obtained from MF1 mice (*n* = 7 to 20). Each symbol represents the mean percentage change in binding  $\pm$  SEM. Asterisks denote values that are significantly different from zero (\*\**P* < 0.01; \*\*\**P* < 0.001; one-sample *t*-test).

**Table 1** Cannabigerol stimulates [<sup>35</sup>S]GTP $\gamma$ S binding to brain membranes obtained from MF1, CB<sub>1</sub><sup>+/-</sup> C57BL/6J and CB<sub>1</sub><sup>-/-</sup> C57BL/6J mice

| Mouse strain                            | Mean EC <sub>50</sub> (95% CL) | Mean E <sub>max</sub> (95% CL) | n       |
|---|--------------------------------|--------------------------------|---------|
| MF1                                     | 0.2 nM (0.006 and 9.0 nM)      | 15.5% (8.5 and 22.5%)          | 7 to 20 |
| CB <sub>1</sub> <sup>+/-</sup> C57BL/6J | 0.04 nM (0.002 and 0.8 nM)     | 17.0% (10.4 and 23.7%)         | 8       |
| CB <sub>1</sub> <sup>-/-</sup> C57BL/6J | 0.17 nM (0.01 and 2.8 nM)      | 11.4% (6.6 and 16.2%)          | 8       |

Mean EC<sub>50</sub> and E<sub>max</sub> values were calculated from data obtained with cannabigerol concentrations of up to 10 nM. CL, confidence limits.




**Figure 3** Mean log concentration-response curves of (A) cannabigerol ( $n = 13$ ), (B) clonidine ( $n = 7$ ) (C) dexmedetomidine ( $n = 5$ ) and (D) maprotiline ( $n = 7$ ) in the MF1 mouse isolated vas deferens constructed in the presence of yohimbine or its vehicle. Each symbol represents the mean value  $\pm$  SEM for inhibition of electrically evoked contractions expressed as a percentage of the amplitude of the twitch response measured immediately before the first addition of cannabigerol, clonidine or maprotiline to the organ bath. Yohimbine or its vehicle was added 30 min before this first addition and all further additions were made at 5 or 15 min intervals (*Methods*). Each log concentration response curve was constructed cumulatively. The mean apparent  $K_b$  value of yohimbine with its 95% confidence limits shown in brackets is 10.1 nM (3.0 and 33.7 nM) against cannabigerol, 14.0 nM (4.9 and 40.6 nM) against clonidine, 8.7 nM (4.3 and 17.8 nM) against dexmedetomidine and 8.2 nM (1.5 and 46.0 nM) against maprotiline. In the absence of yohimbine, electrically evoked contractions were inhibited by cannabigerol, clonidine, dexmedetomidine and maprotiline with mean  $EC_{50}$  values of 72.8 nM (23.4 and 227 nM), 6.3 nM (2.0 and 19.7 nM), 0.24 nM (0.14 and 0.40 nM) and 24.9 nM (5.6 and 111.2 nM) respectively. The 95% confidence limits of these mean values are shown in brackets. The corresponding  $E_{max}$  values are 51.7% (44.3 and 59.1%), 63.0% (54.1 and 71.8%), 85.8% (81.6 and 90.0%) and 58.2% (47.1 and 69.3%) respectively.

concentrations of this  $CB_1$ -selective antagonist that have been found previously to antagonize established  $CB_1$  receptor agonists in this bioassay (Pertwee *et al.*, 1995; Ross *et al.*, 2001). Cannabigerol inhibited electrically evoked contractions of the vas deferens at concentrations below any found to attenuate contractile responses either to the P2 receptor agonist,  $\beta,\gamma$ -methylene ATP, or to the  $\alpha_1$ -adrenoceptor agonist, phenylephrine hydrochloride. Thus it inhibited electrically evoked contractions at concentrations of 100 nM or less (Figure 3A), attenuated contractile responses of the vas deferens to  $\beta,\gamma$ -methylene ATP, at 1  $\mu$ M but not at 100 nM (Figure 4) and did not affect contractions induced by phenylephrine hydrochloride

even at a concentration of 1  $\mu$ M (Figure 4). As electrically evoked contractions of the vas deferens are thought to result from the release of ATP and noradrenaline on to postjunctional P2 receptors and  $\alpha_1$ -adrenoceptors (Introduction), these findings suggest that cannabigerol can inhibit these contractions by acting prejunctionally.

One possibility is that the inhibitory effect of cannabigerol on electrically evoked contractions of the mouse vas deferens is mediated by prejunctional  $\alpha_2$ -adrenoceptors as it is generally accepted that these receptors mediate inhibition of such contractions when activated by endogenously released noradrenaline or by an exogenously added agonist (Pertwee




**Figure 4** Upper panels: mean increases in tension of the MF1 mouse isolated vas deferens induced by  $\beta,\gamma$ -methylene ATP in the presence of DMSO (circles) or cannabigerol (triangles). For the construction of log concentration-response curves,  $\beta,\gamma$ -methylene ATP was first added 30 min (A) after DMSO or 100 nM cannabigerol ( $n = 6$  or  $8$ ) or (B) after DMSO or 1  $\mu$ M cannabigerol ( $n = 8$ ). The asterisk indicates a significant difference between the contractile response to 10  $\mu$ M  $\beta,\gamma$ -methylene ATP in the absence of cannabigerol and the corresponding response in the presence of this cannabinoid ( $P < 0.05$ ; unpaired  $t$ -test). Lower panel: mean increases in tension of the mouse isolated vas deferens induced by 32  $\mu$ M phenylephrine in the absence or presence of 1  $\mu$ M cannabigerol. The two mean values are not significantly different ( $P > 0.05$ ; unpaired  $t$ -test). Additions of phenylephrine were made 30 min after DMSO (open columns) or cannabigerol ( $n = 8$ ). In all panels, mean increases in tension are expressed in grams  $\pm$  SEM. DMSO, dimethyl sulphoxide.

*et al.*, 2005); reviewed in (von Kügelgen and Starke, 1991; Starke, 2001). To test this hypothesis, we investigated whether cannabigerol can be antagonized in the vas deferens by the selective  $\alpha_2$ -adrenoceptor antagonist, yohimbine. We found not only that yohimbine can indeed antagonize cannabigerol-induced inhibition of electrically evoked contractions but also that the potency with which it produces this antagonism is similar to the potency it displays in the same bioassay as an antagonist of clonidine and dexmedetomidine (Figure 3), both of which are established  $\alpha_2$ -adrenoceptor agonists (Newman-Tancredi *et al.*, 1998). Yohimbine also antagonized the inhibition of electrically evoked contractions of the vas deferens induced by maprotiline, an inhibitor of noradrenaline uptake (Barbaccia *et al.*, 1986), with a potency that matched the potency with which it antagonized cannabigerol (Figure 3). It seems likely, therefore, that  $\alpha_2$ -adrenoceptors do indeed mediate

cannabigerol-induced inhibition of electrically evoked contractions of the mouse vas deferens.

#### *Cannabigerol also behaves as an $\alpha_2$ -adrenoceptor agonist in mouse brain membranes*

The results obtained in the vas deferens experiments raised the possibility that cannabigerol-induced stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse brain membranes (Figure 2) might also be  $\alpha_2$ -adrenoceptor-mediated. To investigate this possibility, we first carried out experiments directed at establishing whether dexmedetomidine shares the ability of cannabigerol to stimulate [ $^{35}$ S]GTP $\gamma$ S binding to MF1 mouse brain membranes. We found that this  $\alpha_2$ -adrenoceptor agonist can indeed induce such stimulation and that it is antagonized by yohimbine at 100 nM (Figure 5). This concentration of yohimbine also antagonized cannabigerol-induced




**Figure 5** Mean log concentration-response curves of (A) dexmedetomidine ( $n = 4$ ) and (B) cannabigerol ( $n = 12$ ) constructed in the absence or presence of 100 nM yohimbine. Each symbol represents the mean percentage change in binding of [ $^{35}$ S]GTP $\gamma$ S to MF1 mouse whole brain membranes  $\pm$  SEM. Mean  $EC_{50}$  values of dexmedetomidine and cannabigerol in the absence of yohimbine with 95% confidence limits shown in brackets are 4.3 nM (0.7 and 25.5 nM) and 0.13 nM (0.004 and 4.4 nM) respectively. The corresponding mean  $E_{max}$  values are 26.5% (17.1 and 36.0%) and 14.8% (5.9 and 23.7%) respectively. The right-ward shifts produced by yohimbine in the log concentration response curves of dexmedetomidine and cannabigerol are significant and do not deviate significantly from parallelism ( $P > 0.05$ ). The mean apparent  $K_B$  value of yohimbine for this antagonism, with its 95% confidence limits shown in brackets is 3.9 nM (1.0 and 15.1 nM) against dexmedetomidine and 1.8 nM (0.04 and 90.5 nM) against cannabigerol.

stimulation of [ $^{35}$ S]GTP $\gamma$ S binding, the data obtained indicating a lack of any significant difference between the apparent  $K_B$  values of yohimbine for its antagonism of these two compounds (Figure 5).

#### Cannabigerol and cannabinoid receptors

As cannabigerol is a constituent of cannabis, it was of interest to investigate whether it shares the ability of the plant cannabinoids  $\Delta^9$ -tetrahydrocannabinol,  $\Delta^9$ -tetrahydrocannabivarin and cannabidiol to bind to cannabinoid  $CB_1$  and  $CB_2$  receptors (see Pertwee, 2008). Cannabigerol was able to completely displace [ $^3$ H]CP55940 from specific binding sites both in mouse brain membranes and in CHO-h $CB_2$  cell membranes, its mean  $K_i$  values for this displacement suggesting that it has greater  $CB_1$  than  $CB_2$  receptor affinity (Figure 6). The binding data obtained with brain membranes suggest that although cannabigerol does bind to cannabinoid  $CB_1$  receptors, this is only detectable at concentrations above those at which it stimulates [ $^{35}$ S]GTP $\gamma$ S binding to these membranes (Figures 2 and 5). Further experiments were therefore carried out to establish whether a concentration of cannabigerol that produced a clear reduction in specific binding of [ $^3$ H]CP55940 to brain membranes, also antagonized anandamide or CP55940.


The first set of these experiments showed that anandamide-induced stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse brain membranes was significantly antagonized by cannabigerol at



**Figure 6** Displacement of [ $^3$ H]CP55940 by cannabigerol from specific binding sites on (A) MF1 mouse whole brain membranes ( $n = 4$ ) and (B) CHO-h $CB_2$  cell membranes ( $n = 8$ ). Each symbol represents the mean percent displacement  $\pm$  SEM. Mean  $K_i$  values with 95% confidence limits shown in brackets are (A) 381 nM (231 and 627 nM) for displacement from brain membranes and (B) 2.6  $\mu$ M (1.4 and 4.7  $\mu$ M) for displacement from CHO-h $CB_2$  cell membranes. CHO, Chinese hamster ovary.

10  $\mu$ M (Figure 7A). This it did with a mean apparent  $K_B$  value (Table 2) that is significantly less than its mean  $K_i$  value for displacement of [ $^3$ H]CP55940 from brain membranes (Figure 6). However, in line with the ability of cannabigerol by itself to inhibit [ $^{35}$ S]GTP $\gamma$ S binding to brain membranes at 10  $\mu$ M, it also appeared to produce a downward shift in the log concentration-response curve of anandamide. When this component of cannabigerol-induced antagonism that most likely arises from the ability of this compound to inhibit [ $^{35}$ S]GTP $\gamma$ S binding was excluded, a significant right-ward shift in the log concentration-response curve of anandamide was still apparent (Figure 7C). Importantly, the mean apparent  $K_B$  value calculated from this shift (Table 2) did not differ significantly from the mean  $K_i$  value of cannabigerol for displacement of [ $^3$ H]CP55940 from brain membranes (Figure 6). Also, as this right-ward shift did not deviate significantly from parallelism, it is likely that cannabigerol is a competitive antagonist of anandamide. Neither a right-ward shift nor a downward shift in the log concentration-response curve for anandamide-induced stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to MF1 mouse brain membranes was induced by cannabigerol at 1  $\mu$ M ( $n = 6$ ; data not shown).

Also, 10  $\mu$ M cannabigerol antagonized CP55940-induced stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse brain membranes (Figure 7B). Again, cannabigerol appeared to produce both a right-ward and a downward shift in the log concentration response curve of the agonist. After compensating for the downward shift (Figure 7D) it was found that the mean apparent  $K_B$  value of cannabigerol for antagonism of CP55940 (Table 2) did not differ significantly either from the mean  $K_B$  value of cannabigerol for antagonism of anandamide or from the mean  $K_i$  value of cannabigerol for displacement of [ $^3$ H]CP55940 from brain membranes (Figure 6). As in the anandamide experiments, the right-ward shift induced by cannabigerol in the log concentration-response curve of



**Figure 7** Upper panels: the effect of 10  $\mu\text{M}$  cannabigerol (CBG) on the mean log concentration-response curve of (A) anandamide and (B) CP55940 for stimulation of [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding to mouse brain membranes. Lower panels: the effect of 10  $\mu\text{M}$  cannabigerol (CBG) on the mean log concentration-response curve of (C) anandamide and (D) CP55940 for stimulation of [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding after subtraction of the inhibitory effect induced by 10  $\mu\text{M}$  cannabigerol on [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding in the absence of any other compound ( $30.1 \pm 3.8\%$ ;  $n = 17$ ; Figure 2). This value was subtracted from all values of percent stimulation of [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding by anandamide or CP55490 determined in the presence of cannabigerol. Experiments were performed with MF1 mouse whole brain membranes and stimulation of [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding is expressed as mean percent stimulation  $\pm$  SEM ( $n = 5$  or 6). The right-ward shifts produced by cannabigerol in the log concentration response curves of anandamide and CP55940 do not deviate significantly from parallelism ( $P > 0.05$ ). DMSO, dimethyl sulphoxide.

**Table 2** The mean apparent  $K_{\text{B}}$  values of cannabigerol for antagonism of anandamide- and CP55940-induced stimulation of [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding to MF1 mouse brain membranes

| Agonist    | Antagonist (10 $\mu\text{M}$ ) | Mean apparent $K_{\text{B}}$ (95% CL) | Mean apparent $K_{\text{B}}$ (95% CL) <sup>1</sup> | n |
|------------|--------------------------------|---------------------------------------|--|---|
| Anandamide | Cannabigerol                   | 33.1 nM (13.2 and 82.9 nM)            | 483 nM (162 and 1445 nM)                           | 5 |
| CP55940    | Cannabigerol                   | 53.7 nM (19.4 and 149 nM)             | 936 nM (336 and 2606 nM)                           | 6 |

<sup>1</sup>Calculated after subtraction of the mean inhibitory effect induced by 10  $\mu\text{M}$  cannabigerol on [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  binding in the absence of any other compound (30.1%; Figure 7 legend).


CL, confidence limits.

CP55940 (Figure 7D) did not deviate significantly from parallelism. Taken together, these findings support to the hypothesis that cannabigerol is a CB<sub>1</sub> receptor competitive antagonist, albeit of much lower potency than rimonabant (Thomas *et al.*, 2007).

#### Cannabigerol and 5HT<sub>1A</sub> receptors

There is evidence that some established  $\alpha_2$ -adrenoceptor ligands, including clonidine and yohimbine, target 5-HT<sub>1A</sub> receptors at concentrations above those at which they activate or block  $\alpha_2$ -adrenoceptors (Newman-Tancredi *et al.*,





**Figure 8** Mean log concentration-response curves of *R*-(+)-8-hydroxy-2-(di-*n*-propylamino)tetralin (8-OH-DPAT) constructed in the presence of (A) DMSO or 100 nM WAY100635 ( $n = 7$ ), (B) DMSO or 1 μM cannabigerol ( $n = 10$ ) or (C) DMSO or 10 μM cannabigerol ( $n = 12$ ). Each symbol represents the mean percentage change in binding of [<sup>35</sup>S]GTPγS to MF1 mouse whole brain membranes ± SEM. Neither the right-ward shift produced by WAY100635 in the log concentration response curve of 8-OH-DPAT nor that produced by 1 or 10 μM cannabigerol deviates significantly from parallelism ( $P > 0.05$ ). Mean apparent  $K_B$  values with 95% confidence limits shown in brackets are (A) 1.0 nM (0.5 and 2.3 nM) for WAY100635, (B) 19.6 nM (6.9 and 55.8 nM) for 1 μM cannabigerol and (C) 28.2 nM (7.7 and 102.9 nM) for 10 μM cannabigerol. Panel (D): Schild plot for antagonism of 8-OH-DPAT by 100 nM, 316 nM, 1 μM, 3.16 μM and 10 μM cannabigerol ( $n = 5$  to 12) with the 99% confidence band shown by dotted lines. The mean slope of this best-fit line with 95% confidence limits shown in brackets is 1.1 (1.0 and 1.2) and so does not differ significantly from unity. The mean apparent  $K_B$  value of cannabigerol calculated from this Schild plot, with 95% confidence limits shown in brackets, is 51.9 nM (37.6 and 68.2 nM). DMSO, dimethyl sulphoxide.

1998). This prompted us to investigate whether cannabigerol interacts with 5-HT<sub>1A</sub> receptors at concentrations higher than those at which it stimulates [<sup>35</sup>S]GTPγS binding to MF1 mouse brain membranes.

Initial experiments confirmed that the 5-HT<sub>1A</sub>-selective antagonist, WAY100635 (Forster *et al.*, 1995), did antagonize the stimulatory effect of 8-OH-DPAT, a 5-HT<sub>1A</sub> receptor agonist (Forster *et al.*, 1995), on [<sup>35</sup>S]GTPγS binding to brain membranes (Figure 8A). The mean apparent  $K_B$  value of WAY100635 for this antagonism was 1 nM. Further experiments showed that, at 1 μM, cannabigerol also antagonized 8-OH-DPAT in this bioassay as indicated by the ability of cannabigerol to produce a parallel right-ward shift in the log

concentration response curve of this 5-HT<sub>1A</sub>-selective agonist (Figure 8B). The data from this experiment and from similar experiments performed either with 10 μM cannabigerol (Figure 8C) or with one or other of three lower concentrations of cannabigerol (data not shown) allowed the construction of a Schild plot (Figure 8D), the slope of which was not significantly different from unity. The mean apparent  $K_B$  value of cannabigerol for its antagonism of 8-OH-DPAT, as calculated from this Schild plot, was 51.9 nM.

Cannabigerol (1 μM) was no less effective in producing a parallel right-ward shift in the log concentration response curve of 8-OH-DPAT for stimulating [<sup>35</sup>S]GTPγS binding to brain membranes when these membranes were obtained from

CB<sub>1</sub><sup>+/+</sup> or CB<sub>1</sub><sup>-/-</sup> C57BL/6J mice (data not shown) rather than from MF1 mice. The mean apparent  $K_B$  value of cannabigerol with 95% confidence limits shown in brackets is 19.6 nM (6.9 and 55.8;  $n = 10$ ; Figure 8B) when calculated from the data obtained using MF1 mouse brain membranes and 6.2 nM (2.6 and 14.8 nM;  $n = 11$ ) and 2.3 nM (0.7 and 7.7 nM;  $n = 13$ ) respectively when calculated from the CB<sub>1</sub><sup>+/+</sup> and CB<sub>1</sub><sup>-/-</sup> C57BL/6J mouse brain membrane data.

## Discussion

Results from our initial experiments indicated that cannabigerol exhibited significant potency both as a stimulator of [<sup>35</sup>S]GTPγS binding to mouse brain membranes (Figure 2) and as an inhibitor of electrically evoked contractions of the mouse isolated vas deferens (Figure 3). Neither of these effects appeared to be mediated by cannabinoid CB<sub>1</sub> receptors. Thus, cannabigerol displayed no less potency or efficacy as a stimulator of [<sup>35</sup>S]GTPγS binding to CB<sub>1</sub><sup>-/-</sup> mouse brain membranes than to CB<sub>1</sub><sup>+/+</sup> mouse brain membranes and its inhibitory effect on electrically evoked contractions of the vas deferens was not antagonized by the CB<sub>1</sub>-selective antagonist, rimona-bant, when this was administered at 100 nM, a concentration higher than that required to antagonize the established cannabinoid receptor agonists, CP55940, R-(+)-WIN55212 and Δ<sup>9</sup>-tetrahydrocannabinol, in the same bioassay (Pertwee *et al.*, 1995; Ross *et al.*, 2001). For reasons yet to be established, it was only within a range of very low concentrations that cannabigerol produced concentration-related increases in [<sup>35</sup>S]GTPγS binding to CB<sub>1</sub><sup>+/+</sup> or CB<sub>1</sub><sup>-/-</sup> mouse brain membranes. Thus, within the concentration range 0.01 nM to 1 μM, its log concentration response curve was bell-shaped and indeed, at 10 μM, cannabigerol markedly inhibited [<sup>35</sup>S]GTPγS binding to brain membranes.

It is unlikely that cannabigerol inhibited electrically evoked contractions of the vas deferens by acting postjunctionally to block the actions of the two neurotransmitters that are thought to induce these contractions: ATP acting on post-junctional P2X receptors and noradrenaline acting on post-junctional α<sub>1</sub>-adrenoceptors (von Kugelgen and Starke, 1991; Trendelenburg *et al.*, 2000). Thus, cannabigerol inhibited electrically evoked contractions at concentrations at which it did not significantly affect the amplitude of contractions induced in the vas deferens either by the P2X agonist, β,γ-methylene-ATP, or by the α<sub>1</sub>-adrenoceptor agonist, phenylephrine. Cannabigerol, therefore, differs from another plant cannabinoid, Δ<sup>9</sup>-tetrahydrocannabinol, which in the MF1 mouse vas deferens does reduce the amplitude of contractions induced by 10 μM β,γ-methylene-ATP or 32 μM phenylephrine when it is applied at the lowest of the concentrations at which it inhibits electrically evoked contractions of this tissue (Thomas *et al.*, 2005). The concentrations of β,γ-methylene-ATP and phenylephrine used in these previous experiments with Δ<sup>9</sup>-tetrahydrocannabinol were the same as those used in our cannabigerol experiments.

Results obtained with the selective α<sub>2</sub>-adrenoceptor antagonist, yohimbine, suggest that both the inhibitory effect of cannabigerol on electrically evoked contractions of the vas deferens and its stimulatory effect on [<sup>35</sup>S]GTPγS binding to

mouse brain membranes are mediated by α<sub>2</sub>-adrenoceptors. Thus, yohimbine antagonized both these effects of cannabigerol at a concentration (100 nM) at which it is expected to display selectivity as an α<sub>2</sub>-adrenoceptor antagonist (Newman-Tancredi *et al.*, 1998). Moreover, the mean apparent  $K_B$  values of yohimbine for this antagonism in vas deferens and brain membranes (10.1 and 1.8 nM respectively) were not significantly different either from each other (Results), or from a previously reported  $K_i$  value (5.8 nM) for its binding to human α<sub>2A</sub>-adrenoceptors (Newman-Tancredi *et al.*, 1998). The hypothesis that cannabigerol can activate α<sub>2</sub>-adrenoceptors is also supported by data obtained with yohimbine and two selective α<sub>2</sub>-adrenoceptor agonists, dexmedetomidine and clonidine. These data show that yohimbine possesses similar potency as an antagonist of cannabigerol-, dexmedetomidine- and clonidine-induced inhibition of electrically evoked contractions of the vas deferens (Figure 3) and as an antagonist of cannabigerol and dexmedetomidine, in the [<sup>35</sup>S]GTPγS binding assay (Figure 5). The evidence that cannabigerol targets a prejunctional site to inhibit electrically evoked contractions of the vas deferens (see previous paragraph) and that prejunctional α<sub>2</sub>-adrenoceptors can mediate such contractions lends further support to the hypothesis that cannabigerol can activate α<sub>2</sub>-adrenoceptors. Additional experiments are now required first to establish whether the inhibitory effect of cannabigerol in the vas deferens and its stimulatory effect in the [<sup>35</sup>S]GTPγS binding assay are mediated by α<sub>2A</sub>-, α<sub>2B</sub>- and/or α<sub>2C</sub>-adrenoceptors, and second to investigate why the potency that cannabigerol displays in these two bioassays is so different (Table 1 and Figures 2 and 3). In the meantime, it is noteworthy that there is evidence that electrically evoked neuronal release of noradrenaline from the vas deferens can be inhibited by the activation not only of α<sub>2A</sub>-adrenoceptors but also of α<sub>2C</sub>-adrenoceptors (Scheibner *et al.*, 2001), particularly when the frequency of the electrical stimulation is relatively low as it was in this investigation.

There is also a need for further experiments directed at establishing whether cannabigerol shares the ability of two other plant cannabinoids, Δ<sup>9</sup>-tetrahydrocannabinol and cannabidiol, to inhibit the neuronal uptake of noradrenaline (see Pertwee, 2008). Thus, our experiments showed first that the noradrenaline uptake inhibitor, maprotiline, can suppress electrically evoked contractions of the mouse isolated vas deferens, and second that yohimbine antagonizes cannabigerol and maprotiline in this bioassay with similar potency. It remains possible, therefore, that cannabigerol inhibits the twitch response of the vas deferens at least in part by blocking the reuptake of released noradrenaline in a manner that causes this catecholamine to accumulate at prejunctional α<sub>2</sub>-adrenoceptors and so inhibit both its evoked release and that of ATP. If cannabigerol can indeed inhibit noradrenaline reuptake, this would be consistent with a preliminary report that it increases struggling behaviour in the mouse tail suspension test, an indication that it may possess antidepressant activity (Musty and Deyo, 2006). Importantly though, cannabigerol could not have acted in this way to stimulate [<sup>35</sup>S]GTPγS binding to the mouse brain membranes we used in this investigation, leaving open the possibility that it might be both a potent direct α<sub>2</sub>-adrenoceptor agonist and an

inhibitor of noradrenaline reuptake. It is also noteworthy that it is unlikely that any elevation in extracellular concentration of noradrenaline induced by inhibition of its neuronal uptake would produce much  $\alpha_1$ -adrenoceptor mediated augmentation of electrically evoked contractions of the vas deferens. Thus, there is evidence that under the stimulation conditions used in this investigation, the contraction amplitude of the vas deferens is determined much more by ATP-induced activation of postjunctional P2X purinoceptors than by noradrenaline-induced activation of postjunctional  $\alpha_1$ -adrenoceptors (Pertwee *et al.*, 2002).

Whereas cannabigerol was found to display significantly less potency and efficacy than dexmedetomidine as an inhibitor of electrically evoked contractions of the vas deferens (Figure 3), in the [ $^{35}$ S]GTP $\gamma$ S binding assay in which both compounds displayed high potency but rather low efficacy (Figure 5), no such differences were detected. Why this should be remains to be established, possible explanations being that different  $\alpha_2$ -adrenoceptor subtypes, or indeed different types of receptor, mediate the effects of cannabigerol in brain and vas deferens and/or, as just discussed, that inhibition of noradrenaline reuptake may play a significant part in cannabigerol-induced inhibition of evoked contractions of the vas deferens.

The stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse brain membranes produced by concentrations of cannabigerol in the low nanomolar range does not appear to be cannabinoid CB $_1$  receptor-mediated. However, evidence was obtained that at higher concentrations, this phytocannabinoid can target the CB $_1$  receptor as an antagonist. Thus, in experiments performed with MF1 mouse brain membranes, cannabigerol was found to reduce specific binding of [ $^3$ H]CP55940 to MF1 mouse brain membranes ( $K_i = 381$  nM) and, at 10  $\mu$ M, to antagonize the cannabinoid receptor agonists, anandamide and CP55940, in the [ $^{35}$ S]GTP $\gamma$ S binding assay. No significant antagonism of anandamide was induced by 1  $\mu$ M cannabigerol.

Although the effect of cannabigerol on [ $^{35}$ S]GTP $\gamma$ S binding to MF1 mouse brain membranes is stimulatory at 1, 10 and 100 nM, it was found to be insignificant at 1  $\mu$ M and inhibitory at 10  $\mu$ M. Why the stimulatory effect of cannabigerol on [ $^{35}$ S]GTP $\gamma$ S binding to brain membranes disappears and then changes to an inhibitory effect as its concentration is progressively increased remains to be established. It does seem likely, however, that neither this concentration-dependent loss of the stimulatory effect of cannabigerol nor the inhibitory effect it produces at 10  $\mu$ M are CB $_1$  receptor-mediated, as both effects were also detectable in CB $_1^{-/-}$  C57BL/6J mouse brain membranes. The inhibitory effect produced by 10  $\mu$ M cannabigerol most probably explains why this concentration of this phytocannabinoid seemed to produce downward as well as right-ward shifts in the log concentration-response curves of anandamide and CP55940. When the component of cannabigerol-induced antagonism that seemed to arise from its ability to inhibit [ $^{35}$ S]GTP $\gamma$ S binding to MF1 mouse brain membranes in a seemingly CB $_1$  receptor-independent manner was excluded, significant right-ward shifts in the log concentration response curves of anandamide and CP55940 were still apparent (Figure 8). Importantly, the mean apparent  $K_B$  values calculated from these dextral shifts did not differ sig-

nificantly from the mean  $K_i$  value of cannabigerol for its displacement of [ $^3$ H]CP55940 to MF1 mouse brain membranes, lending further support to the hypothesis that cannabigerol is a CB $_1$  receptor antagonist. Cannabigerol reduced specific binding of [ $^3$ H]CP55940 not only to brain membranes but also to membranes obtained from CHO cells transfected with human CB $_2$  receptors. It did this in manner that suggests it to possess less affinity for CB $_2$  than CB $_1$  receptors. Further experiments are now required to determine whether cannabigerol is a CB $_2$  receptor agonist or antagonist.

Cannabigerol was also found to antagonize the 5-HT $_{1A}$  selective agonist, *R*-(+)-8-hydroxy-2-(di-*n*-propylamino) tetralin (8-OH-DPAT), in the [ $^{35}$ S]GTP $\gamma$ S binding assay in a seemingly competitive manner. This it could do at concentrations above those at which it induced an apparent  $\alpha_2$ -adrenoceptor mediated stimulation of [ $^{35}$ S]GTP $\gamma$ S binding to mouse brain membranes but below the concentration at which it produced detectable antagonism of anandamide and CP55940 in this bioassay. Cannabigerol therefore resembles the selective  $\alpha_2$ -adrenoceptor agonists, clonidine and dexmedetomidine, both of which also target 5-HT $_{1A}$  receptors at concentrations above those at which they activate  $\alpha_2$ -adrenoceptors. However, it differs from these other two compounds in blocking rather than activating the 5-HT $_{1A}$  receptor (Newman-Tancredi *et al.*, 1998). The antagonism of 8-OH-DPAT induced by cannabigerol was presumably CB $_1$  receptor-independent as it was produced with similar potency in experiments performed with CB $_1^{-/-}$  C57BL/6J mouse brain membranes, as in experiments performed with CB $_1^{+/+}$  C57BL/6J mouse brain membranes. Interestingly, in contrast to its effects on the log concentration-response curves of anandamide and CP55940, cannabigerol failed to produce a detectable downward displacement of the 8-OH-DPAT log concentration-response curve when it was administered at a concentration (10  $\mu$ M) that by itself produced a marked inhibition of [ $^{35}$ S]GTP $\gamma$ S binding to brain membranes. The reason for this difference remains to be established. The absence of any such downward displacement does, however, suggest that cannabigerol is a neutral 5-HT $_{1A}$  receptor antagonist that does not interact with this receptor as an inverse agonist.

In conclusion, we have obtained evidence from *in vitro* experiments that cannabigerol is a potent  $\alpha_2$ -adrenoceptor agonist. This was unexpected as the structure of this plant cannabinoid is unlike that of any established  $\alpha_2$ -adrenoceptor ligand and as no other cannabinoid has been reported to behave in this way. We have also obtained evidence that cannabigerol can block 5-HT $_{1A}$  and cannabinoid CB $_1$  receptors albeit with a potency lower than that with which it appears to activate  $\alpha_2$ -adrenoceptors. Further experiments are now required to investigate whether cannabigerol targets any particular subtype of  $\alpha_2$ -adrenoceptor and whether it inhibits the neuronal uptake of noradrenaline. It will also be important to establish first whether cannabigerol possesses high potency and significant efficacy as an  $\alpha_2$ -adrenoceptor agonist when administered *in vivo*, and second whether it displays significant potency *in vivo* as a 5-HT $_{1A}$  receptor antagonist. It would be of interest as well to establish whether cannabigerol can block CB $_1$  receptors *in vivo*, although there is already evidence from experiments with rhesus monkeys that at one dose at least (16.5  $\mu$ g·kg $^{-1}$  i.v.) cannabigerol does not alter the ability

of  $\Delta^9$ -THC to induce *in vivo* effects that are presumably CB<sub>1</sub> receptor-mediated (Mechoulam *et al.*, 1970). In addition, it would be of interest to investigate whether any ability cannabigerol has to activate  $\alpha_2$ -adrenoceptors *in vivo* ceases to be detectable at higher doses as happens *in vitro*. It will also be important to identify which pharmacological actions of cannabigerol are responsible for its reported ability to increase struggling behaviour in the mouse tail suspension test (Musty and Deyo, 2006) or to inhibit human keratinocyte proliferation (Wilkinson and Williamson, 2007), as these effects indicate that cannabigerol may have therapeutic potential as an antidepressant and/or for the treatment of psoriasis. The possibility that cannabigerol has other clinical applications, for example for the production of  $\alpha_2$ -adrenoceptor-mediated analgesia (Tryba and Gehling, 2002; Giovannoni *et al.*, 2009), also merits investigation.

## Acknowledgements

This investigation was supported by grants from GW Pharmaceuticals and the National Institutes of Health (DA-03672).

## Conflict of interest

None.

## References

- Alexander SPH, Mathie A, Peters JA (2008). Guide to Receptors and Channels (GRAC), 3rd edn. *Br J Pharmacol* **153** (Suppl. 2): S1–S209.
- Barbaccia ML, Ravizza L, Costa E (1986). Maprotiline: an antidepressant with an unusual pharmacological profile. *J Pharmacol Exp Ther* **236**: 307–312.
- Breivogel CS, Griffin G, Di Marzo V, Martin BR (2001). Evidence for a new G protein-coupled cannabinoid receptor in mouse brain. *Mol Pharmacol* **60**: 155–163.
- Cascio MG, Gauson LA, Stevenson LA, Ross RA, Pertwee RG (2009). Cannabigerol behaves as a potent alpha-2-adrenoceptor partial agonist. *Symposium on the Cannabinoids. Burlington, Vermont, International Cannabinoid Research Society*, p. 73.
- Cheng Y-C, Prusoff WH (1973). Relationship between the inhibition constant ( $K_i$ ) and the concentration of inhibitor which causes 50 percent inhibition ( $IC_{50}$ ) of an enzymatic reaction. *Biochem Pharmacol* **22**: 3099–3108.
- De Petrocellis L, Vellani V, Schiano-Moriello A, Marini P, Magherini PC, Orlando P *et al.* (2008). Plant-derived cannabinoids modulate the activity of transient receptor potential channels of ankyrin type-1 and melastatin type-8. *J Pharmacol Exp Ther* **325**: 1007–1015.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G *et al.* (1992). Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* **258**: 1946–1949.
- ElSohly MA, Slade D (2005). Chemical constituents of marijuana: the complex mixture of natural cannabinoids. *Life Sci* **78**: 539–548.
- Forster EA, Cliffe IA, Bill DJ, Dover GM, Jones D, Reilly Y *et al.* (1995). A pharmacological profile of the selective silent 5-HT<sub>1A</sub> receptor antagonist, Way-100635. *Eur J Pharmacol* **281**: 81–88.
- Gaoni Y, Mechoulam R (1964). Hashish II. The structure and synthesis of cannabigerol, a new hashish constituent. *Proc Chem Soc (London)*, p. 82.
- Gauson LA, Cascio MG, Thomas A, Ross RA, Pertwee RG (2008). Some pharmacological effects of five little-investigated plant cannabinoids. *Symposium on the Cannabinoids. Burlington, Vermont, International Cannabinoid Research Society*, p. 143.
- Gauson LA, Cascio MG, Ross RA, Pertwee RG (2009). Cannabigerol displays significant potency as a 5-HT<sub>1A</sub> receptor antagonist. *Symposium on the Cannabinoids. Burlington, Vermont, International Cannabinoid Research Society*, p. 25.
- Giovannoni MP, Ghelardini C, Vergelli C, Dal Piaz V (2009).  $\alpha_2$ -agonists as analgesic agents. *Med Res Rev* **29**: 339–368.
- Grunfeld Y, Edery H (1969). Psychopharmacological activity of the active constituents of hashish and some related cannabinoids. *Psychopharmacologia* **14**: 200–210.
- von Kügelgen I, Starke K (1991). Noradrenaline-ATP co-transmission in the sympathetic nervous system. *Trends Pharmacol Sci* **12**: 319–324.
- Kurkinen KMA, Koistinaho J, Laitinen JT (1997). [ $\gamma$ -<sup>35</sup>S]GTP autoradiography allows region-specific detection of muscarinic receptor-dependent G-protein activation in the chick optic tectum. *Brain Res* **769**: 21–28.
- Mechoulam R (2005). Plant cannabinoids: a neglected pharmacological treasure trove. *Br J Pharmacol* **146**: 913–915.
- Mechoulam R, Shani A, Edery H, Grunfeld Y (1970). Chemical basis of hashish activity. *Science* **169**: 611–612.
- Musty RE, Deyo RA (2006). A cannabigerol extract alters behavioral despair in an animal model of depression. *Symposium on the Cannabinoids. Burlington, Vermont, International Cannabinoid Research Society*, p. 32.
- Newman-Tancredi A, Nicolas J-P, Audinot V, Gavaudan S, Verrière L, Touzard M *et al.* (1998). Actions of  $\alpha_2$  adrenoceptor ligands at  $\alpha_{2A}$  and 5-HT<sub>1A</sub> receptors: the antagonist, atipamezole, and the agonist, dexmedetomidine, are highly selective for  $\alpha_{2A}$  adrenoceptors. *Naunyn-Schmiedeberg's Arch Pharmacol* **358**: 197–206.
- Pertwee RG (1997). Pharmacology of cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptors. *Pharmacol Ther* **74**: 129–180.
- Pertwee RG (2008). The diverse CB<sub>1</sub> and CB<sub>2</sub> receptor pharmacology of three plant cannabinoids:  $\Delta^9$ -tetrahydrocannabinol, cannabidiol and  $\Delta^9$ -tetrahydrocannabivarin. *Br J Pharmacol* **153**: 199–215.
- Pertwee RG, Griffin G, Lainton JAH, Huffman JW (1995). Pharmacological characterization of three novel cannabinoid receptor agonists in the mouse isolated vas deferens. *Eur J Pharmacol* **284**: 241–247.
- Pertwee RG, Fernando SR, Griffin G, Ryan W, Razdan RK, Compton DR *et al.* (1996). Agonist-antagonist characterization of 6'-cyanohept-2'-yne- $\Delta^8$ -tetrahydrocannabinol in two isolated tissue preparations. *Eur J Pharmacol* **315**: 195–201.
- Pertwee RG, Ross RA, Craib SJ, Thomas A (2002). (-)-Cannabidiol antagonizes cannabinoid receptor agonists and noradrenaline in the mouse vas deferens. *Eur J Pharmacol* **456**: 99–106.
- Pertwee RG, Thomas A, Stevenson LA, Maor Y, Mechoulam R (2005). Evidence that (-)-7-hydroxy-4'-dimethylheptyl-cannabidiol activates a non-CB<sub>1</sub>, non-CB<sub>2</sub>, non-TRPV1 target in the mouse vas deferens. *Neuropharmacology* **48**: 1139–1146.
- Ross RA, Brockie HC, Stevenson LA, Murphy VL, Templeton F, Makriyannis A *et al.* (1999a). Agonist-inverse agonist characterization at CB<sub>1</sub> and CB<sub>2</sub> cannabinoid receptors of L759633, L759656 and AM630. *Br J Pharmacol* **126**: 665–672.
- Ross RA, Gibson TM, Stevenson LA, Saha B, Crocker P, Razdan RK *et al.* (1999b). Structural determinants of the partial agonist-inverse agonist properties of 6'-azidohept-2'-yne- $\Delta^8$ -tetrahydrocannabinol at cannabinoid receptors. *Br J Pharmacol* **128**: 735–743.
- Ross RA, Gibson TM, Brockie HC, Leslie M, Pashmi G, Craib SJ *et al.* (2001). Structure-activity relationship for the endogenous cannabinoid, anandamide, and certain of its analogues at vanilloid

- receptors in transfected cells and vas deferens. *Br J Pharmacol* **132**: 631–640.
- Scheibner J, Trendelenburg A-U, Hein L, Starke K (2001). Stimulation frequency-noradrenaline release relationships examined in  $\alpha_{2A}$ -,  $\alpha_{2B}$ - and  $\alpha_{2C}$ -adrenoceptor-deficient mice. Naunyn-Schmiedeberg's. *Arch Pharmacol* **364**: 321–328.
- Schlicker E, Kathmann M (2001). Modulation of transmitter release via presynaptic cannabinoid receptors. *Trends Pharmacol Sci* **22**: 565–572.
- Starke K (2001). Presynaptic autoreceptors in the third decade: focus on  $\alpha_2$ -adrenoceptors. *J Neurochem* **78**: 685–693.
- Tallarida RJ, Cowan A, Adler MW (1979). pA<sub>2</sub> and receptor differentiation: a statistical analysis of competitive antagonism. *Life Sci* **25**: 637–654.
- Thomas A, Ross RA, Saha B, Mahadevan A, Razdan RK, Pertwee RG (2004). 6''-Azidohex-2''-yne-cannabidiol: a potential neutral, competitive cannabinoid CB<sub>1</sub> receptor antagonist. *Eur J Pharmacol* **487**: 213–221.
- Thomas A, Stevenson LA, Wease KN, Price MR, Baillie G, Ross RA *et al.* (2005). Evidence that the plant cannabinoid  $\Delta^9$ -tetrahydrocannabinol is a cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptor antagonist. *Br J Pharmacol* **146**: 917–926.
- Thomas A, Baillie GL, Phillips AM, Razdan RK, Ross RA, Pertwee RG (2007). Cannabidiol displays unexpectedly high potency as an antagonist of CB<sub>1</sub> and CB<sub>2</sub> receptor agonists in vitro. *Br J Pharmacol* **150**: 613–623.
- Trendelenburg AU, Cox SL, Schelb V, Klebroff W, Khairallah L, Starke K (2000). Modulation of <sup>3</sup>H-noradrenaline release by presynaptic opioid, cannabinoid and bradykinin receptors and  $\beta$ -adrenoceptors in mouse tissues. *Br J Pharmacol* **130**: 321–330.
- Tryba M, Gehling M (2002). Clonidine – a potent analgesic adjuvant. *Curr Opin Anaesthesiol* **15**: 511–517.
- Wilkinson JD, Williamson EM (2007). Cannabinoids inhibit human keratinocyte proliferation through a non-CB<sub>1</sub>/CB<sub>2</sub> mechanism and have a potential therapeutic value in the treatment of psoriasis. *J Dermatol Sci* **45**: 87–92.