

# **Evolutionary Genetics in the Wild – from Populations to Individuals**

Tuomas Leinonen

Department of Biosciences  
Faculty of Biological and Environmental Sciences  
University of Helsinki  
Finland

Academic dissertation

To be presented for public examination with the permission of the Faculty of Biological and Environmental Sciences of the University of Helsinki in lecture hall 2, Infocentre Korona (Viikinkaari 11) on February 5<sup>th</sup> 2010 at 12 noon.

Supervised by: Acad. Prof. Juha Merilä  
Department of Biosciences  
University of Helsinki, Finland

PhD Jose Manuel Cano Arias  
Department of Biosciences  
University of Helsinki, Finland

Reviewed by: PhD Joost Raeymaekers  
Biology Department  
University of Leuven, Belgium

PhD Folmer Bokma  
Department of Ecology and Environmental  
Science  
Umeå University, Sweden

Examined by: Prof. Michael A. Bell  
Department of Ecology and Evolution  
Stony Brook University, USA

Custos: Prof. Pekka Pamilo  
Department of Biosciences  
University of Helsinki, Finland

ISBN 978-952-92-6854-2 (paperback)  
ISBN 978-952-10-6094-6 (PDF)  
<http://ethesis.helsinki.fi>

Edita  
Helsinki 2010



## Contents

<b>Abstract</b> .....	<b>6</b>
<b>Introduction</b> .....	<b>7</b>
<i>Variation between populations - drift or selection?</i> .....	7
<i>Variation within populations - genetic variance and covariance structure</i> .....	8
<i>Variation between individuals within populations</i> .....	12
<i>Threespine stickleback as a model organism</i> .....	14
<b>Aims of this thesis</b> .....	<b>15</b>
<b>Main results and discussion</b> .....	<b>16</b>
<i>Relative roles of drift and selection in population divergence</i> .....	16
<i>Within population variance and population divergence</i> .....	18
<i>The role of selection in adaptive divergence</i> .....	20
<b>Conclusions and future directions</b> .....	<b>21</b>
<b>Acknowledgements</b> .....	<b>22</b>
<b>Literature cited</b> .....	<b>24</b>

This thesis is based on the following articles, which are referred to in text by their Roman numerals:

- I Leinonen, T., Cano, J.M., Mäkinen, H. & Merilä, J. 2006. Contrasting patterns of body shape and neutral genetic divergence in marine and lake populations of threespine sticklebacks. *Journal of Evolutionary Biology* 19: 1803-1812.
- II Leinonen, T., O'Hara, R.B., Cano, J.M. & Merilä, J. 2008. Comparative studies of quantitative trait and neutral marker divergence: a meta-analysis. *Journal of Evolutionary Biology* 21: 1-17.
- III Leinonen, T., Cano, J.M. & Merilä, J. Genetics of body shape and armour variation in threespine sticklebacks. *Manuscript*.
- IV Leinonen, T., Cano, J.M. & Merilä, J. Genetic basis of sexual dimorphism in the threespine stickleback. *Manuscript*.
- V Loehr, J., Leinonen, T., Herczeg, G., O'Hara, R.B. & Merilä, J. Heritability of fluctuating asymmetry in the threespine stickleback. *Manuscript*.
- VI Leinonen, T., Cano, J.M., Herczeg, G. & Merilä, J. Predation-imposed selection on threespine stickleback (*Gasterosteus aculeatus*) morphology: a test of the refuge usage hypothesis. *Manuscript*.

## Contributions

	I	II	III	IV	V	VI
Concept	TL, JC, JM	JM	TL, JC, JM	TL	GH, JL, JM	TL, JC
Data collection	TL, HM	TL, JC, JM	TL, JC	TL	TL, JL	TL
Analyses	TL, JC	BO, TL	TL	TL	BO, TL	TL, GH
Manuscript preparation	TL, JC, JM	TL, JC, JM	TL, JC, JM	TL, JC, JM	JL, TL, GH, JM	TL, GH, JM

TL: Tuomas Leinonen

JC: José Manuel Cano Arias

JM: Juha Merilä

BO: Robert B. O'Hara

GH: Gábor Herczeg

JL: John Loehr

© Blackwell Publishing (Chapters I and II)

© Authors (Chapters 0, III, IV, V and VI)

Cover by José Manuel Cano Arias

## Abstract

Predicting evolutionary outcomes and reconstructing past evolutionary transitions are among the main goals of evolutionary biology. Ultimately, understanding the mechanisms of evolutionary change will also provide answers to the timely question of whether and how organisms will adapt to changing environmental conditions.

In this thesis, I have investigated the relative roles of natural selection, random genetic drift and genetic correlations in the evolution of complex traits at different levels of organisation – from populations to individuals. I have shown that natural selection has been the driving force behind body shape divergence of marine and freshwater threespine stickleback (*Gasterosteus aculeatus*) populations, while genetic drift may have played a significant role in the more fine scale divergence among isolated freshwater populations. These results are concurrent with the patterns that have emerged in the published studies comparing the relative importance of natural selection and genetic drift as explanations for population divergence in different traits and taxa.

I have also shown that body shape and armour divergence among threespine stickleback populations is likely to be biased by the patterns of genetic variation and covariation. Body shape and armour variation along the most likely direction of evolution – the direction of maximum genetic variance – reflects the general patterns of variation observed wild populations across the distribution range of the threespine stickleback. Conversely, it appears that genetic correlations between the sexes have not imposed significant constraints on the evolution of sexual dimorphism in threespine stickleback body shape and armour.

I have demonstrated that the patterns of evolution seen in the wild can be experimentally recreated to tease out the effects of different selection agents in detail. In addition, I have shown how important it is to take into account the correlative nature of traits, when making interpretations about the effects of natural selection on individual traits. Overall, this thesis provides a demonstration of how considering the relative roles of different mechanism of evolutionary change – at different levels of organisation – can aid in an emergence of a comprehensive picture of how adaptive divergence in wild populations occurs.

## Introduction

Understanding how organisms adapt to changing environments is an integral part of biology, extending from basic ecology and conservation biology to agriculture and medicine. Adaptation to changing environments entails evolutionary change, which is dependent on heritable variation. Evolutionary genetics provides tools for quantifying variation at different levels of resolution, from populations to individual loci, and serves as a platform from which to reach for one of the most important goals of evolutionary biology – predicting and reconstructing evolutionary change in wild populations.

In its very essence, evolution can be defined as a change in phenotypic means, which is a function of genetic drift, natural selection and the matrix of genetic variances and covariances,  $\mathbf{G}$  (Polly 2008). These three components of evolution are intertwined so that population size and  $\mathbf{G}$  determine genetic drift, while an adaptive landscape in terms of fitness determines the strength and direction of selection (Lande 1976, Arnold et al. 2001, Gavrillets 2004). Predicting evolutionary change in wild populations includes estimating all the above three essentials, which in turn requires investigation of variation at different levels of resolution.

### *Variation between populations - drift or selection?*

When starting to uncover the details of variation observed in nature, the foremost issue to consider is that evolution usually takes place on such a long time scale that the observed patterns of phenotypic change could be explained by very little selection or even by genetic drift alone (Lande 1976, Lynch 1988, Kinnison & Hendry 2001; but see Estes & Arnold 2007).

Genetic drift is more likely to play a larger role in populations with small effective population sizes ( $N_e$ ), where the rate of genetic drift is higher than in populations with large  $N_e$ . The higher rate of genetic drift in small populations provides ample scope for non-adaptive differentiation, which in turn decreases the efficiency of natural selection. It can thus be said that the efficiency of natural selection is inversely related to  $N_e$  (e.g. Jones et al. 1968, Frankham et al. 2002, England et al. 2003).

The relative importance of natural selection and genetic drift in explaining evolutionary change has constituted a long-standing debate in evolutionary biology (e.g. Merilä & Crnokrak 2001, McKay & Latta 2002). The debate has remained unresolved, although statistical tools for testing the differentiation have been around for some time (Lande 1976, 1977, Chackraborty & Nei 1982, Lynch & Hill 1986, Lynch 1988, Turelli 1988). The problem with these tools is their inherent assumptions on critical parameters, such as mutational input, time since population divergence and population size, which are often impossible to quantify (e.g. Turelli 1988, Lande 1992, Lynch 1994). However, there is an alternative approach, which is based on comparison of genetic differentiation in neutral marker loci ( $F_{ST}$ ) and quantitative traits ( $Q_{ST}$ ; for details on the calculation of the indices see Box 1). This method has also been available for as long as the first mentioned method (Wright 1951, 1965, Rogers & Harpending 1983, Felsenstein 1986), but for some reason – perhaps partly due to early criticism by Lewontin (1984) on the actual information content of the marker vs. quantitative trait comparisons – it has not been widely adopted prior to the recent

decade or so. Another factor that has possibly hindered the usage of the  $F_{ST}$  vs.  $Q_{ST}$  comparisons, is that estimating the index of quantitative genetic differentiation requires teasing out the genetic component of variance, which requires controlled breeding experiments or known pedigrees of large sample size and a large number of populations (Lynch & Walsh 1998, O'Hara & Merilä 2005).

$F_{ST}$  and  $Q_{ST}$  estimates are based on the relationship of between-population and within-population variance, so that the estimate of population differentiation in a quantitative trait,  $Q_{ST}$ , is analogous to population differentiation at a single locus  $F_{ST}$  (Spitze 1993). There are three possible outcomes of the  $F_{ST}$  vs.  $Q_{ST}$  comparisons with different evolutionary implications (Merilä & Crnokrak 2001). First, if  $Q_{ST}$  is higher than  $F_{ST}$ , directional natural selection must have played a role in the observed population divergence. In this case selection is not favouring similar phenotypes in different populations, which has been the most common outcome in empirical studies (Merilä & Crnokrak 2001). The second possible outcome is that the indices of divergence are equal ( $F_{ST} \approx Q_{ST}$ ). This can be interpreted so that the observed degree of differentiation could have been a product of genetic drift alone. In other words, selection may have been in action, but its effects cannot be distinguished from drift. The third possible outcome,  $F_{ST} > Q_{ST}$  means that the degree of differentiation is less than what would be expected by genetic drift alone. The interpretation of this situation is that natural selection favours similar phenotypes in different populations. In a rare case, this outcome could reflect a situation, where there is not enough genetic variability in the studied populations so that natural selection does not have

any variance to act on (Merilä & Crnokrak 2001).

It must be noted that the approach of comparing the quantitative genetic and neutral genetic differentiation has a number of inherent limitations in both the estimation of  $F_{ST}$  (e.g. Hedrick 1999, Meirmans 2006) and  $Q_{ST}$  (e.g. O'Hara & Merilä 2005, Pujol et al. 2008, Whitlock 2008). Despite these possible problems, the method of comparing  $F_{ST}$  vs.  $Q_{ST}$  estimates provides a good platform for exploring the forces that are in play in phenotypic divergence, and thus, is a natural starting point for the investigation of phenotypic divergence in more detail.

#### *Variation within populations - genetic variance and covariance structure*

Whether the driving force for phenotypic change is natural selection or genetic drift, there will be no evolutionary change if there is no heritable genetic variation. In the simplest case, the interest is on a single trait and evolution (change in the mean of a trait – response,  $R$ ) can be summarised by the breeder's equation:

$$R = h^2S$$

where  $h^2$  denotes the heritability of the trait and  $S$  the selection differential. As the name implies, the origins of the equation lie in animal breeding, where the interest is usually in improving a single trait. However, predicting evolution in wild populations usually involves considering a number of traits simultaneously. In addition, selection rarely, if ever, acts on a single trait alone (Lande 1986), and evolutionary change in a heritable trait is not free to take a direct pathway to its optimal phenotype as imposed by natural selection. The paths of evolution are bounded by the constraints set by the patterns of genetic variation and



development (Bonner 1982, Maynard Smith et al. 1985), which is the reason why it is necessary to have information on genetic variances and covariances of the traits when trying to unravel the details behind phenotypic variation.

The influence of genetic covariance between traits on evolution can be summarised with the multivariate extension of the breeder's equation (Lande 1979, Lande & Arnold 1983):

$$\Delta\mathbf{z} = \mathbf{G}\mathbf{P}^{-1}\mathbf{s}$$

which returns a vector of changes in mean trait values ( $\Delta\mathbf{z}$ ), and where  $\mathbf{G}$  is the matrix of genetic variances and covariances,  $\mathbf{P}$  its phenotypic equivalent and  $\mathbf{s}$  is the vector of selection differentials. To spell it out,  $\mathbf{G}$  describes linkage or pleiotropic connections between traits that might cause correlated patterns of evolution in the traits.  $\mathbf{P}$  depicts phenotypic relations among traits that result from the interplay of the genetic covariance matrix and the developmental processes influenced by the environment.

Estimation of the heritable component of variation requires either controlled breeding experiments in the lab, sampling in the wild so that the pedigree matches a standard design (such as parent-offspring), or using an animal model for a population sampled over multiple generations (reviewed by Roff 2007). Unfortunately, none of the above-mentioned methods are free of problems. Common, and perhaps the most critical problem with all quantitative genetic parameter estimation methods is that they require a rather large sample size (Lynch & Walsh 1998). With a single or a limited number of traits, it is possible to get estimates with reasonable power, but as the number of traits increases, so does the required sample size, approximately as the quartic root of the number of traits (Phillips 1998). In

addition, the power to detect significant heritability decreases when the heritability estimates get smaller (Lynch & Walsh 1998), a situation to be expected when the complexity of traits increases, for example as in life history traits (Price & Schluter 1991, Merilä & Sheldon 1999). As a result, studies of genetic correlations have traditionally been restricted to consider a very limited number of traits. However, the integration of quantitative genetics with geometric morphometric methods (Box 2), has paved the way for quantitative genetic investigation of complex (and comprehensive) traits, such as body shape (Klingenberg & Leamy 2001, Monteiro et al. 2002, Klingenberg & Monteiro 2005, McGuigan et al. 2005).

It must be noted though that single trait cases are not free of problems either. Heritability estimates can easily be misinterpreted, if one does not keep in mind that they are specific to population, environment, generation, age and model used for their estimation (e.g. Hoffmann & Merilä 1999, Charmantier & Garant 2005, Wilson 2008). The issues that undermine the usefulness of heritability in predicting evolution are also reflected in the analysis of multiple traits, which possibly complicates the use of  $\mathbf{G}$  in reconstructing the patterns of evolution that have led to the phenotypic differentiation observed today (e.g. Pigliucci 2006). The usefulness of  $\mathbf{G}$  depends on its stability over time (Turelli 1988) – an issue as yet unresolved (Roff 2000, Stepan et al. 2002, but see Björklund 2004). One solution to this possible problem is to use the  $\mathbf{G}$  of an ancestral population, which unfortunately is rarely possible. Of course, the stability of  $\mathbf{G}$  over time and in different populations depends on the traits in question. For instance, if the genetic covariances result from a pleiotropy among numerous alleles at

**Box 1.** Glossary – abbreviations of quantitative genetic concepts

$N_e$	Effective population size – the average size of a population in terms of individuals that can contribute genes equally to the next generation
FA	Fluctuating asymmetry – small, non directional departures from perfect symmetry
$F_{ST}$	Measure of genetic differentiation in neutral marker genes <sup>1,2</sup> $F_{ST} = \frac{V_b}{(V_b + V_w)}$ $V_b$ = variation between populations $V_w$ = variation within populations
$\mathbf{g}_{max}$	Direction of maximum additive genetic variance or ‘genetic line of least resistance’. The dominant eigenvector of the <b>G</b> matrix <sup>3</sup>
<b>G</b>	Matrix of genetic variances and covariances, calculated based on phenotypic similarity among relatives. $\mathbf{G} = \begin{pmatrix} \sigma^2_{A(1)} & \sigma_{A(1,2)} & \dots & \sigma_{A(1,j)} \\ \sigma_{A(2,1)} & \sigma^2_{A(2)} & \dots & \sigma_{A(2,j)} \\ \dots & \dots & \dots & \dots \\ \sigma_{A(i,1)} & \sigma_{A(i,2)} & \dots & \sigma^2_{A(n)} \end{pmatrix}$ $\sigma^2_{A(i)} =$ variance of trait $i$ $\sigma_{A(i,j)} =$ covariance between traits $i$ and $j$
$h^2$	Heritability – proportion of phenotypic variation in a population that is attributable to genetic variation among individuals $h^2 = \frac{V_A}{V_P}$ $V_A$ = additive genetic variation, $V_P$ = phenotypic variation
<b>P</b>	Matrix of phenotypic variances and covariances (see <b>G</b> for an example)
$Q_{ST}$	Measure of differentiation in quantitative traits <sup>4</sup> $Q_{ST} = \frac{V_b}{(2V_b + V_w)}$ $V_b$ = variation between populations $V_w$ = variation within populations
<b>S</b>	Selection differential – difference between phenotype of selected parents and population mean
<b>s</b>	Vector of selection differentials

References: 1. Wright (1951), 2. Nei (1987), 3. Schluter 1996, 4. Spitze (1993)

**Box 2. A primer to geometric morphometrics**

Geometric morphometrics uses outline or landmark based methods (Adams *et al.* 2004). Landmark analysis relies on biologically definable points, such as points of intersection of bones, which are homologous across all specimens. This makes biological interpretations more holistic, which is the main advantage of landmark-based methods (Rohlf & Marcus 1993). Reyment (2002) defines landmarks as recognizable equivalent points observed on the objects being compared. For each specimen, the landmarks are given coordinates (x and y) on a coordinate plane (coordinates can also be in a three-dimensional plane, when the z coordinate is included). When all the specimens have been given corresponding landmarks, differences in coordinates for each landmark can be compared. This analysis begins with superimposing the landmark configurations of each specimen on each other. Superimposition basically means overlaying one specimen's coordinates on top of another's, so that their homologous landmarks match as closely as possible (Rohlf & Marcus 1993). The aim of superimposition is to remove non-shape variation from a set of data. Non-shape variation is due to changes in translation, rotation, and scale of specimens (Slice *et al.* 1996). The most common of the superimposition methods used today are based on least-squares methods (also called Procrustes analysis; Slice 2001, Adams *et al.* 2004), which estimate the parameters for location and orientation that minimise the sum of squared distances between corresponding points on two configurations (Bookstein 1991). This is done by first translating the centroid of each configuration to the origin and scaling the configurations according to their centroid sizes (Rohlf & Slice 1990). Centroid is essentially the centre of gravity of a specimen as derived from its landmark configuration, and centroid size is the square root of the sums of squared distances of a set of landmarks from their centroid (Slice *et al.* 1996). Superimposition is completed by rotating the configurations so that the squared differences between corresponding landmarks are minimized (Rohlf & Slice 1990). With every specimen the process is iterated to get the mean (reference) shape, which cannot be estimated before superimposition (Adams *et al.* 2004).

After superimposition, the landmark configurations lie on a common coordinate system and the coordinates of landmarks can be used as shape variables (Adams *et al.* 2004). The whole range of multivariate statistics can be used to explore the structure of shape variation. Once the statistical analysis is done, and here lies the beauty of geometric morphometrics, the results can be easily visualised in the space of the original specimens. The most widely used visualisation method today is thin-plate spline (Reyment 2002). Landmarks on a thin-plate spline are thought to lie on an indefinitely thin metal plate. For example, if we had two specimens, one's landmark configuration would lie on a straight plate, while the plate would be bent to fit the configuration of the other one. This new surface should be the least bent of any surface that passes through the specified heights at the locations of the reference landmarks (Bookstein 1991). The difference in the shape of the two specimens can then be expressed as the energy needed to bend the straight plate to its new shape. This is easiest to visualise if a grid is used as a plate. Bending of the grid shows where the variation in specimens lies.

The results of thin-plate spline analyses are usually expressed as warps. Thin-plate spline analysis takes into account the variation at each landmark separately. At each landmark, the difference of a specimen from the mean configuration is called a partial warp. Partial warps are vectors that lie on a coordinate plane defined by principal warps, which in turn are orthogonal shape axes that represent all possible deformations that could occur starting from the reference configuration. When

partial warps are used in a principal component analysis together, so that total shape variation is taken into account, the resulting differences are called relative warps. Scores from the relative warp analysis can be plotted on a grid, where deformation of the grid reflects the variation in the shape of the specimens.

When morphological data are compared with genetic data, it is useful to have an overall measure of shape variance. Analyses based on Procrustes distance have been previously used to assess the magnitude of individual variation, fluctuating asymmetry and measurement error (Klingenberg & McIntyre 1998). Procrustes distance has been also used in quantitative genetics (Monteiro *et al.* 2002), although with some criticism (Klingenberg 2003). Procrustes distance is the square root of the sums of squared differences between corresponding landmarks (Rohlf 1999). It therefore combines all the variation in the landmarks to produce a single measure of overall shape variability between specimens, which can then be compared for example with indices of genetic variability, such as  $F_{ST}$  (see Chapter I).

each locus with a multinormal distribution of effects, the  $\mathbf{G}$  matrix is likely to have remained relatively constant, even under changing selective pressures (Blows & Higgin 2003).

When the  $\mathbf{G}$  matrix is assumed to be constant, genetic covariances are likely to constrain evolution so that phenotypic change is most likely to take place along the direction of maximum additive genetic variance ( $\mathbf{g}_{max}$ ), which in essence is the dominant eigenvector of the  $\mathbf{G}$  matrix (Schluter 1996). However, apart from Schluter's (1996) original tests on evolution along the lines of least resistance, very few studies have found a positive association between the  $\mathbf{g}_{max}$  and phenotypic divergence (but see Arnold & Phillips 1999, Blows & Higgin 2003, McGuigan *et al.* 2005). This indicates that it is possible to break down genetic constraints (i.e. to evolve into a direction different from  $\mathbf{g}_{max}$ ), especially in the long term, and that the constraints themselves are subject to evolution (i.e. the orientation and magnitude of  $\mathbf{G}$  may change with time; Lande 1976, Zeng 1988, Arnold 1992). The impermanence of genetic constraints is also supported by the vast diversity of organisms, even within populations.

### *Variation between individuals within populations*

Within population divergence, namely that between the sexes, offers a way to test the effects of genetic constraints on evolutionary responses. Evolution of sexual dimorphism (SD) involves a puzzle: sex-specific selection acts on genes that are shared by both sexes (Fisher 1930, 1931). In other words, there is a genetic correlation between the sexes ( $r_{g(MF)}$ ), which is likely to slow down the evolution of SD (Lande 1980, Reeve & Fairbairn 2001, Fairbairn *et al.* 2007). Theoretically, in the extreme case, when genetic correlation between the sexes for a given trait is perfect ( $r_{g(MF)} = 1$ ), SD should not evolve. However, even in the presence of high  $r_{g(MF)}$ , SD can still evolve if the sexes differ in amount of genetic variance for a trait in question (Lynch & Walsh 1998).

Simulations have shown that genetic correlations between the sexes constrain the evolution of sexual dimorphism much less than predicted by Lande's (1980) model (Reeve & Fairbairn 2001), which is built around the matrix of genetic variances and covariances between the sexes. Evidence from empirical studies is mixed. A recent meta-analysis of 66 dioecious plant and animal

populations revealed a significant negative association between SD and  $r_{g(MF)}$ , while a survey of 68 studies on hermaphroditic plants revealed mixed results – the relationship between SD and  $r_{g(MF)}$  depended on the species in question (Ashman & Majetic 2006). Studies involving enough traits so that the association between SD and can be tested on the traits within the study are few, and the results from them are also mixed (no correlation: e.g. Cowley et al. 1986, Cowley & Atchley 1988; negative correlation: Ashman 2003, Bonduriansky & Rowe 2005, McDaniel 2005, Fairbairn et al. 2007). The evidence thus indicates that the effect of genetic constraints on the evolution as inferred from sexual dimorphism is still unresolved.


Whether the genomic conflict between sexes imposes severe constraints on the evolution of SD or not, the driving force behind SD is differential selection in males and females. The differential selection between sexes does not have to be sexual selection in its original, strict sense – effects caused by the struggle for mates (Darwin 1871). Disruptive selection between sexes can stem from differences in habitat that lead to differential evolutionary responses from males and females. In fact, similar ecological conditions as those that lie behind adaptive divergence between species or populations can also drive the evolution of SD (Bolnick & Doebeli 2003). Indeed, the ecological differences between the sexes can be larger than those between species (Schoener 1967, 1969, Dayan et al. 1994). It has been suggested that SD and adaptive divergence are interwoven so that when there are no constraints for the evolution of SD, the likelihood of adaptive divergence is restricted (Bolnick & Doebeli 2003). It is thus imperative to consider the differences, ecological and/or genetic, between the sexes, when aiming to

uncover the mechanisms behind adaptive divergence (e.g. Butler et al. 2007).

Sexual dimorphism provides an example of how divergence itself can be adaptive and reduce intraspecific competition within populations. Divergence between sexes can be thought of as a meta-trait, comprising of combined effects of a number of single traits, and thus it should evolve according to the same principles as any other trait. If divergence itself can evolve, it should comprise a heritable component. Indeed, sexual dimorphism of a variety of traits has been found to be heritable, but the estimates of heritability have been restricted to domestic animals or model species (e.g. Cowley et al. 1986, Chapuis et al. 1996, Mignon-Grasteau et al. 1998, David et al. 2003, but see Vandeputte et al. 2007). The published heritability estimates for SD are relatively low (Fig. 1), following the general pattern of decreasing heritability with increasing complexity of traits (Price & Schluter 1991, Merilä & Sheldon 1999).

Strong genetic correlations between the sexes can lead to the situation, where despite of sexual or disruptive selection, male and female phenotypes cannot reach their respective optima. Sexual dimorphism therefore provides one possible solution to the dilemma of how genetic variation is maintained in populations despite of directional selection. Theoretically, if natural selection is driving individuals towards a phenotype that is optimal in terms of fitness, variation around the optimal phenotype should in time diminish. Yet, genetic variation persists despite of strong directional selection (e.g. Barton & Turelli 1989), and even phenotypes that lie outside of the fitness optimum can be heritable. For instance, it has been shown that developmental instability, as inferred from fluctuating asymmetry, can

include a heritable component of variance (Møller & Thornhill 1997).



**Figure 1. Distribution of 61 heritability estimates of sexual dimorphism reported in literature. The heritability estimates were extracted from Araripe et al. (2008), Buvanendran (1969), Chapuis et al. (1996), Cowley et al. (1986), David et al. (2003), Hanrahan & Eisen 1973, Hu et al. (1999), Le Bihan-Duval et al. (1998), Lester et al. (1989), Mignon-Grasteau et al. (1998, 1999, 2004), and Vandeputte et al. (2007).**

#### *Threespine stickleback as a model organism*

The threespine stickleback (*Gasterosteus aculeatus*) has a long history as a model species evolutionary biology. It has traditionally been the pet organism for behavioural ecology, to the extent that it has earned the nickname “white rat of ethology” (Huntingford 2003). Recently, threespine stickleback has become a prominent model in evolutionary genetic research, to the extent that the whole genome of the species has been sequenced.

The threespine stickleback is a small teleost abundant in marine, coastal and freshwater habitats throughout the Northern hemisphere. It is considered to form a species complex with a number of divergent populations (Bell & Foster 1994). The common name of threespine stickleback derives from three dorsal spines, of which the most posterior one is the shortest. There are also two robust pelvic spines that are supported by pelvic skeleton. The sides

of the threespine stickleback are covered to varying extent by bony lateral plates that lie on top of myomeres (body segments). Bony armour is perhaps the most obvious character that varies among threespine stickleback populations across different habitats (Bell & Foster 1994). Bony armour is also the feature from which the species draws its scientific name *Gasterosteus* (*gaster* referring to stomach in Latin and *osseus* meaning bones or bone-like in Greek).

Threespine stickleback evolution is often characterised by divergence in bony armour. The reason being that the reduction in bony armour offers an astonishing example of parallel evolution. The reduction has been observed in numerous independent instances in different parts of the distribution range of the species (reviewed in Bell & Foster 1994). The main trend in distribution of the different lateral plate morphs is that ancestral marine morphs are fully plated, while reduction in the number of lateral plates is evident in landlocked freshwater populations (Bell & Foster 1994, Bell 2001). Reduction of the lateral plates can occur very rapidly on an evolutionary time scale, as has been recently reported by Bell *et al.* (2004). They demonstrated a reduction from fully plated to low plated morph in less than twenty years after colonisation of a freshwater lake. Pelvic girdle reduction is more rare than reduction in the lateral plate number – the complete loss of the pelvic girdle has been reported only in a few populations from recently deglaciated regions (Bell 1987). In evolutionary time scale however, the complete reduction of the pelvic structure in less than 10 000 years is fairly rapid (Shapiro *et al.* 2004). Evolution of other morphological aspects of threespine sticklebacks, such as body shape follows the patterns of bony armour

evolution. Colonisation of freshwater habitats by the ancestral marine threespine sticklebacks has led to repeated patterns of body shape divergence throughout the distribution range of the species (e.g. Walker 1997, Walker & Bell 2000, Kristjansson et al. 2002, Spoljaric & Reimchen 2007, Aguirre et al. 2008, Sharpe et al. 2008, Aguirre 2009).

The advances in the genetic tools available for threespine stickleback research together with the well-established knowledge of basic biology and ecology have led to threespine sticklebacks becoming one of the leading model species for the study of genetic basis of adaptive divergence. Quantitative trait loci as well as individual genes that control divergence in quantitative traits have already been discovered (Peichel et al. 2001, Shapiro et al. 2004, Colosimo et al. 2005, Kimmel et al. 2005, Albert et al. 2008), and changes in the frequencies of individual alleles have been shown to be related to the colonisation of freshwaters by the marine ancestors (Barrett et al. 2008). The advent of increasingly sophisticated methods, such as microarrays for gene expression, is likely to result in an unprecedented increase in the resolution of studies targeting the genetic basis of adaptive divergence of threespine sticklebacks and further our understanding of evolutionary mechanisms to a completely new level.

### **Aims of this thesis**

The main aim of this thesis was to investigate the relative roles of genetic drift, natural selection and genetic constraints in the evolution of complex traits. More specifically, the aim was to find out how the three factors have contributed to the body shape and armour divergence of threespine sticklebacks.

The first aim of this thesis was to quantify the extent and patterns of morphological divergence in Fennoscandian threespine stickleback populations. The next aim was to find out whether the observed patterns of divergence were the result of natural selection or could be explained by genetic drift alone (I), and whether this result conformed with the general trends found in already published studies on the relationship between drift and selection (II). Additionally, since the estimates of quantitative trait divergence ( $Q_{ST}$ ) in the first chapter were based on phenotypic data alone ( $P_{ST}$ ), an additional aim was to explore whether there is a general trend in the published studies that the usage of  $P_{ST}$  as a proxy for  $Q_{ST}$  would result in overestimation of quantitative trait divergence.

Once the driving forces behind the morphological divergence were established, the aim shifted to uncovering the genetic basis of divergence. The aim of Chapter III was to investigate the genetic variation and covariation in body shape and armour traits, and based on these patterns deduce what would be the most likely directions of evolution in these traits. Constraints can have an effect on divergence at the population level, but also within populations in the form of genomic conflict between sexes. The aim of Chapter IV was to find out whether there are constraints imposed by the genomic conflict between the sexes in the ancestral population, and whether these possible constraints are reflected in the patterns of divergence in the derived populations. The aim of Chapter V was to go down one level, to variation within individuals, and to investigate the possibility that a trait reflecting developmental stability could express additive genetic variance. Once the relative roles of genetic drift, natural selection and G were established, it remained to test

whether the observed patterns of divergence could be experimentally recreated (VI), using the knowledge on the mechanisms behind divergence (I, III). The aim of the final chapter was also to find out in detail how genetically correlated traits are affected by selection by a predator.

## Main results and discussion

Specific study questions and the main results of all the chapters are summarised in Table 1. Below I discuss these results and their relevance to the study of evolutionary biology of threespine sticklebacks, and evolution in general.

### *Relative roles of drift and selection in population divergence*

Prior to the present work (I), and the pioneering phylogenetic study by Mäkinen et al. (2006), the extent and patterns of threespine stickleback divergence in Fennoscandia were largely unknown. The results of Chapter I show that body shape divergence among Fennoscandian threespine stickleback populations largely follow the patterns of body shape divergence observed in other parts of the distribution range of the species (e.g. Walker 1997, Walker & Bell 2000, Spoljaric & Reimchen 2007, Aguirre et al. 2008, Aguirre 2009). The largest differentiation was found between threespine sticklebacks from marine and freshwater habitats, but body shape also differed among the lake populations depending on the size of the lake (I). Among the lake populations, the patterns of differentiation in neutral genetic markers were similar to those in body shape differentiation, suggesting that the extent of body shape divergence of Fennoscandian freshwater threespine sticklebacks is what could be expected under drift alone. However, overall and

between habitats, phenotypic differentiation ( $P_{ST}$ ) exceeded the differentiation in neutral markers ( $F_{ST}$ ; Fig. 2), suggesting that directional natural selection has been the main factor influencing divergence among the sea populations, as well as across habitats (Fig. 2). Among the sea populations, despite virtually nonexistent neutral genetic divergence ( $F_{ST}$ ), clear phenotypic divergence was present ( $P_{ST}$ ; Fig. 2), providing evidence that adaptive divergence can take place even in the presence of significant levels of gene flow (Hendry et al. 2002, Hendry & Taylor 2004, Hemmer-Hansen et al. 2007).


Estimation of quantitative trait differentiation using only phenotypic data might result in an overestimation of population divergence due to inclusion of environmental and non-additive genetic effects in the  $P_{ST}$  estimates (e.g. Merilä & Crnokrak 2001, Lee & Frost 2002). In our case however, gross overestimation is unlikely for a number of reasons. First, the conclusions did not change even when assuming that half of the phenotypic variation was additive ( $h^2 = 0.5$ ; Fig. 2). Second, the aspects of body shape showing the most divergence have been demonstrated to have genetic basis (McPhail 1984, Albert et al. 2008, III). Finally, according to the meta-analysis of the published comparisons of quantitative trait and neutral genetic divergence, studies based on data from the wild do not tend to yield higher estimates than  $Q_{ST}$  studies based on common garden data (II). In fact, the meta-analysis also revealed that there were no significant differences in the  $Q_{ST}$  estimates for different trait types (*viz.* morphological, life history or behavioural), contrary to what could have been expected based on how traits with different genetic basis (*i.e.* lower  $h^2$  in traits more closely related to fitness; Mousseau & Roff 1987,



Table 1. Summary of the main study questions and results of the six chapters included in this thesis.

Main study questions	Main results
<p><b>I</b> What is the extent of body shape divergence in the Fennoscandian threespine stickleback populations?</p> <p>Can the observed patterns of divergence be explained by genetic drift alone, or are they the result of natural selection?</p>	<p>The patterns of body shape divergence follow those found elsewhere in the distribution area of threespine sticklebacks; largest differentiation occurring between marine and freshwater populations.</p> <p>Natural selection has been a predominant force behind the observed patterns of body shape divergence, although genetic drift may have contributed to the divergence, especially in the small lake populations.</p>
<p><b>II</b> Does natural selection have a predominant role over genetic drift in the general patterns of quantitative trait divergence?</p> <p>Does divergence in neutral markers reflect divergence in quantitative traits?</p>	<p>Yes. Natural selection has a predominant role as a cause of differentiation in quantitative traits.</p> <p>No. Conclusions on quantitative trait divergence should not be made based solely on divergence in neutral genetic markers. This is especially important in cases in which neutral genetic divergence is low</p>
<p><b>III</b> What is the direction of maximum additive genetic variance (<math>g_{max}</math>) in threespine body shape?</p> <p>Are the directions of additive genetic and phenotypic variance correlated?</p> <p>Are genetic correlations imposing possible constraints on body shape and armour evolution?</p>	<p>Direction of <math>g_{max}</math> reflects the divergence observed in wild populations.</p> <p>Yes. Directions of maximum genetic and phenotypic variance are correlated.</p> <p>Body shape is genetically correlated with armour traits, which should be taken into account when identifying selective factors responsible for evolution of single traits</p>
<p><b>IV</b> Is there a genomic conflict between male and female threespine sticklebacks and is it imposing possible constraints on the evolution of sexual dimorphism?</p> <p>Are the possible constraints reflected in the patterns of sexual dimorphism observed in wild populations?</p>	<p>There is a genomic conflict between males and females, as inferred from the high genetic correlations in body shape and armour traits between the sexes.</p> <p>The effect of genomic conflict is not reflected in the patterns of sexual dimorphism in derived threespine stickleback populations.</p>
<p><b>V</b> Is there a heritable component to the variance in fluctuating asymmetry of lateral plates – a trait reflecting developmental stability?</p>	<p>Yes. This is the first study to find substantial amounts of additive genetic variance in fluctuating asymmetry with appropriate methods and experimental design.</p>
<p><b>VI</b> What are the targets and patterns of predator induced selection in threespine stickleback morphology?</p> <p>Can different anti-predator tactics in pelagic and benthic habitats explain the reduction in the number of lateral plates?</p>	<p>Body size and shape, specifically the length of the caudal peduncle, are both affected by selection by predators.</p> <p>Differing anti-predator tactics in pelagic and benthic habitats can be responsible for the lateral plate number reduction during freshwater colonisation of threespine sticklebacks.</p>

Merilä & Sheldon 1999), and the results of earlier meta-analysis of studies comparing  $Q_{ST}$  and  $F_{ST}$  (Merilä & Crnokrak 2001). The general patterns in studies comparing  $Q_{ST}$  and  $F_{ST}$  estimates suggest that natural selection plays a predominant role in quantitative trait differentiation. It must be noted though that this is the case for those systems that are likely to be known *a priori* to exhibit pronounced phenotypic divergence – they are possibly not a representative sample of systems in respect to degree of population divergence.




**Figure 2.**  $P_{ST}$  values for body shape (the first two principal components) assuming two different heritability ( $h^2$ ) values along with  $F_{ST}$  values. All values are with 95 % Bayesian confidence intervals. Variance in shape is adjusted for variance in sex and body size.

There has been some controversy around the issue whether neutral and quantitative genetic estimates are correlated or not (Merilä & Crnokrak, 2001; Crnokrak & Merilä, 2002; Latta & McKay, 2002; McKay & Latta, 2002). In theory, one index of divergence should not reflect the other (Pearman 2001, McKay & Latta 2002), but certain demographic scenarios, such as habitat fragmentation and reduced gene flow, might influence both  $Q_{ST}$  and  $F_{ST}$  in a similar fashion (Frankham & Weber 2000). This might be one explanation for the lack of difference in  $P_{ST}$  and  $F_{ST}$  estimates among Fennoscandian freshwater

populations of threespine sticklebacks (I). The meta-analysis revealed a positive correlation between  $Q_{ST}$  and  $F_{ST}$  across studies, although with small  $F_{ST}$  values,  $Q_{ST}$  values were highly variable (II). This implies that extreme caution should be exercised before making any conclusions on quantitative genetic differentiation based solely on neutral marker differentiation (e.g. Hendry 2002).

#### *Within population variance and population divergence*

Genetic variation and covariation have a fundamental role in evolutionary change. Without additive genetic variation, there will be no evolution, while covariation between traits means that change in one trait could induce a change in the other. The possible constraining role of the patterns of covariation between traits has led to the prediction that evolution is most likely to take place along the line of least resistance, i.e. to the direction of maximum additive genetic variance ( $\mathbf{g}_{max}$ ; Schluter 1996). It has been fairly well established that threespine stickleback morphological divergence follows similar patterns all over the distribution range of the species (for armour traits: see reviews by Bell & Foster 1994, Bell 2001, Östlund-Nilsson et al. 2007; for body shape: e.g. Walker 1997, Walker & Bell 2000, Kristjansson et al. 2002, Spoljaric & Reimchen 2007, Sharpe et al. 2008, Aguirre et al. 2008, Aguirre 2009, I). If Schluter's (1996) predictions on the evolution along  $\mathbf{g}_{max}$  are correct, the divergence along  $\mathbf{g}_{max}$  of an ancestral population of threespine sticklebacks should be similar to that observed in the wild populations. Chapter III shows that this is the case for threespine stickleback body shape. Body shape divergence in the wild (Walker 1997, Walker & Bell 2000, Kristjansson et al. 2002, Spoljaric &



**Figure 3.** Illustration of body shape divergence along the major axis of genetic variation.

Reimchen 2007, Sharpe et al. 2008, Aguirre et al. 2008, Aguirre 2009, I) and along the  $\mathbf{g}_{\max}$  of the putative ancestral population follow the same patterns; at one end there are threespine sticklebacks with deep bodies, short caudal peduncles, large heads and posterior positioned dorsal spines, pelvis and median fins – typical deep-bodied sticklebacks, while at the other end there are threespine sticklebacks with narrow bodies, long caudal peduncles, small heads and anterior positioned dorsal spines, pelvis and median fins – typical streamlined sticklebacks (Fig. 3).

Directions of maximum additive genetic and phenotypic variance were also correlated, which implies that phenotypic divergence is a fair surrogate for genetic divergence in threespine stickleback body shape (III). This is concordant with results of studies on other taxa and traits (Cheverud 1988, 1996, Roff 1995, Reusch & Blanckenhorn 1998, Waitt & Levin 1998, Reale & Festa-Bianchet 2000, House & Simmons 2005, Åkesson et al. 2007, de Oliveira et al. 2009). The results of Chapter III also show that there are strong negative as well as strong positive genetic correlations between body shape and armour traits. Therefore, disentangling the selective agents responsible for the evolution of any single trait in threespine sticklebacks might be difficult – what may appear to be a direct response to selection on a particular trait (e.g. length of the pelvic girdle) might in fact be a correlated

response to selection on another trait.

Although evolution of threespine stickleback body shape appears to take course along the path dictated by genetic covariances and correlations, this was not the case with sexual dimorphism (IV). Genetic correlations between the sexes ( $r_{g(MF)}$ ) were high and the lines of least resistance to evolution ( $\mathbf{g}_{\max}$ ) had similar directions in males and females, but there was no association between  $r_{g(MF)}$  of the ancestral population and the degree of sexual dimorphism in the derived wild populations. Majority of evidence from published studies points to the direction that  $r_{g(MF)}$  and sexual dimorphism are negatively correlated (Poissant et al. 2010), but the results of Chapter IV add to the few exceptions where no such association has been found (Cowley et al. 1986, Cowley & Atchley 1988, Ashman & Majetic 2006). The lack of association between  $r_{g(MF)}$  and sexual dimorphism also indicates that in threespine sticklebacks, sex-specific selection pressures are strong enough to overcome the genetic constraints, and/or that the sex-limited gene expression can be regulated with relative ease.


Genomic conflict between the sexes is one possible factor that maintains genetic variation despite of directional selection, which theoretically depletes genetic variation. Because of the genetic correlations between male and female traits, it is more difficult for the traits to reach their optimum in terms of fitness. This is also one possible way

in which traits indicative of developmental stability, such as fluctuating asymmetry, can have an additive genetic component of variance. Traditionally, heritability estimates for fluctuating asymmetry have been very low (Leamy 1997, Fuller & Houle 2003), but the estimates have usually suffered from difficulties affecting their accuracy (e.g. Merilä & Björklund 1995, Whitlock 1996, Björklund & Merilä 1997, van Dongen 2006). Chapter V is the first study that has found a substantial amount of additive genetic variance in fluctuating asymmetry ( $h^2 = 0.24$ ) using appropriate methods and experimental design. Selection by predators has been found to favour symmetric threespine sticklebacks (Moodie & Reimchen 1976, Bergstrom & Reimchen 2003), implying that selection on fluctuating asymmetry of lateral plates may have important evolutionary consequences.

#### *The role of selection in adaptive divergence*

With the predominant role of selection in morphological divergence of threespine sticklebacks between marine and freshwater habitats established (I), as well as the possible constraints imposed by the genetic variance-covariance matrix, **G** (III), the final component missing from the breeder's equation was the actual effect of selection on the divergence. Although a number of hypotheses on the effects of abiotic and biotic factors have been advanced to explain the divergence in armour (reviewed by Bell 1995, Reimchen 2000, Bell 2001; see also Marchinko & Schluter 2007, Barrett et al. 2008), as well as in body shape (e.g. Walker 1997, Walker & Bell 2000, Spoljaric & Reimchen 2007, Aguirre et al. 2008, Aguirre 2009), the effect of predation is still included in most of the hypotheses (for armour see

reviews by Reimchen 1994, Bell 2001). The positive effect of predation on presence of lateral plates has been experimentally shown (Moodie et al. 1973, Bell & Haglund 1978, Reimchen 1992, 2000, Vamosi 2002), but the effects of predation on the reduction of plate number has not been experimentally tested, although it has been hypothesised that different antipredation strategies in different habitats lie behind the reduction in lateral plates (Reimchen 1992). Likewise, the role of predation in body shape differentiation of threespine sticklebacks has not been experimentally tested.



**Figure 4. Survival probability (relative fitness) of threespine sticklebacks as a function of number of lateral plates (corrected for body size) in the experiments with (black dots and line) and without (grey dots and line) refuge. Dotted lines denote the 95% confidence intervals.**

Chapter VI provides direct experimental support for Reimchen's (1992) hypothesis that different antipredatory tactics in pelagic and benthic habitats could be responsible for the lateral plate number reduction during freshwater colonisation in threespine sticklebacks. In habitats with refuge, selection favours individuals with low number of lateral plates, while fully plated individuals have a selective advantage in habitats without shelter (Fig. 4). Differences in

body shape also have an effect on survival in the presence of predators (VI). Swimming performance related aspects of body shape, such as the length of the caudal peduncle and body depth (Webb 1982, 1984) appear to be the most important predictors of survival from predators. Taking into account the genetic correlations between body shape and armour traits revealed that spine lengths also have an effect on survival, as expected (Hoogland et al. 1957, Vamosi 2002). It is notable that this effect was not evident when analysing the effects of different traits individually. Therefore, it is imperative to consider genetic correlations between traits when drawing conclusions on the effects of selection on single traits (Lande & Arnold 1983).

### Conclusions and future directions

I have demonstrated how evolution of complex traits, investigated at different levels of resolution can produce a holistic picture of the evolutionary processes. By following the theoretical framework for multivariate trait evolution ( $\Delta\mathbf{z} = \mathbf{GP}^{-1}\mathbf{s}$ ; Lande 1979, Lande & Arnold 1983), I have shown how evolutionary mechanisms can influence adaptive divergence from the level of individuals to populations. Body shape divergence of threespine sticklebacks between populations from marine and freshwater habitats is clearly driven by natural selection, while genetic drift may play significant role in more fine scale divergence, between populations of isolated freshwater habitats. Body shape divergence along the most likely direction of evolution ( $\mathbf{g}_{\max}$ ; Schluter 1996) in the ancestral marine threespine sticklebacks is similar to that observed in wild populations (e.g. Walker & Bell 2000, Spoljaric & Reimchen 2007, I), indicating that

genetic variances and covariances ( $\mathbf{G}$ ) can significantly bias the patterns of evolution.

Genetic conflict, as imposed by high genetic correlations between males and females, appears to be resolved in threespine sticklebacks. Ancestral genetic correlations are not reflected in the patterns of sexual dimorphism in the derived wild populations. The patterns of ancestral sexual dimorphism in body shape reflect the patterns of body shape differentiation observed in threespine stickleback marine-freshwater transition (e.g. Walker & Bell 2000, Spoljaric & Reimchen 2007, I), indicating that ancestral sexual dimorphism could have played a significant role in the adaptive divergence of threespine sticklebacks. Variation in traits within individuals can also diverge and help maintaining genetic variation in a population under directional selection. The significant additive genetic variance component of fluctuating asymmetry provides a demonstration of how phenotypes that are assumed to be outside their fitness optimum can be inherited, and thus, have an influence on evolution.

Finally, I have shown that the patterns of evolution seen in nature can be experimentally recreated to tease out the effects of natural selection on adaptive divergence in detail. I have also provided a demonstration of how important it is to take into account the correlative nature of traits, when interpreting the effects of natural selection. The effects of selection on some traits might be hidden because of selection on genetically correlated traits. Threespine stickleback evolution is characterised by rapid reduction in bony armour, and I have shown how different antipredatory strategies in pelagic and benthic habitats could be responsible for the lateral plate number reduction during freshwater

colonisation. Definitive tests of the effect of predation on the reduction of lateral plates have been thus far lacking, although the effect of marine to freshwater transition on individual genes that affect the number of lateral plates have recently been demonstrated (Barrett et al. 2008). The future challenge lies in trying to establish a basis for adaptive divergence of complex traits, such as body shape, at the level of genes. It is likely though, that instead of changes in genes themselves, changes in gene regulation could play an important role in adaptive divergence of complex traits. For example, changes in gene regulation are more readily achieved than actual changes in the genes themselves in cichlid fishes (Roberts et al. 2009). The recent methodological advances in molecular genetics bring promise on opportunities to take the study of adaptive divergence into a completely new level, especially in threespine sticklebacks with completely sequenced genome.

## Acknowledgements

First of all, I would like to thank my supervisors for giving me the opportunity to do my PhD in EGRU. I am extremely grateful to Cano, who in the beginning believed in me and pushed me to start the project. Thank you for always being available, and at the same time giving me space to do things my way (read: in the last minute or slightly past that). I cannot think of a single occasion, when you would not have answered my emails asking for advice or comments within a few hours. With my way of doing everything usually quite close to the deadlines, your quick responsiveness was really essential for me.

The same really applies to my other supervisor, Juha Merilä, the mastermind behind it all. I am especially grateful for the patience you have showed along the way. I really needed all that time to get the

twists of quantitative genetics into my head.

Although I have only had two official supervisors, I feel that I have also been supervised by a number of other people – an advantage of being part of a large research group. I really owe a big thank you especially to Jussi Alho and Gábor Herczeg. Without Jussi I would still be struggling with the analyses. In addition to writing the R code for half of the chapters, Jussi has always been ready to discuss my work, and go out of his way to help anyone in need. Likewise, Gábor has also been always ready to discuss my work and solve problems, whether theoretical or practical. I am also grateful for your company in the basement, greenhouse, gym, as well as on the field trips to civilised and not so civilised places.

I would also like to thank Hannu Mäkinen for great times in the field, especially in Kevo and Vättern, as well as in the office. In addition to providing me with actual molecular data, Hannu demonstrated with his own example how to survive the stressful phases. In the latter stages of Hannu's PhD project and early days of mine, our office was lucky to have the kindest person on earth, Cim Matsuba. I am sure our whole group is grateful for having had Cim in our office, always ready to discuss anything and help anyone. Cim also contributed enormously to the fish breeding, and made it possible for me to have some time off, by taking care of the fish during many of the weekends, as well as *kurisumasu*.

Another person to whom I owe a special thank you for help with practical issues is Simo Rintakoski, who stained, photographed and measured thousands of fish in addition to being a real handyman in and outside the lab. I owe a big thank you also to Linda Uoti and Ina Naukkarinen for their help with marking hundreds of live fish for the selection experiment.

To be honest, I have probably spent majority of my time in the office asking



questions and bugging people with my problems. The list of people to thank for being there to help me either with theory or practice is really long, but especially Phillip Gienapp, Celine Teplitsky and Bob O'Hara deserve a big thanks for their advice on quantitative genetics as well as statistics. Also, without the effort from John Loehr, the chapter on FA would have never seen daylight. Thank you John also for bringing some life to the group meetings; reminiscing about your stunned face following the silence in your first group meeting still makes me smile. Marika Karjalainen sexed a couple thousand fish for me, for which I am extremely grateful. Erica Leder and Jackie DeFaveri also deserve big thanks for their efforts in sexing hundreds of fish, and Theresa Knopp for helping with collecting the data for the meta-analysis, sharing lifts to Porvoo, and introducing to me the night-life of Tolkkinen. The whole range of EGRU post-docs and students, past and present have offered vital support on which to lean at times of getting stuck.

Perhaps the person, who has suffered the most from my bugging, is Vilppu, who luckily has not found a way yet to escape from the office when (daily or actually hourly) I have had something to moan about. I sincerely thank you for being there, and also for bringing light and some sense of humour into the office. Your company has been vital for keeping myself relatively sane during this project.

The company of the rest of the PhD students and 'other hang-arounds' at the department has also been essential for keeping the stress at a tolerable level. What would be a better way to relieve stress and let out pressure than together complain about work, research group, university administration etc. I want to especially thank Paula, Minttu, Marja, Abi, as well as Kim, Aksu, Jenni, Johan and the rest of the lunch group for your company.

The other stickleback people at the department; Jan, Leon, Ulla and Ulrika also deserve a thank you for accompanying

me to the stickleback meetings, as do Laura, Anti, Paula again, and the rest of the people from Craig Primmer's group in Turku for the good times at the always-so-fun CoE meetings.

A number of people have provided stickleback samples for my work. Thank you Teija Aho, Per-Arne Amundsen, Anders Berglund, Susie Coyle, Piotr Debowski, Jörg Freyhof, Ari Haikonen, Dánjúl Petur Højgaard, Vladimír Kovács, Dmitry Lajus, Arne Levsen, Volker Loeschcke, Pat Monaghan, Arne Nolte, Lennart Persson, Thorsten Reusch, Ari Saura, Per Sjöstrand, Audrius Steponenas, Aleksei Veselov, and Linda Zaveik for providing the study material. I also want to thank Petri Savola and Toivo Mauria for their invaluable assistance in catching pikes. A special thank you also to Briitta Saarela for hospitality during the collection of sticklebacks at Pyhäjoki.

I owe a great debt of gratitude also to the people in the global scientific community who have provided advice in the numerous statistical as well as practical issues. I am especially indebted to Arthur Gilmour, Patrick Phillips, Dean Adams, and Ashley Ward for their help with different softwares and methodological issues.

For helping with the formalities, I would like to thank Maarit Haataja as well as Ilkka Teräs and Veijo Kaitala. I am also grateful for Anni Tonteri for always being ready and happy to help.

I would like to extend my thanks to my friends outside the department, who have been really helpful in taking my mind off work and the studies. I am truly grateful especially to Sampo, Jenni, Jeppe, Janne and Henri, for good times, and for keeping me company while I have been paskiainen joka paikassa.

This work has been financially supported by the LUOVA graduate school, Academy of Finland, and travel grants from the Chancellor of the University of Helsinki.

Finally, and most importantly I want to thank my family for their unconditional love and support. Mom and dad, I really appreciate that you have settled for my extremely short answers on my study subject. I am glad I have always had a sanctuary away from the world of science where to retreat (and where to regress). Thank you also to my sister Aino and brother Simo with his family Emma and Iida for always being there for me when needed. I am also grateful to the Jouppila family for their hospitality during our visits. The weekends and holidays in the countryside have been a true lifeline for me.

I owe the biggest thanks to Sanna and Ilmari for constantly reminding me of the things that really matter. Ilmari, after you came into our life, it has been impossible for me to stress too much about anything; your happiness and laughter make all the worries disappear. It just does not seem possible for you to have a bad day, in which you really take after your mother. Sanna, my love, you have been so kind and understanding with my work that there just are no words kind enough to thank you. I am forever grateful for your love and support during the time when I have been busy with the thesis. You have always remained positive, even when I have tried to sink into despair. Actually, I must also thank the sticklebacks for colonising northern Finland, because that brought me to Kevo to find you.

Thank you Ginza for keeping my feet warm while I have been sitting at my desk and for daily dragging me away from the desk to enjoy the beautiful outdoors.

### Literature cited

Adams, D.C. Rohlf, F.J. & Slice, D.C. 2004. Geometric morphometrics: ten years of progress following the 'revolution'. *Ital. J. Zool.* **71**: 5-16.

- Aguirre, W.E. 2009. Microgeographical diversification of threespine stickleback: Body shape-habitat correlations in a small, ecologically diverse Alaskan drainage. *Biol. J. Linn. Soc.* **98**: 139-151.
- Aguirre, W., E. E. Kaitlyn, M. Kusenda, M. A. Bell. 2008. Phenotypic variation and sexual dimorphism in anadromous threespine stickleback: implications for postglacial adaptive radiation. *Biol. J. Linn. Soc.* **95**: 465-478.
- Åkesson, M., Bensch, S. & Hasselquist, D. 2007. Genetic and phenotypic associations in morphological traits: a long term study of great reed warblers *Acrocephalus arundinaceus*. *J. Avian Biol.* **38**: 58-72.
- Albert, A.Y.K., Sawaya, S., Vines, T.H., Knecht, A.K., Miller, C.T., Summers, B.R., Balabhadra, S., Kingsley, D.M. & Schluter, D. 2008. The genetics of adaptive shape shift in stickleback: Pleiotropy and effect size. *Evolution* **62**: 76-85.
- Araripe, L.O., Yassin, A., Klaczko, L.B., Moreteau, B. & David, J.R. 2008. Divergent abdominal bristle patterns in two distantly related drosophilids: antero-posterior variations and sexual dimorphism in a modular trait. *Genetica* **134**: 211-222.
- Arnold, S. J. 1992. Constraints on phenotypic evolution. *Am. Nat.* **140**: S85-S107.
- Arnold, S.J. & Phillips, P.C. 1999. Hierarchical comparison of genetic variance-covariance matrices. II. coastal-inland divergence in the garter snake, *Thamnophis elegans*. *Evolution* **53**: 1516-1527.
- Ashman T. L. 2003. Constraints on the evolution of males and sexual dimorphism: Field estimates of genetic architecture of reproductive traits in three populations of gynodioecious *Fragaria virginiana*. *Evolution* **57**: 2012-2025.
- Ashman T. L., and C. J. Majetic. 2006. Genetic constraints on floral evolution: a review and evaluation of patterns. *Heredity* **96**: 343-352.
- Barrett, R.D.H., Rogers, S.M. & Schluter, D. 2008. Natural selection on a major armor gene in threespine stickleback. *Science* **322**: 255-257.



- Barton, N.H. & Turelli, M. 1989. Evolutionary quantitative genetics: how little do we know? *Annu. Rev. Genet.* **23**: 337-370.
- Bell, M.A. 1987. Interacting evolutionary constraints in pelvic reduction of threespine sticklebacks, *Gasterosteus aculeatus* (Pisces, Gasterosteidae). *Biol. J. Linn. Soc.* **31**: 347-382.
- Bell, M.A. 1995. Intraspecific systematics of *Gasterosteus aculeatus* populations: Implications for behavioral ecology. *Behaviour* **132**: 1131-1152.
- Bell M. A. 2001. Lateral plate evolution in the threespine stickleback: getting nowhere fast. *Genetica* **112**: 445-461.
- Bell, M.A. & Haglund, T.R. 1978. Selective predation of threespine sticklebacks (*Gasterosteus aculeatus*) by garter snakes. *Evolution* **32**: 304-319.
- Bell, M. A. & Foster, S. A. eds. 1994. *The Evolutionary Biology of Threespine Stickleback*. Oxford University Press. Oxford. 571 pp.
- Bell, M.A. Aguirre, W.E. & Buck, N.J. 2004. Twelve years of contemporary armor evolution in a threespine stickleback population. *Evolution* **58**: 814-824.
- Bergstrom, C.A. & Reimchen, T.E. 2003. Asymmetry in structural defenses: Insights into selective predation in the wild. *Evolution* **57**: 2128-2138.
- Björklund, M. 2004. Constancy of the G matrix in ecological time. *Evolution* **58**: 1157-1164.
- Björklund, M. & Merilä, J. 1997. Why some measurements of fluctuating asymmetry are sensitive to measurement error. *Ann. Zool. Fennici* **34**:133-137.
- Blows, M.W. & Higgie, M. 2003. Genetic constraints on the evolution of mate recognition under natural selection. *Am. Nat.* **161**: 240-253.
- Bolnick, D.I. & Doebeli, M. 2003. Sexual dimorphism and adaptive speciation: Two sides of the same ecological coin. *Evolution* **57**: 2433-2449.
- Bonduriansky R., & Rowe, L. 2005. Intralocus sexual conflict and the genetic architecture of sexually dimorphic traits in *Prochyliza xanthostoma* (Diptera: Piophilidae). *Evolution* **59**: 1965-1975.
- Bonner, J.T. 1982. Evolutionary strategies and developmental constraints in the cellular slime-molds. *Am. Nat.* **119**: 530-552.
- Bookstein, F.L. 1991. *Morphometric Tools for Landmark Data: Geometry and Biology*. Cambridge University Press. Cambridge. 455 pp.
- Butler, M.A., Sawyer, S.A. & Losos, J.B. 2007. Sexual dimorphism and adaptive radiation in *Anolis* lizards. *Nature* **447**: 202-205.
- Buvanendran,V. 1969. Heritability and genetic correlations of sexual dimorphism for 10-week weight in poultry. *Br. Poult. Sci.* **10**: 321-325.
- Chakraborty, R. & Nei, M. 1982. Genetic differentiation of quantitative characters between populations or species .1. mutation and random genetic drift. *Genet. Res.* **39**: 303-314.
- Chapuis H., TixierBoichard, M., Delabrosse, Y. & Ducrocq, V. 1996. Multivariate restricted maximum likelihood estimation of genetic parameters for production traits in three selected turkey strains. *Genet. Sel. Evol.* **28**: 299-317.
- Charmantier, A. & Garant, D. 2005. Environmental quality and evolutionary potential: Lessons from wild populations. *Proc.R. Soc. Lond. B.* **272**: 1415-1425.
- Cheverud, J. M. 1988. A comparison of genetic and phenotypic correlations. *Evolution* **42**:958-968.
- Cheverud, J. M. 1996. Quantitative genetic analysis of cranial morphology in the cotton-top (*Saguinus oedipus*) and saddle-back (*S. fuscicollis*) tamarins. *J. Evol. Biol.* **9**: 5-42.
- Colosimo, P.F., Hosemann, K.E., Balabhadra, S., Villarreal, G., Dickson, M., Grimwood, J., Schmutz, J., Myers, R.M., Schluter, D. & Kingsley, D.M. 2005. Widespread parallel evolution in sticklebacks by repeated fixation of ectodysplasin alleles. *Science* **307**: 1928-1933.
- Cowley, D. E., Atchley, W. R. & Rutledge, J. J. 1986. Quantitative Genetics of *Drosophila melanogaster* .1. Sexual Dimorphism in Genetic-Parameters for Wing Traits. *Genetics* **114**: 549-566.

- Cowley, D. E., & Atchley, W. R. 1988. Quantitative genetics of *Drosophila melanogaster*. 2. Heritabilities and genetic correlations between sexes for head and thorax traits. *Genetics* **119**: 421-433.
- Crnokrak, P. & Merilä, J. 2002. Genetic population divergence: markers and traits. *Trends Ecol. Evol.* **17**: 501.
- Darwin, C. 1871. *The Descent of Man, and Selection in Relation to Sex* (1st ed.). London: John Murray. 409 pp.
- David, J.R., Gibert, P., Mignon-Grasteau, S., Legout, H., Petavy, G., Beaumont, C. & Moreteau, B. 2003. Genetic variability of sexual size dimorphism in a natural population of *Drosophila melanogaster*: An isofemale-line approach. *J. Genet.* **82**: 79-88.
- Dayan, T. & Simberloff, D. 1994. Character displacement, sexual dimorphism, and morphological variation among british and irish mustelids. *Ecology* **75**: 1063-1073.
- de Oliveira, F. B., Porto, A. & Marroig, G. 2009. Covariance structure in the skull of *Catarrhini*: a case of pattern stasis and magnitude evolution. *J. Hum. Evol.* **56**: 417-430.
- England, P.R., Osler, G.H.R., Woodworth, L.M., Montgomery, M.E., Briscoe, D.A. & Frankham, R. 2003. Effects of intense versus diffuse population bottlenecks on microsatellite genetic diversity and evolutionary potential. *Conserv. Genet.* **4**: 595-604.
- Estes, S. & Arnold, S.J. 2007. Resolving the paradox of stasis: Models with stabilizing selection explain evolutionary divergence on all timescales. *Am. Nat.* **169**: 227-244.
- Fairbairn, D.J., Blanckenhorn, W.U. & Szekely, T. (eds). 2007. *Sex, Size and Gender Roles. Evolutionary Studies of Sexual Dimorphism*. Oxford University Press, Oxford. 266 pp.
- Felsenstein, J. 1986. Population differences in quantitative characters and gene-frequencies - a comment. *Am. Nat.* **127**: 731-732.
- Fisher, R. A. 1930. The Evolution of Dominance in Certain Polymorphic Species. *Am. Nat.* **64**: 385-406.
- Fisher, R. A. 1931. The Evolution of Dominance. *Biol. Rev. Camb. Philos. Soc.* **6**: 345-368.
- Frankham, R. & Weber, K.E. 2000. Nature of quantitative genetic variation. In: *Evolutionary Genetics: From Molecules to Morphology* (Singh, R.S., & Crimbas, C.B. eds.), pp. 351-368. Cambridge University Press, Cambridge.
- Frankham, R., Ballou, J.D. & Briscoe, D.A. 2002. *Introduction to Conservation Genetics*. Cambridge University Press. Cambridge. 617 pp.
- Fuller, R.C. & Houle, D. 2003. Inheritance of developmental instability. In: *Developmental Instability: Causes and Consequences* (M. Polak, ed), pp. 157-186. Oxford University Press, Oxford.
- Gavrilets, S. 2004. *Fitness Landscapes and the Origin of Species*. Princeton University Press. Monographs in Population Biology 41. 476 pp.
- Hanrahan, J.P. & Eisen, E.J. 1973. Sexual dimorphism and direct and maternal genetic effects on body-weight in mice. *Theor. Appl. Genet.* **43**: 39-45.
- Hedrick, P.W. 1999. Perspective: Highly variable loci and their interpretation in evolution and conservation. *Evolution* **53**: 313-318.
- Hemmer-Hansen, J., Nielsen, E.E., Frydenberg, J. & Loeschcke, V. 2007. Adaptive divergence in a high gene flow environment: Hsc70 variation in the european flounder (*Platichthys flesus* L.). *Heredity* **99**: 592-600.
- Hendry, A.P. Taylor, E.B. & McPhail, J.D. 2002. Adaptive divergence and the balance between selection and gene flow: lake and stream stickleback in the misty system. *Evolution* **56**: 1199-1216.
- Hendry, A.P. & Taylor, E.B. 2004. How much of the variation in adaptive divergence can be explained by gene flow? - An evaluation using lake-stream stickleback pairs. *Evolution* **58**: 2319-2331.
- Hoffmann, A.A. & Merilä, J. 1999. Heritable variation and evolution under favourable and unfavourable conditions. *Trends Ecol. Evol.* **14**: 96-101.

- Hoogland, R., Morris, D., & Tinbergen, N. 1957. The spines of stickleback (*Gasterosteus* and *Pygosteus*) as a means of defence against predators. *Behaviour* **10**: 205-237.
- House, C. M. & Simmons, L. W. 2005. The evolution of male genitalia: patterns of genetic variation and covariation in the genital sclerites of the dung beetle *Onthophagus taurus*. *J. Evol. Biol.* **18**: 1281-1292.
- Hu, Y.H., Poivey, J.P., Rouvier, R., Wang, C.T. & Tai, C. 1999. Heritabilities and genetic correlations of body weights and feather length in growing Muscovy selected in Taiwan. *Br. Poult. Sci.* **40**: 605-612.
- Huntingford, F.A. 2003. A history of animal behaviour by a partial, ignorant and prejudiced ethologist. *Anim. Behav.* **66**: 409-415.
- Jones, L.P., Frankham, R. & Barker, J.S.F. 1968. Effects of population size and selection intensity in selection for a quantitative character in drosophila .2. long-term response to selection. *Genet. Res.* **12**: 249-266.
- Kimmel, C. B., Ullmann, B., Walker, C., Wilson, C., Currey, M., Phillips, P. C., Bell, M. A., Postlethwait, J. H. & Cresko, W. A. 2005. Evolution and development of facial bone morphology in threespine sticklebacks. *Proc. Natl. Acad. Sci. USA* **102**: 5791-5796.
- Kinnison, M.T. & Hendry, A.P. 2001. The pace of modern life II: From rates of contemporary microevolution to pattern and process. *Genetica* **112**: 145-164.
- Klingenberg, C.P. 2003. Quantitative genetics of geometric shape: heritability and the pitfalls of the univariate approach. *Evolution* **57**: 191-195.
- Klingenberg, C.P. & McIntyre, G.S. 1998. Geometric morphometrics of developmental instability: analyzing patterns of fluctuating asymmetry with Procrustes methods. *Evolution* **52**: 1363-1375.
- Klingenberg, C. P., and L. J. Leamy. 2001. Quantitative genetics of geometric shape in the mouse mandible. *Evolution* **55**:2342-2352.
- Klingenberg, C.P. & Monteiro, L.R. 2005. Distances and directions in multidimensional shape spaces: Implications for morphometric applications. *Syst. Biol.* **54**: 678-688.
- Kristjansson, B. K., Skulason, S. & Noakes, D. L. G. 2002. Morphological segregation of Icelandic threespine stickleback (*Gasterosteus aculeatus* L). *Biol. J. Linn. Soc.* **76**: 247-257.
- Lande, R. 1976. Natural-selection and random genetic drift in phenotypic evolution. *Evolution* **30**: 314-334.
- Lande, R. 1977. Statistical tests for natural-selection on quantitative characters. *Evolution* **31**: 442-444.
- Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain - body size allometry. *Evolution* **33**:402-416.
- Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters. *Evolution* **34**: 292-305.
- Lande, R. 1986. The dynamics of peak shifts and the pattern of morphological evolution. *Paleobiology* **12**: 343-354.
- Lande, R. 1992. Neutral theory of quantitative genetic variance in an island model with local extinction and colonization. *Evolution* **46**: 381-389.
- Lande, R. & Arnold, S.J. 1983. The measurement of selection on correlated characters. *Evolution* **37**: 1210-1226.
- Latta, R.G. & McKay, J.K. 2002. Genetic population divergence: markers and traits. *Trends Ecol. Evol.* **17**: 501-502.
- Le Bihan-Duval, E., Mignon-Grasteau, S., Millet, N. & Beaumont, C. 1998. Genetic analysis of a selection experiment on increased body weight and breast muscle weight as well as on limited abdominal fat weight. *Br. Poult. Sci.* **39**: 346-353.
- Leamy, L.J. 1997. Is developmental stability heritable? *J. Evol. Biol.* **10**: 21-29.
- Lee, C.E. & Frost, B.W. 2002. Morphological stasis in the *Eurythemora affinis* species complex (*Copepoda: Temoridae*). *Hydrobiologia* **480**: 111-128.

- Lester, L.J., Lawson, K.S., Abella, T.A. & Palada, M.S. 1989. Estimated heritability of sex ratio and sexual dimorphism in tilapia. *Aquaculture and Fisheries Management* **20**: 369-380.
- Lewontin, R.C. 1984. Detecting population differences in quantitative characters as opposed to gene-frequencies. *Am. Nat.* **123**: 115-124.
- Lynch, M. 1988. The divergence of neutral quantitative characters among partially isolated populations. *Evolution* **42**: 455-466.
- Lynch, M. 1994. Neutral models of phenotypic evolution. In: *Ecological Genetics* (L.A. Real, ed.), pp. 86-108. Princeton University Press. Princeton, NJ.
- Lynch, M. & Hill, W.G. 1986. Phenotypic evolution by neutral mutation. *Evolution* **40**: 915-935.
- Lynch M. & Walsh, J. B. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates. Sunderland, MA. 980 pp.
- Mäkinen, H. S., Cano, J. M. & Merilä, J. 2006. Genetic relationships among marine and freshwater populations of the European three-spined stickleback (*Gasterosteus aculeatus*) revealed by microsatellites. *Mol. Ecol.* **15**: 1519-1534.
- Marchinko, K.B. & Schluter, D. 2007. Parallel evolution by correlated response: Lateral plate reduction in threespine stickleback. *Evolution* **61**: 1084-1090.
- Maynard Smith, J., Burian, R., Kauffman, S., Alberch, P., Campbell, J., Goodwin, B., Lande, R., Raup, D. & Wolpert, L. 1985. Developmental constraints and evolution. *Q. Rev. Biol.* **60**: 265-87.
- McDaniel, S. F. 2005. Genetic correlations do not constrain the evolution of sexual dimorphism in the moss *Ceratodon purpureus*. *Evolution* **59**: 2353-2361.
- McGuigan, K., Chenoweth, S. F. & Blows, M. W. 2005. Phenotypic divergence along lines of genetic variance. *Am. Nat.* **165**: 32-43.
- McKay, J.K. & Latta, R.G. 2002. Adaptive population divergence: Markers, QTL and traits. *Trends Ecol. Evol.* **17**: 285-291.
- McPhail, J.D. 1984. Ecology and evolution of sympatric sticklebacks (*Gasterosteus*): morphological and genetic evidence for a species pair in Enos Lake, British Columbia. *Can. J. Zool.* **62**: 1402-1408.
- Meirmans, P.G. 2006. Using the AMOVA framework to estimate a standardized genetic differentiation measure. *Evolution* **60**: 2399-2402.
- Merilä, J. & Björklund, M. 1995. Fluctuating asymmetry and measurement error. *Syst. Biol.* **44**: 97-101.
- Merilä, J. & Sheldon, B.C. 1999. Genetic architecture of fitness and nonfitness traits: Empirical patterns and development of ideas. *Heredity* **83**: 103-109.
- Merilä, J. & Crnokrak, P. 2001. Comparison of genetic differentiation at marker loci and quantitative traits. *J. Evol. Biol.* **14**: 892-903.
- Mignon-Grasteau, S., Beaumont, C., Poivey, J.P. & de Rochambeau, H. 1998. Estimation of the genetic parameters of sexual dimorphism of body weight in 'label' chickens and Muscovy ducks. *Gen. Select. Evol.* **30**: 481-491.
- Mignon-Grasteau, S., Beaumont, C., Le Bihan-Duval, E., Poivey, J.P. De Rochambeau, H. & Ricard, F.H. 1999. Genetic parameters of growth curve parameters in male and female chickens. *Br. Poult. Sci.* **40**: 44-51.
- Mignon-Grasteau, S., David, J., Gibert, P., Legout, H., Petavy, G., Moreteau, B. & Beaumont, C. 2004. REML estimates of genetic parameters of sexual dimorphism for wing and thorax length in *Drosophila melanogaster*. *J. Genet.* **83**: 163-170.
- Møller, A.P. & Thornhill, R. 1997. A meta-analysis of the heritability of developmental stability. *J. Evol. Biol.* **10**: 1-16.
- Monteiro, L.R., Diniz-Filho, J.A.F., dos Reis, S.F. & Araujo, E.D. 2002. Geometric estimates of heritability of biological shape. *Evolution* **56**: 563-572.
- Moodie, G.E.E., Mcphail, J.D. & Hagen, D.W. 1973. Experimental demonstration of selective predation on *Gasterosteus aculeatus*. *Behaviour* **47**: 95-105.

- Moodie, G. E. E. & Reimchen, T. E. 1976. Glacial refugia, endemism, and stickleback populations of Queen Charlotte Islands, British Columbia. *Can. Field-Nat.* **90**: 471-474.
- Mousseau, T.A. & Roff, D.A. 1987. Natural-selection and the heritability of fitness components. *Heredity* **59**: 181-197.
- Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia Univ. Press, New York. 512 pp.
- O'Hara, R.B. & Merilä, J. 2005. Bias and precision in  $Q_{ST}$  estimates: Problems and some solutions. *Genetics* **171**: 1331-1339.
- Östlund-Nilsson, S., Mayer, I. & Huntingford, F.A. 2007. (eds.) *Biology of the Three-Spined Stickleback*. CRC Press. Boca Raton, FL. 392 pp.
- Pearman, P.B. 2001. Conservation value of independently evolving units: sacred cow or testable hypothesis? *Cons. Biol.* **15**: 780-783
- Peichel, C. L., Nereng, K. S., Ohgi, K. A., Cole, B. L. E., Colosimo, P. F., Buerkle, C. A., Schluter, D. & Kingsley, D. M. 2001. The genetic architecture of divergence between threespine stickleback species. *Nature* **414**: 901-905.
- Phillips, P.C. 1998. Designing experiments to maximize the power of detecting correlations. *Evolution* **52**: 251-255.
- Pigliucci, M. 2006. Genetic variance-covariance matrices: A critique of the evolutionary quantitative genetics research program. *Biol. Philos.* **21**: 1-23.
- Poissant, J. Wilson, A.J. & Coltman, D.W. Sex-specific variance and the evolution of sexual dimorphism: a systematic review of cross-sex genetic correlations. *Evolution* **64**: 97-107.
- Polly, P.D. 2008. Developmental dynamics and **G**-matrices: Can morphometric spaces be used to model phenotypic evolution? *J. Evol. Biol.* **35**: 83-96.
- Price, T. & Schluter, D. 1991. On the low heritability of life-history traits. *Evolution* **45**: 853-861.
- Reale, D. & Festa-Bianchet, M. 2000. Quantitative genetics of life-history traits in a long-lived wild mammal. *Heredity* **85**: 593-603.
- Reeve, J. P. & Fairbairn, D. J. 2001. Predicting the evolution of sexual size dimorphism. *J. Evol. Biol.* **14**: 244-254.
- Reimchen, T.E. 1992. Injuries on stickleback from attacks by a toothed predator (*Oncorhynchus*) and implications for the evolution of lateral plates. *Evolution* **46**: 1224-1230.
- Reimchen, T.E. 1994. Predators and morphological evolution in threespine stickleback. In: Bell, MA, Foster, SA, eds. *The Evolutionary Biology of the Threespine Stickleback*. Pp. 240–276. Oxford. Oxford University Press.
- Reimchen, T.E. 2000. Predator handling failures of lateral plate morphs in *Gasterosteus aculeatus*: Functional implications for the ancestral plate condition. *Behaviour* **137**: 1081-1096.
- Reusch, T. & Blanckenhorn, W. U. 1998. Quantitative genetics of the dung fly *Sepsis cynipsea*: Cheverud's conjecture revisited. *Heredity* **81**: 111-119.
- Reyment, R.A. 2002. Morphometrics. In: El-Shaarawi, A.H. & Piegorsh, W.W. (eds.) *Encyclopedia of Environmetrics*. Vol. 3. Pp. 1301-1307.
- Roberts, R.B., Ser, J.R. & Kocher, T.D. 2009. Sexual conflict resolved by invasion of a novel sex determiner in Lake Malawi cichlid fishes. *Science* **326**: 998-1001.
- Roff, D. A. 1995. The estimation of genetic correlations from phenotypic correlations - a test of Cheverud's conjecture. *Heredity* **74**: 481-490.
- Roff, D.A. 2000. Trade-offs between growth and reproduction: An analysis of the quantitative genetic evidence. *J. Evol. Biol.* **13**: 434-445.
- Roff, D.A. 2007. A centennial celebration for quantitative genetics. *Evolution* **61**: 1017-1032.
- Rogers, A.R. & Harpending, H.C. 1983. Population-structure and quantitative characters. *Genetics* **105**: 985-1002.
- Rohlf, F.J. 1999. Shape statistics: Procrustes superimpositions and tangent spaces. *J. Classif.* **16**: 197-223.
- Rohlf, F.J. & Slice, D.S. 1990. Extensions of the Procrustes method for the optimal superimposition of landmarks. *Syst. Zool.* **39**: 40-59.
- Rohlf, F.J. & Marcus, L.F. 1993. A revolution in morphometrics. *Trends Ecol. Evol.* **8**: 129-132.

- Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. *Evolution* **50**: 1766-1774.
- Schoener, T.W. 1967. Ecological significance of sexual dimorphism in size in lizard *Anolis cospersus*. *Science* **155**: 474-477.
- Schoener, T.W. 1969. Size patterns in West Indian *Anolis* lizards. I. size and species diversity. *Syst. Zool.* **18**: 386-402.
- Shapiro, M.D., Marks, M.E., Peichel, C.L., Blackman, B.K., Nereng, K.S., Jonsson, B., Schluter, D. & Kingsley, D.M. 2004. Genetic and developmental basis of evolutionary pelvic reduction in threespine sticklebacks. *Nature* **428**: 717-723.
- Sharpe, D. M. T., Räsänen, K., Berner, D., & Hendry, A. P. 2008. Genetic and environmental contributions to the morphology of lake and stream stickleback: implications for gene flow and reproductive isolation. *Evol. Ecol. Res.* **10**: 849-866.
- Slice, D.S. 2001. Landmark coordinates aligned by Procrustes analysis do not lie in Kendall's shape space. *Syst. Biol.* **50**: 141-149.
- Slice, D.S., Bookstein, F.L., Marcus, L.F. & Rohlf, F.J. 1996. Appendix I: A glossary for geometric morphometrics. In: Marcus, L.F., Corti, M., Loy, A. Naylor, G.J.P. & Slice, D.E. (eds.), *Advances In Morphometrics*. Plenum Press. New York. 580 pp.
- Spitze, K. 1993. Population structure in *Daphnia obtusa* - quantitative genetic and allozymic variation. *Genetics* **135**: 367-374.
- Spoljaric, M. A., & Reimchen, T. E. 2007. 10 000 years later: evolution of body shape in Haida Gwaii three-spined stickleback. *J. Fish Biol.* **70**: 1484-1503.
- Steppan, S.J., Phillips, P.C. & Houle, D. 2002. Comparative quantitative genetics: Evolution of the G matrix. *Trends Ecol. Evol.* **17**: 320-327.
- Turelli, M. 1988. Phenotypic evolution, constant covariances, and the maintenance of additive variance. *Evolution* **42**: 1342-1347.
- Vamosi, S.M. 2002. Predation sharpens the adaptive peaks: Survival trade-offs in sympatric sticklebacks. *Ann. Zool. Fennici* **39**: 237-248.
- Van Dongen, S. 2006. Fluctuating asymmetry and developmental instability in evolutionary biology: past, present and future. *J. Evol. Biol.* **19**: 1727-1742.
- Vandeputte, M., Dupont-Nivet, M., Merdy, O., Haffray, P., Chavanne, H. & Chatain, B. 2007. Quantitative genetic determinism of sex-ratio in the European sea bass (*Dicentrarchus labrax* L.). *Aquaculture* **272**: S315-S315.
- Waite, D. E., & Levin, D. A. 1998. Genetic and phenotypic correlations in plants: a botanical test of Cheverud's conjecture. *Heredity* **80**: 310-319.
- Walker, J. A. 1997. Ecological morphology of lacustrine threespine stickleback *Gasterosteus aculeatus* L (Gasterosteidae) body shape. *Biol. J. Linn. Soc.* **61**: 3-50.
- Walker, J.A. & Bell, M.A. 2000. Net evolutionary trajectories of body shape evolution within a microgeographic radiation of threespine sticklebacks (*Gasterosteus aculeatus*). *J. Zool.* **252**: 293-302.
- Webb, P.W. 1982. Locomotor patterns in the evolution of actinopterygian fishes. *Am. Zool.* **22**: 329-342.
- Webb, P.W. 1984. Body form, locomotion and foraging in aquatic vertebrates. *Am. Zool.* **24**: 107-120.
- Wilson, A.J. 2008. Why  $h^2$  does not always equal  $V_A/V_P$ ? *J. Evol. Biol.* **21**: 647-650.
- Whitlock, M. 1996. The heritability of fluctuating asymmetry and the genetic control of developmental stability. *Proc. R. Soc. Lond B* **263**: 849-853.
- Wright, S. 1951. The genetical structure of populations. *Ann. Eugenics* **15**: 323-354.
- Wright, S. 1965. The interpretation of population-structure by F-statistics with special regard to systems of mating. *Evolution* **19**: 395-420.
- Zeng, Z.B. 1988. Long-term correlated response, interpopulation covariation, and interspecific allometry. *Evolution* **42**: 363-374.