

EXAMPLES OF COMPLETE MANIFOLDS OF POSITIVE RICCI CURVATURE WITH NILPOTENT ISOMETRY GROUPS

GUOFANG WEI

It is well known [4] that the isometry group of a complete riemannian manifold M with strictly positive sectional curvature is always compact. This is no longer true in general when M has Ricci curvature $\text{Ric} > 0$. The first example was given in [7] for $\dim M = 4$. In this note we shall prove

THEOREM. *Let L be an n -dimensional simply connected nilpotent Lie group. Then for all sufficiently large p , the product manifold $M^{p+n} = \mathbf{R}^p \times L$ admits complete riemannian metrics with strictly positive Ricci curvature such that the isometry group of M contains L .*

Using a theorem of Malcev [8], we have as an immediate consequence:

COROLLARY. *Every finitely generated torsion-free nilpotent group can be realized as the fundamental group of a complete riemannian manifold with strictly positive Ricci curvature.*

On the other hand, every finitely generated subgroup of the fundamental group of any complete manifold with $\text{Ric} \geq 0$ ($K \geq 0$) is nilpotent (abelian) up to finite index [6, 5, 4].

PROOF OF THE THEOREM. Our construction is inspired by [2]. We first apply an observation in [3, pp. 126–127] to obtain a family of almost flat metrics g_r on L , $0 \leq r < \infty$.

Choose a triangular basis $\{X_1, \dots, X_n\}$ for the Lie algebra l of L , i.e., $[X, X_i] \in l_{i-1}$ whenever $X \in l$, and l_{i-1} is spanned by X_1, \dots, X_{i-1} . For $X = \sum_{i=1}^n a_i X_i$ set $\|X\|^2 = \sum_{i=1}^n h_i^2(r) a_i^2$, where $h_i(r) = (1 + r^2)^{-\alpha_i}$, and $\alpha_n = \alpha > 0$, $2\alpha_i - 4\alpha_{i+1} = 1$, $1 \leq i \leq n-1$. The above norm gives rise to a corresponding almost flat left invariant metric g_r . Then

$$(1) \quad |\text{Ric}_L(X_i)| \leq c(1 + r^2)^{-1},$$

where c is a constant depending on n and the structure constants.

Now we define a warped product metric g on M by

$$g = dr^2 + f^2(r) ds^2 + g_r,$$

where ds^2 is the canonical euclidean metric on the sphere $S^{p-1} \subset \mathbf{R}^p$, $f(r) = r(1 + r^2)^{-1/4}$. g is a complete metric on M , since $f(0) = 0$, $f'(0) = 1$, $f''(0) = 0$, $f(r) > 0$ for $r > 0$, $h_i(r) > 0$ for $r \geq 0$, $h'_i(0) = 0$ for $1 \leq i \leq n$.

Received by the editors December 11, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 53C20; Secondary 57S20.

It is clear that the isometry group of g contains L .

Let $H = \partial/\partial r$ and $U = f(r)^{-1}v$ for a unit tangent vector v of S^{p-1} . Straightforward calculation yields:

$$\begin{aligned} \text{Ric}(H, U) &= 0, \\ \text{Ric}(X_i, H) &= \text{Ric}(X_i, U) = 0, \quad (1 \leq i \leq n), \\ \text{Ric}(X_i, X_j) &= 0, \quad (i \neq j, 1 \leq i, j \leq n). \end{aligned}$$

$$\begin{aligned} \text{Ric}(X_i, X_i) &= -\frac{g''_i}{g_i} - (p-1)\frac{f'g'_i}{fg_i} + \text{Ric}_L(X_i) - \sum_{i \neq j} \frac{g'_i g'_j}{g_i g_j} \\ (2) \quad &\geq \left\{ -2\alpha_i[(2\alpha_i + 1)r^2 - 1] + (p-1)\alpha_i(2 + r^2) \right. \\ &\quad \left. - c(1 + r^2) - \sum_{i \neq j} 4\alpha_i \alpha_j r^2 \right\} / (1 + r^2)^2 \\ &\quad (1 \leq i \leq n). \end{aligned}$$

$$\begin{aligned} \text{Ric}(H, H) &= -\sum_{i=1}^n \frac{g''_i}{g_i} - (p-1)\frac{f''}{f} \\ (3) \quad &= \left\{ -\sum_{i=1}^n 2\alpha_i[(2\alpha_i + 1)r^2 - 1] + (p-1)\frac{r^2 + 6}{4} \right\} / (1 + r^2)^2. \end{aligned}$$

$$(4) \quad \text{Ric}(U, U) = -\frac{f''}{f} + \frac{p-2}{f^2} - (p-2)\left(\frac{f'}{f}\right)^2 - \sum_{i=1}^n \frac{f'g'_i}{fg_i}.$$

Since $1 - (f')^2 \geq 0$, $f'' \leq 0$, we have $\text{Ric}(U, U) > 0$ in (4). Positivity of the Ricci curvature in the equations (2) and (3) follows for p sufficiently large. Observe that every term of the right-hand side decays at a rate of order at least r^{-2} . This completes the proof of the theorem.

REMARK. The smallest p that yields positive Ricci curvature on $M^{p+n} = \mathbf{R}^p \times L$ by means of our construction is quite large in general. For example, in the case of the three-dimensional Heisenberg group $L = H^3$, we have to choose $p > 673$. (With a slightly refined choice of functions, $p > 26$ will already work.) We don't know whether or not p can be chosen much smaller. However, it follows from [1] that necessarily $p \geq 4$ when $L = H^3$.

ACKNOWLEDGMENT. I would like to thank Detlef Gromoll and Michael Anderson for very helpful discussions.

REFERENCES

1. M. Anderson, *On the topology of complete manifolds of nonnegative Ricci curvature* (to appear).
2. L. Bérard Bergery, *Quelques exemples de variétés riemanniennes complètes non compactes à courbure de Ricci positive*, C. R. Acad. Sci. Paris Sér. A-B **302** (1986), 159–161.
3. P. Buser and H. Karcher, *Gromov's almost flat manifolds*, *Astérisque* **81** (1981), 1–148.
4. J. Cheeger and D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, *Ann. of Math. (2)* **96** (1972), 413–443.

5. M. Gromov, *Groups of polynomial growth and expanding maps*, Publ. Math. Inst. Hautes Études Sci. **53** (1981), 53–73.
6. J. Milnor, *A note on curvature and fundamental group*, J. Differential Geom. **2** (1968), 1–7.
7. P. Nabonnand, *Sur les variétés riemanniennes complètes à courbure de Ricci positive*, C. R. Acad. Sci. Paris Sér. A-B **291** (1980), 591–593.
8. M. S. Raghunathan, *Discrete subgroups of Lie groups*, Ergeb. Math. Grenzgeb. Band 68, Springer-Verlag, Berlin and New York, 1972.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK AT STONY BROOK, STONY BROOK, NEW YORK, 11794

