

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Exciton diamagnetic shifts and valley Zeeman effect in monolayer WS₂ and MoS₂ to 65 Tesla A. V. STIER, NHMFL, Los Alamos, K. A. MCCREARY, B. T. JONKER, Naval Research Laboratory, J. KONO, Rice University, S. A. CROOKER, NHMFL, Los Alamos — We report circularly-polarized optical reflection spectroscopy of monolayer WS₂ and MoS₂ at low temperatures (4 K) and in high magnetic fields to 65 T [1]. Both the A and the B exciton transitions exhibit a clear and very similar Zeeman splitting of approximately $-230 \mu\text{eV}/\text{T}$ ($g \simeq -4$), providing the first measurements of the valley Zeeman effect and associated g-factors in monolayer transition-metal disulphides. These results complement and are compared with recent low-field photoluminescence measurements of valley degeneracy breaking in the monolayer diselenides MoSe₂ and WSe₂. Further, the very large magnetic fields used in our studies allows us to observe the small quadratic diamagnetic shifts of the A and B excitons in monolayer WS₂ (0.32 and $0.11 \mu\text{eV}/\text{T}^2$, respectively), from which we calculate exciton radii of 1.53 nm and 1.16 nm. When analyzed within a model of non-local dielectric screening in monolayer semiconductors, these diamagnetic shifts also constrain and provide estimates of the exciton binding energies (410 meV and 470 meV for the A and B excitons, respectively), further highlighting the utility of high magnetic fields for understanding new 2D materials. [1] A. V. Stier et al., submitted, arxiv:1510.07022 (2015)

Andreas Stier
NHMFL, Los Alamos

Date submitted: 05 Nov 2015

Electronic form version 1.4