



Exercise during pregnancy reduces the rate of cesarean section: results of a randomized controlled trial.

Journal:	<i>The Journal of Maternal-Fetal & Neonatal Medicine</i>
Manuscript ID:	DJMF-2011-0799
Manuscript Type:	Original Paper
Date Submitted by the Author:	22-Nov-2011
Complete List of Authors:	Barakat, Ruben; Technical University of Madrid Pelaez, Mireia; Technical University of Madrid Lopez, Carmina; Technical University of Madrid Montejo, Rocio; Hospital Universitario de Fuenlabrada Coteron, Javier; Technical University of Madrid
Keywords:	Delivery, Gestation, Physical activity, Outcome, Labor

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Exercise during pregnancy reduces the rate of cesarean section: results**
6 **of a randomized controlled trial**
7
8
9
10

11
12
13 Authors:

14
15 **Ruben BARAKAT (Corresponding author)**. Faculty of Physical Activity and Sports Sciences.
16 Technical University of Madrid (UPM). +34 91 336 40 20 / barakatruben@gmail.com

17
18 Mireia PELAEZ. Faculty of Physical Activity and Sports Sciences. Technical University of Madrid
19 (UPM).
20

21
22 Carmina LOPEZ. Faculty of Physical Activity and Sports Sciences. Technical University of Madrid
23 (UPM).
24

25 Rocío MONTEJO. Hospital Universitario de Fuenlabrada.
26

27 Javier COTERON. Faculty of Physical Activity and Sports Sciences. Technical University of
28 Madrid (UPM).
29
30
31
32
33
34

35 **Keywords:** Delivery; Gestation; Labor; Outcome; Physical Activity
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ABSTRACT
45
6 Objective:
7

8
9 In this study, the authors assessed the effects of a structured, moderate-intensity
10 exercise program during the entire length of pregnancy on a woman's method of
11 delivery.
12
13
14

15
16
17 Methods:
18

19
20 A randomized controlled trial was conducted with 290 healthy pregnant Caucasian
21 (Spanish) women with a singleton gestation who were randomly assigned to either an
22 exercise ($n=138$) or a control ($n=152$) group. Pregnancy outcomes, including the type of
23 delivery, were measured at the end of the pregnancy.
24
25
26
27
28

29
30 Results:
31

32
33 The percentage of cesarean and instrumental deliveries in the exercise group were
34 lower than in the control group (15.9%, $n=22$; 11.6%, $n=16$ vs. 23%, $n=35$; 19.1%, $n=29$,
35 respectively; $p=0.03$). The overall health status of the newborn as well as other
36 pregnancy outcomes were unaffected.
37
38
39
40
41
42

43
44 Conclusions:
45

46
47 Based on these results, a supervised program of moderate-intensity exercise
48 performed throughout pregnancy was associated with a reduction in the rate of
49 cesarean sections and can be recommended for healthy women in pregnancy.
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Pregnancy is a unique process in which nearly all of the body's control systems are modified to maintain both maternal and fetal homeostasis [1]. With physical exercise (PE) becoming an integral part of many women's lives, the question remains as to whether this exercise has any adverse effects on pregnancy outcomes [2]. In theory, the addition of PE may represent a significant challenge to both the maternal and fetal wellbeing as the dual stresses of pregnancy and exercise may create conflicting physiological demands and thus adversely affect pregnancy outcomes [3].

Research on exercise in pregnancy has increased over the years, and many studies have demonstrated the benefits of PE on maternal and fetal outcomes [4, 5], including positive maternal psychological factors [6, 7] and decreased risks of preeclampsia and gestational diabetes (GD) [8, 9].

Despite the advancement of knowledge on exercise during pregnancy, a general lack of consensus persists regarding the effect of PE on the type of delivery. Scientific evidence from experimental studies on the influence of exercise on the type of delivery is limited and imprecise [10]. Because most studies were observational (with no intervention) and focused on leisure time and occupational physical activity, little information is available about the effects of a supervised exercise program throughout pregnancy on the type of delivery.

Since 2000, the worldwide rate of cesarean delivery has risen even more sharply and is currently over 31 percent. This rate is concerning from a public health perspective because the procedure is not without risk to the mother and fetus [10]. Known risks of cesarean sections include the following: maternal postnatal complications stemming

1
2
3 from major abdominal surgery, placental problems in future pregnancies, neonatal
4
5 respiratory difficulties, poor sucking reflex in the newborn, infection, excessive blood
6
7 loss, maternal respiratory complications, anesthesia reactions and longer
8
9 hospitalizations [13,14].
10
11

12
13 The aim of the present study was to understand the influence of an exercise program
14
15 during pregnancy (all three trimesters) on the type of delivery that ultimately
16
17 occurred. A secondary objective was to determine if exercise could be a protective
18
19 factor against cesarean sections or instrumental delivery methods in women who
20
21 experienced excessive weight gain during pregnancy. We hypothesized that regular
22
23 exercise during pregnancy would be associated with a lower percentage of cesarean
24
25 sections in normal pregnancy and lower cases of instrumental deliveries and cesarean
26
27 sections in pregnant women with excessive weight gain.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

The present study was a randomized controlled clinical trial (RCT) in accordance with the recently published CONSORT guidelines (<http://www.consort-statement.org>) [15].

The study also met the ethical standards in Sport and Exercise Science Research [16].

The research protocol was reviewed and approved by the Ethical Committee of Hospital Universitario de Fuenlabrada (Madrid, Spain) and followed the ethical guidelines outlined in the Declaration of Helsinki (last modified in 2008). The research was performed in the obstetrics department of the Hospital. All participants were informed about the aim and study protocol, and written informed consents were obtained from all participating women.

Study subjects:

Participants were recruited from a hospital database. All the women were healthy with uncomplicated and singleton pregnancies. To allocate participants, a computer-generated list of random numbers was used. The randomization process (sequence generation, allocation concealment and implementation) was conducted for three different study investigators.

Patients were excluded if they had any type of absolute obstetrical contraindication to exercise regarded by the American College of Obstetricians and Gynecologists (ACOG) (e.g. multiple pregnancies or risk for premature labor) [17]. Other exclusion criteria included those women not planning to give birth in the obstetrics department of the study hospital, not receiving medical follow-up throughout the pregnancy,

1
2
3 participating in another physical activity program or having a high level of
4
5 pregestational PE (four or more times per week).
6
7

8 Intervention:
9

10
11 The physical conditioning program included a total of three 40-45 minute sessions per
12
13 week, beginning at the start of the pregnancy (weeks 6–9) until the end of the third
14
15 trimester (weeks 38–39). Thus, there was an average of 85 training sessions planned
16
17 for each participant in the absence of preterm delivery. All subjects used a heart rate
18
19 (HR) monitor (Accurex Plus, Polar Electro OY, Oulu, Finland) during the training
20
21 sessions to ensure that exercise intensity was light to moderate. Their HR was
22
23 consistently kept under 70% of their age-predicted maximum HR value, calculated as
24
25 220 minus age.
26
27
28
29

30
31 Each session included a 25-minute core portion that was preceded and followed by a
32
33 gradual warm-up and cool-down period, both of 7–8 minutes in duration and
34
35 consisting of walking and light, static stretching (to avoid any muscle pains) of most
36
37 muscle groups (upper and lower limbs, neck and trunk muscles).
38
39

40
41 We used exercises covering the major muscle group in the arms and abdomen to
42
43 promote good posture, prevent lower back pain and strengthen the muscles of labor
44
45 and the pelvic floor. We also included an aerobic dance section in every session.
46
47 Exercises involving the Valsalva maneuver, extreme stretching, joint overextension,
48
49 ballistic movements and jumping were specifically avoided. Furthermore, exercises in
50
51 the supine position were performed for no longer than two minutes.
52
53
54
55
56
57
58
59
60

1
2
3 In order to maximize safety, adherence to the training program and its efficacy, all
4 sessions were i) supervised by a qualified fitness specialist (working with groups of 10–
5 12 subjects) with an obstetrician's assistance; ii) accompanied by music; and iii)
6 performed in the Health Care Center in a spacious, well-lit room under favorable
7 environmental conditions (altitude 600 m; temperature=19–21°C; humidity=50–60%).
8 An adequate intake of calories and nutrients was assured before the start of the
9 exercise session.
10
11

12 Measurement of Variables:

13
14 We obtained the type of delivery (normal, instrumental, or cesarean) and pregnancy
15 outcomes from hospital perinatal records including gestational age (days), preterm
16 deliveries (before the completion of 37 weeks gestation), maternal weight gain (kg),
17 blood pressure, one hour oral glucose tolerance test, cases of GD, birth weight/length,
18 pH of the umbilical cord blood and Apgar score.
19

20
21 The following maternal characteristics were obtained by an initial interview at the first
22 prenatal visit: maternal age, parity, smoking habits during pregnancy, occupational
23 activity, time standing per day, time spent on domestic activities and educational level.
24

25
26 We measured weight and height of the mother by standard procedures at the start of
27 the study. Body mass index (BMI) was calculated as weight (in kilograms) divided by
28 height (in meters) squared. Adherence (attendance) to the training program was
29 measured using a checklist at every session.
30
31

32
33 Data analysis:
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Our data (presented in Tables I and II) were analyzed using Student's t-tests for
4
5 independent samples, one-way ANOVA and χ^2 tests.
6
7

8
9 Maternal characteristics of the study sample by group (EG and CG) were reported in
10
11 terms of means and standard deviation (SD), unless otherwise stated. Relative to
12
13 frequencies, χ^2 tests were used. For group comparisons of pregnancy outcomes,
14
15 continuous and nominal data were analyzed with t-tests for unpaired data and χ^2
16
17 tests, respectively. To estimate the effect size, relative risk and 95% confidence
18
19 intervals for the difference were calculated.
20
21

22
23 An interim analysis was conducted to assure the safety of the intervention during the
24
25 trial (n=60). The levels of significance maintained an overall p -value of 0.05 and were
26
27 calculated according to the O'Brien-Fleming stopping boundaries [18].
28
29

30
31 Determination of sample size: To detect a 10% reduction in cesarean sections with a
32
33 two-sided 5% significance level and a power of 80%, a sample size of 125 pregnant per
34
35 group was necessary, given an anticipated dropout rate of 15% [19].
36
37

38
39 Statistical analysis was performed using the Statistical Package for Social Sciences
40
41 software (version 14.0 for Windows; SPSS Inc., Chicago, IL) with the level of statistical
42
43 significance set to <0.05 .
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Out of the 595 women originally contacted at their first prenatal visit, 320 Caucasian, healthy gravid women (aged 31 (SD=4) years) were randomly assigned to either an exercise group (EG, n=160) or a control group (CG, n=160). Of these women, 290 belonged to a low-to-medium socioeconomic class.

A flow diagram of the randomization process of the study participants is shown in Figure 1. Fourteen percent (22/160) and 5% (8/160) of pregnant women in the EG and CG, respectively, were lost to follow-up. We excluded twenty-two women in the EG for the following reasons: discontinued intervention (n=8), risk for premature labor (n=3), pregnancy-induced hypertension (n=2) and personal reasons, such as change of residence or work-related issues (n=9). Eight women were excluded in the CG for pregnancy-induced hypertension (n=2), threat of premature delivery (n=3) and personal reasons (n=3).

Adherence to training and possible adverse effects:

Adherence to the training program in the EG was 87%. In the EG, 138 pregnant completed the program with a high level of satisfaction. Because participating in another exercise program was an exclusion criterion for the CG, this was verified via telephone interview at week 20 and at the end of pregnancy. In the end, 138 women in the EG and 152 women in the CG were included in the analysis.

Figure 1: Flow diagram of the study participants following the CONSORT guidelines (<http://www.consortstatement.org>).

1
2
3 Maternal characteristics: Background variables of the study populations are shown in
4
5 Table 1. There were no significant differences ($p>0.05$) in background variables
6
7 between the two study groups; these background variables included maternal age,
8
9 BMI, parity, previous preterm delivery, previous miscarriages and previous low birth
10
11 weight. Maternal characteristics that could be potentially influence the main study
12
13 outcome (e.g., occupational activity, educational level, smoking during pregnancy,
14
15 standing and housework) were also not significantly different between the groups.
16
17
18
19

20 Table I: Maternal Characteristics: comparison between control and intervention
21
22 groups.
23
24

25 Type of delivery and other pregnancy outcomes:
26
27

28
29 We observed that the percentage of cesarean sections (15.9%, n=22) and instrumental
30
31 deliveries (11.6%, n=16) in the EG was lower than those in the CG (23%, n=35; 19.1%,
32
33 n=29, respectively; $Z=2.73$; $p= 0.03$; $1-\beta=0.65$). We observed a reduced Relative Risk
34
35 (RR) of cesarean section for women in the EG compared with the CG (0.47, 95% CI=
36
37 0.26, 0.82). Nine women delivered before term in the EG (6.5%) and 10 (6.6%) in the
38
39 CG. The mean gestational age was similar in both groups (EG=278.0 (SD=9.9),
40
41 CG=278.3 (SD=10.3) days).
42
43
44
45

46 When other pregnancy outcomes were examined, we found similar results between
47
48 groups for Oral Glucose Tolerance Test, GD and mean blood pressure during
49
50 pregnancy. Maternal weight gain was lower in the EG (11.9 (SD=3.7) kg) compared to
51
52 the CG (13.7 (SD=4.1) Kg) ($p= 0.0001$). We observed no differences between the groups
53
54 in the newborn's overall health status, including Apgar scores at one and five minutes,
55
56 birth weight, birth length and the pH of umbilical cord blood.
57
58
59
60

1
2
3 Table II: Type of delivery and other pregnancy outcomes: comparison between control
4
5 and intervention groups.
6
7

8
9 An additional analysis was conducted to compare the types of deliveries between
10
11 groups of pregnant women with excessive weight gain. There was a lower rate of
12
13 cesarean sections in the EG (16.1%, n=5) compared to the CG (39.7%, n=27), although
14
15 the difference was not statistically significant.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

This study is the first randomized controlled trial that specifically and objectively shows a positive association between the type of delivery and supervised, regular exercise when performed by a large sample of previously sedentary, gravid women throughout the duration of their entire pregnancy. An additional novel intervention of our study was the integration of light resistance, toning, aerobic dance and pelvic floor exercises in the training program, which is not commonly available to pregnant women.

We examined the influence of PE on the delivery method, providing an opportunity to determine the possible influence of a physical activity program during pregnancy (during all three trimesters) on the type of delivery.

The main finding of our study was that supervised resistance training is associated with a lower percentage of cesarean and instrumental deliveries. These findings have clinical relevance given that the cesarean rate is a public health concern because the procedure is not without risks and can adversely affect perinatal outcomes [10, 14].

The overall health status of the baby was unaffected with exercise, as reflected and reinforced by the results of the globally used Apgar score. Moreover, some pregnancy outcomes (e.g., maternal weight gain) improved in the EG, which could explain the protective effect of PE against pregnancy complications.

In additional analyses, our results show a lower percentage of cesarean sections in women with excessive weight gain in the EG. In concordance, many researchers have found that normal maternal BMI and appropriate weight gain during pregnancy is

1
2
3 strongly related to the ability of pregnant women to delivery vaginally without
4
5 difficulty [20-22].
6
7

8
9 In the past, studies on the influence of leisure-time physical activity on the type of
10
11 delivery have generated mixed results. Most of these studies refer to the relationship
12
13 between occupational activity or working conditions and pregnancy outcomes, though
14
15 few clinical trials existed.
16
17

18
19 Our results are in agreement with other study findings. In an experimental study, Clapp
20
21 found that women who continue to exercise during pregnancy compared with
22
23 sedentary pregnant women had a lower incidence of cesarean sections (6 %, n=5 vs.
24
25 30%, n= 13) and instrumental deliveries (6 %, n= 5 vs. 20%, n= 9) [17]. Recently, in an
26
27 observation study, Melzer et al. measured energy expenditure, aerobic fitness and
28
29 sleeping HR in 44 healthy women in late pregnancy and found that the risk for
30
31 operative delivery was lower in active women compared with those that were inactive
32
33 (odds ratio of 3.7) [23].
34
35
36
37

38
39 Bungun et al., by a non-experimental, retrospective study, assessed the association
40
41 between participation in aerobic exercise during the first two trimesters of pregnancy
42
43 and type of delivery in 137 nulliparous women. They found that sedentary women
44
45 (n=93) were 2.05 times more likely to deliver via cesarean section than active women
46
47 (n=44), though this difference was not statistically significant [13].
48
49

50
51 A notable group of studies found no influence of physical activity on the type of
52
53 delivery (5, 24-26). In a recent study, Bø et al. found that exercising at least three times
54
55 per week was not associated with vacuum or forceps deliveries or acute cesarean
56
57 deliveries [11].
58
59
60

1
2
3 In a randomized controlled trial, Cavalcante et al. studied 71 low-risk sedentary
4 pregnant women to evaluate the effectiveness and safety of water aerobics during
5 pregnancy (activity was initiated at 18-20 weeks). They found no difference in the type
6 of delivery between study groups; however, most of the deliveries in the intervention
7 group were normal vaginal deliveries [27].
8
9

10
11
12
13
14
15
16 Bovbjerg and Siega-Riz, by a Pregnancy Risk Assessment Monitoring System, collected
17 data (via questionnaire) on the frequency of exercise during the last three months of
18 pregnancy in 1955 women. They found that exercise was not associated with delivery
19 method [10].
20
21
22
23

24
25
26 In contrast with these investigations, a few studies suggest that exercise during
27 pregnancy may have a negative impact on the type of delivery. Dale, in a study with 33
28 runners and 11 controls, reported a trend noticed in runners; this group tended to fail
29 to progress in labor and delivery, and thus resulted in cesarean sections [12].
30
31
32
33

34
35
36 In a prospective study to evaluate the influence of employment and physical exertion
37 on pregnancy outcomes (quantified by kilocalories expended each day), Magann et al.
38 found an increased proportion of nonelective cesarean sections for women in the
39 higher energy expenditure group (>3000 kcal/day) [28].
40
41
42
43
44
45

46
47 The differences in our results compared to those of other studies can be attributed to
48 the variety of study design, type and intensity of the exercise and duration of the
49 programs used by other studies. In our previous study [5], we observed no influence of
50 exercise on the type of delivery in an RCT performed only during the second and third
51 trimesters of pregnancy. Also, other experimental studies [11, 27] start physical
52 activity programs around 18-20 weeks of gestation, which could be an important factor
53
54
55
56
57
58
59
60

1
2
3 in the difference with the results of the present study. In addition, many observational
4
5 studies derive their results from the information from pregnant women [10, 23, 25] or
6
7 only by the comparison between groups of different exercise intensities [24].
8
9 Moreover, studies that found physical exertion (runners, high workload) was
10
11 associated with abdominal deliveries [12, 28] could demonstrate that exercise during
12
13 pregnancy should be limited to moderate intensity.
14
15

16
17
18 We observe that different studies, usually testing a small sample or based on
19
20 observational designs, have given inconsistent results and conclusions. The main
21
22 contribution of this study is evidence that regular exercise can coexist with pregnancy
23
24 and have positive effects on the type of delivery.
25
26

27
28 In summary, our findings encourage the growing trend in healthy, pregnant women to
29
30 join a physical activity program. This study suggests that light-to-moderate exercise
31
32 can result in improved clinical outcomes for the mother and newborn and serve to be
33
34 an important contribution to public health.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

The authors would like to acknowledge the technical assistance of the Gynaecology and Obstetric Service of Fuenlabrada Hospital of Madrid. This work was partially supported by the Ministerio de Ciencia e Innovación, Spain (DEP2009-08990).

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DECLARATION OF INTEREST

The authors report no Competing of Interest.

For Peer Review Only

REFERENCES

1. Wolfe L, Brenner I, Mottola M. Maternal exercise, fetal well-being and pregnancy outcome. *Exerc Sport Sci Rev.* 1994; 22:145–194.
2. Riemann MK, Kanstrup Hansen IL. Effects on the foetus of exercise in pregnancy. *Scand J Med Sci Sports.* 2000; 10:12-19.
3. Sternfeld B. Physical activity and pregnancy outcome. *Sports Med.* 1997; 23: 33-47.
4. Barakat R, Lucia A, Ruiz JR. Resistance exercise training during pregnancy and newborn's birth size: a randomized controlled trial. *Int J Obes (Lond).* 2009; 33:1048-1057.
5. Barakat R, Ruiz JR, Stirling JR, et al. Type of delivery is not affected by light resistance and toning exercise training during pregnancy: a randomized controlled trial. *Am J Obstet Gynecol.* 2009; 201:590 e 1-6.
6. Barakat R, Pelaez M, Montejo R, et al. Exercise during pregnancy improves maternal health perception: a randomized controlled trial. *Am J Obstet Gynecol.* 2011 May;204(5):402.e1-7.
7. Poudevigne MS, O'Connor PJ. Physical activity and mood during pregnancy. *Med Sci Sports Exerc.* 2005; 37:1374-1380.
8. Mottola MF. The role of exercise in the prevention and treatment of gestational diabetes mellitus. *Curr Sports Med Rep.* 2007; 6:381-386.
9. Sorensen TK, Williams MA, Lee IM, et al. Recreational physical activity during pregnancy and risk of preeclampsia. *Hypertension.* 2003; 41:1273-1280.
10. Bovbjerg ML, Siega-Ruiz AM. Exercise during pregnancy and cesarean delivery: North Carolina PRAMS, 2004-2005. *Birth.* 2009; 36:200-207.

11. Bø K, Fleten C, Nystad W. Effect of antenatal pelvic floor muscle training on labor and birth. *Obstet Gynecol.* 2009; 113:1279-1284.
12. Dale E, Mullinax KM, Bryan DH. Exercise during pregnancy: effects on the fetus. *Can J Appl Sport Sci.* 1982; 7:98-103.
13. Bungum TJ, Peaslee DL, Jackson AW, et al. Exercise during pregnancy and type of delivery in nulliparae. *J Obstet Gynecol Neonatal Nurs.* 2000; 29:258-264.
14. Blanchette H. The rising cesarean delivery rate in america: what are the consequences? *Obstet Gynecol.* 2011 Sep; 118(3):687-690.
15. Schulz KF, Altman DG, Moher D, for the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomized trials. *BMC Medicine.* 2010; 8:18.
16. Harriss DJ, Atkinson G. Ethical Standards in Sport and Exercise Science Research. *Int J Sports Med.* 2009; 30:701-702.
17. Artal R, O'Toole M. Guidelines of the American College of Obstetricians and Gynecologists for exercise during pregnancy and the postpartum period. *Br J Sports Med.* 2003; 37:6-12.
18. Pocock SJ. When to stop a clinical trial. *BMJ.* 1992; 305:235-240.
19. Moher D, Dulberg CS, Wells GA. Statistical power, sample size, and their reporting in randomized controlled trials. *JAMA.* 1994; 272:122-124.
20. Gawade P, Markenson G, Bsat F, Healy A, Pekow P, Plevyak M. Association of gestational weight gain with cesarean delivery rate after labor induction. *J Reprod Med.* 2011 Mar-Apr;56(3-4):95-102.
21. Linné Y. Effects of obesity on women's reproduction and complications during pregnancy. *Obes Rev.* 2004; 5:137-143.

References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
22. Young TK, Woodmansee B. Factors that are associated with cesarean delivery in a large private practice: the importance of prepregnancy body mass index and weight gain. *Am J Obstet Gynecol.* 2002; 187:312-318.
23. Melzer K, Schutz Y, Soehnchen N, et al.. Effects of recommended levels of physical activity on pregnancy outcomes. *Am J Obstet Gynecol.* 2010; 202:266 e1-6.
24. Kardel KR, Kase T. Training in pregnant women: effects on fetal development and birth. *Am J Obstet Gynecol.* 1998; 178:280-286.
25. Sternfeld B, Quesenberry CP Jr, Eskenazi B, et al. Exercise during pregnancy and pregnancy outcome. *Med Sci Sports Exerc.* 1995; 27:634-640.
26. Wong SC, McKenzie DC. Cardiorespiratory fitness during pregnancy and its effect on outcome. *Int J Sports Med.* 1987; 8:79-83.
27. Cavalcante S, Cecatti J, Pereira R, et al. Water aerobics II: maternal body composition and perinatal outcomes after a program for low risk pregnant women. *Reprod Health.* 2009; 6:1.
28. Magann EF, Evans SF, Weitz B, et al. Antepartum, intrapartum, and neonatal significance of exercise on healthy low-risk pregnant working women. *Obstet Gynecol.* 2002; 99:466-472.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

TABLES AND FIGURES

TABLE I: Maternal Characteristics: Comparison Between Control and Exercise Groups.

		Controls (n=152)	Exercise (n=138)	p-value
		Mean (SD) or n/%	Mean (SD) or n/%	
Maternal age (years)		31.7 (4.5)	31.4 (3.2)	0.54
BMI ^a		23.6 (4.0)	24.0 (4.3)	0.45
Parity	0	83/54.6	88/60.9	0.19
	1	60/39.5	46/33.3	
	>1	9/5.9	4/2.9	
Occupational Activity	Active	51/33.6	55/39.9	0.50
	Sedentary	68/44.7	58/42	
	Housewives	33/21.7	25/18.1	
Educational level	College	45/29.6	29/21	0.23
	High school	67/44.1	70/50.7	
Smoked during pregnancy	<High school	40/26.3	39/28.3	0.16
	Yes	29/19.1	18/13	
Standing Hours/day	No	123/80.9	120/87	0.45
	>3 h/day	95/62.5	92/66.7	
Previous Preterm Deliveries	3 or <3 h/day	57/37.5	46/33.3	0.46
	0	146/96.1	130/94.2	
Previous Miscarriages	1 or >1	6/3.9	8/5.8	0.20
	0	104/68.4	107/77.5	
	1	42/27.6	28/20.3	
Previous Low Birth Weight Deliveries	>1	6/3.9	3/2.2	0.56
	0	147/96.7	135/97.8	
Housework Hours/day	1 or >1	5/3.3	3/2.2	0.24
	>2 h/day	81/53.3	83/60.1	
	2 or <2 h/day	71/45.5	55/39.9	

^aBMI at first prenatal medical visit.

TABLE II: Type of Delivery and Other Pregnancy Outcomes—Comparison between Control and Exercise Groups.

		Controls (n=152)	Exercise (n=138)	
		Mean (SD) or n/%	Mean (SD) or n/%	p-value
Type of delivery	Normal	88/57.9	100/72.5	*0.03 $\chi^2 = 6.8$
	Instrumental	29/19.1	16/11.6	Z=2.73 RR=0.47
	Cesarean	35/23	22/15.9	(0.26, 0.82)
Preterm delivery		10/6.6	9/6.5	0.98
Maternal weight gain (kg)		13.7 (4.1)	11.9 (3.7)	*0.0001 **d= 0.56
Gestational age (days)		278.0 (10.3)	278.3 (9.9)	0.81
Apgar Score 1 min.		8.6 (1.3)	8.7 (1.4)	0.34
Apgar Score 5 min.		9.8 (0.8)	9.7 (0.6)	0.22
Birth Weight (g)		3232 (448)	3203 (461)	0.56
Birth Length (cm)		49.5 (2.07)	49.7 (2.06)	0.98
pH Umbilical Cord.		7.26 (0.07)	7.28 (0.08)	0.29
Oral Glucose Tolerance Test (g/dL).		119.2 (28.5)	117.2 (30.7)	0.59
Gestational Diabetes	Yes	12/7.9	6/4.3	0.21
	No	140/92.1	132/95.7	
Blood Pressure ^a	Diastolic	70.8 ± 8.5	69.6 ± 9.6	0.29
	Systolic	115.6 ± 11.7	113.8 ± 15.1	0.25
EWGDP***		Controls (n= 53)	Exercise (n= 31)	
Type of delivery (n/%)	Normal	25/47.2	22/71	
	Instrumental	8/15.1	4/12.9	0.07
	Cesarean	20/37.7	5/16.1	

*Significant level at $p=0.05$ ** Cohen's d

*** Excessive Weight Gain During Pregnancy

^aMean during pregnancy.




Figure 1. Flow chart of the study participants