

## EXISTENCE AND SYMMETRY OF POSITIVE GROUND STATES FOR A DOUBLY CRITICAL SCHRÖDINGER SYSTEM

ZHIJIE CHEN AND WENMING ZOU

ABSTRACT. We study the following doubly critical Schrödinger system:

$$\begin{cases} -\Delta u - \frac{\lambda_1}{|x|^2} u = u^{2^*-1} + \nu \alpha u^{\alpha-1} v^\beta, & x \in \mathbb{R}^N, \\ -\Delta v - \frac{\lambda_2}{|x|^2} v = v^{2^*-1} + \nu \beta u^\alpha v^{\beta-1}, & x \in \mathbb{R}^N, \\ u, v \in D^{1,2}(\mathbb{R}^N), \quad u, v > 0 \text{ in } \mathbb{R}^N \setminus \{0\}, \end{cases}$$

where  $N \geq 3$ ,  $\lambda_1, \lambda_2 \in (0, \frac{(N-2)^2}{4})$ ,  $2^* = \frac{2N}{N-2}$  and  $\alpha > 1, \beta > 1$  satisfying  $\alpha + \beta = 2^*$ . This problem is related to coupled nonlinear Schrödinger equations with critical exponent for Bose-Einstein condensate. For different ranges of  $N, \alpha, \beta$  and  $\nu > 0$ , we obtain positive ground state solutions via some quite different variational methods, which are all radially symmetric. It turns out that the least energy level depends heavily on the relations among  $\alpha, \beta$  and 2. Besides, for sufficiently small  $\nu > 0$ , positive solutions are also obtained via a variational perturbation approach. Note that the Palais-Smale condition cannot hold for any positive energy level, which makes the study via variational methods rather complicated.

### 1. INTRODUCTION

In this paper we consider solitary wave solutions of coupled nonlinear Schrödinger equations, known in the literature as Gross-Pitaevskii equations ([19, 34]):

$$(1.1) \quad \begin{cases} -i \frac{\partial}{\partial t} \Phi_1 = \Delta \Phi_1 - a(x) \Phi_1 + \mu_1 |\Phi_1|^2 \Phi_1 + \nu |\Phi_2|^2 \Phi_1, & x \in \mathbb{R}^N, t > 0, \\ -i \frac{\partial}{\partial t} \Phi_2 = \Delta \Phi_2 - b(x) \Phi_2 + \mu_2 |\Phi_2|^2 \Phi_2 + \nu |\Phi_1|^2 \Phi_2, & x \in \mathbb{R}^N, t > 0, \\ \Phi_j = \Phi_j(x, t) \in \mathbb{C}, & j = 1, 2, \\ \Phi_j(x, t) \rightarrow 0, & \text{as } |x| \rightarrow +\infty, t > 0, j = 1, 2, \end{cases}$$

where  $i$  is the imaginary unit,  $a(x), b(x)$  are potential functions,  $\mu_1, \mu_2 > 0$  and  $\nu \neq 0$  is a coupling constant. System (1.1) appears in many physical problems, especially in nonlinear optics. Physically, the solution  $\Phi_j$  denotes the  $j^{\text{th}}$  component of the beam in Kerr-like photorefractive media (see [3]). The positive constant  $\mu_j$  is for self-focusing in the  $j^{\text{th}}$  component of the beam. The coupling constant  $\nu$  is the interaction between the two components of the beam. Problem (1.1) also arises in the Hartree-Fock theory for a double condensate, i.e., a binary mixture of Bose-Einstein condensates in two different hyperfine states |1) and |2) (see [14]). Physically,  $\Phi_j$  are the corresponding condensate amplitudes, and  $\mu_j$  and  $\nu$  are the

---

Received by the editors December 13, 2012 and, in revised form, July 2, 2013.

2010 *Mathematics Subject Classification*. Primary 35J50, 35J47; Secondary 35B33, 35B09.

This work was supported by NSFC (11025106, 11371212, 11271386) and the Both-Side Tsinghua Fund.

©2014 American Mathematical Society  
 Reverts to public domain 28 years from publication

intraspecies and interspecies scattering lengths. The sign of  $\nu$  determines whether the interactions of states  $|1\rangle$  and  $|2\rangle$  are repulsive or attractive, i.e., the interaction is attractive if  $\nu > 0$ , and the interaction is repulsive if  $\nu < 0$ , where the two states are in strong competition.

To obtain solitary wave solutions of system (1.1), we set  $\Phi_1(x, t) = e^{i\lambda_1 t}u(x)$  and  $\Phi_2(x, t) = e^{i\lambda_2 t}v(x)$ . Write  $V_1(x) = a(x) + \lambda_1$  and  $V_2(x) = b(x) + \lambda_2$  for convenience, and as we are only interested in nonnegative solutions, then system (1.1) is reduced to the following elliptic system:

$$(1.2) \quad \begin{cases} -\Delta u + V_1(x)u = \mu_1 u^3 + \nu uv^2, & x \in \mathbb{R}^N, \\ -\Delta v + V_2(x)v = \mu_2 v^3 + \nu vu^2, & x \in \mathbb{R}^N, \\ u \geq 0, v \geq 0 \text{ in } \mathbb{R}^N, & u(x), v(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty. \end{cases}$$

This Bose-Einstein condensate type system (1.2) is a special case of the following problem:

$$(1.3) \quad \begin{cases} -\Delta u + V_1(x)u = \mu_1 u^{2p-1} + \nu u^{p-1}v^p, & x \in \mathbb{R}^N, \\ -\Delta v + V_2(x)v = \mu_2 v^{2p-1} + \nu v^{p-1}u^p, & x \in \mathbb{R}^N, \\ u \geq 0, v \geq 0 \text{ in } \mathbb{R}^N, & u(x), v(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty, \end{cases}$$

where  $p > 1$  and  $p \leq 2^*/2$  if  $N \geq 3$ , and  $2^* = \frac{2N}{N-2}$  is the critical Sobolev exponent. If  $p = 2$ , then (1.3) turns to be the cubic system (1.2). For further introduction about this problem, readers can also see the survey articles [16, 20], which also contain information about the physical relevance of noncubic nonlinearities (e.g. quintic). For the subcritical case  $p < 2^*/2$ , the existence and multiplicity of solutions to (1.3) have been widely studied under different assumptions on  $V_i$  and  $\nu$ ; see [4, 22, 25–28, 34] and the references therein.

Notice that all the papers mentioned above deal with the subcritical case. To the best of our knowledge, there are no existence results for (1.3) in the critical case  $2p = 2^*$  in the literature.

In this paper, we study (1.3) in critical case where  $N \geq 3$  and  $2p = 2^*$ . In this case, if  $V_i(x) = \lambda_i$  are nonzero constants with the same sign, then by Pohozaev identity, we easily conclude that any solution  $(u, v)$  of (1.3) satisfies  $\int_{\mathbb{R}^N} \lambda_1 u^2 + \lambda_2 v^2 = 0$ , and so  $(u, v) = (0, 0)$ . Hence we do not consider the case  $V_i(x) = \lambda_i$  here, and in the sequel we assume that  $V_i(x) = -\frac{\lambda_i}{|x|^2}$  are Hardy type potentials. The Hardy type potentials, which arise in several physical contexts (e.g., in nonrelativistic quantum mechanics, molecular physics, quantum cosmology, and linearization of combustion models), do not belong to Kato’s class, so they cannot be regarded as a lower order perturbation term. In particular, any nontrivial solutions of (1.3) with  $V_i(x) = -\frac{\lambda_i}{|x|^2}$  are singular at  $x = 0$ . For the sake of simplicity, in the sequel we assume that  $\mu_1 = \mu_2 = 1$ . Then, to study (1.3) with  $2p = 2^*$ ,  $V_i(x) = -\frac{\lambda_i}{|x|^2}$  and  $\mu_1 = \mu_2 = 1$ , we turn to study the following general problem:

$$(1.4) \quad \begin{cases} -\Delta u - \frac{\lambda_1}{|x|^2}u = u^{2^*-1} + \nu \alpha u^{\alpha-1}v^\beta, & x \in \mathbb{R}^N, \\ -\Delta v - \frac{\lambda_2}{|x|^2}v = v^{2^*-1} + \nu \beta u^\alpha v^{\beta-1}, & x \in \mathbb{R}^N, \\ u, v \in D^{1,2}(\mathbb{R}^N), & u, v > 0 \text{ in } \mathbb{R}^N \setminus \{0\}, \end{cases}$$

where  $N \geq 3$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$ ,  $\Lambda_N := \frac{(N-2)^2}{4}$ ,

$$(1.5) \quad \alpha > 1, \quad \beta > 1, \quad \alpha + \beta = 2^*,$$

and  $D^{1,2}(\mathbb{R}^N)$  is the completion of  $C_0^\infty(\mathbb{R}^N)$  with respect to the norm

$$\|u\| := \left( \int_{\mathbb{R}^N} |\nabla u|^2 dx \right)^{1/2}.$$

Note that if  $\alpha = \beta = p = 2^*/2$ , then (1.4) turns to be (1.3) with  $V_i(x) = -\frac{\lambda_i}{|x|^2}$  and  $\mu_1 = \mu_2 = 1$ . The mathematical interest in system (1.4) relies on their double criticality, due to the fact that both the exponent of the nonlinearities (which is critical in the sense of the Sobolev embedding) and the singularities share the same order of homogeneity as the Laplacian. *The main goal of this paper is to study the existence and radial symmetry of ground state solutions to system (1.4)*, where the ground state solution is defined in Definition 1.1 below.

Recall that  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$ ; from Hardy's inequality

$$(1.6) \quad \Lambda_N \int_{\mathbb{R}^N} \frac{u^2}{|x|^2} dx \leq \int_{\mathbb{R}^N} |\nabla u|^2 dx, \quad \forall u \in D^{1,2}(\mathbb{R}^N),$$

we see that  $\|\cdot\|_{\lambda_i}, i = 1, 2$ , are equivalent norms to  $\|\cdot\|$ , where

$$(1.7) \quad \|u\|_{\lambda_i}^2 := \int_{\mathbb{R}^N} |\nabla u|^2 - \frac{\lambda_i}{|x|^2} u^2 dx.$$

Denote the norm of  $L^p(\mathbb{R}^N)$  by  $|u|_p = \left(\int_{\mathbb{R}^N} |u|^p dx\right)^{\frac{1}{p}}$ . The case of a single equation has been deeply investigated in the literature. In particular, by [33], the problem

$$(1.8) \quad \begin{cases} -\Delta u - \frac{\lambda_i}{|x|^2} u = u^{2^*-1}, & x \in \mathbb{R}^N, \\ u(x) \in D^{1,2}(\mathbb{R}^N), & u > 0 \text{ in } \mathbb{R}^N \setminus \{0\} \end{cases}$$

has exactly a one-dimensional  $C^2$  manifold of positive solutions given by

$$(1.9) \quad Z_i = \left\{ z_\mu^i(x) = \mu^{-\frac{N-2}{2}} z_1^i\left(\frac{x}{\mu}\right), \quad \mu > 0 \right\},$$

where

$$(1.10) \quad z_1^i(x) = \frac{A(N, \lambda_i)}{|x|^{a_{\lambda_i}} \left(1 + |x|^2 - \frac{4a_{\lambda_i}}{N-2}\right)^{\frac{N-2}{2}}},$$

$a_{\lambda_i} = \frac{N-2}{2} - \sqrt{\frac{(N-2)^2}{4} - \lambda_i}$  and  $A(N, \lambda_i) = \frac{N(N-2-2a_{\lambda_i})^2}{N-2}$ . Moreover, all positive solutions of (1.8) satisfy

$$(1.11) \quad S(\lambda_i) := \inf_{u \in D^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\|u\|_{\lambda_i}^2}{|u|_{2^*}^2} = \frac{\|z_\mu^i\|_{\lambda_i}^2}{|z_\mu^i|_{2^*}^2} = \left(1 - \frac{4\lambda_i}{(N-2)^2}\right)^{\frac{N-1}{N}} S,$$

and

$$(1.12) \quad I_{\lambda_i}(z_\mu^i) = \frac{1}{N} \|z_\mu^i\|_{\lambda_i}^2 = \frac{1}{N} |z_\mu^i|_{2^*}^{2^*} = \frac{1}{N} S(\lambda_i)^{N/2},$$

where  $S$  is the sharp constant of  $D^{1,2}(\mathbb{R}^N) \hookrightarrow L^{2^*}(\mathbb{R}^N)$ ,

$$(1.13) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq S \left( \int_{\mathbb{R}^N} |u|^{2^*} dx \right)^{\frac{2}{2^*}},$$

and

$$(1.14) \quad I_{\lambda_i}(u) := \frac{1}{2} \|u\|_{\lambda_i}^2 - \frac{1}{2^*} \int_{\mathbb{R}^N} |u|^{2^*} dx, \quad i = 1, 2;$$

see [33]. There are also many papers working on related equations with a Hardy type potential and a critical nonlinearity; we refer the readers to [2, 15, 29] and the references therein.

We call a solution  $(u, v)$  of (1.4) *nontrivial* if both  $u \not\equiv 0$  and  $v \not\equiv 0$ ; we call a solution  $(u, v)$  *positive* if both  $u > 0$  and  $v > 0$  in  $\mathbb{R}^N \setminus \{0\}$ ; we call a solution  $(u, v)$  *semi-trivial* if  $(u, v)$  is a type of  $(u, 0)$  or  $(0, v)$ .

One of the difficulties in the study of (1.4) is that it has semi-trivial solutions  $(z_\mu^1, 0)$  and  $(0, z_\mu^2)$ . Here, we are only interested in nontrivial solutions of (1.4). Define  $\mathbb{D} := D^{1,2}(\mathbb{R}^N) \times D^{1,2}(\mathbb{R}^N)$  with the norm

$$\|(u, v)\|_{\mathbb{D}}^2 := \|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2.$$

Then nontrivial solutions of (1.4) can be found as nontrivial critical points of the  $C^1$  functional  $J_\nu : \mathbb{D} \rightarrow \mathbb{R}$ , where

$$J_\nu(u, v) := I_{\lambda_1}(u) + I_{\lambda_2}(v) - \nu \int_{\mathbb{R}^N} |u|^\alpha |v|^\beta dx.$$

Another difficulty is the failure of the Palais-Smale condition, which makes the study of (1.4) very tough. Since (1.4) is invariant under the transformation  $(u(x), v(x)) \mapsto (\mu^{\frac{N-2}{2}} u(\mu x), \mu^{\frac{N-2}{2}} v(\mu x))$ , where  $\mu > 0$ , it is easy to see that the Palais-Smale condition (*PS* condition for short) cannot hold for any energy level  $c > 0$ . In fact, assume by contradiction that the *PS* condition holds for some  $c > 0$ , and let  $(u_n, v_n)$  be a  $(PS)_c$  sequence; that is,  $J_\nu(u_n, v_n) \rightarrow c$  and  $J'_\nu(u_n, v_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then up to a subsequence, we may assume that  $(u_n, v_n) \rightarrow (u, v)$  strongly in  $\mathbb{D}$ . Define  $(\tilde{u}_n(x), \tilde{v}_n(x)) := (n^{\frac{N-2}{2}} u_n(nx), n^{\frac{N-2}{2}} v_n(nx))$ ; then it is easy to check that  $(\tilde{u}_n, \tilde{v}_n)$  is also a  $(PS)_c$  sequence and  $(\tilde{u}_n, \tilde{v}_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ . Since the  $(PS)_c$  condition holds, we have  $(\tilde{u}_n, \tilde{v}_n) \rightarrow (0, 0)$  strongly in  $\mathbb{D}$ , which contradicts with  $c > 0$ .

**Definition 1.1.** We say a solution  $(u_0, v_0)$  of (1.4) is a ground state solution if  $(u_0, v_0)$  is nontrivial and  $J_\nu(u_0, v_0) \leq J_\nu(u, v)$  for any other nontrivial solution  $(u, v)$  of (1.4).

To obtain ground state solutions of (1.4), as in [22], we define

$$(1.15) \quad \mathcal{N}_\nu := \left\{ (u, v) \in \mathbb{D} : u \not\equiv 0, v \not\equiv 0, \|u\|_{\lambda_1}^2 = \int_{\mathbb{R}^N} (|u|^{2^*} + \nu\alpha |u|^\alpha |v|^\beta), \right. \\ \left. \|v\|_{\lambda_2}^2 = \int_{\mathbb{R}^N} (|v|^{2^*} + \nu\beta |u|^\alpha |v|^\beta) \right\}.$$

Then any nontrivial solution of (1.4) has to belong to  $\mathcal{N}_\nu$ . Take  $\varphi, \psi \in C_0^\infty(\mathbb{R}^N)$  with  $\varphi, \psi \not\equiv 0$  and  $supp(\varphi) \cap supp(\psi) = \emptyset$ ; then there exist  $t_1, t_2 > 0$  such that  $(t_1\varphi, t_2\psi) \in \mathcal{N}_\nu$  for any  $\nu \neq 0$ . So  $\mathcal{N}_\nu \neq \emptyset$ . We set

$$(1.16) \quad c_\nu := \inf_{(u,v) \in \mathcal{N}_\nu} J_\nu(u, v) = \inf_{(u,v) \in \mathcal{N}_\nu} \frac{1}{N} (\|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2).$$

By (1.11) we have

$$(1.17) \quad \|u\|_{\lambda_i}^2 \geq S(\lambda_i) |u|_{2^*}^2, \quad \forall u \in D^{1,2}(\mathbb{R}^N), \quad i = 1, 2.$$

Then it is easy to see that  $c_\nu > 0$  for all  $\nu$ . Moreover, if  $(u_0, v_0)$  is a non-trivial solution satisfying  $J_\nu(u_0, v_0) = c_\nu$ , then  $(u_0, v_0)$  is a ground state solution. Our first result is concerned with ground state solutions with energy below  $\frac{1}{N} \min\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ .

**Theorem 1.1.** *Assume that  $N \geq 3$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$  and (1.5) holds.*

- (1) *If  $\nu < 0$ , then  $c_\nu \equiv \frac{1}{N}S(\lambda_1)^{N/2} + \frac{1}{N}S(\lambda_2)^{N/2}$ , and  $c_\nu$  cannot be attained.*
- (2) *Let*

$$(1.18) \quad \nu_0 := \frac{1}{2^*} \left[ \left( 1 + \max \left\{ \frac{\Lambda_N - \lambda_1}{\Lambda_N - \lambda_2}, \frac{\Lambda_N - \lambda_2}{\Lambda_N - \lambda_1} \right\} \right)^{\frac{2^*}{2}} - 2 \right] > 0;$$

*then for all  $\nu > \nu_0$ , (1.4) has a positive ground state solution  $(u_\nu, v_\nu) \in \mathbb{D}$ , which is radially symmetric and satisfies*

$$(1.19) \quad J_\nu(u_\nu, v_\nu) = c_\nu < \frac{1}{N} \min \left\{ S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2} \right\}.$$

- (3) *If one of the conditions*

- (C<sub>1</sub>)  $N \geq 5$  and  $\max\{\alpha, \beta\} < 2$ ,
- (C<sub>2</sub>)  $\lambda_1 \leq \lambda_2$  and  $\alpha < 2$ ,
- (C<sub>3</sub>)  $\lambda_2 \leq \lambda_1$  and  $\beta < 2$

*holds, then for all  $\nu > 0$ , (1.4) has a positive ground state solution  $(u_\nu, v_\nu) \in \mathbb{D}$ , which is radially symmetric and satisfies (1.19).*

Now we want to obtain ground state solutions with energy above the value  $\frac{1}{N} \max\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ , which seems much more interesting to us. To this goal, by Theorem 1.1 (2)-(3) we have to assume that  $\min\{\alpha, \beta\} \geq 2$  and  $\nu > 0$  is small. In this case, since  $4 \leq \alpha + \beta = 2^*$ , then  $N = 3$  or  $N = 4$ . Moreover, if  $N = 4$ , then we must have  $\alpha = \beta = 2$ . Note that if  $N = 4$  and  $\alpha = \beta = 2$ , then (1.4) turns to be the cubic system

$$(1.20) \quad \begin{cases} -\Delta u - \frac{\lambda_1}{|x|^2} u = u^3 + 2\nu uv^2, & x \in \mathbb{R}^4, \\ -\Delta v - \frac{\lambda_2}{|x|^2} v = v^3 + 2\nu u^2 v, & x \in \mathbb{R}^4, \\ u, v \in D^{1,2}(\mathbb{R}^4), \quad u, v > 0 \text{ in } \mathbb{R}^4 \setminus \{0\}, \end{cases}$$

which is just the Bose-Einstein condensate type system (1.2) with  $V_i(x) = -\frac{\lambda_i}{|x|^2}$  in the critical case  $N = 4$ . Note that  $\Lambda_4 = 1$ . Then we have the following results.

**Theorem 1.2.** *Assume that  $N = 4$ ,  $\alpha = \beta = 2$  and  $\lambda_1, \lambda_2 \in (0, 1)$ . Define*

$$(1.21) \quad \nu_1 := \min \frac{1}{2} \left\{ \frac{1 - \lambda_1}{1 - \lambda_2}, \frac{1 - \lambda_2}{1 - \lambda_1}, \frac{(1 - \lambda_1)^{\frac{3}{4}}(1 - \lambda_2)^{\frac{3}{4}}}{(1 - \lambda_1)^{\frac{3}{2}} + (1 - \lambda_2)^{\frac{3}{2}}} \right\}.$$

*Then for any  $\nu \in (0, \nu_1)$ , (1.20) has a positive ground state solution  $(u_\nu, v_\nu) \in \mathbb{D}$ , which satisfies*

$$(1.22) \quad c_\nu = J_\nu(u_\nu, v_\nu) \rightarrow \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2), \quad \text{as } \nu \rightarrow 0.$$

**Theorem 1.3.** *Assume that  $N = 3$ ,  $\alpha + \beta = 2^*$ ,  $\alpha \geq 2$ ,  $\beta \geq 2$  and  $\lambda_1, \lambda_2 \in (0, \Lambda_3)$ . Then there exists  $\tilde{\nu}_1 > 0$  such that for any  $\nu \in (0, \tilde{\nu}_1)$ , (1.4) has a positive ground state solution  $(u_\nu, v_\nu) \in \mathbb{D}$ , which satisfies*

$$(1.23) \quad J_\nu(u_\nu, v_\nu) \rightarrow \frac{1}{3} \left( S(\lambda_1)^{3/2} + S(\lambda_2)^{3/2} \right), \quad \text{as } \nu \rightarrow 0.$$

*Remark 1.1.* (1) (1.22)-(1.23) yield  $c_\nu > \frac{1}{N} \max\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$  for  $\nu > 0$  small appropriately. That is, we obtain positive ground state solutions with energy above  $\frac{1}{N} \max\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ , so the case  $\min\{\alpha, \beta\} \geq 2$  is completely different from the cases studied in Theorem 1.1-(3). As we can see in the following sections, the case  $\min\{\alpha, \beta\} \geq 2$  is much more complicated (see Theorem 2.1 for an example). Besides, if  $\lambda_1 = \lambda_2$ , then Theorem 1.2 will be improved by Theorem 7.1 in Section 7.

(2) The case  $N = 3$  is much tougher than the case  $N = 4$ , and we cannot give an accurate definition of  $\tilde{v}_1$  in Theorem 1.3 as (1.21) unfortunately. As we will see in the following sections, the idea of proving Theorem 1.2 takes full use of the fact that  $\alpha = \beta = 2$  and thus cannot be used in the case  $N = 3$ . Meanwhile, the idea of proving Theorem 1.3 is quite different and more general.

As we will see in Section 2, the radial symmetry of ground state solutions obtained in Theorem 1.1 is an easy corollary of the Schwartz symmetrization. However, the Schwartz symmetrization cannot be used to prove the radial symmetry of ground state solutions obtained in Theorems 1.2 and 1.3. Here, to get the radial symmetry of solutions obtained in Theorems 1.2 and 1.3, we will use the moving planes method. Precisely, we have the following result.

**Theorem 1.4.** *Assume that  $N = 3$  or  $N = 4$ ,  $\alpha + \beta = 2^*$ ,  $\alpha \geq 2$ ,  $\beta \geq 2$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$  and  $\nu > 0$ . Then any positive solution of (1.4) is radially symmetric with respect to the origin. Therefore, ground state solutions  $(u_\nu, v_\nu)$  obtained in Theorems 1.2 and 1.3 are radially symmetric.*

There are some other special cases, such as the case in which  $N = 3, 4, 5$ ,  $1 < \alpha < 2 \leq \beta$ ,  $\alpha + \beta = 2^*$ ,  $\lambda_2 < \lambda_1$  and  $\nu > 0$  sufficiently small, where we have no idea whether the ground state solutions exist or not. This remains to be an interesting open question. Here we can obtain positive solutions for these cases if  $\nu > 0$  is sufficiently small. Precisely, we have the following result, which plays a crucial role in the proof of Theorem 1.3.

**Theorem 1.5.** *Assume that  $N \geq 3$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$  and (1.5) hold. Then there exists  $\nu_2 > 0$  such that for any  $\nu \in (0, \nu_2]$ , (1.4) has a positive solution  $(u_\nu, v_\nu) \in \mathbb{D}$ , which is radially symmetric with respect to the origin and satisfies*

$$(1.24) \quad J_\nu(u_\nu, v_\nu) < \frac{1}{N} \left( S(\lambda_1)^{\frac{N}{2}} + S(\lambda_2)^{\frac{N}{2}} \right).$$

We should mention that Abdellaoui, Felli and Peral [1] studied the following class of weakly coupled nonlinear elliptic equations:

$$(1.25) \quad \begin{cases} -\Delta u - \frac{\lambda_1}{|x|^2} u = u^{2^*-1} + \nu h(x) \alpha u^{\alpha-1} v^\beta, & x \in \mathbb{R}^N, \\ -\Delta v - \frac{\lambda_2}{|x|^2} v = v^{2^*-1} + \nu h(x) \beta u^\alpha v^{\beta-1}, & x \in \mathbb{R}^N, \\ u, v \in D^{1,2}(\mathbb{R}^N), \quad u, v > 0 \text{ in } \mathbb{R}^N \setminus \{0\}. \end{cases}$$

Note that if  $h(x) \equiv 1$ , then (1.25) turns to be (1.4). For the case (1.5), they assumed the following condition on  $h(x)$ :

**(H<sub>1</sub>)**  $h \in L^\infty(\mathbb{R}^N)$ ,  $h \geq 0$ ,  $h \not\equiv 0$ ,  $h$  is continuous in a neighborhood of 0 and  $\infty$ , and  $h(0) = \lim_{|x| \rightarrow \infty} h(x) = 0$ .

Then they proved some existence results of ground state solutions for (1.25) in the case  $\nu > 0$  (see Section 4 in [1]). We should point out that, under condition  $(\mathbf{H}_1)$ , the Palais-Smale condition holds for energy level  $c$  with

$$(1.26) \quad c < \frac{1}{N} \min \left\{ S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2} \right\}$$

(see [1, Lemmas 4.1 and 4.3]), which plays a crucial role in obtaining ground state solutions in [1]. Therefore, *problem (1.4) is completely different from (1.25)*. Moreover, there are no results about the existence of ground state solutions to (1.25) with energy above  $\frac{1}{N} \max\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$  in [1]. It was only pointed out in [1, Remark 4.6] that for the case where  $h(x) \equiv 1$ ,  $\nu > 0$ ,  $\alpha + \beta = 2^*$  and  $\lambda_1 = \lambda_2 = \lambda$ , it is easy to construct, by a direct computation, positive solutions to (1.4) of the form  $(\phi, c\phi)$ ,  $c > 0$ . We remark that whether these solutions  $(\phi, c\phi)$  are ground state solutions is not known in [1], and there are not any conclusions about (1.4) for the general case  $\lambda_1 \neq \lambda_2$  in [1].

The rest of this paper proves these theorems, and we give some notation here. In the sequel, we denote positive constants (possibly different in different places) by  $C, C_1, C_2, \dots$ , and  $B(x, r) := \{y \in \mathbb{R}^N : |x - y| < r\}$ . Denote  $B_r := B(0, r)$  for convenience. The paper is organized as follows.

We give the proof of Theorem 1.1 in Section 2, where we will use the concentration-compactness principle from Lions ([23, 24]) and some ideas from [1]. The proof of Theorem 1.2 is given in Section 3, where we will borrow some ideas from [33] and the authors' paper [12].

In Section 4, we will prove Theorem 1.5 via a perturbation method, where we will use some ideas from Byeon and Jeanjean [9]. In order to construct a spike solution of the nonlinear elliptic problem

$$-\varepsilon^2 \Delta u + V(x)u = f(u), \quad u \in H^1(\mathbb{R}^N),$$

for a general subcritical nonlinearity  $f(u)$  and sufficiently small  $\varepsilon > 0$ , Byeon and Jeanjean [9] developed a new variational approach. We will mainly follow this variational approach to prove Theorem 1.5. Note that this approach cannot be used directly, and we need some crucial modifications for our proof. For example, we will define a special mountain-pass value  $a_\nu$ , where all paths are required to be bounded in  $\mathbb{D}$  by the same constant which is independent of  $\nu$ . This special  $a_\nu$  is essential to our proof. Moreover, we should point out that the lack of compactness is the main difficulty because of the failure of the  $(PS)$  condition of (1.4), and especially because  $Z_i, i = 1, 2$ , are not compact in  $\mathbb{D}$ .

In Section 5, we will prove Theorem 1.3 with the help of Theorem 1.5. Here, quite different ideas are needed compared to those of proving Theorem 1.2 in Section 3.

In Section 6, we will prove Theorem 1.4 via the moving planes method. The moving planes method has been used by many authors to prove symmetry and monotonicity of positive solutions to various nonlinear elliptic problems; we refer the readers to [10, 11, 17, 18] and the references therein.

Finally, by following some arguments from the authors' papers [12, 13], we will give some remarks for the special case  $\lambda_1 = \lambda_2$ ,  $\alpha = \beta = 2^*/2$  and  $\nu > 0$  in Section 7, where some uniqueness results about the ground state solutions will be obtained; see Theorems 7.1 and 7.2. In the authors' papers [12, 13], we studied the following

Bose-Einstein condensation system for the critical case:

$$(1.27) \quad \begin{cases} -\Delta u + \lambda_1 u = \mu_1 u^{2^*-1} + \nu u^{\frac{2^*}{2}-1} v^{\frac{2^*}{2}}, & x \in \Omega, \\ -\Delta v + \lambda_2 v = \mu_2 v^{2^*-1} + \nu v^{\frac{2^*}{2}-1} u^{\frac{2^*}{2}}, & x \in \Omega, \\ u \geq 0, v \geq 0 \text{ in } \Omega, \quad u = v = 0 \text{ on } \partial\Omega. \end{cases}$$

Here,  $\Omega \subset \mathbb{R}^N (N \geq 4)$  is a smooth bounded domain and  $\mu_1, \mu_2 > 0$  (the special case  $N = 4$  was studied in [12], and the general case  $N \geq 5$  in [13]). When  $\nu = 0$ , (1.27) turns to be the well-known Brezis-Nirenberg problem ([8]). Thanks to the celebrated idea from Brezis and Nirenberg [8], we can show that the (PS) condition of (1.27) holds for some ranges of energy level (see [12, 13] for details). Therefore, problem (1.4) is also completely different from (1.27). Fortunately, some ideas of studying (1.27) in [12, 13] can be used in this paper.

## 2. PROOF OF THEOREM 1.1

### 2.1. The case $\nu < 0$ .

**Lemma 2.1.** *If  $c_\nu$  is attained by a couple  $(u, v) \in \mathcal{N}_\nu$ , then this couple is a critical point of  $J_\nu$ , provided  $\nu < 0$ .*

*Proof.* This proof is standard. Let  $\nu < 0$ . Assume that  $(u, v) \in \mathcal{N}_\nu$  such that  $c_\nu = J_\nu(u, v)$ . Define

$$G_1(u, v) = \|u\|_{\lambda_1}^2 - \int_{\mathbb{R}^N} (|u|^{2^*} + \nu\alpha|u|^\alpha|v|^\beta),$$

$$G_2(u, v) = \|v\|_{\lambda_2}^2 - \int_{\mathbb{R}^N} (|v|^{2^*} + \nu\beta|u|^\alpha|v|^\beta).$$

Then there exist two Lagrange multipliers  $K_1, K_2 \in \mathbb{R}$  such that

$$(2.1) \quad J'_\nu(u, v) + K_1 G'_1(u, v) + K_2 G'_2(u, v) = 0.$$

Testing (2.1) with  $(u, 0)$  and  $(0, v)$  respectively, we conclude from  $(u, v) \in \mathcal{N}_\nu$  that

$$\begin{aligned} & \left( (2^* - 2)|u|_{2^*}^{2^*} + \alpha(2 - \alpha)|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) K_1 - \alpha\beta|\nu| \left( \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) K_2 = 0, \\ & \left( (2^* - 2)|v|_{2^*}^{2^*} + \beta(2 - \beta)|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) K_2 - \alpha\beta|\nu| \left( \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) K_1 = 0. \end{aligned}$$

Recall that  $|u|_{2^*}^{2^*} > \alpha|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta$  and  $|v|_{2^*}^{2^*} > \beta|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta$ ; we see from  $\alpha + \beta = 2^*$  that

$$\begin{aligned} & \left( (2^* - 2)|u|_{2^*}^{2^*} + \alpha(2 - \alpha)|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) \\ & \times \left( (2^* - 2)|v|_{2^*}^{2^*} + \beta(2 - \beta)|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right) > \left( \alpha\beta|\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right)^2. \end{aligned}$$

From this we deduce that  $K_1 = K_2 = 0$ , and so  $J'_\nu(u, v) = 0$ . □

**Lemma 2.2.** *Let  $\nu < 0$ . For any  $(u, v) \in \mathbb{D}$  with  $u \not\equiv 0$  and  $v \not\equiv 0$ , if*

$$(2.2) \quad \left( \int_{\mathbb{R}^N} |u|^{2^*} \right)^\alpha \left( \int_{\mathbb{R}^N} |v|^{2^*} \right)^\beta > \alpha^\alpha \beta^\beta \left( |\nu| \int_{\mathbb{R}^N} |u|^\alpha|v|^\beta \right)^{2^*},$$

*then there exist  $t_1 > 0, s_1 > 0$ , such that  $(t_1 u, s_1 v) \in \mathcal{N}_\nu$ .*



*Proof.* For simplicity, we denote

$$A_1 = \|u\|_{\lambda_1}^2, \quad B_1 = |u|_{2^*}^2, \quad C = |\nu| \int_{\mathbb{R}^N} |u|^\alpha |v|^\beta, \quad A_2 = \|v\|_{\lambda_2}^2, \quad B_2 = |v|_{2^*}^2.$$

Recall definition (1.15) of  $\mathcal{N}_\nu$ ; we see that  $(tu, sv) \in \mathcal{N}_\nu$  for  $t, s > 0$  is equivalent to  $t, s > 0$  satisfying

$$(2.3) \quad A_1 t^{2-\alpha} = B_1 t^\beta - \alpha C s^\beta, \quad A_2 s^{2-\beta} = B_2 s^\alpha - \beta C t^\alpha.$$

If  $C = 0$ , then it is trivial to see that (2.3) has a solution  $(t_1, s_1)$  with  $t_1, s_1 > 0$ . So we may assume that  $C > 0$ . Then the equation  $A_1 t^{2-\alpha} = B_1 t^\beta - \alpha C s^\beta$  is equivalent to

$$s = g(t) := \left( \frac{B_1 t^\beta - A_1 t^{2-\alpha}}{\alpha C} \right)^{1/\beta} > 0, \quad t > t_0 := \left( \frac{A_1}{B_1} \right)^{\frac{1}{2-\alpha}}.$$

Therefore, it suffices to prove that

$$(2.4) \quad A_2 \left( \frac{B_1 t^\beta - A_1 t^{2-\alpha}}{\alpha C} \right)^{\frac{2-\beta}{\beta}} - B_2 \left( \frac{B_1 t^\beta - A_1 t^{2-\alpha}}{\alpha C} \right)^{\frac{\alpha}{\beta}} + \beta C t^\alpha = 0$$

has a solution  $t > t_0$ . Note that (2.4) is equivalent to

$$f(t) := A_2 \left( \frac{B_1 - A_1 t^{2-2^*}}{\alpha C} \right)^{\frac{2-\beta}{\beta}} + t^{2^*-2} \left[ \beta C - B_2 \left( \frac{B_1 - A_1 t^{2-2^*}}{\alpha C} \right)^{\frac{\alpha}{\beta}} \right] = 0.$$

Note that (2.2) implies that

$$\beta C - B_2 \left( \frac{B_1}{\alpha C} \right)^{\frac{\alpha}{\beta}} < 0;$$

then it is easy to check that  $\lim_{t \searrow t_0} f(t) > 0$  and  $\lim_{t \rightarrow +\infty} f(t) = -\infty$ . So there exists  $t_1 > t_0 > 0$  such that  $f(t_1) = 0$ . Let  $s_1 = g(t_1)$ ; then  $s_1 > 0$  and  $(t_1 u, s_1 v) \in \mathcal{N}_\nu$ . This completes the proof.  $\square$

*Proof of Theorem 1.1-(1).* Fix any  $\nu < 0$ . Recall (1.9)-(1.12). It is easy to see that  $z_\mu^2 \rightharpoonup 0$  weakly in  $D^{1,2}(\mathbb{R}^N)$  and so  $(z_\mu^2)^\beta \rightharpoonup 0$  weakly in  $L^{2^*/\beta}(\mathbb{R}^N)$  as  $\mu \rightarrow +\infty$ . That is,

$$\lim_{\mu \rightarrow +\infty} |\nu| \int_{\mathbb{R}^N} (z_1^1)^\alpha (z_\mu^2)^\beta dx = 0.$$

Then (2.2) holds for  $(z_1^1, z_\mu^2)$  when  $\mu > 0$  sufficiently large, and so there exist  $t_\mu, s_\mu > 0$  such that  $(t_\mu z_1^1, s_\mu z_\mu^2) \in \mathcal{N}_\nu$ . Denote  $F_\mu := |\nu| \int_{\mathbb{R}^N} (z_1^1)^\alpha (z_\mu^2)^\beta dx$ . Then

$$(2.5) \quad t_\mu^2 S(\lambda_1)^{\frac{N}{2}} = t_\mu^{2^*} S(\lambda_1)^{\frac{N}{2}} - \alpha t_\mu^\alpha s_\mu^\beta F_\mu, \quad s_\mu^2 S(\lambda_2)^{\frac{N}{2}} = s_\mu^{2^*} S(\lambda_2)^{\frac{N}{2}} - \beta t_\mu^\alpha s_\mu^\beta F_\mu.$$

Assume that, up to a subsequence,  $t_\mu \rightarrow +\infty$  as  $\mu \rightarrow \infty$ . Then by

$$\beta(t_\mu^{2^*} - t_\mu^2) S(\lambda_1)^{\frac{N}{2}} = \alpha(s_\mu^{2^*} - s_\mu^2) S(\lambda_2)^{\frac{N}{2}}$$

we also have  $s_\mu \rightarrow +\infty$ . Then

$$t_\mu^{2^*} - t_\mu^2 \geq \frac{1}{2} t_\mu^{2^*}, \quad s_\mu^{2^*} - s_\mu^2 \geq \frac{1}{2} s_\mu^{2^*} \quad \text{for } \mu \text{ large enough.}$$

Combining this with (2.5) we see that

$$\left( \frac{t_\mu}{s_\mu} \right)^\beta \leq 2\alpha S(\lambda_1)^{-\frac{N}{2}} F_\mu \rightarrow 0, \quad \left( \frac{s_\mu}{t_\mu} \right)^\alpha \leq 2\beta S(\lambda_2)^{-\frac{N}{2}} F_\mu \rightarrow 0, \quad \text{as } \mu \rightarrow +\infty,$$

a contradiction. Therefore,  $t_\mu$  and  $s_\mu$  are uniformly bounded. Then by (2.5) and  $F_\mu \rightarrow 0$  as  $\mu \rightarrow \infty$ , we get that  $\lim_{\mu \rightarrow +\infty} (t_\mu, s_\mu) = (1, 1)$ . Note that  $(t_\mu z_1^1, s_\mu z_\mu^2) \in \mathcal{N}_\nu$ , and we see from (1.12) and (1.16) that

$$\begin{aligned} c_\nu &\leq J_\nu(t_\mu z_1^1, s_\mu z_\mu^2) = \frac{1}{N} (t_\mu^2 \|z_1^1\|_{\lambda_1}^2 + s_\mu^2 \|z_\mu^2\|_{\lambda_2}^2) \\ &= \frac{1}{N} (t_\mu^2 S(\lambda_1)^{N/2} + s_\mu^2 S(\lambda_2)^{N/2}). \end{aligned}$$

Letting  $\mu \rightarrow +\infty$ , we get that  $c_\nu \leq \frac{1}{N} (S(\lambda_1)^{N/2} + S(\lambda_2)^{N/2})$ . On the other hand, for any  $(u, v) \in \mathcal{N}_\nu$ , we see from  $\nu < 0$  and (1.17) that

$$\|u\|_{\lambda_1}^2 \leq \int_{\mathbb{R}^N} |u|^{2^*} dx \leq S(\lambda_1)^{-2^*/2} \|u\|_{\lambda_1}^{2^*},$$

and so  $\|u\|_{\lambda_1}^2 \geq S(\lambda_1)^{N/2}$ . Similarly,  $\|v\|_{\lambda_2}^2 \geq S(\lambda_2)^{N/2}$ . Combining these with (1.16), we get that  $c_\nu \geq \frac{1}{N} (S(\lambda_1)^{N/2} + S(\lambda_2)^{N/2})$ . Hence,

$$(2.6) \quad c_\nu = \frac{1}{N} (S(\lambda_1)^{N/2} + S(\lambda_2)^{N/2}).$$

Now, assume that  $c_\nu$  is attained by some  $(u, v) \in \mathcal{N}_\nu$ ; then  $(|u|, |v|) \in \mathcal{N}_\nu$  and  $J_\nu(|u|, |v|) = c_\nu$ . By Lemma 2.1, we know that  $(|u|, |v|)$  is a nontrivial solution of (1.4). By the maximum principle, we may assume that  $u > 0, v > 0$  in  $\mathbb{R}^N \setminus \{0\}$ , and so  $\int_{\mathbb{R}^N} u^\alpha v^\beta dx > 0$ . Then

$$\|u\|_{\lambda_1}^2 < \int_{\mathbb{R}^N} |u|^{2^*} dx \leq S(\lambda_1)^{-2^*/2} \|u\|_{\lambda_1}^{2^*}.$$

Therefore, it is easy to see that  $c_\nu = J_\nu(u, v) > \frac{1}{N} (S(\lambda_1)^{N/2} + S(\lambda_2)^{N/2})$ , which is a contradiction. This completes the proof. □

**2.2. The case  $\nu > 0$ .** In this subsection, we let  $\nu > 0$ . Define

$$(2.7) \quad c'_\nu := \inf_{(u,v) \in \mathcal{N}'_\nu} J_\nu(u, v),$$

where

$$(2.8) \quad \mathcal{N}'_\nu := \left\{ (u, v) \in D \setminus \{(0, 0)\} : J'_\nu(u, v)(u, v) = 0 \right\}.$$

Note that  $\mathcal{N}_\nu \subset \mathcal{N}'_\nu$ , so  $c'_\nu \leq c_\nu$ . By (1.17) it is easy to prove that  $c'_\nu > 0$ . Moreover, it is standard to prove that

$$\begin{aligned} (2.9) \quad c'_\nu &= \inf_{(u,v) \in \mathbb{D} \setminus \{(0,0)\}} \max_{t>0} J_\nu(tu, tv) \\ &= \inf_{(u,v) \in \mathbb{D} \setminus \{(0,0)\}} \frac{1}{N} \left[ \frac{\|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2}{\left( \int_{\mathbb{R}^N} |u|^{2^*} + 2^* \nu |u|^\alpha |v|^\beta + |v|^{2^*} \right)^{\frac{2}{2^*}}} \right]^{\frac{N}{2}}. \end{aligned}$$

Define  $E(u, v) := |\nabla u|^2 + |\nabla v|^2 - \frac{\lambda_1}{|x|^2} |u|^2 - \frac{\lambda_2}{|x|^2} |v|^2$  and  $F(u, v) := |u|^{2^*} + 2^* \nu |u|^\alpha |v|^\beta + |v|^{2^*}$  for simplicity; then

$$(2.10) \quad \int_{\mathbb{R}^N} E(u, v) dx \geq (Nc'_\nu)^{\frac{2}{N}} \left( \int_{\mathbb{R}^N} F(u, v) dx \right)^{\frac{2}{2^*}}, \quad \forall (u, v) \in \mathbb{D}.$$

The following lemma is the counterpart of the Brezis-Lieb Lemma ([7]) for  $(u, v)$ , and the idea of its proof comes from [7] (see also [35, Lemma 1.32]).

**Lemma 2.3.** *Let  $\Omega \subset \mathbb{R}^N$  be an open set and  $(u_n, v_n)$  be a bounded sequence in  $L^{2^*}(\Omega) \times L^{2^*}(\Omega)$ . If  $(u_n, v_n) \rightarrow (u, v)$  almost everywhere in  $\Omega$ , then*

$$\lim_{n \rightarrow \infty} \int_{\Omega} (|u_n|^\alpha |v_n|^\beta - |u_n - u|^\alpha |v_n - v|^\beta) \, dx = \int_{\Omega} |u|^\alpha |v|^\beta \, dx.$$

*Proof.* Fatou Lemma yields

$$\int_{\Omega} |u|^{2^*} \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |u_n|^{2^*} < \infty, \quad \int_{\Omega} |v|^{2^*} \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |v_n|^{2^*} < \infty.$$

Recall that  $\alpha, \beta$  satisfy (1.5). For any  $a_1, a_2, b_1, b_2 \in \mathbb{R}$  and  $\varepsilon > 0$ , we deduce from the mean value theorem and Young’s inequality that

$$\begin{aligned} & \left| |a_1 + a_2|^\alpha |b_1 + b_2|^\beta - |a_1|^\alpha |b_1|^\beta \right| \\ & \leq \left| |a_1 + a_2|^\alpha - |a_1|^\alpha \right| |b_1 + b_2|^\beta + |a_1|^\alpha \left| |b_1 + b_2|^\beta - |b_1|^\beta \right| \\ & \leq C \left[ (|a_1| + |a_2|)^{\alpha-1} (|b_1| + |b_2|)^\beta |a_2| + |a_1|^\alpha (|b_1| + |b_2|)^{\beta-1} |b_2| \right] \\ & \leq C\varepsilon \left[ (|a_1| + |a_2|)^{2^*} + (|b_1| + |b_2|)^{2^*} \right] + C\varepsilon^{1-2^*} \left( |a_2|^{2^*} + |b_2|^{2^*} \right) \\ & \leq C\varepsilon \left( |a_1|^{2^*} + |a_2|^{2^*} + |b_1|^{2^*} + |b_2|^{2^*} \right) + C\varepsilon^{1-2^*} \left( |a_2|^{2^*} + |b_2|^{2^*} \right). \end{aligned}$$

Denote  $\omega_n = u_n - u$  and  $\sigma_n = v_n - v$ . Then

$$\begin{aligned} f_n^\varepsilon & := \left[ \left| |u_n|^\alpha |v_n|^\beta - |\omega_n|^\alpha |\sigma_n|^\beta - |u|^\alpha |v|^\beta \right| \right. \\ & \quad \left. - C\varepsilon \left( |\omega_n|^{2^*} + |u|^{2^*} + |\sigma_n|^{2^*} + |v|^{2^*} \right) \right]_+ \\ & \leq |u|^\alpha |v|^\beta + C\varepsilon^{1-2^*} \left( |u|^{2^*} + |v|^{2^*} \right), \end{aligned}$$

and so the dominated convergence theorem yields  $\int_{\Omega} f_n^\varepsilon \, dx \rightarrow 0$  as  $n \rightarrow \infty$ . Note that

$$\left| |u_n|^\alpha |v_n|^\beta - |\omega_n|^\alpha |\sigma_n|^\beta - |u|^\alpha |v|^\beta \right| \leq f_n^\varepsilon + C\varepsilon \left( |\omega_n|^{2^*} + |u|^{2^*} + |\sigma_n|^{2^*} + |v|^{2^*} \right),$$

so we obtain

$$\limsup_{n \rightarrow \infty} \int_{\Omega} \left| |u_n|^\alpha |v_n|^\beta - |\omega_n|^\alpha |\sigma_n|^\beta - |u|^\alpha |v|^\beta \right| \leq C\varepsilon.$$

Since  $C > 0$  is independent of  $\varepsilon > 0$ , the proof is complete. □

The following lemma is the counterpart of Lions’ concentration-compactness principle ([23, 24]) for problem (1.4).

**Lemma 2.4.** *Let  $(u_n, v_n) \in \mathbb{D}$  be a sequence such that*

$$(2.11) \quad \begin{cases} (u_n, v_n) \rightharpoonup (u, v) & \text{weakly in } \mathbb{D}, \\ (u_n, v_n) \rightarrow (u, v) & \text{almost everywhere on } \mathbb{R}^N, \\ E(u_n - u, v_n - v) \rightharpoonup \mu & \text{in the sense of measures,} \\ F(u_n - u, v_n - v) \rightharpoonup \rho & \text{in the sense of measures.} \end{cases}$$

Define

$$\begin{aligned}
 \mu_\infty &:= \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{|x| \geq R} E(u_n, v_n) \, dx, \\
 \rho_\infty &:= \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{|x| \geq R} F(u_n, v_n) \, dx.
 \end{aligned}
 \tag{2.12}$$

Then it follows that

$$\|\mu\| \geq (Nc'_\nu)^{\frac{2}{N}} \|\rho\|^{\frac{2}{2^*}},
 \tag{2.13}$$

$$\mu_\infty \geq (Nc'_\nu)^{\frac{2}{N}} \rho_\infty^{\frac{2}{2^*}},
 \tag{2.14}$$

$$\limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} E(u_n, v_n) \, dx = \int_{\mathbb{R}^N} E(u, v) \, dx + \|\mu\| + \mu_\infty,
 \tag{2.15}$$

$$\limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n, v_n) \, dx = \int_{\mathbb{R}^N} F(u, v) \, dx + \|\rho\| + \rho_\infty.
 \tag{2.16}$$

Moreover, if  $(u, v) = (0, 0)$  and  $\|\mu\| = (Nc'_\nu)^{\frac{2}{N}} \|\rho\|^{\frac{2}{2^*}}$ , then  $\mu$  and  $\rho$  are concentrated at a single point.

*Proof.* In this proof we mainly follow the argument of [35, Lemma 1.40]. First we assume  $(u, v) = (0, 0)$ . For any  $h \in C_0^\infty(\mathbb{R}^N)$ , we see from (2.10) that

$$\int_{\mathbb{R}^N} E(hu_n, hv_n) \, dx \geq (Nc'_\nu)^{\frac{2}{N}} \left( \int_{\mathbb{R}^N} |h|^{2^*} F(u_n, v_n) \, dx \right)^{\frac{2}{2^*}}.
 \tag{2.17}$$

Since  $u_n \rightarrow 0, v_n \rightarrow 0$  in  $L^2_{loc}(\mathbb{R}^N)$ , we have

$$\int_{\mathbb{R}^N} E(hu_n, hv_n) \, dx - \int_{\mathbb{R}^N} |h|^2 E(u_n, v_n) \, dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then by letting  $n \rightarrow \infty$  in (2.17), we obtain

$$\int_{\mathbb{R}^N} |h|^2 \, d\mu \geq (Nc'_\nu)^{\frac{2}{N}} \left( \int_{\mathbb{R}^N} |h|^{2^*} \, d\rho \right)^{\frac{2}{2^*}};
 \tag{2.18}$$

that is, (2.13) holds.

For  $R > 1$ , let  $\psi_R \in C^1(\mathbb{R}^N)$  be such that  $0 \leq \psi_R \leq 1, \psi_R(x) = 1$  for  $|x| \geq R + 1$  and  $\psi_R(x) = 0$  for  $|x| \leq R$ . Then we see from (2.10) that

$$\int_{\mathbb{R}^N} E(\psi_R u_n, \psi_R v_n) \, dx \geq (Nc'_\nu)^{\frac{2}{N}} \left( \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) \, dx \right)^{\frac{2}{2^*}}.$$

Since  $u_n \rightarrow 0, v_n \rightarrow 0$  in  $L^2_{loc}(\mathbb{R}^N)$ , then

$$\begin{aligned}
 &\limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx \\
 &\geq (Nc'_\nu)^{\frac{2}{N}} \limsup_{n \rightarrow \infty} \left( \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) \, dx \right)^{\frac{2}{2^*}}.
 \end{aligned}
 \tag{2.19}$$

Note that  $\int_{|x| \geq R+1} F(u_n, v_n) \leq \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) \leq \int_{|x| \geq R} F(u_n, v_n)$ , so

$$\rho_\infty = \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) \, dx.
 \tag{2.20}$$

On the other hand,

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx \\ &= \limsup_{n \rightarrow \infty} \int_{|x| \geq R+1} E(u_n, v_n) \, dx + \limsup_{n \rightarrow \infty} \int_{R \leq |x| \leq R+1} |\psi_R|^2 E(u_n, v_n) \, dx \\ &= \limsup_{n \rightarrow \infty} \int_{|x| \geq R+1} E(u_n, v_n) \, dx + \limsup_{n \rightarrow \infty} \int_{R \leq |x| \leq R+1} |\psi_R|^2 (|\nabla u_n|^2 + |\nabla v_n|^2) \, dx \\ &\geq \limsup_{n \rightarrow \infty} \int_{|x| \geq R+1} E(u_n, v_n) \, dx. \end{aligned}$$

Letting  $R \rightarrow \infty$  we see that  $\mu_\infty \leq \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx$ . Similarly,

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx \\ &= \limsup_{n \rightarrow \infty} \int_{|x| \geq R} E(u_n, v_n) \, dx - \liminf_{n \rightarrow \infty} \int_{R \leq |x| \leq R+1} (1 - |\psi_R|^2) E(u_n, v_n) \, dx \\ &\leq \limsup_{n \rightarrow \infty} \int_{|x| \geq R} E(u_n, v_n) \, dx. \end{aligned}$$

Letting  $R \rightarrow \infty$  we see that  $\mu_\infty \geq \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx$ . Hence

$$(2.21) \quad \mu_\infty = \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^2 E(u_n, v_n) \, dx.$$

Then (2.14) follows directly from (2.19), (2.20) and (2.21).

Assume moreover that  $\|\mu\| = (Nc'_\nu)^{\frac{2}{N}} \|\rho\|^{\frac{2}{2^*}}$ . Then by Hölder inequality and (2.18), we have

$$\int_{\mathbb{R}^N} |h|^{2^*} \, d\rho \leq (Nc'_\nu)^{-\frac{2}{N-2}} \|\mu\|^{\frac{2}{N-2}} \int_{\mathbb{R}^N} |h|^{2^*} \, d\mu, \quad \forall h \in C_0^\infty(\mathbb{R}^N).$$

From this we deduce that  $\rho = (Nc'_\nu)^{-\frac{2}{N-2}} \|\mu\|^{\frac{2}{N-2}} \mu$ . So  $\mu = (Nc'_\nu)^{\frac{2}{N}} \|\rho\|^{-\frac{2}{N}} \rho$ , and we see from (2.18) that

$$\|\rho\|^{\frac{2}{N}} \left( \int_{\mathbb{R}^N} |h|^{2^*} \, d\rho \right)^{\frac{2}{2^*}} \leq \int_{\mathbb{R}^N} |h|^2 \, d\rho, \quad \forall h \in C_0^\infty(\mathbb{R}^N).$$

That is, for each open set  $\Omega$ , we have  $\rho(\Omega)^{\frac{2}{2^*}} \rho(\mathbb{R}^N)^{\frac{2}{N}} \leq \rho(\Omega)$ . Therefore,  $\rho$  is concentrated at a single point.

For the general case, we denote  $\omega_n = u_n - u$  and  $\sigma_n = v_n - v$ ; then  $(\omega_n, \sigma_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ . From the Brezis-Lieb Lemma ([7]) and Lemma 2.3, we obtain for nonnegative  $h \in C_0(\mathbb{R}^N)$  that

$$\begin{aligned} \int_{\mathbb{R}^N} hE(u, v) \, dx &= \lim_{n \rightarrow \infty} \left( \int_{\mathbb{R}^N} hE(u_n, v_n) \, dx - \int_{\mathbb{R}^N} hE(\omega_n, \sigma_n) \, dx \right), \\ \int_{\mathbb{R}^N} hF(u, v) \, dx &= \lim_{n \rightarrow \infty} \left( \int_{\mathbb{R}^N} hF(u_n, v_n) \, dx - \int_{\mathbb{R}^N} hF(\omega_n, \sigma_n) \, dx \right), \end{aligned}$$

so

(2.22)

$$E(u_n, v_n) \rightharpoonup E(u, v) + \mu, \quad F(u_n, v_n) \rightharpoonup F(u, v) + \rho, \quad \text{in the sense of measures.}$$

Inequality (2.13) follows from the corresponding one for  $(\omega_n, \sigma_n)$ . From the Brezis-Lieb Lemma ([7]) and Lemma 2.3 again, it is easy to prove that

$$\begin{aligned} \mu_\infty &:= \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{|x| \geq R} E(\omega_n, \sigma_n) \, dx, \\ \rho_\infty &:= \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{|x| \geq R} F(\omega_n, \sigma_n) \, dx, \end{aligned}$$

and so inequality (2.14) follows from the corresponding one for  $(\omega_n, \sigma_n)$ . For any  $R > 1$ , we deduce from (2.22) that

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n, v_n) \\ &= \limsup_{n \rightarrow \infty} \left( \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) + \int_{\mathbb{R}^N} (1 - |\psi_R|^{2^*}) F(u_n, v_n) \right) \\ &= \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\psi_R|^{2^*} F(u_n, v_n) + \int_{\mathbb{R}^N} (1 - |\psi_R|^{2^*}) F(u, v) + \int_{\mathbb{R}^N} (1 - |\psi_R|^{2^*}) \, d\rho. \end{aligned}$$

Letting  $R \rightarrow \infty$ , we see from (2.20) that (2.16) holds. The proof of (2.15) is similar. This completes the proof.  $\square$

**Lemma 2.5.** *Let  $\nu > 0$ . Then (1.4) has a solution  $(u, v) \in \mathbb{D} \setminus \{(0, 0)\}$  (maybe semi-trivial) such that  $J_\nu(u, v) = c'_\nu$  and  $u, v \geq 0$  are radially symmetric with respect to the origin. Moreover, if  $c'_\nu < \frac{1}{N} \min\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ , then  $(u, v) \in \mathbb{D}$  is a positive ground state solution of (1.4), and  $c_\nu = c'_\nu = J_\nu(u, v)$ .*

*Proof.* For  $(u, v) \in \mathcal{N}'_\nu$  with  $u \geq 0, v \geq 0$ , we denote by  $(u^*, v^*)$  its Schwartz symmetrization. Then by the properties of Schwartz symmetrization (see [21] for example), we see from  $\lambda_1, \lambda_2, \nu > 0$  that

$$\int_{\mathbb{R}^N} (|\nabla u^*|^2 + |\nabla v^*|^2 - \frac{\lambda_1}{|x|^2} |u^*|^2 - \frac{\lambda_2}{|x|^2} |v^*|^2) \leq \int_{\mathbb{R}^N} (|u^*|^{2^*} + 2^* \nu |u^*|^\alpha |v^*|^\beta + |v^*|^{2^*}).$$

Therefore, there exists  $0 < t^* \leq 1$  such that  $(t^* u^*, t^* v^*) \in \mathcal{N}'_\nu$ , and then

$$\begin{aligned} J_\nu(t^* u^*, t^* v^*) &= \frac{1}{N} (t^*)^2 (\|u^*\|_{\lambda_1}^2 + \|v^*\|_{\lambda_2}^2) \\ (2.23) \qquad \qquad &\leq \frac{1}{N} (\|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2) = J_\nu(u, v). \end{aligned}$$

Therefore, we may take a minimizing sequence  $(\tilde{u}_n, \tilde{v}_n) \in \mathcal{N}'_\nu$  of  $c'_\nu$  such that  $(\tilde{u}_n, \tilde{v}_n) = (\tilde{u}_n^*, \tilde{v}_n^*)$  and  $J_\nu(\tilde{u}_n, \tilde{v}_n) \rightarrow c'_\nu$  as  $n \rightarrow \infty$ . Define the Lévy concentration functions

$$Q_n(R) := \sup_{y \in \mathbb{R}^N} \int_{B(y, R)} F(\tilde{u}_n, \tilde{v}_n) \, dx.$$

Since  $\tilde{u}_n, \tilde{v}_n \geq 0$  are radially symmetric nonincreasing, one has that  $Q_n(R) = \int_{B(0, R)} F(\tilde{u}_n, \tilde{v}_n) \, dx$ . Then there exists  $R_n > 0$  such that

$$Q_n(R_n) = \int_{B(0, R_n)} F(\tilde{u}_n, \tilde{v}_n) \, dx = \frac{1}{2} \int_{\mathbb{R}^N} F(\tilde{u}_n, \tilde{v}_n) \, dx.$$

Define

$$(u_n(x), v_n(x)) := \left( R_n^{\frac{N-2}{2}} \tilde{u}_n(R_n x), R_n^{\frac{N-2}{2}} \tilde{v}_n(R_n x) \right).$$

Then by a direct computation, we see that  $(u_n, v_n) \in \mathcal{N}'_\nu$ ,  $J_\nu(u_n, v_n) \rightarrow c'_\nu$ ,  $u_n, v_n \geq 0$  are radially symmetric nonincreasing, and

$$(2.24) \quad \int_{B(0,1)} F(u_n, v_n) dx = \frac{1}{2} \int_{\mathbb{R}^N} F(u_n, v_n) dx = \sup_{y \in \mathbb{R}^N} \int_{B(y,1)} F(u_n, v_n) dx.$$

From (2.23) we know that  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}$ . Then passing to a subsequence, there exist  $(u, v) \in \mathbb{D}$  and finite measures  $\mu, \rho$  such that (2.11) holds. Define  $\mu_\infty, \rho_\infty$  as in (2.12); then by Lemma 2.4 we see that (2.13)-(2.16) hold. Note that

$$(2.25) \quad \|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2 = \int_{\mathbb{R}^N} F(u_n, v_n) dx \rightarrow Nc'_\nu, \quad \text{as } n \rightarrow \infty.$$

We conclude from (2.13)-(2.16) and (2.10) that

$$\begin{aligned} Nc'_\nu &= \int_{\mathbb{R}^N} F(u, v) dx + \|\rho\| + \rho_\infty, \\ Nc'_\nu &\geq (Nc'_\nu)^{\frac{2}{N}} \left[ \left( \int_{\mathbb{R}^N} F(u, v) dx \right)^{\frac{2}{2^*}} + \|\rho\|^{\frac{2}{2^*}} + \rho_\infty^{\frac{2}{2^*}} \right]. \end{aligned}$$

Therefore,  $\int_{\mathbb{R}^N} F(u, v) dx$ ,  $\|\rho\|$  and  $\rho_\infty$  are equal either to 0 or to  $Nc'_\nu$ . By (2.24)-(2.25), we have  $\rho_\infty \leq \frac{1}{2}Nc'_\nu$ , so  $\rho_\infty = 0$ . If  $\|\rho\| = Nc'_\nu$ , then one has that  $\int_{\mathbb{R}^N} F(u, v) dx = 0$ , and so  $(u, v) = (0, 0)$ . Moreover, since  $\|\mu\| \leq Nc'_\nu$ , we deduce from (2.13) that  $\|\mu\| = (Nc'_\nu)^{\frac{2}{N}} \|\rho\|^{\frac{2}{2^*}}$ . Then Lemma 2.4 implies that  $\rho$  is concentrated at a single point  $z$ , and we see from (2.24)-(2.25) that

$$\frac{1}{2}Nc'_\nu = \lim_{n \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B(y,1)} F(u_n, v_n) \geq \lim_{n \rightarrow \infty} \int_{B(z,1)} F(u_n, v_n) = \|\rho\|,$$

a contradiction. Therefore,  $\int_{\mathbb{R}^N} F(u, v) dx = Nc'_\nu$ . Since  $\|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2 \leq Nc'_\nu$ , we deduce from (2.10) and (2.25) that

$$Nc'_\nu = \|u\|_{\lambda_1}^2 + \|v\|_{\lambda_2}^2 = \int_{\mathbb{R}^N} F(u, v) dx = \lim_{n \rightarrow \infty} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2),$$

that is,  $(u_n, v_n) \rightarrow (u, v)$  strongly in  $\mathbb{D}$ ,  $(u, v) \in \mathcal{N}'_\nu$  and  $J_\nu(u, v) = c'_\nu$ . Recall that  $c'_\nu > 0$ , so  $(u, v) \neq (0, 0)$ . By definition (2.5) of  $\mathcal{N}'_\nu$  and using the Lagrange multiplier method, it is standard to prove that  $J'_\nu(u, v) = 0$ , so  $(u, v)$  is a solution of (1.4). Moreover,  $u, v \geq 0$  are radially symmetric.

Now, assume that  $c'_\nu < \frac{1}{N} \min\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ . Then it is easy to prove that both  $u \not\equiv 0$  and  $v \not\equiv 0$ , that is,  $(u, v) \in \mathcal{N}_\nu$ , and so  $J_\nu(u, v) = c'_\nu = c_\nu$ . Hence,  $(u, v)$  is a ground state solution of (1.4). By the maximum principle,  $u, v > 0$  in  $\mathbb{R}^N \setminus \{0\}$  and are radially symmetric. This completes the proof.  $\square$

Since  $(z_\mu^1, 0)$  and  $(0, z_\mu^2)$  belong to  $\mathcal{N}'_\nu$ , thus  $c'_\nu \leq \frac{1}{N} \min\{S(\lambda_1)^{\frac{N}{2}}, S(\lambda_2)^{\frac{N}{2}}\}$  always holds. However, the following result says that the conclusion

$$c'_\nu < \frac{1}{N} \min\{S(\lambda_1)^{\frac{N}{2}}, S(\lambda_2)^{\frac{N}{2}}\}$$

cannot always hold unfortunately.

**Theorem 2.1.** *Assume that  $\alpha, \beta \geq 2$ . Then there exists  $\tilde{\nu} > 0$  such that for all  $\nu \in (0, \tilde{\nu})$  there holds*

$$c'_\nu = \frac{1}{N} \min \left\{ S(\lambda_1)^{\frac{N}{2}}, S(\lambda_2)^{\frac{N}{2}} \right\}.$$

Moreover  $c'_\nu$  is achieved by and only by

$$\begin{cases} (0, \pm z_\mu^2), & \mu > 0, & \text{if } \lambda_1 < \lambda_2, \\ (\pm z_\mu^1, 0), & \mu > 0, & \text{if } \lambda_1 > \lambda_2, \\ (0, \pm z_\mu^2), (\pm z_\mu^1, 0), & \mu > 0, & \text{if } \lambda_1 = \lambda_2. \end{cases}$$

*Proof.* Thanks to Lemma 2.5, this proof is completely the same as that of [1, Theorem 3.4], and we omit the details here. □

*Proof of Theorem 1.1 (2)-(3).* Let  $\nu > 0$ . By Lemma 2.5 and Theorem 2.1 we know that we have to require further assumptions on  $\alpha, \beta$  and  $\nu$  to obtain positive ground state solutions with energy below  $\frac{1}{N} \min\{S(\lambda_1)^{\frac{N}{2}}, S(\lambda_2)^{\frac{N}{2}}\}$ .

Denote

$$d_1 := \frac{\Lambda_N - \lambda_2}{\Lambda_N - \lambda_1}, \quad d_2 := \frac{\Lambda_N - \lambda_1}{\Lambda_N - \lambda_2}.$$

Recall (1.18); we let  $\nu > \nu_0$ . Then

$$(2.26) \quad 1 + \max\{d_1, d_2\} < (2 + 2^* \nu)^{\frac{2}{2^*}}.$$

Without loss of generality, we may assume that  $\lambda_1 \leq \lambda_2$ . Then (1.11) yields  $S(\lambda_2) \leq S(\lambda_1)$ . By Hardy inequality (1.6) we have  $\|u\|_{\lambda_1}^2 \leq d_2 \|u\|_{\lambda_2}^2$  for all  $u \in D^{1,2}(\mathbb{R}^N)$ . Then we deduce from (1.11), (2.9) and (2.26) that

$$\begin{aligned} c'_\nu &\leq \frac{1}{N} \left[ \frac{\|z_\mu^2\|_{\lambda_1}^2 + \|z_\mu^2\|_{\lambda_2}^2}{\left(\int_{\mathbb{R}^N} |z_\mu^2|^{2^*} + 2^* \nu |z_\mu^2|^{2^*} + |z_\mu^2|^{2^*}\right)^{\frac{2}{2^*}}} \right]^{\frac{N}{2}} \\ &\leq \frac{1}{N} \left[ \frac{1 + d_2}{(2 + 2^* \nu)^{\frac{2}{2^*}}} \cdot \frac{\|z_\mu^2\|_{\lambda_2}^2}{\left(\int_{\mathbb{R}^N} |z_\mu^2|^{2^*}\right)^{\frac{2}{2^*}}} \right]^{\frac{N}{2}} \\ &< \frac{1}{N} \left[ \frac{\|z_\mu^2\|_{\lambda_2}^2}{\left(\int_{\mathbb{R}^N} |z_\mu^2|^{2^*}\right)^{\frac{2}{2^*}}} \right]^{\frac{N}{2}} = \frac{1}{N} S(\lambda_2)^{\frac{N}{2}} \\ &= \frac{1}{N} \min \left\{ S(\lambda_1)^{\frac{N}{2}}, S(\lambda_2)^{\frac{N}{2}} \right\}. \end{aligned}$$

Hence, conclusion (2) follows from Lemma 2.5.

Repeating the proof of [1, Theorem 2.2 (iii)-(iv)] with minor modifications, we can show that if  $\alpha < 2$ , then for all  $\mu > 0$ ,  $(0, z_\mu^2)$  is a saddle point for  $J_\nu$  in  $\mathcal{N}'_\nu$ , and so  $c'_\nu < \frac{1}{N} S(\lambda_2)^{N/2}$ ; if  $\beta < 2$ , then for all  $\mu > 0$ ,  $(z_\mu^1, 0)$  is a saddle point for  $J_\nu$  in  $\mathcal{N}'_\nu$ , and so  $c'_\nu < \frac{1}{N} S(\lambda_1)^{N/2}$ . Meanwhile, by (1.11) we see that  $\lambda_1 < \lambda_2$  implies  $S(\lambda_1) > S(\lambda_2)$  and  $\lambda_1 > \lambda_2$  implies  $S(\lambda_1) < S(\lambda_2)$ . Then under any one condition of **(C<sub>1</sub>)**, **(C<sub>2</sub>)** and **(C<sub>3</sub>)**, we have that  $c'_\nu < \frac{1}{N} \min\{S(\lambda_1)^{N/2}, S(\lambda_2)^{N/2}\}$ , and so conclusion (3) follows from Lemma 2.5. This completes the proof. □



3. PROOF OF THEOREM 1.2: THE CASE  $N = 4$

In this section, we assume that  $N = 4$ ,  $\alpha = \beta = 2$  and  $\lambda_1, \lambda_2 \in (0, 1)$ . By Theorem 2.1 we know that the ideas of proving Theorem 1.1 cannot be used here, and we need to use a different approach, which is much more complicated. This approach will take full use of the fact  $\alpha = \beta = 2$ , and so cannot be used in the case  $N = 3$  unfortunately.

3.1. **The special case**  $\lambda_1 = \lambda_2 = 0$ . Consider the following problem:

$$(3.1) \quad \begin{cases} -\Delta u = u^3 + 2\nu uv^2, & x \in \mathbb{R}^4, \\ -\Delta v = v^3 + 2\nu vu^2, & x \in \mathbb{R}^4, \\ u, v \in D^{1,2}(\mathbb{R}^4), & u, v > 0 \text{ in } \mathbb{R}^4. \end{cases}$$

For  $\varepsilon > 0$  and  $y \in \mathbb{R}^4$ , we consider the Aubin-Talenti instanton [6, 32]  $U_{\varepsilon,y} \in D^{1,2}(\mathbb{R}^4)$  defined by

$$(3.2) \quad U_{\varepsilon,y}(x) := \frac{2\sqrt{2}\varepsilon}{\varepsilon^2 + |x - y|^2}.$$

Then  $U_{\varepsilon,y}$  satisfies  $-\Delta u = u^3$  in  $\mathbb{R}^4$  and

$$(3.3) \quad \int_{\mathbb{R}^4} |\nabla U_{\varepsilon,y}|^2 dx = \int_{\mathbb{R}^4} |U_{\varepsilon,y}|^4 dx = S^2.$$

Furthermore,  $\{U_{\varepsilon,y} : \varepsilon > 0, y \in \mathbb{R}^4\}$  contains all positive solutions of the equation  $-\Delta u = u^3$  in  $\mathbb{R}^4$ . Note that (3.1) has semi-trivial solutions  $(U_{\varepsilon,y}, 0)$  and  $(0, U_{\varepsilon,y})$ . Here we are only interested in nontrivial solutions of (3.1), which can be found as nontrivial critical points of the  $C^2$  functional  $L_\nu : \mathbb{D} \rightarrow \mathbb{R}$ , where

$$(3.4) \quad L_\nu(u, v) = \frac{1}{2} (\|u\|^2 + \|v\|^2) - \frac{1}{4} \int_{\mathbb{R}^4} (u^4 + 4\nu u^2 v^2 + v^4).$$

**Definition 3.1.** We say a solution  $(u_0, v_0)$  of (3.1) is a ground state solution if  $(u_0, v_0)$  is nontrivial and  $L_\nu(u_0, v_0) \leq L_\nu(u, v)$  for any other nontrivial solution  $(u, v)$  of (3.1).

Define the general Nehari manifold of (3.4) as

$$\mathcal{M}_\nu := \left\{ (u, v) \in \mathbb{D} : u \neq 0, v \neq 0, \int_{\mathbb{R}^4} |\nabla u|^2 = \int_{\mathbb{R}^4} (u^4 + 2\nu u^2 v^2), \int_{\mathbb{R}^4} |\nabla v|^2 = \int_{\mathbb{R}^4} (v^4 + 2\nu u^2 v^2) \right\}.$$

Then any nontrivial solution of (3.1) has to belong to  $\mathcal{M}_\nu$ . Similarly as  $\mathcal{N}_\nu$ , we see that  $\mathcal{M}_\nu \neq \emptyset$ . We set

$$(3.5) \quad m_\nu := \inf_{(u,v) \in \mathcal{M}_\nu} L_\nu(u, v) = \inf_{(u,v) \in \mathcal{M}_\nu} \frac{1}{4} \int_{\mathbb{R}^4} (|\nabla u|^2 + |\nabla v|^2) dx.$$

By Sobolev inequality (1.13), it is easily seen that  $m_\nu > 0$  for all  $\nu$ . Moreover, if  $(u_0, v_0)$  is a nontrivial solution of (3.1) satisfying  $L_\nu(u_0, v_0) = m_\nu$ , then  $(u_0, v_0)$  is a ground state solution. Then we have the following result, which will play a crucial role in the proof of Theorem 1.2. Part of this result comes from the authors' paper [12].

**Theorem 3.1.** *Let  $\nu > 0$ .*

- (1) *If  $\nu \neq 1/2$ , then for any  $\varepsilon > 0$ ,  $y \in \mathbb{R}^4$ ,  $((1 + 2\nu)^{-1/2}U_{\varepsilon,y}, (1 + 2\nu)^{-1/2}U_{\varepsilon,y})$  is a positive ground state solution of (3.1), with*

$$(3.6) \quad m_\nu = L_\nu \left( (1 + 2\nu)^{-1/2}U_{\varepsilon,y}, (1 + 2\nu)^{-1/2}U_{\varepsilon,y} \right) = \frac{1}{2(1 + 2\nu)}S^2.$$

*Moreover, the set  $\{(1 + 2\nu)^{-1/2}U_{\varepsilon,y}, (1 + 2\nu)^{-1/2}U_{\varepsilon,y} : \varepsilon > 0, y \in \mathbb{R}^4\}$  contains all positive ground state solutions of (3.1).*

- (2) *If  $\nu = 1/2$ , then for any  $\varepsilon > 0$ ,  $y \in \mathbb{R}^4$ ,  $\theta \in (0, \pi/2)$ ,  $(\sin \theta U_{\varepsilon,y}, \cos \theta U_{\varepsilon,y})$  is a ground state solution of (3.1), and  $m_{1/2} = \frac{1}{4}S^2$ . Moreover, the set  $\{(\sin \theta U_{\varepsilon,y}, \cos \theta U_{\varepsilon,y}) : \varepsilon > 0, y \in \mathbb{R}^4, \theta \in (0, \pi/2)\}$  contains all positive ground state solutions of (3.1).*

*Proof.* (1) Let  $\nu > 0$  and  $\nu \neq 1/2$ . Then this result follows directly from [12, Theorem 1.5 and Theorem 4.1].

(2) Let  $\nu = 1/2$ . Firstly, note that for any  $\theta \in (0, \pi/2)$ ,  $(\sin \theta U_{\varepsilon,y}, \cos \theta U_{\varepsilon,y})$  is a positive solution of (3.1) when  $\nu = 1/2$ , so

$$(3.7) \quad m_{1/2} \leq L_{1/2}(\sin \theta U_{\varepsilon,y}, \cos \theta U_{\varepsilon,y}) = \frac{1}{4}S^2.$$

Secondly, take any  $(u, v) \in \mathcal{M}_{1/2}$ . If

$$(3.8) \quad \left( \int_{\mathbb{R}^4} u^2 v^2 dx \right)^2 = \int_{\mathbb{R}^4} u^4 dx \int_{\mathbb{R}^4} v^4 dx,$$

then by Hölder’s inequality we may assume that  $v = Cu$  for some constant  $C \neq 0$ . Recall the definition of  $\mathcal{M}_\nu$ . We see from (1.13) that

$$\int_{\mathbb{R}^4} |\nabla u|^2 = (1 + C^2) \int_{\mathbb{R}^4} u^4 \leq (1 + C^2)S^{-2} \left( \int_{\mathbb{R}^4} |\nabla u|^2 \right)^2,$$

so  $\int_{\mathbb{R}^4} |\nabla u|^2 \geq (1 + C^2)^{-1}S^2$  and then

$$L_{1/2}(u, v) = L_{1/2}(u, Cu) \geq \frac{1}{4}S^2.$$

If (3.8) does not hold, then

$$\left( \int_{\mathbb{R}^4} u^2 v^2 dx \right)^2 < \int_{\mathbb{R}^4} u^4 dx \int_{\mathbb{R}^4} v^4 dx,$$

and it is easy to prove that for any  $\nu \in (0, 1/2)$ , there exist  $t_\nu, s_\nu > 0$  such that  $(\sqrt{t_\nu}u, \sqrt{s_\nu}v) \in \mathcal{M}_\nu$  and  $(t_\nu, s_\nu) \rightarrow (1, 1)$  as  $\nu \rightarrow 1/2$ . This implies that

$$L_{1/2}(u, v) = \lim_{\nu \nearrow 1/2} L_\nu(\sqrt{t_\nu}u, \sqrt{s_\nu}v) \geq \lim_{\nu \nearrow 1/2} m_\nu = \frac{1}{4}S^2.$$

Therefore, for any  $(u, v) \in \mathcal{M}_{1/2}$ , we have  $L_{1/2}(u, v) \geq \frac{1}{4}S^2$ , and so  $m_{1/2} \geq \frac{1}{4}S^2$ . Combining this with (3.7), we see that  $(\sin \theta U_{\varepsilon,y}, \cos \theta U_{\varepsilon,y})$  is a ground state solution of (3.1) when  $\nu = 1/2$ .

Now assume that  $(u, v)$  is any positive ground state solution of (3.1) when  $\nu = 1/2$ . Then we deduce from (1.13) that

$$S|u|_4^2 \leq \|u\|^2 = |u|_4^4 + \int_{\mathbb{R}^4} u^2 v^2 dx \leq |u|_4^4 + |u|_4^2 |v|_4^2,$$

that is,  $|u|_4^2 + |v|_4^2 \geq S$ . Meanwhile, since  $L_{1/2}(u, v) = m_{1/2} = \frac{1}{4}S^2$ , we have

$$S^2 = \|u\|^2 + \|v\|^2 \geq S|u|_4^2 + S|v|_4^2,$$

that is,  $|u|_4^2 + |v|_4^2 \leq S$ . So  $|u|_4^2 + |v|_4^2 = S$ , that is,

$$S|u|_4^2 = \|u\|^2 = |u|_4^4 + \int_{\mathbb{R}^4} u^2 v^2 = |u|_4^4 + |u|_4^2 |v|_4^2.$$

First this means that  $v = Cu$  for some  $C > 0$ . Second, combining (1.13) with  $\|u\|^2 = S|u|_4^2$ , it is well known that  $u = C_1 U_{\varepsilon, y}$  for some  $C_1 > 0$ ,  $\varepsilon > 0$  and  $y \in \mathbb{R}^4$  (see [6, 32] for example). Hence,  $(u, v) = (C_1 U_{\varepsilon, y}, C_2 U_{\varepsilon, y})$  for some  $C_1, C_2 > 0$ . Then  $L_{1/2}(u, v) = \frac{1}{4}S^2$  yields that  $C_1^2 + C_2^2 = 1$ . Therefore, there exists  $\theta \in (0, \pi/2)$  such that  $C_1 = \sin \theta$  and  $C_2 = \cos \theta$ . This completes the proof.  $\square$

**3.2. The general case**  $\lambda_1, \lambda_2 \in (0, 1)$ . Recall the definition (1.21) of  $\nu_1$ . We have the following important energy estimate, and the idea of the proof comes from the authors' paper [12].

**Lemma 3.1.** *For any  $\nu \in (0, \nu_1)$ , there holds*

$$c_\nu < \min \left\{ \frac{1}{4}S(\lambda_1)^2 + \frac{1}{4}S(\lambda_2)^2, \quad m_\nu \right\}.$$

*Proof.* Define

$$(3.9) \quad G(u, v) := \begin{pmatrix} \int_{\mathbb{R}^4} u^4 dx & 2\nu \int_{\mathbb{R}^4} u^2 v^2 dx \\ 2\nu \int_{\mathbb{R}^4} u^2 v^2 dx & \int_{\mathbb{R}^4} v^4 dx \end{pmatrix}.$$

When  $\det G(u, v) > 0$ , the inverse matrix of  $G(u, v)$  is

$$(3.10) \quad G^{-1}(u, v) := \frac{1}{\det G(u, v)} \begin{pmatrix} \int_{\mathbb{R}^4} v^4 dx & -2\nu \int_{\mathbb{R}^4} u^2 v^2 dx \\ -2\nu \int_{\mathbb{R}^4} u^2 v^2 dx & \int_{\mathbb{R}^4} u^4 dx \end{pmatrix}.$$

Assume  $\nu \in (0, \nu_1)$ . Obviously, one has that  $2\nu < 1$ , and so  $\det G(z_1^1, z_1^2) > 0$ . Recall that  $\|z_1^i\|_{\lambda_i}^2 = |z_1^i|_4^4 = 4S(\lambda_i)^2, i = 1, 2$ ; we see that  $(\sqrt{t_0}z_1^1, \sqrt{s_0}z_1^2) \in \mathcal{N}_\nu$  for some  $t_0 > 0, s_0 > 0$  is equivalent to

$$(3.11) \quad \begin{pmatrix} t_0 \\ s_0 \end{pmatrix} := G^{-1}(z_1^1, z_1^2) \begin{pmatrix} |z_1^1|_4^4 \\ |z_1^2|_4^4 \end{pmatrix} = \frac{1}{\det G(z_1^1, z_1^2)} \begin{pmatrix} |z_1^2|_4^4 (|z_1^1|_4^4 - 2\nu \int_{\mathbb{R}^4} (z_1^1)^2 (z_1^2)^2) \\ |z_1^1|_4^4 (|z_1^2|_4^4 - 2\nu \int_{\mathbb{R}^4} (z_1^1)^2 (z_1^2)^2) \end{pmatrix} > \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Here and in the following,  $\begin{pmatrix} a \\ b \end{pmatrix} > \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  means both  $a > 0$  and  $b > 0$ . Meanwhile, (1.11) yields  $S_{\lambda_i} = (1 - \lambda_i)^{3/4}S$ , so we deduce from (1.21) that

$$2\nu < \min \left\{ \frac{1 - \lambda_1}{1 - \lambda_2}, \frac{1 - \lambda_2}{1 - \lambda_1} \right\} \leq \min \left\{ \frac{S(\lambda_1)}{S(\lambda_2)}, \frac{S(\lambda_2)}{S(\lambda_1)} \right\}.$$

Then

$$2\nu \int_{\mathbb{R}^4} (z_1^1)^2 (z_1^2)^2 < \min \left\{ \frac{S(\lambda_1)}{S(\lambda_2)}, \frac{S(\lambda_2)}{S(\lambda_1)} \right\} |z_1^1|_4^2 |z_1^2|_4^2 = \min \left\{ |z_1^1|_4^4, |z_1^2|_4^4 \right\}.$$

So (3.11) holds and  $(\sqrt{t_0}z_1^1, \sqrt{s_0}z_1^2) \in \mathcal{N}_\nu$  for  $(t_0, s_0)$  defined in (3.11). Then

$$\begin{aligned} c_\nu &\leq J_\nu(\sqrt{t_0}z_1^1, \sqrt{s_0}z_1^2) = \frac{t_0}{4}\|z_1^1\|_{\lambda_1}^2 + \frac{s_0}{4}\|z_1^2\|_{\lambda_2}^2 \\ &= \frac{t_0}{4} \int_{\mathbb{R}^4} (z_1^1)^4 + \frac{s_0}{4} \int_{\mathbb{R}^4} (z_1^2)^4 \\ &< \frac{t_0}{4} \int_{\mathbb{R}^4} \left( (z_1^1)^4 + 2\nu(z_1^1)^2(z_1^2)^2 \right) + \frac{s_0}{4} \int_{\mathbb{R}^4} \left( (z_1^2)^4 + 2\nu(z_1^1)^2(z_1^2)^2 \right) \\ &= \frac{1}{4}\|z_1^1\|_{\lambda_1}^2 + \frac{1}{4}\|z_1^2\|_{\lambda_2}^2 = \frac{1}{4}S(\lambda_1)^2 + \frac{1}{4}S(\lambda_2)^2. \end{aligned}$$

Hence  $c_\nu < \frac{1}{4}S(\lambda_1)^2 + \frac{1}{4}S(\lambda_2)^2$ . It remains to prove  $c_\nu < m_\nu$ . Take  $y_0 \in \mathbb{R}^4$  such that  $|y_0| = 2$ . Let  $\psi \in C_0^\infty(B(y_0, 1), \mathbb{R})$  be a function with  $0 \leq \psi \leq 1$ ,  $\psi \equiv 1$  for  $x \in B(y_0, 1/2)$ . Recall  $U_{\varepsilon, y_0}$  in (3.2) and (3.3); we define  $U_\varepsilon := \psi U_{\varepsilon, y_0}$ . Then by [8] or [35, Lemma 1.46], we have the inequalities

$$\begin{aligned} \int_{\mathbb{R}^4} |\nabla U_\varepsilon|^2 &= S^2 + O(\varepsilon^2), \quad \int_{\mathbb{R}^4} |U_\varepsilon|^4 = S^2 + O(\varepsilon^4), \\ \int_{\mathbb{R}^4} \frac{|U_\varepsilon|^2}{|x|^2} dx &\geq \frac{1}{9} \int_{B(y_0, 1)} |U_\varepsilon|^2 \geq C\varepsilon^2 |\ln \varepsilon| + O(\varepsilon^2), \end{aligned}$$

where  $C$  is a positive constant. Recalling that  $\lambda_1, \lambda_2 > 0$ , we have

$$\begin{aligned} J_\nu(\sqrt{t}U_\varepsilon, \sqrt{s}U_\varepsilon) &= \frac{1}{2}t \int_{\mathbb{R}^4} \left( |\nabla U_\varepsilon|^2 - \frac{\lambda_1}{|x|^2} U_\varepsilon^2 \right) + \frac{1}{2}s \int_{\mathbb{R}^4} \left( |\nabla U_\varepsilon|^2 - \frac{\lambda_2}{|x|^2} U_\varepsilon^2 \right) \\ &\quad - \frac{1}{4}(t^2 + 4\nu ts + s^2) \int_{\mathbb{R}^4} U_\varepsilon^4 \\ &\leq \frac{1}{2}(t + s) (S^2 - C\varepsilon^2 |\ln \varepsilon| + O(\varepsilon^2)) \\ (3.12) \quad &\quad - \frac{1}{4}(t^2 + 4\nu ts + s^2) (S^2 + O(\varepsilon^4)). \end{aligned}$$

Denote

$$A_\varepsilon = S^2 - C\varepsilon^2 |\ln \varepsilon| + O(\varepsilon^2), \quad B_\varepsilon = S^2 + O(\varepsilon^4);$$

then  $0 < A_\varepsilon < B_\varepsilon$  and  $A_\varepsilon < S^2$  for  $\varepsilon > 0$  small enough. Consider

$$f_\varepsilon(t, s) := \frac{1}{2}A_\varepsilon(t + s) - \frac{1}{4}B_\varepsilon(t^2 + 4\nu ts + s^2);$$

then it is easy to see that there exists  $t_\varepsilon, s_\varepsilon > 0$  such that

$$f_\varepsilon(t_\varepsilon, s_\varepsilon) = \max_{t, s > 0} f_\varepsilon(t, s).$$

By  $\frac{\partial}{\partial t} f_\varepsilon(t, s)|_{(t_\varepsilon, s_\varepsilon)} = \frac{\partial}{\partial s} f_\varepsilon(t, s)|_{(t_\varepsilon, s_\varepsilon)} = 0$ , we see that

$$t_\varepsilon = s_\varepsilon = \frac{A_\varepsilon}{(1 + 2\nu)B_\varepsilon}.$$

Then it follows from (3.6) and (3.12) that

$$\begin{aligned}
 \max_{t,s>0} J_\nu(\sqrt{t}U_\varepsilon, \sqrt{s}U_\varepsilon) &\leq \max_{t,s>0} f_\varepsilon(t, s) = f_\varepsilon(t_\varepsilon, s_\varepsilon) \\
 &= \frac{1}{2(1+2\nu)} \frac{A_\varepsilon^2}{B_\varepsilon} < \frac{1}{2(1+2\nu)} A_\varepsilon \\
 (3.13) \quad &< \frac{S^2}{2(1+2\nu)} = m_\nu \quad \text{holds for } \varepsilon \text{ small enough.}
 \end{aligned}$$

Similarly as above, we have  $\det G(U_\varepsilon, U_\varepsilon) > 0$ . Moreover,  $(\sqrt{\tilde{t}_\varepsilon}U_\varepsilon, \sqrt{\tilde{s}_\varepsilon}U_\varepsilon) \in \mathcal{N}_\nu$  for some  $\tilde{t}_\varepsilon > 0, \tilde{s}_\varepsilon > 0$  is equivalent to

$$(3.14) \quad \begin{pmatrix} \tilde{t}_\varepsilon \\ \tilde{s}_\varepsilon \end{pmatrix} = \frac{|U_\varepsilon|_4^4}{\det G(U_\varepsilon, U_\varepsilon)} \begin{pmatrix} \|U_\varepsilon\|_{\lambda_1}^2 - 2\nu\|U_\varepsilon\|_{\lambda_2}^2 \\ \|U_\varepsilon\|_{\lambda_2}^2 - 2\nu\|U_\varepsilon\|_{\lambda_1}^2 \end{pmatrix} > \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

On the other hand, by (1.6) we have

$$\begin{aligned}
 \|U_\varepsilon\|_{\lambda_1}^2 - 2\nu\|U_\varepsilon\|_{\lambda_2}^2 &= (1-2\nu) \int_{\mathbb{R}^4} |\nabla U_\varepsilon|^2 - (\lambda_1 - 2\nu\lambda_2) \int_{\mathbb{R}^4} \frac{U_\varepsilon^2}{|x|^2} \\
 &\geq (1-2\nu) \int_{\mathbb{R}^4} \frac{U_\varepsilon^2}{|x|^2} - (\lambda_1 - 2\nu\lambda_2) \int_{\mathbb{R}^4} \frac{U_\varepsilon^2}{|x|^2} \\
 &= [(1-\lambda_1) - 2\nu(1-\lambda_2)] \int_{\mathbb{R}^4} \frac{U_\varepsilon^2}{|x|^2} > 0.
 \end{aligned}$$

Similarly,  $\|U_\varepsilon\|_{\lambda_2}^2 - 2\nu\|U_\varepsilon\|_{\lambda_1}^2 > 0$ . Hence, (3.14) holds and  $(\sqrt{\tilde{t}_\varepsilon}U_\varepsilon, \sqrt{\tilde{s}_\varepsilon}U_\varepsilon) \in \mathcal{N}_\nu$  for  $(\tilde{t}_\varepsilon, \tilde{s}_\varepsilon)$  defined in (3.14). Then we see from (3.13) that

$$c_\nu \leq J_\nu(\sqrt{\tilde{t}_\varepsilon}U_\varepsilon, \sqrt{\tilde{s}_\varepsilon}U_\varepsilon) \leq \max_{t,s>0} J_\nu(\sqrt{t}U_\varepsilon, \sqrt{s}U_\varepsilon) < m_\nu.$$

This completes the proof. □

**Lemma 3.2.** *Assume that  $\nu \in (0, \nu_1)$ . Then there exist  $C_2 > C_1 > 0$  such that for any  $(u, v) \in \mathcal{N}_\nu$  with  $J_\nu(u, v) \leq \frac{1}{4}S(\lambda_1)^2 + \frac{1}{4}S(\lambda_2)^2$ , there holds*

$$(3.15) \quad C_1 \leq \int_{\mathbb{R}^4} u^4 dx, \int_{\mathbb{R}^4} v^4 dx \leq C_2.$$

*Proof.* Take any  $(u, v) \in \mathcal{N}_\nu$  with  $J_\nu(u, v) \leq \frac{1}{4}S(\lambda_1)^2 + \frac{1}{4}S(\lambda_2)^2$ . By (1.17) and Hölder’s inequality, one has

$$\begin{aligned}
 S(\lambda_1)|u|_4^2 &\leq \|u\|_{\lambda_1}^2 = \int_{\mathbb{R}^4} (u^4 + 2\nu u^2 v^2) \leq |u|_4^4 + 2\nu|u|_4^2|v|_4^2, \\
 S(\lambda_2)|v|_4^2 &\leq \|v\|_{\lambda_2}^2 = \int_{\mathbb{R}^4} (v^4 + 2\nu u^2 v^2) \leq |v|_4^4 + 2\nu|u|_4^2|v|_4^2.
 \end{aligned}$$

Therefore, there exists  $C_2 > 0$  such that  $\int_{\mathbb{R}^4} u^4, \int_{\mathbb{R}^4} v^4 \leq C_2$ . Moreover,

$$(3.16) \quad |u|_4^2 + 2\nu|v|_4^2 \geq S(\lambda_1),$$

$$(3.17) \quad 2\nu|u|_4^2 + |v|_4^2 \geq S(\lambda_2),$$

$$(3.18) \quad S(\lambda_1)|u|_4^2 + S(\lambda_2)|v|_4^2 \leq S(\lambda_1)^2 + S(\lambda_2)^2.$$

Recall that  $S(\lambda_i) = (1 - \lambda_i)^{3/4}S$ . Since  $\nu \in (0, \nu_1)$  and  $\nu_1$  is defined in (1.21), by (3.16) and (3.18) we have

$$(3.19) \quad |u|_4^2 \geq \frac{S(\lambda_1)S(\lambda_2) - 2\nu(S(\lambda_1)^2 + S(\lambda_2)^2)}{S(\lambda_2) - 2\nu S(\lambda_1)} > 0,$$

and by (3.17) and (3.18) we have

$$|v|_4^2 \geq \frac{S(\lambda_1)S(\lambda_2) - 2\nu(S(\lambda_1)^2 + S(\lambda_2)^2)}{S(\lambda_1) - 2\nu S(\lambda_2)} > 0.$$

This completes the proof. □

The following lemma is motivated by [33], and some ideas of the proof come from [30].

**Lemma 3.3.** *Assume that  $\nu \in (0, \nu_1)$ . Let  $(u_n, v_n) \in \mathcal{N}_\nu$  be a minimizing sequence of  $c_\nu$ , and  $(u_n, v_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ . Then for any  $r > 0$  and for every  $\varepsilon \in (-r, 0) \cup (0, r)$ , there exists  $\rho \in (\varepsilon, 0) \cup (0, \varepsilon)$  such that, up to a subsequence,*

$$(3.20) \quad \text{either } \int_{B_{r+\rho}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0 \text{ or } \int_{\mathbb{R}^N \setminus B_{r+\rho}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0.$$

*Proof.* Without loss of generality, we only consider the case  $\varepsilon \in (0, r)$  (the proof for the case  $\varepsilon \in (-r, 0)$  is similar). Since  $(u_n, v_n) \in \mathcal{N}_\nu$  is a minimizing sequence of  $c_\nu$ , then  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}$ . Moreover, by Lemmas 3.1 and 3.2 we may assume that  $(u_n, v_n)$  satisfies (3.15) for all  $n \in \mathbb{N}$ .

*Step 1.* We prove (3.20) by further assuming that  $J'_\nu(u_n, v_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

In the following, some arguments are borrowed from [30] (see also [31, Lemma III.3.3] or [33, Proposition 5.2]). Denote  $\mathbb{S}$  as the unit sphere of  $\mathbb{R}^4$ . Since

$$\int_r^{r+\varepsilon} d\rho \int_{\rho\mathbb{S}} (|\nabla u_n|^2 + |\nabla v_n|^2) = \int_{r \leq |x| \leq r+\varepsilon} (|\nabla u_n|^2 + |\nabla v_n|^2)$$

is bounded, we can find  $\rho \in (0, \varepsilon)$  such that

$$\int_{(r+\rho)\mathbb{S}} (|\nabla u_n|^2 + |\nabla v_n|^2) \leq \frac{3}{\varepsilon} \int_{r \leq |x| \leq r+\varepsilon} (|\nabla u_n|^2 + |\nabla v_n|^2)$$

holds for infinitely many  $n$ 's. Therefore, as  $H^1((r + \rho)\mathbb{S})$  is compactly embedded into  $H^{1/2}((r + \rho)\mathbb{S})$ , up to a subsequence we can assume that  $u_n \rightarrow u, v_n \rightarrow v$  strongly in  $H^{1/2}((r + \rho)\mathbb{S})$ . On the other hand, by the continuity of the embedding  $H^1(B_{r+\rho}) \hookrightarrow H^{1/2}((r + \rho)\mathbb{S})$  and by the weak convergence to  $(0, 0)$  of  $(u_n, v_n)$ , we deduce that  $(u, v) = (0, 0)$ , that is,  $u_n \rightarrow 0$  and  $v_n \rightarrow 0$  strongly in  $H^{1/2}((r + \rho)\mathbb{S})$ . Let  $w_{i,n}, i = 1, 2$ , be the solutions to the Dirichlet problems

$$(3.21) \quad \begin{cases} \Delta w_{1,n} = 0 & \text{in } B_{r+\varepsilon} \setminus B_{r+\rho}, \\ w_{1,n} = 0 & \text{on } (r + \varepsilon)\mathbb{S}, \\ w_{1,n} = u_n & \text{on } (r + \rho)\mathbb{S}, \end{cases} \quad \begin{cases} \Delta w_{2,n} = 0 & \text{in } B_{r+\rho} \setminus B_{r-\varepsilon}, \\ w_{2,n} = 0 & \text{on } (r - \varepsilon)\mathbb{S}, \\ w_{2,n} = u_n & \text{on } (r + \rho)\mathbb{S}, \end{cases}$$

and let  $\sigma_{i,n}, i = 1, 2$  be the solutions to the Dirichlet problems

$$(3.22) \quad \begin{cases} \Delta \sigma_{1,n} = 0 & \text{in } B_{r+\varepsilon} \setminus B_{r+\rho}, \\ \sigma_{1,n} = 0 & \text{on } (r + \varepsilon)\mathbb{S}, \\ \sigma_{1,n} = v_n & \text{on } (r + \rho)\mathbb{S}, \end{cases} \quad \begin{cases} \Delta \sigma_{2,n} = 0 & \text{in } B_{r+\rho} \setminus B_{r-\varepsilon}, \\ \sigma_{2,n} = 0 & \text{on } (r - \varepsilon)\mathbb{S}, \\ \sigma_{2,n} = v_n & \text{on } (r + \rho)\mathbb{S}. \end{cases}$$

By continuity of the inverse Laplace operator from  $H^{1/2}(\partial\Omega)$  to  $H^1(\Omega)$ , it follows from the above discussion that  $w_{1,n} \rightarrow 0, \sigma_{1,n} \rightarrow 0$  strongly in  $H^1(B_{r+\varepsilon} \setminus B_{r+\rho})$  and  $w_{2,n} \rightarrow 0, \sigma_{2,n} \rightarrow 0$  strongly in  $H^1(B_{r+\rho} \setminus B_{r-\varepsilon})$ . Define

$$(3.23) \quad u_{1,n}(x) = \begin{cases} u_n(x) & \text{if } x \in B_{r+\rho}, \\ w_{1,n} & \text{if } x \in B_{r+\varepsilon} \setminus B_{r+\rho}, \\ 0 & \text{elsewhere,} \end{cases}$$

$$(3.24) \quad v_{1,n}(x) = \begin{cases} v_n(x) & \text{if } x \in B_{r+\rho}, \\ \sigma_{1,n} & \text{if } x \in B_{r+\varepsilon} \setminus B_{r+\rho}, \\ 0 & \text{elsewhere,} \end{cases}$$

$$(3.25) \quad u_{2,n}(x) = \begin{cases} 0 & \text{if } x \in B_{r-\varepsilon}, \\ w_{2,n} & \text{if } x \in B_{r+\rho} \setminus B_{r-\varepsilon}, \\ u_n(x) & \text{elsewhere,} \end{cases}$$

$$(3.26) \quad v_{2,n}(x) = \begin{cases} 0 & \text{if } x \in B_{r-\varepsilon}, \\ \sigma_{2,n} & \text{if } x \in B_{r+\rho} \setminus B_{r-\varepsilon}, \\ v_n(x) & \text{elsewhere.} \end{cases}$$

Then it is easy to see that

$$(3.27) \quad \|u_n\|_{\lambda_1}^2 = \|u_{1,n}\|_{\lambda_1}^2 + \|u_{2,n}\|_{\lambda_1}^2 + o(1),$$

$$(3.28) \quad \|v_n\|_{\lambda_2}^2 = \|v_{1,n}\|_{\lambda_2}^2 + \|v_{2,n}\|_{\lambda_2}^2 + o(1).$$

Moreover, we can easily obtain

$$(3.29) \quad J'_\nu(u_{1,n}, v_{1,n})(u_{1,n}, 0) = J'_\nu(u_n, v_n)(u_{1,n}, 0) + o(1) = o(1),$$

$$(3.30) \quad J'_\nu(u_{1,n}, v_{1,n})(0, v_{1,n}) = J'_\nu(u_n, v_n)(0, v_{1,n}) + o(1) = o(1),$$

$$(3.31) \quad J'_\nu(u_{2,n}, v_{2,n})(u_{2,n}, 0) = J'_\nu(u_n, v_n)(u_{2,n}, 0) + o(1) = o(1),$$

$$(3.32) \quad J'_\nu(u_{2,n}, v_{2,n})(0, v_{2,n}) = J'_\nu(u_n, v_n)(0, v_{2,n}) + o(1) = o(1).$$

Then we claim that

$$(3.33) \quad \text{either } \lim_{n \rightarrow \infty} (\|u_{1,n}\|^2 + \|v_{1,n}\|^2) = 0 \text{ or } \lim_{n \rightarrow \infty} (\|u_{2,n}\|^2 + \|v_{2,n}\|^2) = 0.$$

In fact, if (3.33) does not hold, then up to a subsequence,

$$(3.34) \quad \text{both } \lim_{n \rightarrow \infty} (\|u_{1,n}\|^2 + \|v_{1,n}\|^2) > 0 \text{ and } \lim_{n \rightarrow \infty} (\|u_{2,n}\|^2 + \|v_{2,n}\|^2) > 0.$$

We have the following several cases.

*Case 1.* Up to a subsequence, both  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 > 0$ .

Since norms  $\|\cdot\|_{\lambda_i}, i = 1, 2$ , are equivalent to  $\|\cdot\|$ , and (3.29)-(3.30) yield

$$(3.35) \quad \|u_{1,n}\|_{\lambda_1}^2 = |u_{1,n}|_4^4 + 2\nu \int_{\mathbb{R}^4} u_{1,n}^2 v_{1,n}^2 + o(1),$$

$$(3.36) \quad \|v_{1,n}\|_{\lambda_2}^2 = |v_{1,n}|_4^4 + 2\nu \int_{\mathbb{R}^4} u_{1,n}^2 v_{1,n}^2 + o(1),$$

hence, both  $\liminf_{n \rightarrow \infty} |u_{1,n}|_4^4 > 0$  and  $\liminf_{n \rightarrow \infty} |v_{1,n}|_4^4 > 0$ . Since  $2\nu < 2\nu_1 \leq 1$ , by Hölder's inequality we have

$$\liminf_{n \rightarrow \infty} \left[ |u_{1,n}|_4^4 |v_{1,n}|_4^4 - \left( 2\nu \int_{\mathbb{R}^4} u_{1,n}^2 v_{1,n}^2 \right)^2 \right] > 0.$$

Combining this with (3.35)-(3.36), it is easy to prove that there exist  $t_n, s_n > 0$  such that  $(\sqrt{t_n}u_{1,n}, \sqrt{s_n}v_{1,n}) \in \mathcal{N}_\nu$  and  $(t_n, s_n) \rightarrow (1, 1)$ , and so we conclude from (3.27)-(3.28) and (3.34) that

$$\begin{aligned} c_\nu &= \lim_{n \rightarrow \infty} J_\nu(u_n, v_n) = \lim_{n \rightarrow \infty} \frac{1}{4}(\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \\ &= \lim_{n \rightarrow \infty} \frac{1}{4}(t_n\|u_{1,n}\|_{\lambda_1}^2 + s_n\|v_{1,n}\|_{\lambda_2}^2) + \lim_{n \rightarrow \infty} \frac{1}{4}(\|u_{2,n}\|_{\lambda_1}^2 + \|v_{2,n}\|_{\lambda_2}^2) \\ &> \lim_{n \rightarrow \infty} \frac{1}{4}(t_n\|u_{1,n}\|_{\lambda_1}^2 + s_n\|v_{1,n}\|_{\lambda_2}^2) \\ &= \lim_{n \rightarrow \infty} J_\nu(\sqrt{t_n}u_{1,n}, \sqrt{s_n}v_{1,n}) \geq c_\nu, \end{aligned}$$

a contradiction. So Case 1 is impossible.

*Case 2.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 = 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 > 0$ .

Then (3.36) yields that

$$\|v_{1,n}\|_{\lambda_2}^2 = |v_{1,n}|_4^4 + o(1) \leq S(\lambda_2)^{-2}\|v_{1,n}\|_{\lambda_2}^4 + o(1),$$

and so  $\lim_{n \rightarrow \infty} \|v_{1,n}\|_{\lambda_2}^2 \geq S(\lambda_2)^2$ . By (3.15) we have

$$\liminf_{n \rightarrow \infty} \|u_{2,n}\|^2 = \liminf_{n \rightarrow \infty} \|u_n\|^2 - \lim_{n \rightarrow \infty} \|u_{1,n}\|^2 > 0.$$

If up to a subsequence,  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 > 0$ , then we can get a contradiction just as in Case 1. Therefore,  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 = 0$ . Then similarly as above, we can deduce from (3.31) that  $\lim_{n \rightarrow \infty} \|u_{2,n}\|_{\lambda_1}^2 \geq S(\lambda_1)^2$ . Then

$$\begin{aligned} c_\nu &= \lim_{n \rightarrow \infty} \frac{1}{4}(\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \\ &= \lim_{n \rightarrow \infty} \frac{1}{4}(\|u_{2,n}\|_{\lambda_1}^2 + \|v_{1,n}\|_{\lambda_2}^2) \geq \frac{1}{4}(S(\lambda_1)^2 + S(\lambda_2)^2), \end{aligned}$$

a contradiction with Lemma 3.1. So Case 2 is impossible.

*Case 3.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 = 0$ .

By a similar argument as in Case 2, we get a contradiction. So Case 3 is impossible.

Since none of Cases 1, 2 and 3 is true, we see that (3.34) is impossible, that is, (3.33) holds. Recall the definition of  $(u_{i,n}, v_{i,n})$ ; (3.20) follows directly from (3.33). This completes the proof of Step 1.

*Step 2.* We prove (3.20) without assuming that  $J'_\nu(u_n, v_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

By the Ekeland variational principle (see [31, Theorem 5.1] for example), there exists a sequence  $\{(\tilde{u}_n, \tilde{v}_n)\} \in \mathcal{N}_\nu$  such that

$$(3.37) \quad J_\nu(\tilde{u}_n, \tilde{v}_n) \leq J_\nu(u_n, v_n), \quad \|(u_n, v_n) - (\tilde{u}_n, \tilde{v}_n)\| \leq \frac{1}{n},$$

$$(3.38) \quad J_\nu(u, v) \geq J_\nu(\tilde{u}_n, \tilde{v}_n) - \frac{1}{n}\|(\tilde{u}_n, \tilde{v}_n) - (u, v)\|, \quad \forall (u, v) \in \mathcal{N}_\nu.$$

Here,  $\|(u, v)\| := (\int_{\mathbb{R}^4} (|\nabla u|^2 + |\nabla v|^2) dx)^{1/2}$  is also a norm of  $\mathbb{D}$ , which is equivalent to  $\|(u, v)\|_{\mathbb{D}}$ . Recall that  $(u_n, v_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ ; by (3.37) we also have  $J_\nu(\tilde{u}_n, \tilde{v}_n) \rightarrow c_\nu$  and  $(\tilde{u}_n, \tilde{v}_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ . Moreover, by Lemma 3.1 we may assume that  $(\tilde{u}_n, \tilde{v}_n)$  satisfies (3.15) for all  $n \in \mathbb{N}$ . Then by repeating the proof



of [12, Theorem 1.3 (1)-(2)], we can prove that  $J'_\nu(\tilde{u}_n, \tilde{v}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Hence Step 1 yields that (3.20) holds for  $(\tilde{u}_n, \tilde{v}_n)$ . Combining this with (3.37), we see that (3.20) holds for  $(u_n, v_n)$ . This completes the proof.  $\square$

*Proof of Theorem 1.2.* Fix any  $\nu \in (0, \nu_1)$ . Take a sequence  $(\bar{u}_n, \bar{v}_n) \in \mathcal{N}_\nu$  such that  $J_\nu(\bar{u}_n, \bar{v}_n) \rightarrow c_\nu$  as  $n \rightarrow \infty$ . Recall that  $E(u, v) = |\nabla u|^2 + |\nabla v|^2 - \frac{\lambda_1}{|x|^2}|u|^2 - \frac{\lambda_2}{|x|^2}|v|^2$ ; there exists  $R_n > 0$  such that

$$\int_{B_{R_n}} E(\bar{u}_n, \bar{v}_n) = \int_{\mathbb{R}^N \setminus B_{R_n}} E(\bar{u}_n, \bar{v}_n) = \frac{1}{2}(\|\bar{u}_n\|_{\lambda_1}^2 + \|\bar{v}_n\|_{\lambda_2}^2).$$

Define

$$(\tilde{u}_n(x), \tilde{v}_n(x)) := \left( R_n^{\frac{N-2}{2}} \bar{u}_n(R_n x), R_n^{\frac{N-2}{2}} \bar{v}_n(R_n x) \right).$$

Then by a direct computation, we see that  $(\tilde{u}_n, \tilde{v}_n) \in \mathcal{N}_\nu$  and  $J_\nu(\tilde{u}_n, \tilde{v}_n) \rightarrow c_\nu$ . Moreover,

$$(3.39) \quad \int_{B_1} E(\tilde{u}_n, \tilde{v}_n) = \int_{\mathbb{R}^N \setminus B_1} E(\tilde{u}_n, \tilde{v}_n) = \frac{1}{2}(\|\tilde{u}_n\|_{\lambda_1}^2 + \|\tilde{v}_n\|_{\lambda_2}^2) \rightarrow 2c_\nu > 0.$$

By the Ekeland variational principle (see [31, Theorem 5.1] for example), there exists a sequence  $\{(u_n, v_n)\} \in \mathcal{N}_\nu$  such that

$$(3.40) \quad J_\nu(u_n, v_n) \leq J_\nu(\tilde{u}_n, \tilde{v}_n), \quad \|(u_n, v_n) - (\tilde{u}_n, \tilde{v}_n)\| \leq \frac{1}{n},$$

$$(3.41) \quad J_\nu(u, v) \geq J_\nu(u_n, v_n) - \frac{1}{n}\|(u, v) - (u_n, v_n)\|, \quad \forall (u, v) \in \mathcal{N}_\nu.$$

Similarly as in Step 2 in the proof of Lemma 3.3, we have that  $J_\nu(u_n, v_n) \rightarrow c_\nu$  and  $J'_\nu(u_n, v_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Moreover, (3.39) and (3.40) yield that

$$(3.42) \quad \lim_{n \rightarrow \infty} \int_{B_1} E(u_n, v_n) = \lim_{n \rightarrow \infty} \int_{B_1} E(\tilde{u}_n, \tilde{v}_n) = 2c_\nu,$$

$$(3.43) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N \setminus B_1} E(u_n, v_n) = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N \setminus B_1} E(\tilde{u}_n, \tilde{v}_n) = 2c_\nu.$$

Note that  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}$ . Then up to a subsequence, we assume that  $(u_n, v_n) \rightharpoonup (u, v)$  weakly in  $\mathbb{D}$ . Then  $J'_\nu(u, v) = 0$ .

*Case 1.*  $(u, v) \equiv (0, 0)$ .

Then we can apply Lemma 3.3 twice with  $r = 1$  and  $\varepsilon = \pm 1/4$  respectively, and there exist  $\rho^+ \in (0, 1/4)$  and  $\rho^- \in (-1/4, 0)$  such that the alternative (3.20) holds. By (3.42)-(3.43) we can rule out all possibilities other than

$$(3.44) \quad \int_{B_{1+\rho^-}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0 \quad \text{and} \quad \int_{\mathbb{R}^N \setminus B_{1+\rho^+}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0.$$

Now let  $\eta \in C_0^\infty(\mathbb{R}^4)$  such that  $0 \leq \eta \leq 1$ ,  $\eta(x) = 1$  for  $|x| \in [3/4, 5/4]$  and  $\eta(x) = 0$  for  $|x| \notin [1/2, 3/2]$ . Recall that  $(u_n, v_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ , so  $u_n, v_n \rightarrow 0$  strongly in  $L^2_{loc}(\mathbb{R}^4)$ . Combining this with (3.44), we obtain that

$$\|(\eta u_n) - u_n\| \rightarrow 0, \quad \|(\eta v_n) - v_n\| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

By Hardy inequality (1.6), we have

$$\begin{aligned} \int_{\mathbb{R}^4} \frac{u_n^2}{|x|^2} &= \int_{\mathbb{R}^4} \frac{(1-\eta)^2 u_n^2}{|x|^2} + \int_{\mathbb{R}^4} \frac{(2-\eta)\eta u_n^2}{|x|^2} \\ &\leq \|(\eta u_n) - u_n\|^2 + 8 \int_{1/2 \leq |x| \leq 3/2} u_n^2 = o(1). \end{aligned}$$

Similarly,  $\int_{\mathbb{R}^4} \frac{v_n^2}{|x|^2} = o(1)$ . Therefore, we see from  $(u_n, v_n) \in \mathcal{N}_\nu$  that

$$\begin{aligned} \int_{\mathbb{R}^4} |\nabla u_n|^2 &= \int_{\mathbb{R}^4} u_n^4 + 2\nu \int_{\mathbb{R}^4} u_n^2 v_n^2 + o(1), \\ \int_{\mathbb{R}^4} |\nabla v_n|^2 &= \int_{\mathbb{R}^4} v_n^4 + 2\nu \int_{\mathbb{R}^4} u_n^2 v_n^2 + o(1). \end{aligned}$$

From Lemma 3.2 we may assume that  $|u_n|_4^4, |v_n|_4^4 \geq C > 0$ , where  $C$  is independent of  $n$ . Since  $2\nu < 2\nu' \leq 1$ , then it is easy to prove that there exist  $t_n, s_n > 0$  such that  $(\sqrt{t_n}u_n, \sqrt{s_n}v_n) \in \mathcal{M}_\nu$  and  $(t_n, s_n) \rightarrow (1, 1)$  as  $n \rightarrow \infty$ . Hence,

$$\begin{aligned} c_\nu &= \lim_{n \rightarrow \infty} J_\nu(u_n, v_n) = \lim_{n \rightarrow \infty} \frac{1}{4} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \\ &= \lim_{n \rightarrow \infty} \frac{1}{4} \left( t_n \int_{\mathbb{R}^4} |\nabla u_n|^2 + s_n \int_{\mathbb{R}^4} |\nabla v_n|^2 \right) \\ &= \lim_{n \rightarrow \infty} L_\nu(\sqrt{t_n}u_n, \sqrt{s_n}v_n) \geq m_\nu, \end{aligned}$$

a contradiction with Lemma 3.1. So Case 1 is impossible.

*Case 2.* Either  $u \equiv 0, v \not\equiv 0$  or  $u \not\equiv 0, v \equiv 0$ .

Without loss of generality, we assume that  $u \not\equiv 0, v \equiv 0$ . Note that  $J'_\nu(u, v)(u, 0) = 0$  yields

$$\|u\|_{\lambda_1}^2 = |u|_4^4 \leq S(\lambda_1)^{-2} \|u\|_{\lambda_1}^4,$$

which implies  $\|u\|_{\lambda_1}^2 \geq S(\lambda_1)^2$ .

*Case 2.1.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_n - u\| > 0$ .

Denote  $w_n = u_n - u$ . Note that  $(u_n, v_n) \in \mathcal{N}_\nu$ . Then by the Brezis-Lieb Lemma ([7]) and Lemma 2.3 we conclude that

$$\begin{aligned} \|w_n\|_{\lambda_1}^2 &= \int_{\mathbb{R}^4} w_n^4 + 2\nu \int_{\mathbb{R}^4} w_n^2 v_n^2 + o(1), \\ \|v_n\|_{\lambda_2}^2 &= \int_{\mathbb{R}^4} v_n^4 + 2\nu \int_{\mathbb{R}^4} w_n^2 v_n^2 + o(1). \end{aligned}$$

Similarly as above, it is easy to prove that there exist  $t_n, s_n > 0$  such that  $(\sqrt{t_n}w_n, \sqrt{s_n}v_n) \in \mathcal{N}_\nu$  and  $(t_n, s_n) \rightarrow (1, 1)$  as  $n \rightarrow \infty$ . Hence,

$$\begin{aligned} c_\nu &= \lim_{n \rightarrow \infty} J_\nu(u_n, v_n) = \lim_{n \rightarrow \infty} \frac{1}{4} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \\ &= \frac{1}{4} \|u\|_{\lambda_1}^2 + \lim_{n \rightarrow \infty} \frac{1}{4} (t_n \|w_n\|_{\lambda_1}^2 + s_n \|v_n\|_{\lambda_2}^2) \\ &> \lim_{n \rightarrow \infty} J_\nu(\sqrt{t_n}w_n, \sqrt{s_n}v_n) \geq c_\nu, \end{aligned}$$

a contradiction. So Case 2.1 is impossible.

*Case 2.2.*  $u_n \rightarrow u$  strongly in  $D^{1,2}(\mathbb{R}^N)$ .

Then  $u_n^2 \rightarrow u^2$  strongly in  $L^2(\mathbb{R}^4)$ . Recall that  $v_n \rightarrow 0$  in  $D^{1,2}(\mathbb{R}^4)$ , so  $v_n^2 \rightarrow 0$  weakly in  $L^2(\mathbb{R}^4)$ , which easily implies

$$\int_{\mathbb{R}^4} u_n^2 v_n^2 \leq \int_{\mathbb{R}^4} u^2 v_n^2 + \int_{\mathbb{R}^4} |u_n^2 - u^2| v_n^2 = o(1).$$

Then we have

$$\|v_n\|_{\lambda_2}^2 = |v_n|_4^4 + o(1) \leq S(\lambda_2)^{-2} \|v_n\|_{\lambda_2}^4 + o(1).$$

Since Lemma 3.2 yields  $\lim_{n \rightarrow \infty} \|v_n\|_{\lambda_2}^2 > 0$ , then  $\lim_{n \rightarrow \infty} \|v_n\|_{\lambda_2}^2 \geq S(\lambda_2)^2$ , and

$$\begin{aligned} c_\nu &= \lim_{n \rightarrow \infty} J_\nu(u_n, v_n) = \lim_{n \rightarrow \infty} \frac{1}{4} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \\ &= \frac{1}{4} \|u\|_{\lambda_1}^2 + \lim_{n \rightarrow \infty} \frac{1}{4} \|v_n\|_{\lambda_2}^2 \geq \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2), \end{aligned}$$

a contradiction with Lemma 3.1. So Case 2.2 is impossible, and so Case 2 is impossible.

Since neither Case 1 nor Case 2 is true, we obtain that  $u \neq 0$  and  $v \neq 0$ . Since  $J'_\nu(u, v) = 0$ , then  $(u, v) \in \mathcal{N}_\nu$ . Meanwhile, Fatou's Lemma implies that

$$c_\nu \leq J_\nu(u, v) \leq \liminf_{n \rightarrow \infty} J_\nu(u_n, v_n) = c_\nu,$$

so  $J_\nu(u, v) = c_\nu$ . Then  $(|u|, |v|) \in \mathcal{N}_\nu$  and  $J_\nu(|u|, |v|) = c_\nu$ . Since  $2\nu < 1$  and  $\alpha = \beta = 2$ , then by repeating the proof of Lemma 2.1, we can prove that  $J'_\nu(|u|, |v|) = 0$ . By the maximum principle,  $|u|, |v| > 0$  in  $\mathbb{R}^4 \setminus \{0\}$ . Therefore,  $(|u|, |v|)$  is a positive ground state solution of (1.20).

To finish the proof, it suffices to prove  $c_\nu \rightarrow \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2)$  as  $\nu \rightarrow 0$ . From the above argument, we may assume that  $(u_\nu, v_\nu)$  is a positive ground state solution of (1.20) with  $c_\nu = J_\nu(u_\nu, v_\nu)$  for any  $\nu \in (0, \nu_1)$ . Since  $c_\nu < \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2)$ , we see that  $(u_\nu, v_\nu)$  are uniformly bounded in  $\mathbb{D}$ . Then

$$\|u_\nu\|_{\lambda_1}^2 = |u_\nu|_4^4 + 2\nu \int_{\mathbb{R}^4} u_\nu^2 v_\nu^2 \leq S(\lambda_1)^{-2} \|u\|_{\lambda_1}^4 + O(\nu).$$

From (3.19) we see that  $\liminf_{\nu \rightarrow 0} \|u_\nu\|_{\lambda_1}^2 > 0$ , so  $\liminf_{\nu \rightarrow 0} \|u_\nu\|_{\lambda_1}^2 \geq S(\lambda_1)^2$ . Similarly, we can prove that  $\liminf_{\nu \rightarrow 0} \|v_\nu\|_{\lambda_2}^2 \geq S(\lambda_2)^2$ , and so  $\liminf_{\nu \rightarrow 0} c_\nu \geq \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2)$ . That is,  $\lim_{\nu \rightarrow 0} c_\nu = \frac{1}{4} (S(\lambda_1)^2 + S(\lambda_2)^2)$ . This completes the proof.  $\square$

#### 4. PROOF OF THEOREM 1.5: A VARIATIONAL PERTURBATION APPROACH

In this section, we give the proof of Theorem 1.5, and this result will be used in the proof of Theorem 1.3. Assume that  $N \geq 3$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$  and (1.5) hold. Let  $\nu > 0$ . To obtain positive solutions of (1.4), we consider the following modified problem:

$$(4.1) \quad \begin{cases} -\Delta u - \frac{\lambda_1}{|x|^2} u - u_+^{2^*-1} = \nu \alpha u_+^{\alpha-1} v_+^\beta, & x \in \mathbb{R}^N, \\ -\Delta v - \frac{\lambda_2}{|x|^2} v - v_+^{2^*-1} = \nu \beta u_+^\alpha v_+^{\beta-1}, & x \in \mathbb{R}^N, \\ u(x), v(x) \in D^{1,2}(\mathbb{R}^N), \end{cases}$$

where  $u_\pm(x) := \max\{\pm u(x), 0\}$  and so does  $v_\pm$ . The associated energy functional of (4.1) is

$$(4.2) \quad \bar{J}_\nu(u, v) := \frac{1}{2} \|u\|_{\lambda_1}^2 + \frac{1}{2} \|v\|_{\lambda_2}^2 - \frac{1}{2^*} \int_{\mathbb{R}^N} (u_+^{2^*} + v_+^{2^*}) - \nu \int_{\mathbb{R}^N} u_+^\alpha v_+^\beta.$$

Then it is standard to prove that  $\bar{J} \in C^1(\mathbb{D}, \mathbb{R})$ . Define

$$C_{0,r}^\infty(\mathbb{R}^N) := \{u \in C_0^\infty(\mathbb{R}^N) : u \text{ is radially symmetric}\},$$

$$D_r^{1,2}(\mathbb{R}^N) := \{u \in D^{1,2}(\mathbb{R}^N) : u \text{ is radially symmetric}\},$$

and  $\mathbb{D}_r := D_r^{1,2}(\mathbb{R}^N) \times D_r^{1,2}(\mathbb{R}^N)$ . Then  $\mathbb{D}_r$  is a subspace of  $\mathbb{D}$  with norm  $\|\cdot\|_{\mathbb{D}}$ . In this section, we consider the functional  $\bar{J}_\nu$  restricted to  $\mathbb{D}_r$ . By Palais’s Symmetric Criticality Principle, any critical points of  $\bar{J}_\nu : \mathbb{D}_r \rightarrow \mathbb{R}$  are radially symmetric solutions of (4.1).

Without loss of generality, we assume that  $S(\lambda_1) \leq S(\lambda_2)$ . Then we see from (1.12) that

$$(4.3) \quad S(\lambda_1)^{N/4} = \|z_\mu^1\|_{\lambda_1} \leq \|z_\mu^2\|_{\lambda_2} = S(\lambda_2)^{N/4}, \quad \forall \mu > 0.$$

Define

$$P_{\lambda_i}(u) := \begin{cases} \frac{|u|_{2^*}^2}{\|u\|_{\lambda_i}^2}, & \text{if } u \in D^{1,2}(\mathbb{R}^N) \setminus \{0\}, \\ 0, & \text{if } u = 0, \end{cases} \quad i = 1, 2.$$

By (1.17) it is easy to prove that  $P_{\lambda_i} \in C(D^{1,2}(\mathbb{R}^N), \mathbb{R})$ . Note that  $P_{\lambda_i}(u) = 1$  is equivalent to  $I'_{\lambda_i}(u)u = 0$ . Then by (1.11)-(1.14) it is easy to check that

$$(4.4) \quad M_i := \frac{1}{N}S(\lambda_i)^{N/2} = \inf_{\substack{u \in D^{1,2}(\mathbb{R}^N) \\ P_{\lambda_i}(u)=1}} I_{\lambda_i}(u), \quad i = 1, 2.$$

By (1.12)-(1.14) we have

$$(4.5) \quad I_{\lambda_i}(tz_1^i) = \frac{t^2}{2}\|z_1^i\|_{\lambda_i}^2 - \frac{t^{2^*}}{2^*}|z_1^i|_{2^*}^{2^*} = \left(\frac{t^2}{2} - \frac{t^{2^*}}{2^*}\right)\|z_1^i\|_{\lambda_i}^2, \quad i = 1, 2.$$

Note that

$$(4.6) \quad I_{\lambda_i}(z_1^i) = \max_{t>0} I_{\lambda_i}(tz_1^i) = M_i, \quad i = 1, 2, \quad M_1 \leq M_2.$$

It is easily seen that there exist  $0 < t_0 < 1 < t_1$  such that

$$(4.7) \quad I_{\lambda_i}(tz_1^i) \leq M_1/4 \quad \text{for } t \in (0, t_0] \cup [t_1, \infty), \quad i = 1, 2.$$

Define

$$\tilde{\gamma}_i(t) := tz_1^i \text{ for } 0 \leq t \leq t_1, \quad i = 1, 2; \quad \tilde{\gamma}(t, s) := (\tilde{\gamma}_1(t), \tilde{\gamma}_2(s)).$$

Then  $\tilde{\gamma}(t, s) \in \mathbb{D}_r$  for all  $(t, s)$  and there exists a constant  $\mathcal{C} > 0$  such that

$$(4.8) \quad \max_{(t,s) \in [0,t_1] \times [0,t_1]} \|\tilde{\gamma}(t, s)\|_{\mathbb{D}} \leq \mathcal{C}.$$

Denote  $Q := [0, t_1] \times [0, t_1]$  for convenience. For  $\nu \geq 0$ , we define

$$a_\nu := \inf_{\gamma \in \Gamma} \max_{(t,s) \in Q} \bar{J}_\nu(\gamma(t, s)), \quad d_\nu := \max_{(t,s) \in Q} \bar{J}_\nu(\tilde{\gamma}(t, s)),$$

where

$$(4.9) \quad \Gamma := \left\{ \gamma \in C(Q, \mathbb{D}_r) : \max_{(t,s) \in Q} \|\gamma(t, s)\|_{\mathbb{D}} \leq 2S(\lambda_2)^{N/4} + \mathcal{C}, \right. \\ \left. \gamma(t, s) = \tilde{\gamma}(t, s) \text{ for } (t, s) \in Q \setminus (t_0, t_1) \times (t_0, t_1) \right\}.$$

The definition of  $a_\nu$  is different from the definitions of usual mountain-pass values (cf. [5]). All paths in  $\Gamma$  are required to be uniformly bounded in  $\mathbb{D}$  by  $2S(\lambda_2)^{N/4} + \mathcal{C}$ , which will play a crucial role in the proof of Lemma 4.1 below.

**Lemma 4.1.** *There hold  $d_\nu < d_0$  for  $\nu > 0$  and  $\lim_{\nu \rightarrow 0+} a_\nu = \lim_{\nu \rightarrow 0+} d_\nu = a_0 = d_0 = M_1 + M_2$ .*

*Proof.* Note that  $z_1^i > 0$  in  $\mathbb{R}^N \setminus \{0\}$ ; we have

$$d_0 = \max_{(t,s) \in Q} \bar{J}_0(\tilde{\gamma}(t, s)) = \max_{t \in (0,t_1)} I_{\lambda_1}(tz_1^1) + \max_{s \in (0,t_1)} I_{\lambda_2}(sz_1^2).$$

Then by (4.5)-(4.6) we see that

$$(4.10) \quad d_0 = \bar{J}_0(\tilde{\gamma}(1, 1)) = M_1 + M_2, \quad \bar{J}_0(\tilde{\gamma}(t, s)) < d_0 \text{ for } (t, s) \in Q \setminus \{(1, 1)\}.$$

Fix any  $\nu > 0$ . Note that there exists  $(t_\nu, s_\nu) \in Q \setminus \{(0, 0)\}$  such that  $d_\nu = \bar{J}_\nu(\tilde{\gamma}(t_\nu, s_\nu))$ . If  $(t_\nu, s_\nu) = (1, 1)$ , then we see from (4.10) that

$$d_\nu = \bar{J}_\nu(\tilde{\gamma}(1, 1)) = \bar{J}_0(\tilde{\gamma}(1, 1)) - \nu \int_{\mathbb{R}^N} (z_1^1)^\alpha (z_1^2)^\beta dx < d_0;$$

if  $(t_\nu, s_\nu) \neq (1, 1)$ , then we deduce from (4.10) again that

$$d_\nu = \bar{J}_\nu(\tilde{\gamma}(t_\nu, s_\nu)) = \bar{J}_0(\tilde{\gamma}(t_\nu, s_\nu)) - \nu t_\nu^\alpha s_\nu^\beta \int_{\mathbb{R}^N} (z_1^1)^\alpha (z_1^2)^\beta dx < d_0,$$

that is,  $d_\nu < d_0$  for any  $\nu > 0$ . Note that  $\tilde{\gamma} \in \Gamma$ ; we have  $a_\nu \leq d_\nu$ , that is

$$(4.11) \quad \liminf_{\nu \rightarrow 0+} a_\nu \leq \liminf_{\nu \rightarrow 0+} d_\nu, \quad \limsup_{\nu \rightarrow 0+} a_\nu \leq \limsup_{\nu \rightarrow 0+} d_\nu \leq d_0 \quad \text{and} \quad a_0 \leq d_0.$$

On the other hand, for any  $\gamma(t, s) = (\gamma_1(t, s), \gamma_2(t, s)) \in \Gamma$ , we define  $\Upsilon(\gamma) : [t_0, t_1]^2 \rightarrow \mathbb{R}^2$  by

$$\Upsilon(\gamma)(t, s) := (P_{\lambda_1}(\gamma_1(t, s)) - 1, P_{\lambda_2}(\gamma_2(t, s)) - 1).$$

By the definitions of  $P_{\lambda_i}$  and  $\tilde{\gamma}$ , it is easily seen that

$$\Upsilon(\tilde{\gamma})(t, s) = (t^{2^*-2} - 1, s^{2^*-2} - 1).$$

Then  $\deg(\Upsilon(\tilde{\gamma}), [t_0, t_1]^2, (0, 0)) = 1$ . By (4.9) we see that for any  $(t, s) \in \partial([t_0, t_1]^2)$ ,  $\Upsilon(\gamma)(t, s) = \Upsilon(\tilde{\gamma})(t, s) \neq (0, 0)$ . Therefore,  $\deg(\Upsilon(\gamma), [t_0, t_1]^2, (0, 0))$  is well defined and

$$\deg(\Upsilon(\gamma), [t_0, t_1]^2, (0, 0)) = \deg(\Upsilon(\tilde{\gamma}), [t_0, t_1]^2, (0, 0)) = 1.$$

Then there exists  $(t_2, s_2) \in [t_0, t_1]^2$  such that  $\Upsilon(\gamma)(t_2, s_2) = (0, 0)$ , that is,  $P_{\lambda_1}(\gamma_1(t_2, s_2)) = 1$  and  $P_{\lambda_2}(\gamma_2(t_2, s_2)) = 1$ . Combining these with (4.4), we have

$$(4.12) \quad \begin{aligned} \max_{(t,s) \in Q} \bar{J}_0(\gamma(t, s)) &\geq \bar{J}_0(\gamma(t_2, s_2)) \geq J_0(\gamma(t_2, s_2)) \\ &= I_{\lambda_1}(\gamma_1(t_2, s_2)) + I_{\lambda_2}(\gamma_2(t_2, s_2)) \\ &\geq M_1 + M_2 = d_0. \end{aligned}$$

Therefore,  $a_0 \geq d_0$ . By (4.11) one gets that  $a_0 = d_0$ .

Assume by contradiction that  $\liminf_{\nu \rightarrow 0+} a_\nu < d_0$ . Then there exists  $\varepsilon > 0$ ,  $\nu_n \rightarrow 0+$  and  $\gamma_n = (\gamma_{n,1}, \gamma_{n,2}) \in \Gamma$  such that

$$\max_{(t,s) \in Q} J_{\nu_n}(\gamma_n(t, s)) \leq d_0 - 2\varepsilon.$$

Recall that  $\alpha + \beta = 2^*$ . By (4.9) and Hölder's inequality, there exists  $n_0$  large enough such that

$$\max_{(t,s) \in Q} \nu_n \left| \int_{\mathbb{R}^N} (\gamma_{n,1}(t, s)_+)^{\alpha} (\gamma_{n,2}(t, s)_+)^{\beta} dx \right| \leq C\nu_n \leq \varepsilon, \quad \forall n \geq n_0,$$

and so

$$a_0 \leq \max_{(t,s) \in Q} \bar{J}_0(\gamma_n(t,s)) \leq \max_{(t,s) \in Q} \bar{J}_{\nu_n}(\gamma_n(t,s)) + \varepsilon \leq d_0 - \varepsilon, \quad \forall n \geq n_0,$$

a contradiction with  $a_0 = d_0$ . Therefore,  $\liminf_{\nu \rightarrow 0^+} a_\nu \geq d_0$ . Combining this with (4.11), we complete the proof.  $\square$

Recall (1.9); we define  $X := Z_1 \times Z_2 \subset \mathbb{D}_r$  and

$$(4.13) \quad \begin{aligned} X^\delta &:= \{(u,v) \in \mathbb{D}_r : \text{dist}((u,v), X) \leq \delta\}, \quad \bar{J}_\nu^d := \{(u,v) \in \mathbb{D}_r : \bar{J}_\nu(u,v) \leq d\}; \\ \delta &:= \min \left\{ \frac{1}{2}, \frac{1}{16} S(\lambda_1)^{N/4}, \frac{C}{4} \right\}. \end{aligned}$$

Here  $\text{dist}((u,v), X) := \inf\{\|(u-\varphi, v-\psi)\|_{\mathbb{D}} : (\varphi, \psi) \in X\}$ . Define

$$\|\bar{J}'_\nu(u,v)\| := \sup \left\{ \bar{J}'_\nu(u,v)(\psi, \phi) : (\psi, \phi) \in \mathbb{D}_r, \|( \psi, \phi)\|_{\mathbb{D}} = 1 \right\}.$$

**Lemma 4.2.** *Recall  $\delta$  in (4.13). Then there exist  $0 < \sigma < 1$  and  $\nu_3 \in (0, 1)$  such that  $\|\bar{J}'_\nu(u,v)\| \geq \sigma$  holds for any  $(u,v) \in \bar{J}_\nu^{d_\nu} \cap (X^\delta \setminus X^{\delta/2})$  and  $\nu \in (0, \nu_3]$ .*

*Proof.* Assume by contradiction that there exist  $\nu_n \rightarrow 0^+$  and  $(u_n, v_n) \in \bar{J}_{\nu_n}^{d_{\nu_n}} \cap (X^\delta \setminus X^{\delta/2})$  such that  $\|\bar{J}'_{\nu_n}(u_n, v_n)\| \rightarrow 0$ . Then there exist  $\mu_{i,n} > 0, i = 1, 2, n \in \mathbb{N}$ , such that

$$\left\| (u_n, v_n) - \left( z_{\mu_{1,n}}^1, z_{\mu_{2,n}}^2 \right) \right\|_{\mathbb{D}} \leq 2\delta, \quad \forall n \in \mathbb{N}.$$

Hence  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}_r$ , and up to a subsequence, we may assume that  $(u_n, v_n) \rightharpoonup (\bar{u}, \bar{v})$  weakly in  $\mathbb{D}_r$ . However, since  $(z_\mu^1, z_\mu^2) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}_r$  as  $\mu \rightarrow \infty$ , we know that  $X$  is not compact in  $\mathbb{D}_r$ . So it seems very difficult for us to show that  $\bar{u} \not\equiv 0$  and  $\bar{v} \not\equiv 0$ . To overcome this difficulty, let us define

$$\tilde{u}_n(x) := \mu_{1,n}^{\frac{N-2}{2}} u_n(\mu_{1,n}x), \quad \tilde{v}_n(x) := \mu_{2,n}^{\frac{N-2}{2}} v_n(\mu_{2,n}x).$$

Note that  $\|\cdot\|_{\lambda_i}, i = 1, 2$  are invariant with respect to the transformation  $u(\cdot) \mapsto \mu^{-\frac{N-2}{2}} u(\frac{\cdot}{\mu})$  for all  $\mu > 0$ . Therefore,

$$\|\tilde{u}_n - z_1^1\|_{\lambda_1} = \|u_n - z_{\mu_{1,n}}^1\|_{\lambda_1} \leq 2\delta, \quad \|\tilde{v}_n - z_1^2\|_{\lambda_2} = \|v_n - z_{\mu_{2,n}}^2\|_{\lambda_2} \leq 2\delta.$$

This means that  $(\tilde{u}_n, \tilde{v}_n)$  are uniformly bounded in  $\mathbb{D}_r$ . Up to a subsequence, we may assume that  $(\tilde{u}_n, \tilde{v}_n) \rightharpoonup (\tilde{u}, \tilde{v})$  weakly in  $\mathbb{D}_r \cap L^{2^*}(\mathbb{R}^N) \times L^{2^*}(\mathbb{R}^N)$ . Then we have  $\|\tilde{u} - z_1^1\|_{\lambda_1} \leq \liminf_{n \rightarrow \infty} \|\tilde{u}_n - z_1^1\|_{\lambda_1} \leq 2\delta$ . Combining this with (4.3) and (4.13), we get that  $\tilde{u} \not\equiv 0$ . Similarly,  $\tilde{v} \not\equiv 0$ .

Take any  $\tilde{\phi} \in C_{0,r}^\infty(\mathbb{R}^N)$  such that  $\|\tilde{\phi}\|_{\lambda_1} = 1$ ; we define

$$(4.14) \quad \phi_n(x) := \mu_{1,n}^{-\frac{N-2}{2}} \tilde{\phi} \left( \frac{x}{\mu_{1,n}} \right).$$

Then  $\|\phi_n\|_{\lambda_1} = \|\tilde{\phi}\|_{\lambda_1} = 1$ . Since  $\nu_n \rightarrow 0$ , by Hölder's inequality and the Sobolev inequality, we easily obtain that

$$\lim_{n \rightarrow \infty} \left| \nu_n \alpha \int_{\mathbb{R}^N} (u_n)_+^{\alpha-1} (v_n)_+^\beta \phi_n dx \right| = 0.$$

Therefore,

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \bar{J}'_{\nu_n}(u_n, v_n)(\phi_n, 0) \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \nabla u_n \nabla \phi_n - \frac{\lambda_1}{|x|^2} u_n \phi_n - (u_n)_+^{2^*-1} \phi_n \, dx \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \nabla \tilde{u}_n \nabla \tilde{\phi} - \frac{\lambda_1}{|x|^2} \tilde{u}_n \tilde{\phi} - (\tilde{u}_n)_+^{2^*-1} \tilde{\phi} \, dx \\ &= \int_{\mathbb{R}^N} \nabla \tilde{u} \nabla \tilde{\phi} - \frac{\lambda_1}{|x|^2} \tilde{u} \tilde{\phi} - \tilde{u}_+^{2^*-1} \tilde{\phi} \, dx \text{ holds for any } \tilde{\phi} \in C_{0,r}^\infty(\mathbb{R}^N), \end{aligned}$$

that is,  $-\Delta \tilde{u} - \frac{\lambda_1}{|x|^2} \tilde{u} = \tilde{u}_+^{2^*-1}$  and  $\tilde{u} \in D_r^{1,2}(\mathbb{R}^N)$ . By testing this equation with  $\tilde{u}_-$  we see that  $\tilde{u} \geq 0$ . By the maximum principle, one has that  $\tilde{u} > 0$  in  $\mathbb{R}^N \setminus \{0\}$ , that is,  $\tilde{u}$  is a positive solution of (1.8) with  $i = 1$ . Then by (1.8)-(1.9) we get that  $\tilde{u} \in Z_1$ . Similarly, we may prove that  $\tilde{v} \in Z_2$ , that is,  $(\tilde{u}, \tilde{v}) \in X$ . Recall that  $\bar{J}_{\nu_n}(u_n, v_n) \leq d_{\nu_n}$  and  $\alpha + \beta = 2^*$ . We deduce from Lemma 4.1 that

$$\begin{aligned} M_1 + M_2 &\geq \lim_{n \rightarrow \infty} \left( \bar{J}_{\nu_n}(u_n, v_n) - \frac{1}{2^*} \bar{J}'_{\nu_n}(u_n, v_n)(u_n, v_n) \right) \\ &= \frac{1}{N} \lim_{n \rightarrow \infty} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) = \frac{1}{N} \lim_{n \rightarrow \infty} (\|\tilde{u}_n\|_{\lambda_1}^2 + \|\tilde{v}_n\|_{\lambda_2}^2) \\ &\geq \frac{1}{N} \|\tilde{u}\|_{\lambda_1}^2 + \frac{1}{N} \|\tilde{v}\|_{\lambda_2}^2 = M_1 + M_2, \end{aligned}$$

which implies that all inequalities above are identities, so  $\bar{J}_{\nu_n}(u_n, v_n) \rightarrow M_1 + M_2$  and  $(\tilde{u}_n, \tilde{v}_n) \rightarrow (\tilde{u}, \tilde{v}) \in X$  strongly in  $\mathbb{D}_r$ . Then

$$\|(\tilde{u}_n, \tilde{v}_n) - (\tilde{u}, \tilde{v})\|_{\mathbb{D}} \leq \delta/4, \quad \text{for } n \text{ large enough,}$$

and so

$$\|(u_n, v_n) - (\bar{u}_n, \bar{v}_n)\|_{\mathbb{D}} \leq \delta/4, \quad \text{for } n \text{ large enough,}$$

where

$$(\bar{u}_n(x), \bar{v}_n(x)) := \left( \mu_{1,n}^{-\frac{N-2}{2}} \tilde{u} \left( \frac{x}{\mu_{1,n}} \right), \mu_{2,n}^{-\frac{N-2}{2}} \tilde{v} \left( \frac{x}{\mu_{2,n}} \right) \right) \in X.$$

This contradicts with  $(u_n, v_n) \notin X^{\delta/2}$  for any  $n$ . This completes the proof. □

**Lemma 4.3.** *There exists  $\nu_4 \in (0, \nu_3]$  and  $\varepsilon > 0$  such that for any  $\nu \in (0, \nu_4]$ ,*

$$\bar{J}_\nu(\tilde{\gamma}(t, s)) \geq a_\nu - \varepsilon \quad \text{implies that } \tilde{\gamma}(t, s) \in X^{\delta/2}.$$

*Proof.* Assume by contradiction that there exist  $\nu_n \rightarrow 0$ ,  $\varepsilon_n \rightarrow 0$  and  $(t_n, s_n) \in Q$  such that

$$(4.15) \quad \bar{J}_{\nu_n}(\tilde{\gamma}(t_n, s_n)) \geq a_{\nu_n} - \varepsilon_n \quad \text{and} \quad \tilde{\gamma}(t_n, s_n) \notin X^{\delta/2}, \quad \forall n \in \mathbb{N}.$$

Passing to a subsequence, we may assume that  $(t_n, s_n) \rightarrow (\tilde{t}, \tilde{s}) \in Q$ . Then by Lemma 4.1 and letting  $n \rightarrow \infty$  in (4.15), we have

$$J_0(\tilde{\gamma}(\tilde{t}, \tilde{s})) \geq \lim_{n \rightarrow \infty} a_{\nu_n} = M_1 + M_2.$$

Combining this with (4.10), we obtain that  $(\tilde{t}, \tilde{s}) = (1, 1)$ . Hence,

$$\lim_{n \rightarrow \infty} \|\tilde{\gamma}(t_n, s_n) - \tilde{\gamma}(1, 1)\|_{\mathbb{D}} = 0.$$

However,  $\tilde{\gamma}(1, 1) = (z_1^1, z_1^2) \in X$ , which is a contradiction with (4.15). □

Let

$$(4.16) \quad \varepsilon_0 := \min \left\{ \frac{\varepsilon}{2}, \frac{M_1}{4}, \frac{1}{8} \delta \sigma^2 \right\},$$

where  $\delta, \sigma$  are seen in Lemma 4.2. By Lemma 4.1 there exists  $\nu_2 \in (0, \nu_4]$  such that

$$(4.17) \quad |a_\nu - d_\nu| < \varepsilon_0, \quad |a_\nu - (M_1 + M_2)| < \varepsilon_0, \quad \forall \nu \in (0, \nu_2].$$

**Lemma 4.4.** *For any fixed  $\nu \in (0, \nu_2]$ , there exists  $\{(u_n, v_n)\}_{n=1}^\infty \subset X^\delta \cap \bar{J}_\nu^{d_\nu}$  such that*

$$\left\| \bar{J}'_\nu(u_n, v_n) \right\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

*Proof.* Fix any  $\nu \in (0, \nu_2]$ . Assume by contradiction that there exists  $0 < l(\nu) < 1$  such that  $\|\bar{J}'_\nu(u, v)\| \geq l(\nu)$  on  $X^\delta \cap \bar{J}_\nu^{d_\nu}$ . Then there exists a pseudo-gradient vector field  $T_\nu$  in  $\mathbb{D}_r$  which is defined on a neighborhood  $Z_\nu \subset \mathbb{D}_r$  of  $X^\delta \cap \bar{J}_\nu^{d_\nu}$  (cf. [31]), such that for any  $(u, v) \in Z_\nu$ , there holds

$$\begin{aligned} \|T_\nu(u, v)\|_{\mathbb{D}} &\leq 2 \min\{1, \|\bar{J}'_\nu(u, v)\|\}, \\ \bar{J}'_\nu(u, v)(T_\nu(u, v)) &\geq \min\{1, \|\bar{J}'_\nu(u, v)\|\} \|\bar{J}'_\nu(u, v)\|. \end{aligned}$$

Let  $\eta_\nu$  be a Lipschitz continuous function on  $\mathbb{D}_r$  such that  $0 \leq \eta_\nu \leq 1$ ,  $\eta_\nu \equiv 1$  on  $X^\delta \cap \bar{J}_\nu^{d_\nu}$  and  $\eta_\nu \equiv 0$  on  $\mathbb{D}_r \setminus Z_\nu$ . Let  $\xi_\nu$  be a Lipschitz continuous function on  $\mathbb{R}$  such that  $0 \leq \xi_\nu \leq 1$ ,  $\xi_\nu(l) \equiv 1$  if  $|l - a_\nu| \leq \frac{\varepsilon}{2}$  and  $\xi_\nu(l) \equiv 0$  if  $|l - a_\nu| \geq \varepsilon$ . Let

$$e_\nu(u, v) = \begin{cases} -\eta_\nu(u, v) \xi_\nu(\bar{J}_\nu(u, v)) T_\nu(u, v) & \text{if } (u, v) \in Z_\nu, \\ 0 & \text{if } (u, v) \in \mathbb{D}_r \setminus Z_\nu. \end{cases}$$

Then there exists a global solution  $\psi_\nu : \mathbb{D}_r \times [0, +\infty) \rightarrow \mathbb{D}_r$  to the following initial value problem:

$$\begin{cases} \frac{d}{d\theta} \psi_\nu(u, v, \theta) = e_\nu(\psi_\nu(u, v, \theta)), \\ \psi_\nu(u, v, 0) = (u, v). \end{cases}$$

It is easy to see from Lemma 4.2 and (4.16)-(4.17) that  $\psi_\nu$  has the following properties:

- (1)  $\psi_\nu(u, v, \theta) = (u, v)$  if  $\theta = 0$  or  $(u, v) \in \mathbb{D}_r \setminus Z_\nu$  or  $|\bar{J}_\nu(u, v) - a_\nu| \geq \varepsilon$ ;
- (2)  $\left\| \frac{d}{d\theta} \psi_\nu(u, v, \theta) \right\|_{\mathbb{D}} \leq 2$ ;
- (3)  $\frac{d}{d\theta} \bar{J}_\nu(\psi_\nu(u, v, \theta)) = \bar{J}'_\nu(\psi_\nu(u, v, \theta))(e_\nu(\psi_\nu(u, v, \theta))) \leq 0$ ;
- (4)  $\frac{d}{d\theta} J_\nu(\psi_\nu(u, v, \theta)) \leq -l(\nu)^2$  if  $\psi_\nu(u, v, \theta) \in X^\delta \cap (\bar{J}_\nu^{d_\nu} \setminus \bar{J}_\nu^{a_\nu - \varepsilon/2})$ ;
- (5)  $\frac{d}{d\theta} J_\nu(\psi_\nu(u, v, \theta)) \leq -\sigma^2$  if  $\psi_\nu(u, v, \theta) \in (X^\delta \setminus X^{\delta/2}) \cap (\bar{J}_\nu^{d_\nu} \setminus \bar{J}_\nu^{a_\nu - \varepsilon/2})$ .

*Step 1.* For any  $(t, s) \in Q$ , we claim that there exists  $\theta_{t,s} \in [0, +\infty)$  such that  $\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}) \in \bar{J}_\nu^{a_\nu - \varepsilon_0}$ , where  $\varepsilon_0$  is seen in (4.16).

Assume by contradiction that there exists  $(t, s) \in Q$  such that

$$\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta)) > a_\nu - \varepsilon_0, \quad \forall \theta \geq 0.$$

Note that  $\varepsilon_0 < \varepsilon$ ; we see from Lemma 4.3 that  $\tilde{\gamma}(t, s) \in X^{\delta/2}$ . Note that  $\bar{J}_\nu(\tilde{\gamma}(t, s)) \leq d_\nu < a_\nu + \varepsilon_0$ ; we see from the property (3) that

$$a_\nu - \varepsilon_0 < \bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta)) \leq d_\nu < a_\nu + \varepsilon_0, \quad \forall \theta \geq 0.$$



This implies  $\xi_\nu(\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta))) \equiv 1$ . If  $\psi_\nu(\tilde{\gamma}(t, s), \theta) \in X^\delta$  for all  $\theta \geq 0$ , then  $\eta_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta)) \equiv 1$ , and  $\|\bar{J}'_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta))\| \geq l(\nu)$  for all  $\theta > 0$ . Then we see from property (4) that

$$\bar{J}_\nu \left( \psi_\nu \left( \tilde{\gamma}(t, s), \frac{\varepsilon}{l(\nu)^2} \right) \right) \leq a_\nu + \frac{\varepsilon}{2} - \int_0^{\frac{\varepsilon}{l(\nu)^2}} l(\nu)^2 dt \leq a_\nu - \frac{\varepsilon}{2},$$

a contradiction. Thus, there exists  $\theta_{t,s} > 0$  such that  $\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}) \notin X^\delta$ . Since  $\tilde{\gamma}(t, s) \in X^{\delta/2}$ , so there exists  $0 < \theta_{t,s}^1 < \theta_{t,s}^2 \leq \theta_{t,s}$  such that  $\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^1) \in \partial X^{\delta/2}$ ,  $\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^2) \in \partial X^\delta$  and  $\psi_\nu(\tilde{\gamma}(t, s), \theta) \in X^\delta \setminus X^{\delta/2}$  for all  $\theta \in (\theta_{t,s}^1, \theta_{t,s}^2)$ .

Then by Lemma 4.2 we have  $\|\bar{J}'_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta))\| \geq \sigma$  for all  $\theta \in (\theta_{t,s}^1, \theta_{t,s}^2)$ . Then using property (2) we have

$$\delta/2 \leq \|\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^2) - \psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^1)\|_{\mathbb{D}} \leq 2|\theta_{t,s}^2 - \theta_{t,s}^1|,$$

that is,  $\theta_{t,s}^2 - \theta_{t,s}^1 \geq \delta/4$ . This implies from (4.16) and property (5) that

$$\begin{aligned} \bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^2)) &\leq \bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^1)) + \int_{\theta_{t,s}^1}^{\theta_{t,s}^2} \frac{d}{d\theta} \bar{J}_\nu(\psi_\nu(u, v, \theta)) d\theta \\ &< a_\nu + \varepsilon_0 - \sigma^2(\theta_{t,s}^2 - \theta_{t,s}^1) \leq a_\nu + \varepsilon_0 - \frac{1}{4}\delta\sigma^2 \\ &\leq a_\nu - \varepsilon_0, \end{aligned}$$

which is a contradiction.

By Step 1 we can define  $T(t, s) := \inf\{\theta \geq 0 : \bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta)) \leq a_\nu - \varepsilon_0\}$  and let  $\gamma(t, s) := \psi_\nu(\tilde{\gamma}(t, s), T(t, s))$ . Then  $\bar{J}_\nu(\gamma(t, s)) \leq a_\nu - \varepsilon_0$  for all  $(t, s) \in Q$ .

*Step 2.* We shall prove that  $\gamma(t, s) \in \Gamma$ .

For any  $(t, s) \in Q \setminus (t_0, t_1) \times (t_0, t_1)$ , by (4.6)-(4.7) and (4.16)-(4.17) we have

$$\begin{aligned} \bar{J}_\nu(\tilde{\gamma}(t, s)) &\leq \bar{J}_0(\tilde{\gamma}(t, s)) = I_{\lambda_1}(\tilde{\gamma}_1(t)) + I_{\lambda_2}(\tilde{\gamma}_2(s)) \\ &\leq M_1/4 + M_2 \leq M_1 + M_2 - 3\varepsilon_0 < a_\nu - \varepsilon_0, \end{aligned}$$

which implies that  $T(t, s) = 0$ , and so  $\gamma(t, s) = \tilde{\gamma}(t, s)$ . From the definition of  $\Gamma$  in (4.9), it suffices to prove that  $\|\gamma(t, s)\|_{\mathbb{D}} \leq 2S(\lambda_2)^{N/4} + \mathcal{C}$  for all  $(t, s) \in Q$  and  $T(t, s)$  is continuous with respect to  $(t, s)$ .

For any  $(t, s) \in Q$ , if  $\bar{J}_\nu(\tilde{\gamma}(t, s)) \leq a_\nu - \varepsilon_0$ , we have  $T(t, s) = 0$ , and so  $\gamma(t, s) = \tilde{\gamma}(t, s)$ , and by (4.8) we see that  $\|\gamma(t, s)\|_{\mathbb{D}} \leq \mathcal{C} < 2S(\lambda_2)^{N/4} + \mathcal{C}$ .

If  $\bar{J}_\nu(\tilde{\gamma}(t, s)) > a_\nu - \varepsilon_0$ , then  $\tilde{\gamma}(t, s) \in X^{\delta/2}$  and

$$a_\nu - \varepsilon_0 < \bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta)) \leq d_\nu < a_\nu + \varepsilon_0, \quad \forall \theta \in [0, T(t, s)).$$

This implies  $\xi_\nu(\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta))) \equiv 1$  for  $\theta \in [0, T(t, s))$ . If  $\psi_\nu(\tilde{\gamma}(t, s), T(t, s)) \notin X^\delta$ , then there exists  $0 < \theta_{t,s}^1 < \theta_{t,s}^2 < T(t, s)$  as above. Then we can prove that  $\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), \theta_{t,s}^2)) \leq a_\nu - \varepsilon_0$  as above, which contradicts the definition of  $T(t, s)$ . Therefore,  $\gamma(t, s) = \psi_\nu(\tilde{\gamma}(t, s), T(t, s)) \in X^\delta$ . Then there exists  $(u, v) \in X$  such that  $\|\gamma(t, s) - (u, v)\|_{\mathbb{D}} \leq 2\delta \leq \mathcal{C}$ . By (4.3) we have

$$\|\gamma(t, s)\|_{\mathbb{D}} \leq \|(u, v)\|_{\mathbb{D}} + \mathcal{C} \leq 2S(\lambda_2)^{N/4} + \mathcal{C}.$$

To prove the continuity of  $T(t, s)$ , we fix any  $(\tilde{t}, \tilde{s}) \in Q$ . Assume  $\bar{J}_\nu(\gamma(\tilde{t}, \tilde{s})) < a_\nu - \varepsilon_0$  first. Then  $T(\tilde{t}, \tilde{s}) = 0$  from the definition of  $T(t, s)$ . So  $\bar{J}_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s})) < a_\nu - \varepsilon_0$ .

By the continuity of  $\tilde{\gamma}$ , there exists  $\tau > 0$  such that for any  $(t, s) \in (\tilde{t} - \tau, \tilde{t} + \tau) \times (\tilde{s} - \tau, \tilde{s} + \tau) \cap Q$ , there holds  $\bar{J}_\nu(\tilde{\gamma}(t, s)) < a_\nu - \varepsilon_0$ , that is,  $T(t, s) = 0$  for any  $(t, s) \in (\tilde{t} - \tau, \tilde{t} + \tau) \times (\tilde{s} - \tau, \tilde{s} + \tau) \cap Q$ , and so  $T$  is continuous at  $(\tilde{t}, \tilde{s})$ . Now we assume that  $\bar{J}_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s})) = a_\nu - \varepsilon_0$ . Then from the previous proof we see that  $\gamma(\tilde{t}, \tilde{s}) = \psi_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s}), T(\tilde{t}, \tilde{s})) \in X^\delta$ , and so

$$\left\| \bar{J}'_\nu(\psi_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s}), T(\tilde{t}, \tilde{s}))) \right\| \geq l(\nu) > 0.$$

Then for any  $\omega > 0$ , we have  $\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s}), T(\tilde{t}, \tilde{s}) + \omega)) < a_\nu - \varepsilon_0$ . By the continuity of  $\psi_\nu$  and  $\tilde{\gamma}$ , there exists  $\tau > 0$  such that for any  $(t, s) \in (\tilde{t} - \tau, \tilde{t} + \tau) \times (\tilde{s} - \tau, \tilde{s} + \tau) \cap Q$ , we have  $\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(t, s), T(\tilde{t}, \tilde{s}) + \omega)) < a_\nu - \varepsilon_0$ , so  $T(t, s) \leq T(\tilde{t}, \tilde{s}) + \omega$ . It follows that

$$0 \leq \limsup_{(t,s) \rightarrow (\tilde{t}, \tilde{s})} T(t, s) \leq T(\tilde{t}, \tilde{s}).$$

If  $T(\tilde{t}, \tilde{s}) = 0$ , we have  $\lim_{(t,s) \rightarrow (\tilde{t}, \tilde{s})} T(t, s) = T(\tilde{t}, \tilde{s})$  immediately. If  $T(\tilde{t}, \tilde{s}) > 0$ , then for any  $0 < \omega < T(\tilde{t}, \tilde{s})$  we similarly have  $\bar{J}_\nu(\psi_\nu(\tilde{\gamma}(\tilde{t}, \tilde{s}), T(\tilde{t}, \tilde{s}) - \omega)) > a_\nu - \varepsilon_0$ . By the continuity of  $\psi_\nu$  and  $\tilde{\gamma}$  again, we easily obtain that

$$\liminf_{(t,s) \rightarrow (\tilde{t}, \tilde{s})} T(t, s) \geq T(\tilde{t}, \tilde{s}).$$

Thus  $T$  is continuous at  $(\tilde{t}, \tilde{s})$ . This completes the proof of Step 2.

Now, we have proved that  $\gamma(t, s) \in \Gamma$  and  $\max_{(t,s) \in Q} \bar{J}_\nu(\gamma(t, s)) \leq a_\nu - \varepsilon_0$ , which contradicts the definition of  $a_\nu$ . This completes the proof.  $\square$

*Proof of Theorem 1.5.* Fix any  $\nu \in (0, \nu_2]$ . By Lemma 4.4 there exists  $\{(u_n, v_n)\}_{n=1}^\infty \subset X^\delta \cap \bar{J}_\nu^{d_\nu}$  such that

$$\left\| \bar{J}'_\nu(u_n, v_n) \right\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Note that there exist  $\mu_{i,n} > 0, i = 1, 2, n \in \mathbb{N}$ , such that

$$(4.18) \quad \left\| (u_n, v_n) - \left( z_{\mu_{1,n}}^1, z_{\mu_{2,n}}^2 \right) \right\|_{\mathbb{D}} \leq 2\delta, \quad \forall n \in \mathbb{N}.$$

By (4.3),  $\{(u_n, v_n), n \geq 1\}$  are uniformly bounded in  $\mathbb{D}_r$ . Up to a subsequence, we may assume that  $(u_n, v_n) \rightharpoonup (u, v)$  weakly in  $\mathbb{D}_r$ . As pointed out before, since  $X$  is not compact in  $\mathbb{D}_r$ , it seems very difficult to prove that  $u \neq 0$  and  $v \neq 0$ . To overcome this difficulty, let us define

$$(4.19) \quad \begin{aligned} \tilde{u}_n(x) &:= \mu_{1,n}^{\frac{N-2}{2}} u_n(\mu_{1,n}x), & \tilde{v}_n(x) &:= \mu_{1,n}^{\frac{N-2}{2}} v_n(\mu_{1,n}x); \\ \bar{u}_n(x) &:= \mu_{2,n}^{\frac{N-2}{2}} u_n(\mu_{2,n}x), & \bar{v}_n(x) &:= \mu_{2,n}^{\frac{N-2}{2}} v_n(\mu_{2,n}x). \end{aligned}$$

Then by a direct computation, we see that  $\|\bar{J}'_\nu(\tilde{u}_n, \tilde{v}_n)\| \rightarrow 0$  and  $\|\bar{J}'_\nu(\bar{u}_n, \bar{v}_n)\| \rightarrow 0$  as  $n \rightarrow \infty$ . Moreover,

$$\|\tilde{u}_n - z_1^1\|_{\lambda_1} \leq 2\delta, \quad \|\bar{v}_n - z_1^2\|_{\lambda_2} \leq 2\delta.$$

Up to a subsequence, we may assume that  $(\tilde{u}_n, \tilde{v}_n) \rightharpoonup (\tilde{u}, \tilde{v})$  and  $(\bar{u}_n, \bar{v}_n) \rightharpoonup (\bar{u}, \bar{v})$  weakly in  $\mathbb{D}_r \cap L^{2^*}(\mathbb{R}^N) \times L^{2^*}(\mathbb{R}^N)$ . Then  $\bar{J}'_\nu(\tilde{u}, \tilde{v}) = 0$  and  $\bar{J}'_\nu(\bar{u}, \bar{v}) = 0$ . Moreover, as in the proof of Lemma 4.2, we get that  $\tilde{u} \neq 0$  and  $\bar{v} \neq 0$ .

Now we claim that either  $\tilde{v} \neq 0$  or  $\bar{u} \neq 0$ .

Assume by contradiction that both  $\tilde{v} \equiv 0$  and  $\bar{u} \equiv 0$ . Then  $\tilde{v}_n \rightharpoonup 0$  weakly in  $D_r^{1,2}(\mathbb{R}^N) \cap L^{2^*}(\mathbb{R}^N)$ . Hence it is easy to prove that  $(\tilde{v}_n)_+^{2^*-1} \rightharpoonup 0$  in  $L^{\frac{2^*}{2^*-1}}(\mathbb{R}^N)$ .

Take any  $\tilde{\phi} \in C_{0,r}^\infty(\mathbb{R}^N)$  such that  $\|\tilde{\phi}\|_{\lambda_1} = 1$ , and  $\phi_n$  is defined in (4.14). Then we see from Hölder's inequality and  $\alpha + \beta = 2^*$  that

$$(4.20) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \left| \nu \alpha \int_{\mathbb{R}^N} (u_n)_+^{\alpha-1} (v_n)_+^\beta \phi_n \, dx \right| = \lim_{n \rightarrow \infty} \left| \nu \alpha \int_{\mathbb{R}^N} (\tilde{u}_n)_+^{\alpha-1} (\tilde{v}_n)_+^\beta \tilde{\phi} \, dx \right| \\ & \leq \lim_{n \rightarrow \infty} C \left( \int_{\mathbb{R}^N} (\tilde{u}_n)_+^{2^*-1} |\tilde{\phi}| \, dx \right)^{\frac{\alpha-1}{2^*-1}} \left( \int_{\mathbb{R}^N} (\tilde{v}_n)_+^{2^*-1} |\tilde{\phi}| \, dx \right)^{\frac{\beta}{2^*-1}} = 0. \end{aligned}$$

Combining this with  $\bar{J}'_\nu(u_n, v_n)(\phi_n, 0) \rightarrow 0$ , we may repeat the proof of Lemma 4.2 and then get that  $\tilde{u} \in Z_1$ . Similarly, we can prove that  $\bar{v} \in Z_2$ . Then

$$(4.21) \quad \begin{aligned} d_\nu & \geq \lim_{n \rightarrow \infty} \left( \bar{J}_\nu(u_n, v_n) - \frac{1}{2^*} \bar{J}'_\nu(u_n, v_n)(u_n, v_n) \right) \\ & = \frac{1}{N} \lim_{n \rightarrow \infty} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) = \frac{1}{N} \lim_{n \rightarrow \infty} (\|\tilde{u}_n\|_{\lambda_1}^2 + \|\bar{v}_n\|_{\lambda_2}^2) \\ & \geq \frac{1}{N} \|\tilde{u}\|_{\lambda_1}^2 + \frac{1}{N} \|\bar{v}\|_{\lambda_2}^2 = M_1 + M_2 = d_0 > d_\nu, \end{aligned}$$

a contradiction. So either  $\tilde{v} \not\equiv 0$  or  $\bar{u} \not\equiv 0$ . Without loss of generality, we may assume that  $\tilde{v} \not\equiv 0$ . Note that  $\tilde{u} \not\equiv 0$  and  $\bar{J}'_\nu(\tilde{u}, \tilde{v}) = 0$ . Then by testing (4.1) with  $\tilde{u}_-$  and  $\tilde{v}_-$ , we see from Hardy's inequality (1.6) that  $\tilde{u} \geq 0$  and  $\tilde{v} \geq 0$ . By the maximum principle, one has that  $\tilde{u} > 0$  and  $\tilde{v} > 0$  in  $\mathbb{R}^N \setminus \{0\}$ . Hence,  $(\tilde{u}, \tilde{v})$  is a positive solution of (1.4), which is radially symmetric. Moreover,

$$\begin{aligned} d_\nu & \geq \lim_{n \rightarrow \infty} \left( \bar{J}_\nu(u_n, v_n) - \frac{1}{2^*} \bar{J}'_\nu(u_n, v_n)(u_n, v_n) \right) \\ & = \frac{1}{N} \lim_{n \rightarrow \infty} (\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) = \frac{1}{N} \lim_{n \rightarrow \infty} (\|\tilde{u}_n\|_{\lambda_1}^2 + \|\tilde{v}_n\|_{\lambda_2}^2) \\ & \geq \frac{1}{N} \|\tilde{u}\|_{\lambda_1}^2 + \frac{1}{N} \|\tilde{v}\|_{\lambda_2}^2 = J_\nu(\tilde{u}, \tilde{v}), \end{aligned}$$

that is,  $J_\nu(\tilde{u}, \tilde{v}) \leq d_\nu < d_0 = \frac{1}{N}(S(\lambda_1)^{N/2} + S(\lambda_2)^{N/2})$ . This completes the proof. □

### 5. PROOF OF THEOREM 1.3: THE CASE $N = 3$

In this section, we give the proof of Theorem 1.3. Assume that  $N = 3$ ,  $\alpha + \beta = 2^* = 6$ ,  $\alpha \geq 2$ ,  $\beta \geq 2$ ,  $\lambda_1, \lambda_2 \in (0, \Lambda_3)$  and  $\nu > 0$ . Recall from (1.11) that  $S(\lambda_i) < S$ ; we take a  $\varepsilon_0 \in (0, 1/2)$  such that

$$(5.1) \quad \begin{cases} \frac{\max\{S(\lambda_1)^{\frac{3}{2}}, S(\lambda_2)^{\frac{3}{2}}\}}{1-\varepsilon_0} \leq S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \leq (2 - 2\varepsilon_0)S^{\frac{3}{2}}, \\ (1 - 2\varepsilon_0)S(\lambda_2)^{\frac{3}{2}} \geq \varepsilon_0 S(\lambda_1)^{\frac{3}{2}}, \quad (1 - 2\varepsilon_0)S(\lambda_1)^{\frac{3}{2}} \geq \varepsilon_0 S(\lambda_2)^{\frac{3}{2}}. \end{cases}$$

Define

$$K_\nu := \{(u, v) \in \mathbb{D} : u \not\equiv 0, v \not\equiv 0, J'_\nu(u, v) = 0\}$$

as the set of nontrivial critical points of  $J_\nu$ , and

$$(5.2) \quad b_\nu := \inf_{(u,v) \in K_\nu} J_\nu(u, v).$$

By Theorem 1.5 we see that for any  $\nu \in (0, \nu_2]$ ,  $K_\nu \neq \emptyset$ ,  $b_\nu$  is well defined and

$$b_\nu < \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right).$$

Note that  $K_\nu \subset \mathcal{N}_\nu$ , so  $b_\nu \geq c_\nu > 0$ . Define

$$(5.3) \quad \mathcal{C}_0 := \max \left\{ \left[ S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right]^{\frac{\alpha}{6}}, \left[ S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right]^{\frac{\beta}{6}} \right\}.$$

Then for any  $(u, v) \in \mathcal{N}_\nu$  with  $J_\nu(u, v) < \frac{1}{3}(S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}})$ , since

$$\int_{\mathbb{R}^3} (u^6 + v^6 + 6\nu|u|^\alpha|v|^\beta) = 3J_\nu(u, v) < S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}},$$

we have

$$(5.4) \quad |u|_6^\alpha \leq \mathcal{C}_0, \quad |v|_6^\beta \leq \mathcal{C}_0.$$

Then we see from Hölder’s inequality and (1.17) that

$$S(\lambda_1)|u|_6^2 \leq \|u\|_{\lambda_1}^2 = |u|_6^6 + \nu\alpha \int_{\mathbb{R}^3} |u|^\alpha|v|^\beta \leq |u|_6^6 + \nu\alpha\mathcal{C}_0|u|_6^\alpha,$$

that is, we can obtain

$$(5.5) \quad \begin{cases} S(\lambda_1)|u|_6^2 \leq |u|_6^6 + \nu\alpha\mathcal{C}_0|u|_6^\alpha, \\ S(\lambda_2)|v|_6^2 \leq |v|_6^6 + \nu\beta\mathcal{C}_0|v|_6^\beta. \end{cases}$$

The following result was introduced in [1].

**Lemma 5.1** (see [1, Lemma 3.3]). *Let  $N \geq 3$ ,  $A, B > 0$ , and  $\theta \geq 2$  be fixed. For any  $\nu > 0$ , let*

$$S_\nu := \left\{ \sigma > 0 : A\sigma^{\frac{2}{2^*}} \leq \sigma + \nu B\sigma^{\frac{\theta}{2^*}} \right\}.$$

*Then for any  $\varepsilon > 0$ , there exists  $\nu_1 > 0$  depending only on  $\varepsilon, A, B, \theta$  and  $N$ , such that*

$$\inf S_\nu \geq (1 - \varepsilon)A^{\frac{N}{2}} \quad \text{for all } 0 < \nu < \nu_1.$$

Recall that  $\alpha, \beta \geq 2$  and  $2^* = 6$ . From Lemma 5.1, we have the following result trivially.

**Lemma 5.2.** *Recall  $\varepsilon_0$  in (5.1). Then there exists  $\tilde{\nu}_1 \in (0, \nu_2]$  such that for any  $\nu \in (0, \tilde{\nu}_1)$  there hold*

$$(5.6) \quad S(\lambda_1)\sigma^{\frac{1}{3}} \leq \sigma + \nu\alpha\mathcal{C}_0\sigma^{\frac{\alpha}{6}}, \quad \sigma > 0 \Rightarrow \sigma \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}},$$

$$(5.7) \quad S(\lambda_2)\sigma^{\frac{1}{3}} \leq \sigma + \nu\beta\mathcal{C}_0\sigma^{\frac{\beta}{6}}, \quad \sigma > 0 \Rightarrow \sigma \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}},$$

$$(5.8) \quad S\sigma^{\frac{1}{3}} \leq \sigma + \nu\alpha\mathcal{C}_0\sigma^{\frac{\alpha}{6}}, \quad \sigma > 0 \Rightarrow \sigma \geq (1 - \varepsilon_0)S^{\frac{3}{2}},$$

$$(5.9) \quad S\sigma^{\frac{1}{3}} \leq \sigma + \nu\beta\mathcal{C}_0\sigma^{\frac{\beta}{6}}, \quad \sigma > 0 \Rightarrow \sigma \geq (1 - \varepsilon_0)S^{\frac{3}{2}}.$$

The following lemma is the counterpart of Lemma 3.3 for the case  $N = 3$ . The first part of the proof is similar to that of Lemma 3.3, so we do not give the details, but the latter part of the proof is quite different.

**Lemma 5.3.** *Assume that  $\nu \in (0, \tilde{\nu}_1)$ . Let  $(u_n, v_n) \in K_\nu$  be a minimizing sequence of  $b_\nu$ , and  $(u_n, v_n) \rightharpoonup (0, 0)$  weakly in  $\mathbb{D}$ . Then for any  $r > 0$  and for every  $\varepsilon \in (-r, 0) \cup (0, r)$ , there exists  $\rho \in (\varepsilon, 0) \cup (0, \varepsilon)$  such that, up to a subsequence,*

$$(5.10) \quad \text{either } \int_{B_{r+\rho}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0 \text{ or } \int_{\mathbb{R}^N \setminus B_{r+\rho}} (|\nabla u_n|^2 + |\nabla v_n|^2) \rightarrow 0.$$

*Proof.* Fix any  $\nu \in (0, \tilde{\nu}_1)$ . Without loss of generality, we only consider the case  $\varepsilon \in (0, r)$  (the proof for the case  $\varepsilon \in (-r, 0)$  is similar). Since  $(u_n, v_n) \in K_\nu$  is a minimizing sequence of  $b_\nu$ , we may assume that  $J_\nu(u_n, v_n) < \frac{1}{3}(S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}})$ , and so  $(u_n, v_n)$  satisfy (5.4)-(5.5) for all  $n$ . Then by (5.6)-(5.7) of Lemma 5.2 we have

$$(5.11) \quad |u_n|_6^6 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad |v_n|_6^6 \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}, \quad \forall n \in \mathbb{N}.$$

Note that  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}$  and  $J'_\nu(u_n, v_n) = 0$ . Then by repeating the argument of Lemma 3.3 with trivial modifications and using the same notation  $w_{i,n}, \sigma_{i,n}, u_{i,n}, v_{i,n}$  with the same definitions as (3.21)-(3.26), there exists  $\rho \in (0, \varepsilon)$  such that  $u_{i,n}, v_{i,n}, i = 1, 2$ , satisfy (3.27)-(3.32). Moreover, we can prove that

$$(5.12) \quad |u_n|_6^6 = |u_{1,n}|_6^6 + |u_{2,n}|_6^6 + o(1), \quad |v_n|_6^6 = |v_{1,n}|_6^6 + |v_{2,n}|_6^6 + o(1).$$

Now we claim that

$$(5.13) \quad \text{either } \lim_{n \rightarrow \infty} (\|u_{1,n}\|^2 + \|v_{1,n}\|^2) = 0 \text{ or } \lim_{n \rightarrow \infty} (\|u_{2,n}\|^2 + \|v_{2,n}\|^2) = 0.$$

In fact, if (5.13) does not hold, then up to a subsequence,

$$(5.14) \quad \text{both } \lim_{n \rightarrow \infty} (\|u_{1,n}\|^2 + \|v_{1,n}\|^2) > 0 \text{ and } \lim_{n \rightarrow \infty} (\|u_{2,n}\|^2 + \|v_{2,n}\|^2) > 0.$$

We have the following several cases.

*Case 1.* Up to a subsequence, both  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 > 0$ .

Recall that norms  $\|\cdot\|_{\lambda_i}, i = 1, 2$ , are equivalent to  $\|\cdot\|$ . Note that (3.29)-(3.30) yield

$$(5.15) \quad \|u_{1,n}\|_{\lambda_1}^2 = |u_{1,n}|_6^6 + \nu\alpha \int_{\mathbb{R}^3} |u_{1,n}|^\alpha |v_{1,n}|^\beta + o(1),$$

$$(5.16) \quad \|v_{1,n}\|_{\lambda_2}^2 = |v_{1,n}|_6^6 + \nu\beta \int_{\mathbb{R}^3} |u_{1,n}|^\alpha |v_{1,n}|^\beta + o(1).$$

Hence, both  $A_1 := \liminf_{n \rightarrow \infty} |u_{1,n}|_6^6 > 0$  and  $B_1 := \liminf_{n \rightarrow \infty} |v_{1,n}|_6^6 > 0$ . Since  $(u_n, v_n)$  satisfy (5.4) for all  $n$ , by (5.12) and letting  $n \rightarrow \infty$  in (5.15)-(5.16), similarly as in (5.5) we can prove that

$$(5.17) \quad S(\lambda_1)A_1^{\frac{1}{3}} \leq A_1 + \nu\alpha C_0 A_1^{\frac{\alpha}{6}}, \quad S(\lambda_2)B_1^{\frac{1}{3}} \leq B_1 + \nu\beta C_0 B_1^{\frac{\beta}{6}}.$$

Then by (5.6)-(5.7) of Lemma 5.2 we have

$$(5.18) \quad A_1 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad B_1 \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}.$$

*Case 1.1.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{2,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 > 0$ .

Then similarly as above, we can prove that

$$(5.19) \quad A_2 := \liminf_{n \rightarrow \infty} |u_{2,n}|_6^6 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad B_2 := \liminf_{n \rightarrow \infty} |v_{2,n}|_6^6 \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}.$$

Combining this with (5.18), (5.1) and (5.12), we deduce that

$$\begin{aligned} b_\nu &= \lim_{n \rightarrow \infty} J_\nu(u_n, v_n) = \lim_{n \rightarrow \infty} \frac{1}{3} \left( |u_n|_6^6 + |v_n|_6^6 + 6\nu \int_{\mathbb{R}^3} |u_n|^\alpha |v_n|^\beta \right) \\ &\geq \lim_{n \rightarrow \infty} \frac{1}{3} (|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3} (A_1 + B_1 + A_2 + B_2) \\ &\geq \frac{2 - 2\varepsilon_0}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 1.1 is impossible.

*Case 1.2.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{2,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 = 0$ .

Then  $\int_{\mathbb{R}^3} |u_{2,n}|^\alpha |v_{2,n}|^\beta \rightarrow 0$  as  $n \rightarrow \infty$ , so (3.31) yields

$$S(\lambda_1) |u_{2,n}|_6^2 \leq \|u_{2,n}\|_{\lambda_1}^2 = |u_{2,n}|_6^6 + o(1),$$

and so  $A_2 \geq S(\lambda_1)^{3/2}$ . Then we conclude from (5.18), (5.1) and (5.12) that

$$\begin{aligned} b_\nu &\geq \lim_{n \rightarrow \infty} \frac{1}{3} (|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3} (A_1 + B_1 + A_2) \\ &\geq \frac{1 - \varepsilon_0}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) + \frac{1}{3} S(\lambda_1)^{\frac{3}{2}} \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 1.2 is impossible.

*Case 1.3.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{2,n}\|^2 = 0$  and  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 > 0$ .

Then (3.32) yields

$$S(\lambda_2) |v_{2,n}|_6^2 \leq \|v_{2,n}\|_{\lambda_2}^2 = |v_{2,n}|_6^6 + o(1),$$

and so  $B_2 \geq S(\lambda_2)^{3/2}$ . Then we conclude from (5.18), (5.1) and (5.12) that

$$\begin{aligned} b_\nu &\geq \lim_{n \rightarrow \infty} \frac{1}{3} (|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3} (A_1 + B_1 + B_2) \\ &\geq \frac{1 - \varepsilon_0}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) + \frac{1}{3} S(\lambda_2)^{\frac{3}{2}} \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 1.3 is impossible.

Since none of Cases 1.1-1.3 is true, Case 1 is impossible.

*Case 2.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 = 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 > 0$ .

Then similarly as in Case 1.3, we have  $B_1 \geq S(\lambda_2)^{3/2}$ . Moreover, we see from (5.11) and (5.12) that

$$\liminf_{n \rightarrow \infty} |u_{2,n}|_6^6 = \liminf_{n \rightarrow \infty} |u_n|_6^6 - \lim_{n \rightarrow \infty} |u_{1,n}|_6^6 > 0.$$

*Case 2.1.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{2,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 > 0$ .

Then similarly as above, we see that (5.19) holds, and so

$$\begin{aligned} b_\nu &\geq \lim_{n \rightarrow \infty} \frac{1}{3}(|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3}(B_1 + A_2 + B_2) \\ &\geq \frac{1 - \varepsilon_0}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) + \frac{1}{3} S(\lambda_2)^{\frac{3}{2}} \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 2.1 is impossible.

*Case 2.2.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{2,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{2,n}\|^2 = 0$ .

Then similarly as in Case 1.2, we have  $A_2 \geq S(\lambda_1)^{3/2}$ , and so

$$\begin{aligned} b_\nu &\geq \lim_{n \rightarrow \infty} \frac{1}{3}(|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3}(B_1 + A_2) \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 2.2 is impossible.

Since neither Case 2.1 nor Case 2.2 is true, Case 2 is impossible.

*Case 3.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_{1,n}\|^2 > 0$  and  $\lim_{n \rightarrow \infty} \|v_{1,n}\|^2 = 0$ .

By a similar argument as in Case 2, we get a contradiction. So Case 3 is impossible.

Since none of Cases 1, 2 and 3 are true, we see that (5.14) is impossible, that is, (5.13) holds. Recall the definitions (3.23)-(3.26) of  $(u_{i,n}, v_{i,n})$ , (5.10) follows directly from (5.13). This completes the proof.  $\square$

*Proof of Theorem 1.3.* Fix any  $\nu \in (0, \tilde{\nu}_1)$ . Take a sequence  $(\bar{u}_n, \bar{v}_n) \in K_\nu$  such that  $J_\nu(\bar{u}_n, \bar{v}_n) \rightarrow b_\nu$  as  $n \rightarrow \infty$ . Recall that  $E(u, v) = |\nabla u|^2 + |\nabla v|^2 - \frac{\lambda_1}{|x|^2}|u|^2 - \frac{\lambda_2}{|x|^2}|v|^2$  and there exists  $R_n > 0$  such that

$$\int_{B_{R_n}} E(\bar{u}_n, \bar{v}_n) = \int_{\mathbb{R}^N \setminus B_{R_n}} E(\bar{u}_n, \bar{v}_n) = \frac{1}{2}(\|\bar{u}_n\|_{\lambda_1}^2 + \|\bar{v}_n\|_{\lambda_2}^2).$$

Define

$$(u_n(x), v_n(x)) := \left( R_n^{\frac{N-2}{2}} \bar{u}_n(R_n x), R_n^{\frac{N-2}{2}} \bar{v}_n(R_n x) \right).$$

Then by a direct computation, we see that  $(u_n, v_n) \in K_\nu$  and  $J_\nu(u_n, v_n) \rightarrow b_\nu$ . Moreover,

$$(5.20) \quad \int_{B_1} E(u_n, v_n) = \int_{\mathbb{R}^N \setminus B_1} E(u_n, v_n) = \frac{1}{2}(\|u_n\|_{\lambda_1}^2 + \|v_n\|_{\lambda_2}^2) \rightarrow \frac{3}{2}b_\nu > 0.$$

Besides, we may assume that  $J_\nu(u_n, v_n) < \frac{1}{3}(S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}})$ , and so  $(u_n, v_n)$  satisfy (5.4)-(5.5) for all  $n$ . Then by (5.6)-(5.7) of Lemma 5.2 we have

$$(5.21) \quad |u_n|_6^6 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad |v_n|_6^6 \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}, \quad \forall n \in \mathbb{N}.$$

Note that  $(u_n, v_n)$  are uniformly bounded in  $\mathbb{D}$ . Then up to a subsequence, we assume that  $(u_n, v_n) \rightharpoonup (u, v)$  weakly in  $\mathbb{D}$ . Then  $J'_\nu(u_n, v_n) = 0$  implies  $J'_\nu(u, v) = 0$ .

*Step 1.* We show that both  $u \not\equiv 0$  and  $v \not\equiv 0$ , that is,  $(u, v) \in K_\nu$ . Moreover,  $J_\nu(u, v) = b_\nu$ .

*Case 1.*  $(u, v) \equiv (0, 0)$ .

Then we can apply Lemma 5.3 twice with  $r = 1$  and  $\varepsilon = \pm 1/4$  respectively, and there exist  $\rho^+ \in (0, 1/4)$  and  $\rho^- \in (-1/4, 0)$  such that the alternative (5.10) holds. Then by repeating the argument of Case 1 in the proof of Theorem 1.2 with trivial modifications, we get that  $\int_{\mathbb{R}^3} \frac{u_n^2}{|x|^2} = o(1)$ ,  $\int_{\mathbb{R}^3} \frac{v_n^2}{|x|^2} = o(1)$ , and so

$$S|u_n|_6^2 \leq \int_{\mathbb{R}^3} |\nabla u_n|^2 = \int_{\mathbb{R}^3} u_n^6 + \nu\alpha \int_{\mathbb{R}^3} |u_n|^\alpha |v_n|^\beta + o(1),$$

$$S|v_n|_6^2 \leq \int_{\mathbb{R}^3} |\nabla v_n|^2 = \int_{\mathbb{R}^3} v_n^6 + \nu\beta \int_{\mathbb{R}^3} |u_n|^\alpha |v_n|^\beta + o(1).$$

Denote  $A = \liminf_{n \rightarrow \infty} |u_n|_6^6$  and  $B = \liminf_{n \rightarrow \infty} |v_n|_6^6$ ; then (5.21) yields  $A > 0$  and  $B > 0$ . Then by Hölder’s inequality it is easy to prove that

$$SA^{\frac{1}{3}} \leq A + \nu\alpha C_0 A^{\frac{\alpha}{6}}, \quad SB^{\frac{1}{3}} \leq B + \nu\beta C_0 B^{\frac{\beta}{6}}.$$

Then by (5.8)-(5.9) of Lemma 5.2 we have

$$A \geq (1 - \varepsilon_0)S^{\frac{3}{2}}, \quad B \geq (1 - \varepsilon_0)S^{\frac{3}{2}}.$$

So we conclude from (5.1) that

$$b_\nu \geq \lim_{n \rightarrow \infty} \frac{1}{3}(|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3}(A + B) \geq \frac{2 - 2\varepsilon_0}{3}S^{\frac{3}{2}}$$

$$\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu,$$

a contradiction. So Case 1 is impossible.

*Case 2.* Either  $u \equiv 0, v \neq 0$  or  $u \neq 0, v \equiv 0$ .

Without loss of generality, we assume that  $u \neq 0, v \equiv 0$ . We see from  $J'_\nu(u, v)(u, 0) = 0$  that

$$S(\lambda_1)|u|_6^2 \leq \|u\|_{\lambda_1}^2 = |u|_6^6,$$

which implies  $|u|_6^6 \geq S(\lambda_1)^{3/2}$ .

*Case 2.1.* Up to a subsequence,  $\lim_{n \rightarrow \infty} \|u_n - u\| > 0$ .

Denote  $w_n = u_n - u$ . Note that  $J'_\nu(u_n, v_n) = 0$ . Then by the Brezis-Lieb Lemma ([7]) and Lemma 2.3 we conclude that

$$\|w_n\|_{\lambda_1}^2 = \int_{\mathbb{R}^3} w_n^6 + \nu\alpha \int_{\mathbb{R}^3} |w_n|^\alpha |v_n|^\beta + o(1),$$

$$\|v_n\|_{\lambda_2}^2 = \int_{\mathbb{R}^3} v_n^6 + \nu\beta \int_{\mathbb{R}^3} |w_n|^\beta |v_n|^\beta + o(1).$$

Denote  $C = \liminf_{n \rightarrow \infty} |w_n|_6^6$ ; then  $C > 0$ . Then by Hölder’s inequality it is easy to prove that

$$S(\lambda_1)C^{\frac{1}{3}} \leq C + \nu\alpha C_0 C^{\frac{\alpha}{6}}, \quad S(\lambda_2)B^{\frac{1}{3}} \leq B + \nu\beta C_0 B^{\frac{\beta}{6}}.$$

Then by (5.6)-(5.7) of Lemma 5.2 we have

$$C \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad B \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}.$$



So we conclude from (5.1) that

$$\begin{aligned} b_\nu &\geq \lim_{n \rightarrow \infty} \frac{1}{3}(|u_n|_6^6 + |v_n|_6^6) = \frac{1}{3}|u|_6^6 + \lim_{n \rightarrow \infty} \frac{1}{3}(|w_n|_6^6 + |v_n|_6^6) \\ &\geq \frac{1}{3}S(\lambda_1)^{\frac{3}{2}} + \frac{1}{3}(B + C) \geq \frac{1}{3}S(\lambda_1)^{\frac{3}{2}} + \frac{1 - \varepsilon_0}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So Case 2.1 is impossible.

Case 2.2.  $u_n \rightarrow u$  strongly in  $D^{1,2}(\mathbb{R}^3)$ .

Then  $u_n \rightarrow u$  strongly in  $L^6(\mathbb{R}^3)$ . Recall that  $v_n \rightarrow 0$  in  $D^{1,2}(\mathbb{R}^3)$ ; up to a subsequence,  $u_n \rightarrow u$  and  $v_n \rightarrow 0$  almost everywhere in  $\mathbb{R}^3$ . So Lemma 2.3 yields

$$\int_{\mathbb{R}^3} |u_n|^\alpha |v_n|^\beta = \int_{\mathbb{R}^3} |u_n - u|^\alpha |v_n|^\beta + o(1) = o(1).$$

Then we have

$$S(\lambda_2)|v_n|_6^2 \leq \|v_n\|_{\lambda_2}^2 = |v_n|_6^6 + o(1),$$

so  $B \geq S(\lambda_2)^{3/2}$ , and we conclude from (5.1) that

$$b_\nu \geq \lim_{n \rightarrow \infty} \frac{1}{3}(|u_n|_6^6 + |v_n|_6^6) \geq \frac{1}{3}(|u|_6^6 + B) \geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu,$$

a contradiction. So Case 2.2 is impossible, and thus Case 2 is impossible.

Since neither Case 1 nor Case 2 is true, we obtain that  $u \neq 0$  and  $v \neq 0$ . Since  $J'_\nu(u, v) = 0$ , thus  $(u, v) \in K_\nu$ . Then

$$b_\nu \leq J_\nu(u, v) = \frac{1}{3} \|(u, v)\|_{\mathbb{D}}^2 \leq \liminf_{n \rightarrow \infty} \frac{1}{3} \|(u_n, v_n)\|_{\mathbb{D}}^2 = \liminf_{n \rightarrow \infty} J_\nu(u_n, v_n) = b_\nu,$$

so  $J_\nu(u, v) = b_\nu$ , and  $(u_n, v_n) \rightarrow (u, v)$  strongly in  $\mathbb{D}$ . Then (5.21) implies that

$$(5.22) \quad |u|_6^6 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}, \quad |v|_6^6 \geq (1 - \varepsilon_0)S(\lambda_2)^{\frac{3}{2}}.$$

Step 2. We show that neither  $u$  or  $v$  is sign-changing, so  $(|u|, |v|)$  is a positive ground state solution of (1.4).

Assume by contradiction that  $u_+ \neq 0$  and  $u_- \neq 0$ . By  $J'_\nu(u, v)(u_\pm, 0) = 0$  we obtain

$$S(\lambda_1)|u_\pm|_6^2 \leq \|u_\pm\|_{\lambda_1}^2 = \int_{\mathbb{R}^3} u_\pm^6 + \nu\alpha \int_{\mathbb{R}^3} |u_\pm|^\alpha |v|^\beta.$$

Then

$$S(\lambda_1)|u_\pm|_6^2 \leq |u_\pm|_6^6 + \nu\alpha C_0 |u_\pm|_6^\alpha.$$

By (5.6) of Lemma 5.2 we have

$$|u_\pm|_6^6 \geq (1 - \varepsilon_0)S(\lambda_1)^{\frac{3}{2}}.$$

So we conclude from (5.1) and (5.22) that

$$\begin{aligned} b_\nu &\geq \frac{1}{3}(|u|_6^6 + |v|_6^6) \geq \frac{2 - 2\varepsilon_0}{3}S(\lambda_1)^{\frac{3}{2}} + \frac{1 - \varepsilon_0}{3}S(\lambda_2)^{\frac{3}{2}} \\ &\geq \frac{1}{3} \left( S(\lambda_1)^{\frac{3}{2}} + S(\lambda_2)^{\frac{3}{2}} \right) > b_\nu, \end{aligned}$$

a contradiction. So  $u$  is not sign-changing. Similarly,  $v$  is not sign-changing. That is,  $(|u|, |v|)$  is a solution of  $J_\nu$ . By the maximum principle, we see that  $|u| > 0$  and  $|v| > 0$  in  $\mathbb{R}^3 \setminus \{0\}$ . Since  $J_\nu(|u|, |v|) = b_\nu$ , then  $(|u|, |v|)$  is a positive ground state solution of (1.4).

*Step 3.* We show that  $b_\nu \rightarrow \frac{1}{3} (S(\lambda_1)^{3/2} + S(\lambda_2)^{3/2})$  as  $\nu \rightarrow 0$ .

From the above argument, we may assume that  $(u_\nu, v_\nu)$  is a positive ground state solution of (1.4) with  $b_\nu = J_\nu(u_\nu, v_\nu)$  for any  $\nu \in (0, \tilde{\nu}_1)$ . The rest of the argument is similar to that in the proof of Theorem 1.2, and we omit the details. This completes the proof.  $\square$

6. PROOF OF THEOREM 1.4: THE MOVING PLANES METHOD

In this section, we will use the moving planes method to prove Theorem 1.4. In the sequel, we assume that  $N = 3$  or  $N = 4$ ,  $\alpha + \beta = 2^*$ ,  $\alpha \geq 2$ ,  $\beta \geq 2$  and  $\lambda_1, \lambda_2 \in (0, \Lambda_N)$ . Fix any  $\nu > 0$ . Let  $(u, v)$  be any a positive solution of (1.4). For  $\lambda < 0$  we consider the reflection

$$x = (x_1, x_2, \dots, x_N) \mapsto x^\lambda = (2\lambda - x_1, x_2, \dots, x_N),$$

where  $x \in \Sigma^\lambda := \{x \in \mathbb{R}^N : x_1 < \lambda\}$ . Define  $u^\lambda(x) := u(x^\lambda)$  and  $v^\lambda(x) := v(x^\lambda)$ ; then

$$u(x) = u^\lambda(x), \quad v(x) = v^\lambda(x), \quad \text{for } x \in \partial\Sigma^\lambda = \{x \in \mathbb{R}^N : x_1 = \lambda\}.$$

Define  $w^\lambda(x) := u^\lambda(x) - u(x)$  and  $\sigma^\lambda(x) := v^\lambda(x) - v(x)$  for  $x \in \Sigma^\lambda$ . Then

$$(6.1) \quad w^\lambda(x) = \sigma^\lambda(x) = 0, \quad \forall x \in \partial\Sigma^\lambda.$$

Recall that  $(u, v)$  satisfies (1.4). Thus we have that

$$(6.2) \quad \begin{aligned} -\Delta w^\lambda(x) &= \frac{\lambda_1}{|x|^2} w^\lambda(x) + a_1^\lambda(x) w^\lambda(x) + a_2^\lambda(x) \sigma^\lambda(x) + \lambda_1 \left( \frac{1}{|x^\lambda|^2} - \frac{1}{|x|^2} \right) u^\lambda(x) \\ &\geq \frac{\lambda_1}{|x|^2} w^\lambda(x) + a_1^\lambda(x) w^\lambda(x) + a_2^\lambda(x) \sigma^\lambda(x) \end{aligned}$$

holds in  $\Sigma^\lambda \setminus \{0^\lambda\}$ , where

$$(6.3) \quad a_1^\lambda := \frac{(u^\lambda)^{2^*-1} - u^{2^*-1}}{u^\lambda - u} + \nu \alpha v^\beta \frac{(u^\lambda)^{\alpha-1} - u^{\alpha-1}}{u^\lambda - u} \geq 0,$$

$$(6.4) \quad a_2^\lambda := \nu \alpha (u^\lambda)^{\alpha-1} \frac{(v^\lambda)^\beta - v^\beta}{v^\lambda - v} \geq 0.$$

Similarly,

$$(6.5) \quad -\Delta \sigma^\lambda(x) \geq \frac{\lambda_2}{|x|^2} \sigma^\lambda(x) + b_1^\lambda(x) \sigma^\lambda(x) + b_2^\lambda(x) w^\lambda(x)$$

holds in  $\Sigma^\lambda \setminus \{0^\lambda\}$ , where

$$(6.6) \quad b_1^\lambda := \frac{(v^\lambda)^{2^*-1} - v^{2^*-1}}{v^\lambda - v} + \nu \beta u^\alpha \frac{(v^\lambda)^{\beta-1} - v^{\beta-1}}{v^\lambda - v} \geq 0,$$

$$(6.7) \quad b_2^\lambda := \nu \beta (v^\lambda)^{\beta-1} \frac{(u^\lambda)^\alpha - u^\alpha}{u^\lambda - u} \geq 0.$$

Define

$$\Omega_1^\lambda := \{x \in \Sigma^\lambda : w^\lambda(x) < 0\}, \quad \Omega_2^\lambda := \{x \in \Sigma^\lambda : \sigma^\lambda(x) < 0\}.$$

Since  $u, v \in L^{2^*}(\mathbb{R}^N)$  and  $\Omega_i^\lambda \subset \Sigma^\lambda$ , there exists  $\lambda_0 < 0$  such that for any  $\lambda \leq \lambda_0$ , we have

$$(6.8) \quad \left\| (2^* - 1)u^{2^*-2} + \nu\alpha(\alpha - 1)u^{\alpha-2}v^\beta \right\|_{L^{N/2}(\Omega_1^\lambda)} \leq \frac{1}{4} \left( 1 - \frac{\lambda_1}{\Lambda_N} \right) S,$$

$$(6.9) \quad \left\| (2^* - 1)v^{2^*-2} + \nu\beta(\beta - 1)u^\alpha v^{\beta-2} \right\|_{L^{N/2}(\Omega_2^\lambda)} \leq \frac{1}{4} \left( 1 - \frac{\lambda_2}{\Lambda_N} \right) S,$$

$$(6.10) \quad (\nu\alpha\beta)^2 \|u^{\alpha-1}v^{\beta-1}\|_{L^{N/2}(\Omega_1^\lambda \cap \Omega_2^\lambda)}^2 \leq \frac{1}{16} \left( 1 - \frac{\lambda_1}{\Lambda_N} \right) \left( 1 - \frac{\lambda_2}{\Lambda_N} \right) S^2.$$

*Step 1.* We claim that for any  $\lambda \leq \lambda_0$ , both  $w^\lambda > 0$  and  $\sigma^\lambda > 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ .

Fix any  $\lambda \leq \lambda_0$ . Define  $w_-^\lambda := \max\{-w^\lambda, 0\}$  and  $\sigma_-^\lambda := \max\{-\sigma^\lambda, 0\}$ ; then  $w_-^\lambda, \sigma_-^\lambda \in D^{1,2}(\mathbb{R}^N)$ . Testing (6.2) with  $w_-^\lambda$  and using Hölder’s inequality and Hardy’s inequality (1.6), we obtain

$$\begin{aligned} \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 &\leq \int_{\Omega_1^\lambda} \frac{\lambda_1}{|x|^2} |w_-^\lambda|^2 + \int_{\Omega_1^\lambda} a_1^\lambda |w_-^\lambda|^2 + \int_{\Omega_1^\lambda \cap \Omega_2^\lambda} a_2^\lambda w_-^\lambda \sigma_-^\lambda \\ &\leq \frac{\lambda_1}{\Lambda_N} \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 + \|a_1^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)}^2 \\ &\quad + \|a_2^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda \cap \Omega_2^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)} \|\sigma_-^\lambda\|_{L^{2^*}(\Omega_2^\lambda)}. \end{aligned}$$

When  $\theta \geq 1$ , we see from the mean value theorem that

$$\frac{s^\theta - t^\theta}{s - t} \leq \theta t^{\theta-1}, \quad \forall 0 < s < t.$$

Recall that  $u^\lambda < u$  in  $\Omega_1^\lambda$  and  $v^\lambda < v$  in  $\Omega_2^\lambda$ . Since  $\alpha \geq 2$  and  $\beta \geq 2$ , we see from (6.3)-(6.4) and (6.6)-(6.7) that

$$(6.11) \quad \begin{aligned} a_1^\lambda &\leq (2^* - 1)u^{2^*-2} + \nu\alpha(\alpha - 1)u^{\alpha-2}v^\beta, \quad \text{in } \Omega_1^\lambda, \\ a_2^\lambda, b_2^\lambda &\leq \nu\alpha\beta u^{\alpha-1}v^{\beta-1}, \quad \text{in } \Omega_1^\lambda \cap \Omega_2^\lambda, \\ b_1^\lambda &\leq (2^* - 1)v^{2^*-2} + \nu\beta(\beta - 1)u^\alpha v^{\beta-2}, \quad \text{in } \Omega_2^\lambda. \end{aligned}$$

Then we see from (6.8) and (1.13) that

$$\|a_1^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)}^2 \leq \frac{1}{4} \left( 1 - \frac{\lambda_1}{\Lambda_N} \right) \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2.$$

From above we obtain

$$(6.12) \quad \frac{3}{4} \left( 1 - \frac{\lambda_1}{\Lambda_N} \right) \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 \leq \|a_2^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda \cap \Omega_2^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)} \|\sigma_-^\lambda\|_{L^{2^*}(\Omega_2^\lambda)}.$$

Similarly, testing (6.5) with  $\sigma_-^\lambda$  we can prove that

$$(6.13) \quad \frac{3}{4} \left( 1 - \frac{\lambda_2}{\Lambda_N} \right) \int_{\Omega_2^\lambda} |\nabla \sigma_-^\lambda|^2 \leq \|b_2^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda \cap \Omega_2^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)} \|\sigma_-^\lambda\|_{L^{2^*}(\Omega_2^\lambda)}.$$

Let  $|\Omega|$  denotes the Lebesgue measure of  $\Omega$  in  $\mathbb{R}^N$ . If  $|\Omega_1^\lambda \cap \Omega_2^\lambda| > 0$ , then

$$(6.14) \quad \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 > 0, \quad \int_{\Omega_2^\lambda} |\nabla \sigma_-^\lambda|^2 > 0.$$

Combining this with (6.11)-(6.13), (1.13) and (6.10), we get

$$\begin{aligned} 0 &< \frac{9}{16} \left(1 - \frac{\lambda_1}{\Lambda_N}\right) \left(1 - \frac{\lambda_2}{\Lambda_N}\right) \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 \int_{\Omega_2^\lambda} |\nabla \sigma_-^\lambda|^2 \\ &\leq \|a_2^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda \cap \Omega_2^\lambda)} \|b_2^\lambda\|_{L^{\frac{N}{2}}(\Omega_1^\lambda \cap \Omega_2^\lambda)} \|w_-^\lambda\|_{L^{2^*}(\Omega_1^\lambda)}^2 \|\sigma_-^\lambda\|_{L^{2^*}(\Omega_2^\lambda)}^2 \\ &\leq \frac{1}{16} \left(1 - \frac{\lambda_1}{\Lambda_N}\right) \left(1 - \frac{\lambda_2}{\Lambda_N}\right) \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 \int_{\Omega_2^\lambda} |\nabla \sigma_-^\lambda|^2, \end{aligned}$$

a contradiction. Hence  $|\Omega_1^\lambda \cap \Omega_2^\lambda| = 0$ , and so (6.12)-(6.13) yield

$$(6.15) \quad \int_{\Omega_1^\lambda} |\nabla w_-^\lambda|^2 \leq 0, \quad \int_{\Omega_2^\lambda} |\nabla \sigma_-^\lambda|^2 \leq 0.$$

This implies that  $|\Omega_1^\lambda| = 0$  and  $|\Omega_2^\lambda| = 0$ . That is, both  $w^\lambda \geq 0$  and  $\sigma^\lambda \geq 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ . If  $w^\lambda \equiv 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ , then we see from (6.2) that

$$-\Delta w^\lambda(x) \geq \lambda_1 \left(\frac{1}{|x^\lambda|^2} - \frac{1}{|x|^2}\right) u^\lambda(x) > 0 \text{ in } \Sigma^\lambda \setminus \{0^\lambda\},$$

a contradiction. So  $w^\lambda \not\equiv 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ . Then by the maximum principle, we conclude that  $w^\lambda > 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ . Similarly,  $\sigma^\lambda > 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ .

*Step 2.* Define  $\lambda^* = \sup \{\bar{\lambda} < 0 : w^\lambda > 0, \sigma^\lambda > 0 \text{ in } \Sigma^\lambda \setminus \{0^\lambda\}, \forall \lambda < \bar{\lambda}\}$ . Then we claim that  $\lambda^* = 0$ .

Assume by contradiction that  $\lambda^* < 0$ . Clearly we have both  $w^{\lambda^*} \geq 0$  and  $\sigma^{\lambda^*} \geq 0$  in  $\Sigma^{\lambda^*} \setminus \{0^{\lambda^*}\}$ . By a similar argument as in Step 1, in fact we have both  $w^{\lambda^*} > 0$  and  $\sigma^{\lambda^*} > 0$  in  $\Sigma^{\lambda^*} \setminus \{0^{\lambda^*}\}$ . Take  $\varepsilon > 0$  such that

$$\varepsilon < \frac{1}{2} \min \left\{ \frac{1}{4} \left(1 - \frac{\lambda_1}{\Lambda_N}\right) S, \frac{1}{4} \left(1 - \frac{\lambda_2}{\Lambda_N}\right) S, \frac{1}{16} \left(1 - \frac{\lambda_1}{\Lambda_N}\right) \left(1 - \frac{\lambda_2}{\Lambda_N}\right) S^2 \right\}.$$

Then there exists a small  $\delta_1 \in (0, |\lambda^*|)$  such that for any  $\lambda \in [\lambda^*, \lambda^* + \delta_1)$ , there hold

$$(6.16) \quad \left\| (2^* - 1)u^{2^*-2} + \nu\alpha(\alpha - 1)u^{\alpha-2}v^\beta \right\|_{L^{N/2}(\Sigma^\lambda \setminus \Sigma^{\lambda^*})} \leq \varepsilon,$$

$$(6.17) \quad \left\| (2^* - 1)v^{2^*-2} + \nu\beta(\beta - 1)u^\alpha v^{\beta-2} \right\|_{L^{N/2}(\Sigma^\lambda \setminus \Sigma^{\lambda^*})} \leq \varepsilon,$$

$$(6.18) \quad (\nu\alpha\beta)^2 \left\| u^{\alpha-1}v^{\beta-1} \right\|_{L^{N/2}(\Sigma^\lambda \setminus \Sigma^{\lambda^*})}^2 \leq \varepsilon.$$

Meanwhile, since  $w^{\lambda^*} > 0$  and  $\sigma^{\lambda^*} > 0$  in  $\Sigma^{\lambda^*} \setminus \{0^{\lambda^*}\}$ , by convergence almost everywhere and thereby in the measure sense of  $(w^\lambda, v^\lambda) \rightarrow (w^{\lambda^*}, v^{\lambda^*})$  in  $\Sigma^{\lambda^*}$ , we have that

$$\begin{aligned} \lim_{\lambda \rightarrow \lambda^*} \left\| (2^* - 1)u^{2^*-2} + \nu\alpha(\alpha - 1)u^{\alpha-2}v^\beta \right\|_{L^{N/2}(\Omega_1^\lambda \cap \Sigma^{\lambda^*})} &= 0, \\ \lim_{\lambda \rightarrow \lambda^*} \left\| (2^* - 1)v^{2^*-2} + \nu\beta(\beta - 1)u^\alpha v^{\beta-2} \right\|_{L^{N/2}(\Omega_2^\lambda \cap \Sigma^{\lambda^*})} &= 0, \\ \lim_{\lambda \rightarrow \lambda^*} (\nu\alpha\beta)^2 \left\| u^{\alpha-1}v^{\beta-1} \right\|_{L^{N/2}(\Omega_1^\lambda \cap \Omega_2^\lambda \cap \Sigma^{\lambda^*})}^2 &= 0. \end{aligned}$$

Then there exists  $\delta_2 \in (0, \delta_1)$  such that for any  $\lambda \in [\lambda^*, \lambda^* + \delta_2)$ , there hold

$$\begin{aligned} & \left\| (2^* - 1)u^{2^*-2} + \nu\alpha(\alpha - 1)u^{\alpha-2}v^\beta \right\|_{L^{N/2}(\Omega_1^\lambda \cap \Sigma^{\lambda^*})} \leq \varepsilon, \\ & \left\| (2^* - 1)v^{2^*-2} + \nu\beta(\beta - 1)u^\alpha v^{\beta-2} \right\|_{L^{N/2}(\Omega_2^\lambda \cap \Sigma^{\lambda^*})} \leq \varepsilon, \\ & (\nu\alpha\beta)^2 \left\| u^{\alpha-1}v^{\beta-1} \right\|_{L^{N/2}(\Omega_1^\lambda \cap \Omega_2^\lambda \cap \Sigma^{\lambda^*})}^2 \leq \varepsilon. \end{aligned}$$

Recall that  $\Omega_i^\lambda \subset \Sigma^\lambda$ . Combining these with (6.16)-(6.18), we see that (6.8)-(6.10) hold for any  $\lambda \in [\lambda^*, \lambda^* + \delta_2)$ . Then repeating the proof of Step 1, we conclude that for any  $\lambda \in [\lambda^*, \lambda^* + \delta_2)$ ,  $w^\lambda > 0$  and  $\sigma^\lambda > 0$  in  $\Sigma^\lambda \setminus \{0^\lambda\}$ , which contradicts the definition of  $\lambda^*$ . Therefore  $\lambda^* = 0$ .

*Step 3.* We claim that both  $u$  and  $v$  are radially symmetric with respect to the origin.

With the help of Steps 1 and 2, this argument is standard. Since  $\lambda^* = 0$ , then we can carry out the above procedure in the opposite direction, namely moving the parallel planes in the negative  $x_1$  direction from positive infinity. Then they must stop at the origin again, and so we get the symmetry of both  $u$  and  $v$  with respect to 0 in the  $x_1$  direction by combining the two inequalities obtained in the two opposite directions. Since the direction can be chosen arbitrarily, we conclude that both  $u$  and  $v$  are radially symmetric with respect to the origin. This completes the proof of Theorem 1.4.  $\square$

7. UNIQUENESS RESULTS FOR THE SPECIAL CASES  $\lambda_1 = \lambda_2$  AND  $\alpha = \beta = \frac{2^*}{2}$

When  $\lambda_1 = \lambda_2$ , some uniqueness results about ground state solutions of (1.27) were obtained by the authors in [12, 13]. We remark that, by using the same ideas as in [12, 13], these results also hold for problem (1.4) if we assume  $N \geq 4$ ,  $\lambda_1 = \lambda_2$  and  $\alpha = \beta = \frac{2^*}{2}$ . First we consider the case  $N = 4$ ; then we have the following result, which improves Theorem 1.2 in case  $\lambda_1 = \lambda_2$ .

**Theorem 7.1.** *Assume that  $N = 4$ ,  $\lambda_1 = \lambda_2 \in (0, 1)$ ,  $\alpha = \beta = 2$  and  $\nu > 0$ .*

- (1) *If  $\nu \neq 1/2$ , then for any  $\mu > 0$ ,  $((1 + 2\nu)^{-1/2}z_\mu^1, (1 + 2\nu)^{-1/2}z_\mu^1)$  is a ground state solution of (1.4), with*

$$(7.1) \quad c_\nu = J_\nu \left( (1 + 2\nu)^{-1/2}z_\mu^1, (1 + 2\nu)^{-1/2}z_\mu^1 \right) = \frac{1}{2(1 + 2\nu)} S(\lambda_1)^2.$$

*Moreover, the set  $\{((1 + 2\nu)^{-1/2}z_\mu^1, (1 + 2\nu)^{-1/2}z_\mu^1) : \mu > 0\}$  contains all positive ground state solutions of (1.4).*

- (2) *If  $\nu = 1/2$ , then for any  $\mu > 0$  and  $\theta \in (0, \pi/2)$ ,  $(\sin \theta z_\mu^1, \cos \theta z_\mu^1)$  is a ground state solution of (1.4) and  $c_{1/2} = \frac{1}{4}S(\lambda_1)^2$ . Moreover, the set  $\{(\sin \theta z_\mu^1, \cos \theta z_\mu^1) : \mu > 0, \theta \in (0, \pi/2)\}$  contains all positive ground state solutions of (1.4).*

*Proof.* (1) This result can be obtained by repeating the proofs of [12, Theorem 1.1 and Theorem 1.2] with trivial modifications. We omit the details.

(2) This result can be obtained by repeating the proofs of Theorem 3.1-(2) with trivial modifications. We omit the details.  $\square$

*Remark 7.1.* As pointed out in the Introduction, by [33] we know that  $Z_i$  contains all positive solutions of (1.8). Here, for the case where  $N = 4$ ,  $\lambda_1 = \lambda_2 \in (0, 1)$ ,  $\alpha = \beta = 2$ ,  $\nu > 0$  and  $\nu \neq 1/2$ , we conjecture that the set  $\{(1 + 2\nu)^{-1/2} z_\mu^1, (1 + 2\nu)^{-1/2} z_\mu^1) : \mu > 0\}$  contains all positive solutions of (1.4).

Now we consider the case  $N \geq 5$ . Denote  $p = \frac{2^*}{2}$  for simplicity. Consider

$$(7.2) \quad \begin{cases} k^{p-1} + p\nu k^{\frac{p}{2}-1} l^{\frac{p}{2}} = 1, \\ p\nu k^{\frac{p}{2}} l^{\frac{p}{2}-1} + l^{p-1} = 1, \\ k > 0, \quad l > 0. \end{cases}$$

Let  $\nu > 0$ . By a direct computation, it was proved in [13, Lemma 2.1] that there exists  $(k_0, l_0)$ , such that

$$(7.3) \quad (k_0, l_0) \text{ satisfies (7.2) and } k_0 = \min\{k : (k, l) \text{ is a solution of (7.2)}\}.$$

Then we have the following uniqueness result.

**Theorem 7.2.** *Assume that  $N \geq 5$ ,  $\lambda_1 = \lambda_2 \in (0, \Lambda_N)$  and  $\alpha = \beta = p = \frac{2^*}{2}$ . If  $\nu \geq \frac{2}{N}$ , then for any  $\mu > 0$ ,  $(\sqrt{k_0} z_\mu^1, \sqrt{l_0} z_\mu^1)$  is a positive ground state solution of (1.4). Moreover, the set  $\{(\sqrt{k_0} z_\mu^1, \sqrt{l_0} z_\mu^1) : \mu > 0\}$  contains all positive ground state solutions of (1.4).*

*Proof.* This result can be obtained by repeating the proofs of [13, Theorem 1.1 and Theorem 1.2] with trivial modifications. We omit the details.  $\square$

#### ACKNOWLEDGEMENT

The authors wish to thank the anonymous referee very much for the careful reading and valuable comments.

#### REFERENCES

- [1] Boumediene Abdellaoui, Veronica Felli, and Ireneo Peral, *Some remarks on systems of elliptic equations doubly critical in the whole  $\mathbb{R}^N$* , Calc. Var. Partial Differential Equations **34** (2009), no. 1, 97–137, DOI 10.1007/s00526-008-0177-2. MR2448311 (2010d:35067)
- [2] Boumediene Abdellaoui, Ireneo Peral, and Veronica Felli, *Existence and multiplicity for perturbations of an equation involving a Hardy inequality and the critical Sobolev exponent in the whole of  $\mathbb{R}^N$* , Adv. Differential Equations **9** (2004), no. 5-6, 481–508. MR2099969 (2006d:35062)
- [3] N. Akhmediev and A. Ankiewicz, *Partially coherent solitons on a finite background*, Phys. Rev. Lett. **82** (1999), 2661-2664.
- [4] Antonio Ambrosetti and Eduardo Colorado, *Standing waves of some coupled nonlinear Schrödinger equations*, J. Lond. Math. Soc. (2) **75** (2007), no. 1, 67–82, DOI 10.1112/jlms/jdl020. MR2302730 (2008f:35369)
- [5] Antonio Ambrosetti and Paul H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis **14** (1973), 349–381. MR0370183 (51 #6412)
- [6] Thierry Aubin, *Problèmes isopérimétriques et espaces de Sobolev* (French), J. Differential Geometry **11** (1976), no. 4, 573–598. MR0448404 (56 #6711)
- [7] Haïm Brézis and Elliott Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc. **88** (1983), no. 3, 486–490, DOI 10.2307/2044999. MR699419 (84e:28003)
- [8] Haïm Brézis and Louis Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), no. 4, 437–477, DOI 10.1002/cpa.3160360405. MR709644 (84h:35059)

- [9] Jaeyoung Byeon and Louis Jeanjean, *Standing waves for nonlinear Schrödinger equations with a general nonlinearity*, Arch. Ration. Mech. Anal. **185** (2007), no. 2, 185–200, DOI 10.1007/s00205-006-0019-3. MR2317788 (2008g:35049)
- [10] Wenxiong Chen and Congming Li, *An integral system and the Lane-Emden conjecture*, Discrete Contin. Dyn. Syst. **24** (2009), no. 4, 1167–1184, DOI 10.3934/dcds.2009.24.1167. MR2505697 (2010d:35068)
- [11] Wenxiong Chen, Congming Li, and Biao Ou, *Classification of solutions for an integral equation*, Comm. Pure Appl. Math. **59** (2006), no. 3, 330–343, DOI 10.1002/cpa.20116. MR2200258 (2006m:45007a)
- [12] Zhijie Chen and Wenming Zou, *Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent*, Arch. Ration. Mech. Anal. **205** (2012), no. 2, 515–551, DOI 10.1007/s00205-012-0513-8. MR2947540
- [13] Zhijie Chen and Wenming Zou, *Positive least energy solutions and phase separation for coupled Schrödinger equations with critical exponent: Higher dimensional case*, Calc. Var. Partial Differential Equations, to appear, DOI 10.1007/s00526-014-0717-x.
- [14] B. Esry, C. Greene, J. Burke, and J. Bohn, *Hartree-Fock theory for double condensates*, Phys. Rev. Lett. **78** (1997), 3594–3597.
- [15] Veronica Felli and Angela Pistoia, *Existence of blowing-up solutions for a nonlinear elliptic equation with Hardy potential and critical growth*, Comm. Partial Differential Equations **31** (2006), no. 1-3, 21–56, DOI 10.1080/03605300500358145. MR2209748 (2006k:35090)
- [16] D. J. Frantzeskakis, *Dark solitons in atomic Bose-Einstein condensates: from theory to experiments*, J. Phys. A **43** (2010), no. 21, 213001, 68, DOI 10.1088/1751-8113/43/21/213001. MR2644602 (2011m:82041)
- [17] B. Gidas, Wei Ming Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), no. 3, 209–243. MR544879 (80h:35043)
- [18] B. Gidas, Wei Ming Ni, and L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbf{R}^n$* , Mathematical analysis and applications, Part A, Adv. in Math. Suppl. Stud., vol. 7, Academic Press, New York, 1981, pp. 369–402. MR634248 (84a:35083)
- [19] D. Hall, M. Matthews, J. Ensher, C. Wieman, and E. Cornell, *Dynamics of component separation in a binary mixture of Bose-Einstein condensates*, Phys. Rev. Lett. **81** (1998), 1539–1542.
- [20] Yu. S. Kivshar and B. Luther-Davies, *Dark optical solitons: physics and applications*, Physics Reports **298** (1998), 81–197.
- [21] Elliott H. Lieb and Michael Loss, *Analysis*, Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 1997. MR1415616 (98b:00004)
- [22] Tai-Chia Lin and Juncheng Wei, *Ground state of  $N$  coupled nonlinear Schrödinger equations in  $\mathbf{R}^n$ ,  $n \leq 3$* , Comm. Math. Phys. **255** (2005), no. 3, 629–653, DOI 10.1007/s00220-005-1313-x. MR2135447 (2006g:35044)
- [23] P.-L. Lions, *The concentration-compactness principle in the calculus of variations. The limit case. I*, Rev. Mat. Iberoamericana **1** (1985), no. 1, 145–201, DOI 10.4171/RMI/6. MR834360 (87c:49007)
- [24] P.-L. Lions, *The concentration-compactness principle in the calculus of variations. The limit case. II*, Rev. Mat. Iberoamericana **1** (1985), no. 2, 45–121, DOI 10.4171/RMI/12. MR850686 (87j:49012)
- [25] Zhaoli Liu and Zhi-Qiang Wang, *Multiple bound states of nonlinear Schrödinger systems*, Comm. Math. Phys. **282** (2008), no. 3, 721–731, DOI 10.1007/s00220-008-0546-x. MR2426142 (2009k:58022)
- [26] L. A. Maia, E. Montefusco, and B. Pellacci, *Positive solutions for a weakly coupled nonlinear Schrödinger system*, J. Differential Equations **229** (2006), no. 2, 743–767, DOI 10.1016/j.jde.2006.07.002. MR2263573 (2007h:35070)
- [27] Alessio Pomponio, *Coupled nonlinear Schrödinger systems with potentials*, J. Differential Equations **227** (2006), no. 1, 258–281, DOI 10.1016/j.jde.2005.09.002. MR2233961 (2007e:35263)
- [28] Boyan Sirakov, *Least energy solitary waves for a system of nonlinear Schrödinger equations in  $\mathbf{R}^n$* , Comm. Math. Phys. **271** (2007), no. 1, 199–221, DOI 10.1007/s00220-006-0179-x. MR2283958 (2007k:35477)

- [29] Didier Smets, *Nonlinear Schrödinger equations with Hardy potential and critical nonlinearities*, Trans. Amer. Math. Soc. **357** (2005), no. 7, 2909–2938 (electronic), DOI 10.1090/S0002-9947-04-03769-9. MR2139932 (2006f:35102)
- [30] Michael Struwe, *A global compactness result for elliptic boundary value problems involving limiting nonlinearities*, Math. Z. **187** (1984), no. 4, 511–517, DOI 10.1007/BF01174186. MR760051 (86k:35046)
- [31] Michael Struwe, *Variational methods*, 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 34, Springer-Verlag, Berlin, 1996. Applications to nonlinear partial differential equations and Hamiltonian systems. MR1411681 (98f:49002)
- [32] Giorgio Talenti, *Best constant in Sobolev inequality*, Ann. Mat. Pura Appl. (4) **110** (1976), 353–372. MR0463908 (57 #3846)
- [33] Susanna Terracini, *On positive entire solutions to a class of equations with a singular coefficient and critical exponent*, Adv. Differential Equations **1** (1996), no. 2, 241–264. MR1364003 (97b:35057)
- [34] Susanna Terracini and Gianmaria Verzini, *Multipulse phases in  $k$ -mixtures of Bose-Einstein condensates*, Arch. Ration. Mech. Anal. **194** (2009), no. 3, 717–741, DOI 10.1007/s00205-008-0172-y. MR2563622 (2011a:35109)
- [35] Michel Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications, 24, Birkhäuser Boston Inc., Boston, MA, 1996. MR1400007 (97h:58037)

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* [chenzhijie1987@sina.com](mailto:chenzhijie1987@sina.com)

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* [wzou@math.tsinghua.edu.cn](mailto:wzou@math.tsinghua.edu.cn)