

 Open access • Journal Article • DOI:10.1016/J.NONRWA.2021.103405

Existence and uniqueness of maximal strong solution of a 1D blood flow in a network of vessels — [Source link](#)

Debayan Maity, Jean-Pierre Raymond, Arnab Roy

Institutions: Tata Institute of Fundamental Research, University of Toulouse, Academy of Sciences of the Czech Republic

Published on: 01 Feb 2022 - Nonlinear Analysis-real World Applications (Pergamon)

Topics: Robin boundary condition, Nonlinear system, Uniqueness and Partial differential equation

Related papers:

- [Long Time Existence of Solutions to an Elastic Flow of Networks](#)
- [Uniqueness of weak solutions for a pseudo-parabolic equation modeling two phase flow in porous media](#)
- [Existence and Uniqueness for an Exponentially Derived Switching Scheme](#)
- [Existence of a solution for generalized Forchheimer flow in porous media with minimal regularity conditions](#)
- [Solutions to muscle fiber equations and their long time behaviour](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/existence-and-uniqueness-of-maximal-strong-solution-of-a-1d-4ld4tu6fct>



HAL
open science

Existence and uniqueness of maximal strong solution of a 1d blood flow in a network of vessels

Debayan Maity, Jean-Pierre Raymond, Arnab Roy

► To cite this version:

Debayan Maity, Jean-Pierre Raymond, Arnab Roy. Existence and uniqueness of maximal strong solution of a 1d blood flow in a network of vessels. 2020. hal-02912208

HAL Id: hal-02912208

<https://hal.archives-ouvertes.fr/hal-02912208>

Preprint submitted on 5 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXISTENCE AND UNIQUENESS OF MAXIMAL STRONG SOLUTION OF A 1D BLOOD FLOW IN A NETWORK OF VESSELS

DEBAYAN MAITY, JEAN-PIERRE RAYMOND, AND ARNAB ROY

ABSTRACT. We study the well-posedness of a system of one-dimensional partial differential equations modeling blood flows in a network of vessels with viscoelastic walls. We prove the existence and uniqueness of maximal strong solution for this type of hyperbolic/parabolic model. We also prove a stability estimate under suitable nonlinear Robin boundary conditions.

1. INTRODUCTION

In this paper, we consider a one dimensional blood flow model in a network of vessels with viscoelastic walls. In each vessel (of length 1 in nondimensional variables), the cross sectional area $A(x, t)$ of the vessel at the axial coordinate $x \in I = (0, 1)$ and at time $t > 0$, the flow rate $Q(x, t)$, and the average internal pressure $P(x, t)$, over a cross section, satisfy the mass conservation and momentum balance equations:

$$\begin{aligned} \frac{\partial A}{\partial t} + \frac{\partial Q}{\partial x} &= 0, \quad x \in I, t > 0, \\ \frac{\partial Q}{\partial t} + \frac{\partial}{\partial x} \left(\frac{Q^2}{A} \right) + \frac{A}{\rho} \frac{\partial P}{\partial x} &= -k_f \frac{Q}{A}, \quad x \in I, t > 0, \end{aligned} \tag{1.1}$$

where ρ is the fluid density, assumed to be constant, and k_f is the friction coefficient per unit length. To close the system we need a constitutive law connecting the pressure P to the cross-sectional area A . In the Kelvin-Voigt model, the pressure law (or vessel law) is given by:

$$P = P_{\text{ext}} + \frac{\beta}{A_0} \left(\sqrt{A} - \sqrt{A_0} \right) + \frac{\nu}{A_0} \frac{\partial}{\partial t} (\sqrt{A}), \tag{1.2}$$

where P_{ext} denotes the constant external pressure, A_0 denotes the reference cross-sectional area, ν is a viscoelastic coefficient depending on the thickness h of the vessel, the coefficient β is related to the vessel stiffness, and is defined by $\beta = \frac{\sqrt{\pi} h E}{1 - \sigma^2}$, where E is the Young's modulus and σ is the Poisson's ratio. The system (1.1)-(1.2) has to be completed by initial and boundary conditions.

For simplicity, we analyze models corresponding to the vessel law (1.2), but the results of the paper can be adapted to more general vessel laws as those considered in [17].

When we substitute the pressure law (1.2) in (1.1)₂, by taking (1.1)₁ into account, we obtain the following system

$$\begin{aligned} \frac{\partial A}{\partial t} + \frac{\partial Q}{\partial x} &= 0, \quad x \in I, t > 0, \\ \frac{\partial Q}{\partial t} + \frac{\partial}{\partial x} \left(\frac{Q^2}{A} \right) + \frac{\beta \sqrt{A}}{2A_0 \rho} \frac{\partial A}{\partial x} + \frac{\nu}{4A_0 A^{1/2} \rho} \frac{\partial A}{\partial x} \frac{\partial Q}{\partial x} - \frac{\nu \sqrt{A}}{2A_0 \rho} \frac{\partial^2 Q}{\partial x^2} &= -k_f \frac{Q}{A}, \quad x \in I, t > 0. \end{aligned} \tag{1.3}$$

Date: August 5, 2020.

2010 Mathematics Subject Classification. 35M10, 35Q35, 76D03, 76Z05, 92C35.

Key words and phrases. Blood flow model, viscoelastic vessels, one-dimensional model, arterial network, strong solutions, maximal-in-time solutions, uniqueness of solution, fluid-structure interaction.

The first author was partially supported by Department of Atomic Energy, Government of India, under project no. 12-R & D-TFR-5.01-0520 and by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 694126-DYCON). The second author is partially supported by the ANR-Project HANUMAN 18-CE45-0014-01. The third author is supported by the Czech Science Foundation (GAČR) project GA19-04243S. The Institute of Mathematics, CAS is supported by RVO:67985840.

When the viscoelastic coefficient ν is equal to zero, the system corresponding to (1.3) can be written as a quasilinear hyperbolic system (see, e.g., [6, 18, 23, 10, 26]). The diffusive effect, induced by the viscous term $-\frac{\nu\sqrt{A}}{2A_0\rho}\frac{\partial^2 Q}{\partial x^2}$ when $\nu > 0$, makes the system of hyperbolic/parabolic nature. Even if, in blood flow models, the hyperbolic nature of system (1.3) is dominant, because the viscous term $-\frac{\nu\sqrt{A}}{2A_0\rho}\frac{\partial^2 Q}{\partial x^2}$ is small compared to the other terms, this additional viscous term plays a role in numerical simulations [19], in estimation problems [7, 14], and when data coming from numerical models are compared with *in vivo* data [3, 4].

The viscoelastic behavior of vessels has been observed in several experimental studies [1, 27]. Several studies demonstrate that the incorporation of viscoelastic tube laws allows more physiological predictions than those obtained with elastic laws, because blood pressure and vessel deformation are often overestimated by 1D elastic models [22, 21, 25]. For the analysis of other viscoelastic models we refer to [20, 22, 7].

In the numerical approximations of system (1.3), the viscous term is often considered as a viscous correction in a quasilinear hyperbolic system, and therefore the viscous term is taken into account as a source term [18, 19]. Splitting methods are other numerical strategies, consisting of solving alternatively an hyperbolic system and a parabolic equation, see [17, 24].

From the well-posedness point of view, the existence of global-in-time regular solutions under some smallness conditions, or local-in-time regular solutions, for the quasilinear hyperbolic system corresponding to $\nu = 0$, is studied in [6] and in [10] in a single vessel. The coupling of a quasilinear hyperbolic system with a Windkessel type boundary condition is considered in [9]. As far as we know, similar results in the viscous case, when $\nu > 0$, are not known. Another viscoelastic model is derived in [5] for a single vessel, but not for a network.

The goal of this paper is to prove the existence and uniqueness of maximal strong solution of a system modeling a blood flow in a network of vessels, corresponding to (1.3) in each vessel. In this paper, due to its length, we do not study outflow boundary conditions of Windkessel type. But several results of the present paper may be extended to such models. This will be studied in a forthcoming paper.

Before studying a general network, for clarity, we introduce the first results for a binary vascular bifurcation as represented in Figure 1. For $i = 1, 2, 3$, (A_i, Q_i) satisfies (1.1) with the pressure law defined in (1.2). More precisely, we consider the following system:

$$\begin{aligned} &\text{For } i \in \{1, 2, 3\}, (A_i, Q_i) \text{ satisfies} \\ &\frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0, \quad t \in (0, T), x \in I, \\ &\frac{\partial Q_i}{\partial t} + \frac{\partial}{\partial x} \left(\frac{Q_i^2}{A_i} \right) + \frac{A_i}{\rho} \frac{\partial P_i}{\partial x} = -k_f \frac{Q_i}{A_i}, \quad t \in (0, T), x \in I, \\ &P_i = P_{\text{ext}} + \frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) + \frac{\nu}{A_{i,0}} \frac{\partial}{\partial t} (\sqrt{A_i}), \end{aligned} \tag{1.4}$$

where $A_{i,0}$ denotes the reference sectional area of the i -th vessel.

At the branching point, the balance of rate flows and the continuity of total pressures read as follows

$$\begin{aligned} Q_1(1, t) &= Q_2(0, t) + Q_3(0, t), \quad t \geq 0, \\ P_1(1, t) + \frac{\rho}{2} \frac{Q_1^2}{A_1^2}(1, t) &= P_2(0, t) + \frac{\rho}{2} \frac{Q_2^2}{A_2^2}(0, t) = P_3(0, t) + \frac{\rho}{2} \frac{Q_3^2}{A_3^2}(0, t), \quad t \geq 0. \end{aligned} \tag{1.5}$$

The above system is completed by the following initial conditions

$$A_i(x, 0) = A_i^0(x), \quad Q_i(x, 0) = Q_i^0(x), \quad x \in I, \tag{1.6}$$

and boundary conditions

$$Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \geq 0. \tag{1.7}$$

Using the expression of P_i in (1.4)₃, we want to eliminate all the terms involving P_i . Observe that using (1.4)₁, we can rewrite P_i as

$$P_i = P_{\text{ext}} + \frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \frac{\partial Q_i}{\partial x}.$$




FIGURE 1

By differentiating the above pressure law with respect to x , we obtain

$$\frac{\partial P_i}{\partial x} = \frac{\beta}{2A_{i,0}\sqrt{A_i}} \frac{\partial A_i}{\partial x} - \frac{\partial}{\partial x} \left(\frac{\nu}{2A_{i,0}\sqrt{A_i}} \frac{\partial Q_i}{\partial x} \right).$$

Using the above expression, we can write the system (1.4)–(1.7), in an arbitrary time interval (t_0, t_1) , in the following form:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, (A_i, Q_i) \text{ satisfies} \\ \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0, \quad t \in (t_0, t_1), \quad x \in I, \\ \frac{\rho}{A_i} \frac{\partial Q_i}{\partial t} + \frac{\rho}{A_i} \left(\frac{2Q_i}{A_i} \frac{\partial Q_i}{\partial x} - \frac{Q_i^2}{A_i^2} \frac{\partial A_i}{\partial x} \right) + \frac{\beta}{2A_{i,0}\sqrt{A_i}} \frac{\partial A_i}{\partial x} - \frac{\partial}{\partial x} \left(\frac{\nu}{2A_{i,0}\sqrt{A_i}} \frac{\partial Q_i}{\partial x} \right) \\ \quad = -k_f \rho \frac{Q_i}{A_i^2}, \quad t \in (t_0, t_1), \quad x \in I, \\ Q_1(1, t) = Q_2(0, t) + Q_3(0, t), \quad t \in (t_0, t_1), \\ Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \in (t_0, t_1), \\ \left[-\frac{\nu}{2A_{1,0}\sqrt{A_1}} \frac{\partial Q_1}{\partial x} + \frac{\beta}{A_{1,0}} \left(\sqrt{A_1} - \sqrt{A_{1,0}} \right) + \frac{1}{2} \rho \frac{Q_1^2}{A_1^2} \right] \Big|_{x=1} \\ \quad = \left[-\frac{\nu}{2A_{2,0}\sqrt{A_2}} \frac{\partial Q_2}{\partial x} + \frac{\beta}{A_{2,0}} \left(\sqrt{A_2} - \sqrt{A_{2,0}} \right) + \frac{1}{2} \rho \frac{Q_2^2}{A_2^2} \right] \Big|_{x=0} \\ \quad = - \left[\frac{\nu}{2A_{3,0}\sqrt{A_3}} \frac{\partial Q_3}{\partial x} + \frac{\beta}{A_{3,0}} \left(\sqrt{A_3} - \sqrt{A_{3,0}} \right) + \frac{1}{2} \rho \frac{Q_3^2}{A_3^2} \right] \Big|_{x=0}, \quad t \in (t_0, t_1), \\ A_i(x, t_0) = A_i^0(x), \quad Q_i(x, t_0) = Q_i^0(x), \quad x \in I. \end{array} \right. \quad (1.8)$$

We shall need to consider system (1.8) over a time interval (t_0, t_1) , and not necessarily over a fixed time interval $(0, T)$, to prove the existence of maximal strong solutions (see Section 5.1).

From now on, to simplify the notation we set

$$A = (A_i)_{i=1}^3, \quad A^0 = (A_i^0)_{i=1}^3, \quad \bar{A} = (\bar{A}_i)_{i=1}^3, \quad Q = (Q_i)_{i=1}^3, \quad Q^0 = (Q_i^0)_{i=1}^3, \quad (A, Q) = (A_i, Q_i)_{i=1}^3.$$

For $-\infty < t_0 < t_1 < \infty$, we look for solutions to system (1.8) in the space

$$\mathcal{E}(t_0, t_1) = \left\{ (A, Q) \mid A \in [H^1(t_0, t_1; H^1(I))]^3, \quad Q \in [L^2(t_0, t_1; H^2(I)) \cap H^1(t_0, t_1; L^2(I))]^3 \right\}, \quad (1.9)$$

equipped with the norm

$$\begin{aligned} \|(A, Q)\|_{\mathcal{E}(t_0, t_1)} = & \sum_{i=1}^3 \left(\|A_i\|_{H^1(t_0, t_1; H^1(0,1))} + \|A_i\|_{L^\infty(t_0, t_1; H^1(0,1))} + \|Q_i\|_{L^2(t_0, t_1; H^2(0,1))} \right. \\ & \left. + \|Q_i\|_{H^1(t_0, t_1; L^2(0,1))} + \|Q_i\|_{L^\infty(t_0, t_1; H^1(0,1))} \right). \end{aligned} \quad (1.10)$$

Moreover, for $\bar{A} \in [H^1(I)]^3$, $A \in [H^1(t_0, t_1; H^1(I))]^3$, we introduce the following quantities

$$\gamma_{\bar{A}} = \min \left\{ \bar{A}(x) \mid x \in \bar{I} \right\} > 0, \quad (1.11)$$

$$\gamma_A(t_0, t_1) = \min \left\{ A(x, t) \mid x \in \bar{I}, t \in [t_0, t_1] \right\} > 0. \quad (1.12)$$

We also introduce the spaces

$$E_{\bar{A}}(t_0, t_1) = \left\{ A \in [H^1(t_0, t_1; H^1(I))]^3 \mid A(t_0) = \bar{A} \right\}, \quad (1.13)$$

and, for $\gamma > 0$,

$$E_{\bar{A}}(t_0, t_1; \gamma) = \left\{ A \in E_{\bar{A}}(t_0, t_1) \mid \gamma_A(t_0, t_1) \geq \gamma \right\}. \quad (1.14)$$

Definition 1.1. We say that a pair (A, Q) is a strong solution to system (1.8) over the time interval $[0, T]$ when $(A, Q) \in \mathcal{E}(0, T)$, $\gamma_A(0, T) > 0$, (A, Q) satisfies (1.8)₂₋₄ in the sense of distributions in $I \times (0, T)$ and (1.8)₅₋₉ in the sense of traces.

We say that (A, Q) is a maximal strong solution to system (1.8) over the time interval $[0, T_m)$ when either $T_m = \infty$, or $T_m < \infty$ and, for all $0 < T < T_m$, (A, Q) is a strong solution to system (1.8) over the time interval $[0, T]$, and when

$$\lim_{T \rightarrow T_m} \left(\|(A, Q)\|_{\mathcal{E}(0, T)} + \max \left\{ |A_i(x, T)|^{-1} \mid 1 \leq i \leq 3, x \in [0, 1] \right\} \right) = \infty. \quad (1.15)$$

We are now in a position to state the main result of the paper in the case of the simple network represented in Figure 1.

Theorem 1.2. Let us assume that, for $i = 1, 2, 3$, $A_i^0 > 0$, $A^0 \in [H^1(I)]^3$, $Q^0 \in [H^1(I)]^3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$, and that the following compatibility conditions are satisfied

$$\begin{aligned} Q_1^0(1) &= Q_2^0(0) + Q_3^0(0), \\ Q_1^0(0) &= h_1(0), \quad Q_2^0(1) = h_2(0), \quad Q_3^0(1) = h_3(0). \end{aligned} \quad (1.16)$$

Then, the system (1.8) admits a unique maximal strong solution over $[0, T_m)$, for some $T_m > 0$. Both the solution and the maximal time of existence T_m are unique.

Remark 1.3. In the above theorem, we state the existence of a unique maximal solution for the system (1.4)–(1.6) with Dirichlet boundary conditions on the flow rate (1.7). Later on, we shall prove the existence of maximal unique solution when the Dirichlet boundary conditions are replaced by nonlinear Dirichlet boundary conditions (see Theorem 5.3) and nonlinear Robin boundary conditions (see Theorem 6.2). These nonlinear Robin boundary conditions, which approximate Dirichlet boundary conditions, allow us to prove an energy estimate satisfied by the corresponding solutions (see Proposition 6.3). In Theorem 5.3, the nonlinear Dirichlet boundary conditions are introduced to take into account boundary conditions on the velocity.

To study the system (1.8), in Section 2, we rewrite it in the form of a linear system in which the nonlinear terms are collected in source terms. The existence and regularity results for the associated nonhomogeneous linear system are obtained in Section 3. The nonlinear terms are estimated in Section 4 and Theorem 1.2 is proved in Section 5 with the Banach fixed point Theorem. In Section 6, by adapting results obtained in [10] for the quasilinear hyperbolic system corresponding to our model when $\nu = 0$, we prove an energy identity satisfied by strong solutions of system (1.4)–(1.7). This energy identity is not sufficient to obtain a stability estimate because of the nonhomogeneous Dirichlet boundary conditions in (1.7). We are able to prove that if, in (1.7), we replace the classical nonhomogeneous Dirichlet boundary conditions by nonlinear Robin boundary conditions, the associated nonlinear system admits a unique maximal strong solution, and that this solution satisfies a stability estimate (see Theorem 6.2 and Proposition 6.3). We generalize the previous results to general networks in Section 7.

2. REFORMULATION OF THE NONLINEAR SYSTEM

In order to prove the existence of solutions to system (1.8), we rewrite it in the form of a linearized system in which the nonlinear terms of system (1.8) are right hand side terms in the equations. For that, for an arbitrary $\bar{A} \in [H^1(I)]^3$ satisfying $\gamma_{\bar{A}} > 0$, we introduce the coefficients

$$L_{\bar{A}}^i = \frac{\rho}{\bar{A}_i}, \quad N_{\bar{A}}^i = \frac{\nu}{2\bar{A}_{i,0}\sqrt{\bar{A}_i}}, \quad i = 1, 2, 3, \quad (2.1)$$

and, for $i = 1, 2, 3$, we define the nonlinear terms $F_{\bar{A}}^i$ and $G_{\bar{A}}^i$ by

$$\begin{aligned} F_{\bar{A}}^i(A_i, Q_i) &= -k_f \rho \frac{Q_i}{\bar{A}_i^2} - \frac{2\rho Q_i}{\bar{A}_i^2} \frac{\partial Q_i}{\partial x} + \frac{\rho Q_i^2}{\bar{A}_i^3} \frac{\partial A_i}{\partial x} - \rho \left(\frac{1}{\bar{A}_i} - \frac{1}{\bar{A}_i} \right) \frac{\partial Q_i}{\partial t} \\ &+ \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2\bar{A}_{i,0}\sqrt{\bar{A}_i}} - \frac{\nu}{2\bar{A}_{i,0}\sqrt{\bar{A}_i}} \right) \frac{\partial Q_i}{\partial x} \right] - \frac{\beta}{2\bar{A}_{i,0}\sqrt{\bar{A}_i}} \frac{\partial A_i}{\partial x} - \frac{1}{4\bar{A}_{i,0}\bar{A}_i^{3/2}} \frac{\partial A_i}{\partial x} \frac{\partial Q_i}{\partial x}, \end{aligned} \quad (2.2)$$

and

$$\begin{aligned} G_{\bar{A}}^1(A_1, Q_1) &= \left[\frac{\beta}{\bar{A}_{1,0}} \left(\sqrt{A_1} - \sqrt{\bar{A}_1} \right) + \frac{\nu}{2\bar{A}_{1,0}} \frac{\partial Q_1}{\partial x} \left(\frac{1}{\sqrt{\bar{A}_1}} - \frac{1}{\sqrt{A_1}} \right) + \frac{1}{2} \rho \frac{Q_1^2}{\bar{A}_1^2} \right] \Big|_{x=1}, \\ G_{\bar{A}}^i(A_i, Q_i) &= \left[\frac{\beta}{\bar{A}_{i,0}} \left(\sqrt{A_i} - \sqrt{\bar{A}_i} \right) + \frac{\nu}{2\bar{A}_{i,0}} \frac{\partial Q_i}{\partial x} \left(\frac{1}{\sqrt{\bar{A}_i}} - \frac{1}{\sqrt{A_i}} \right) + \frac{1}{2} \rho \frac{Q_i^2}{\bar{A}_i^2} \right] \Big|_{x=0}, \quad i = 2, 3. \end{aligned} \quad (2.3)$$

Moreover, the constant terms are defined by

$$\begin{aligned} g_{\bar{A}}^1 &= \left[\frac{\beta}{\bar{A}_{1,0}} \left(\sqrt{\bar{A}_1} - \sqrt{A_{1,0}} \right) \right] \Big|_{x=1}, \\ g_{\bar{A}}^i &= \left[\frac{\beta}{\bar{A}_{i,0}} \left(\sqrt{\bar{A}_i} - \sqrt{A_{i,0}} \right) \right] \Big|_{x=0}, \quad i = 2, 3. \end{aligned} \quad (2.4)$$

From now on, to simplify the presentation, we are going to choose $(t_0, t_1) = (0, T)$, but all the results can be adapted to the case when $(t_0, t_1) \neq (0, T)$. With the nonlinear terms introduced above, the system (1.8) can now be rewritten as:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, (A_i, Q_i) \text{ satisfies} \\ \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0, \quad t \in (0, T), \quad x \in I, \\ L_{A^0}^i \frac{\partial Q_i}{\partial t} - \frac{\partial}{\partial x} \left(N_{A^0}^i \frac{\partial Q_i}{\partial x} \right) = F_{A^0}^i(A_i, Q_i), \quad t \in (0, T), \quad x \in I, \\ Q_1(1, t) = Q_2(0, t) + Q_3(0, t), \quad t \in (0, T), \\ Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \in (0, T), \\ -N_{A^0}^1 \frac{\partial Q_1}{\partial x}(1, t) + g_{A^0}^1 + G_{A^0}^1(A_1, Q_1) = -N_{A^0}^2 \frac{\partial Q_2}{\partial x}(0, t) + g_{A^0}^2 + G_{A^0}^2(A_2, Q_2) \\ \quad = -N_{A^0}^3 \frac{\partial Q_3}{\partial x}(0, t) + g_{A^0}^3 + G_{A^0}^3(A_3, Q_3), \quad t \in (0, T), \\ A_i(x, 0) = A_i^0(x), \quad Q_i(x, 0) = Q_i^0(x), \quad x \in I. \end{array} \right. \quad (2.5)$$

In the above system, we use the nonlinear terms and the coefficients corresponding to $\bar{A} = A^0$. But in Section 5.1, we shall need to study system (1.8) over $(0, \hat{T})$, and next over (\hat{T}, τ) . This is why it is important to express the dependence of the nonlinear terms and the coefficients on A^0 , or on $A(\hat{T})$.

3. STUDY OF A LINEAR MODEL

The proof of Theorem 1.2 relies on the Banach fixed point Theorem. The idea is to replace the nonlinear terms in (2.5) by given source terms f_i and g_i . Throughout this section, we assume that $\bar{A} \in [H^1(I)]^3$ and $\gamma_{\bar{A}} > 0$ ($\gamma_{\bar{A}}$ is defined in (1.11)). To study system (2.5), we are going to establish regularity results for the following linear system:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, (A_i, Q_i) \text{ satisfies} \\ \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0 \quad t \in (0, T), x \in I, \\ L_{\bar{A}}^i \frac{\partial Q_i}{\partial t} - \frac{\partial}{\partial x} \left(N_{\bar{A}}^i \frac{\partial Q_i}{\partial x} \right) = f_i \quad t \in (0, T), x \in I, \\ Q_1(1, t) = Q_2(0, t) + Q_3(0, t) \quad t \in (0, T), \\ Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \in (0, T), \\ -N_{\bar{A}}^1 \frac{\partial Q_1}{\partial x}(1, t) + g_1(t) = -N_{\bar{A}}^2 \frac{\partial Q_2}{\partial x}(0, t) + g_2(t) = -N_{\bar{A}}^3 \frac{\partial Q_3}{\partial x}(0, t) + g_3(t) \quad t \in (0, T), \\ A_i(x, 0) = A_i^0(x), \quad Q_i(x, 0) = Q_i^0(x) \quad x \in I. \end{array} \right. \quad (3.1)$$

Let us remark that the above linear system can be solved ‘‘in cascades’’: The equations (3.1)_{3–6} satisfied by Q_i , with the initial condition for Q_i in (3.1)₇, can be solved independently of A_i . Once we have obtained the regularity of Q_i , we can easily get the regularity of A_i from (3.1)₂. Thus, at first we concentrate in solving the following system with homogeneous boundary conditions:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, Q_i \text{ satisfies} \\ L_{\bar{A}}^i \frac{\partial Q_i}{\partial t} - \frac{\partial}{\partial x} \left(N_{\bar{A}}^i \frac{\partial Q_i}{\partial x} \right) = f_i \quad t \in (0, T), x \in I, \\ Q_1(1, t) = Q_2(0, t) + Q_3(0, t) \quad t \in (0, T), \\ Q_1(0, t) = Q_2(1, t) = Q_3(1, t) = 0 \quad t \in (0, T), \\ -N_{\bar{A}}^1 \frac{\partial Q_1}{\partial x}(1, t) = -N_{\bar{A}}^2 \frac{\partial Q_2}{\partial x}(0, t) = -N_{\bar{A}}^3 \frac{\partial Q_3}{\partial x}(0, t) \quad t \in (0, T), \\ Q_i(x, 0) = Q_i^0(x) \quad x \in I. \end{array} \right. \quad (3.2)$$

3.1. Study of system (3.2). We first study the following stationary system:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, Q_i \text{ satisfies} \\ -\frac{\partial}{\partial x} \left(N_{\bar{A}}^i \frac{\partial Q_i}{\partial x} \right) = f_i \quad \text{in } I, \\ Q_1(1) = Q_2(0) + Q_3(0), \\ Q_1(0) = Q_2(1) = Q_3(1) = 0, \\ -N_{\bar{A}}^1 \frac{dQ_1}{dx}(1) = -N_{\bar{A}}^2 \frac{dQ_2}{dx}(0) = -N_{\bar{A}}^3 \frac{dQ_3}{dx}(0). \end{array} \right. \quad (3.3)$$

We introduce the space

$$V = \left\{ (Q_1, Q_2, Q_3) \in [H^1(I)]^3 \mid Q_1(0) = 0 = Q_2(1) = Q_3(1), \quad Q_1(1) = Q_2(0) + Q_3(0) \right\},$$

and the bilinear form ‘ a ’ defined on $V \times V$ by

$$a(Q, \Phi) = \sum_{i=1}^3 \int_0^1 N_{\bar{A}}^i \frac{dQ_i}{dx} \frac{d\Phi_i}{dx}, \quad \Phi = (\Phi_1, \Phi_2, \Phi_3).$$

Lemma 3.1. *A function $Q \in (H^2(I))^3$ satisfies (3.3) if and only if it is solution of the following variational problem*

$$\text{Determine } Q \in V \quad \text{such that} \quad a(Q, \Phi) = \sum_{i=1}^3 \int_0^1 f_i \Phi_i \, dx \quad \text{for all } \Phi \in V. \quad (3.4)$$

Proof. If $Q \in (H^2(I))^3$ is solution of (3.3), then an integration by parts gives

$$a(Q, \Phi) + \sum_{i=1}^3 \left[-N_A^i \frac{dQ_i}{dx}(1) \Phi_i(1) + N_A^i \frac{dQ_i}{dx}(0) \Phi_i(0) \right] = \sum_{i=1}^3 \int_0^1 f_i \Phi_i dx.$$

Using the condition $\Phi_1(0) = 0 = \Phi_2(1) = \Phi_3(1)$ and the Neumann boundary conditions (3.3)₅ satisfied by (Q_1, Q_2, Q_3) , we have

$$\begin{aligned} & \sum_{i=1}^3 \left[-N_A^i \frac{dQ_i}{dx}(1) \Phi_i(1) + N_A^i \frac{dQ_i}{dx}(0) \Phi_i(0) \right] \\ &= -N_A^2 \frac{dQ_2}{dx}(0) \Phi_1(1) + N_A^2 \frac{dQ_2}{dx}(0) \Phi_2(0) + N_A^3 \frac{dQ_3}{dx}(0) \Phi_3(0) \\ &= \left[-N_A^2 \frac{dQ_2}{dx}(0) + N_A^3 \frac{dQ_3}{dx}(0) \right] \Phi_3(0) = 0. \end{aligned}$$

Thus, Q is a solution to the variational problem (3.4).

The converse statement can be proved in a classical way by first recovering equation (3.3)₂, and next the regularity of $(Q_i)_{i=1}^3$ and the boundary conditions (3.3)₃₋₅. The proof is complete. \blacksquare

Proposition 3.2. *Let us assume that, for $i = 1, 2, 3$, $f_i \in L^2(I)$. Then the system (3.3) admits a unique solution $Q \in (H^2(I))^3$ and*

$$\sum_{i=1}^3 \|Q_i\|_{H^2(I)} \leq C \sum_{i=1}^3 \|f_i\|_{L^2(I)}. \quad (3.5)$$

Proof. The bilinear form a is continuous and coercive on $V \times V$. The existence of a unique weak solution to the variational problem (3.4) follows from the Lax-Milgram Lemma. The end of proof is classical. \blacksquare

We introduce the unbounded operator $(\mathcal{A}, \mathcal{D}(\mathcal{A}))$ in $[L^2(I)]^3$ defined by

$$\begin{aligned} \mathcal{D}(\mathcal{A}) &= \{Q \in V \cap [H^2(I)]^3 \mid Q = (Q_1, Q_2, Q_3) \text{ satisfies (3.3)}_5\}, \\ \mathcal{A}Q &= (\mathcal{A}_1 Q_1, \mathcal{A}_2 Q_2, \mathcal{A}_3 Q_3) \quad \text{where} \\ \mathcal{A}_i Q_i &= -\frac{\partial}{\partial x} \left(N_A^i \frac{\partial Q_i}{\partial x} \right) \quad \text{for } i = 1, 2, 3. \end{aligned}$$

The following proposition is an easy consequence of Proposition 3.2 and of its proof.

Proposition 3.3. *The operator $(\mathcal{A}, \mathcal{D}(\mathcal{A}))$ is the infinitesimal generator of a strongly continuous analytic semi-group on $[L^2(I)]^3$. Moreover, $(\mathcal{A}, \mathcal{D}(\mathcal{A}))$ is a self-adjoint operator with compact resolvent in $[L^2(I)]^3$.*

To define weak solutions to system (3.2), we introduce the following spaces

$$\begin{aligned} H(0, T) &= [H^1(0, T; L^2(I))]^3, \\ W(0, T) &= \left\{ Q \in L^2(0, T; V) \mid \frac{dQ}{dt} \in L^2(0, T; V') \right\}, \end{aligned}$$

where V' is the dual of V with respect to the pivot space $(L^2(I))^3$.

Definition 3.4. *We say that $Q \in W(0, T)$ is a weak solution to (3.2) if the following conditions hold*

$$\sum_{i=1}^3 \left(\left\langle L_A^i \frac{dQ_i}{dt}(t), \xi_i \right\rangle_{V', V} + \int_0^1 N_A^i \frac{\partial Q_i}{\partial x} \frac{\partial \xi_i}{\partial x} dx \right) = \sum_{i=1}^3 \int_0^1 f_i \xi_i dx \quad \text{for all } \xi = (\xi_i)_{i=1}^3 \in V, \quad (3.6)$$

$$Q_i(\cdot, 0) = Q_i^0. \quad (3.7)$$

The main result of this subsection is the following:

Theorem 3.5. *Let $f_i \in L^2(0, T; L^2(I))$ and $Q^0 \in V$. Then there exists a unique weak solution $Q \in W(0, T)$ to (3.2). Moreover, we have*

$$Q_i \in L^2(0, T; H^2(I)) \cap H^1(0, T; L^2(I)) \cap L^\infty(0, T; H^1(I)), \quad (3.8)$$

and

$$\begin{aligned} \sum_{i=1}^3 \left[\|Q_i\|_{L^\infty(0, T; H^1(I))} + \|Q_i\|_{L^2(0, T; H^2(I))} + \|Q_i\|_{H^1(0, T; L^2(I))} \right] \\ \leq C \sum_{i=1}^3 (\|f_i\|_{L^2(0, T; L^2(I))} + \|Q_i^0\|_{H^1(I)}), \quad (3.9) \end{aligned}$$

where the constant C depends on L_i, N_i , but is independent of T .

Proof. Due to Proposition 3.3, there exists a Hilbertian basis in $(L^2(I))^3$, namely $(\xi_{1,k}, \xi_{2,k}, \xi_{3,k})_{k \in \mathbb{N}^*}$, constituted of eigenfunctions of \mathcal{A} , such that $(\xi_{1,k}, \xi_{2,k}, \xi_{3,k})$ belongs to $\mathcal{D}(\mathcal{A})$ for all $k \in \mathbb{N}^*$. Using that basis, we can follow the lines of the proof of [8, Theorem 4, Chapter 7] to prove the existence of a unique weak solution to system (3.2) in $W(0, T)$.

The estimate in (3.9) can be proved as in [8, Theorem 5, Chapter 7]. \blacksquare

3.2. Study of system (3.1). We want to study the regularity of solutions of the linear problem (3.1).

Lemma 3.6. *Let us assume that, for $i = 1, 2, 3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$, $g_i \in H_{\text{loc}}^{1/4}([0, \infty))$, $Q_i^0 \in H^1(I)$, and the compatibility conditions (1.16) are satisfied.*

Then, for $i = 1, 2, 3$, there exists a function $Q_i \in L_{\text{loc}}^2([0, \infty); H^2(I)) \cap H_{\text{loc}}^1([0, \infty); L^2(I)) \cap L_{\text{loc}}^\infty([0, \infty); H^1(I))$ such that

$$\begin{aligned} Q_1(0, t) = h_1(t), \quad Q_1(1, t) = 0, \quad \frac{\partial Q_1}{\partial x}(1, t) = g_1(t), \quad Q_1(\cdot, 0) = Q_1^0, \\ Q_2(0, t) = 0, \quad Q_2(1, t) = h_2(t), \quad \frac{\partial Q_2}{\partial x}(0, t) = g_2(t), \quad Q_2(\cdot, 0) = Q_2^0, \\ Q_3(0, t) = 0, \quad Q_3(1, t) = h_3(t), \quad \frac{\partial Q_3}{\partial x}(0, t) = g_3(t), \quad Q_3(\cdot, 0) = Q_3^0, \quad \text{for all } t \geq 0. \end{aligned} \quad (3.10)$$

Moreover, there exists a constant $C > 0$, independent of $T > 0$, such that, for $i = 1, 2, 3$, we have

$$\|Q_i\|_{L^2(0, T; H^2(I)) \cap H^1(0, T; L^2(I)) \cap L^\infty(0, T; H^1(I))} \leq C (\|g_i\|_{H^{1/4}(0, T)} + \|h_i\|_{H^{3/4}(0, T)} + \|Q_i^0\|_{H^1(I)}). \quad (3.11)$$

Proof. The proof of the above lemma follows from [15, Theorem 2.3, page 18]. \blacksquare

We prove the following theorem:

Theorem 3.7. *Let us assume that, for $i = 1, 2, 3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$ and $g_i \in H_{\text{loc}}^{1/4}([0, \infty))$, $f_i \in L^2(0, T; L^2(I))$, and the following compatibility conditions are satisfied*

$$Q_1^0(1) = Q_2^0(0) + Q_3^0(0), \quad Q_1^0(0) = h_1(0), \quad Q_2^0(1) = h_2(0), \quad Q_3^0(1) = h_3(0). \quad (3.12)$$

Then, the system (3.1) admits a unique solution (A, Q) satisfying

$$\begin{aligned} \frac{\partial A_i}{\partial t} \in L^\infty(0, T; L^2(I)), \quad A_i \in H^1(0, T; H^1(I)), \\ Q_i \in L^2(0, T; H^2(I)) \cap H^1(0, T; L^2(I)) \cap L^\infty(0, T; H^1(I)), \quad i = 1, 2, 3. \end{aligned} \quad (3.13)$$

Furthermore, there exists a constant $C_{\bar{A}}$, depending on \bar{A} but independent of T , such that

$$\begin{aligned} \sum_{i=1}^3 \left[\left\| \frac{\partial A_i}{\partial t} \right\|_{L^\infty(0,T;L^2(I))} + \|A_i\|_{H^1(0,T;H^1(I))} + \|A_i\|_{L^\infty(0,T;H^1(I))} \right. \\ \left. + \|Q_i\|_{L^\infty(0,T;H^1(I))} + \|Q_i\|_{L^2(0,T;H^2(I))} + \|Q_i\|_{H^1(0,T;L^2(I))} \right] \\ \leq C_{\bar{A}} \sum_{i=1}^3 (\|f_i\|_{L^2(0,T;L^2(I))} + \|g_i\|_{H^{1/4}(0,T)} + \|h_i\|_{H^{3/4}(0,T)} + \|A_i^0\|_{H^1(I)} + \|Q_i^0\|_{H^1(I)}), \end{aligned} \quad (3.14)$$

for all $T > 0$.

Proof. We first study the existence and uniqueness of a solution (Q_1, Q_2, Q_3) in $[L^2(0, T; H^2(I))]^3 \cap [H^1(0, T; L^2(I))]^3 \cap [L^\infty(0, T; H^1(I))]^3$ to system (3.1)₃₋₇. Let $(\tilde{Q}_1, \tilde{Q}_2, \tilde{Q}_3)$ be a solution to (3.10) satisfying (3.11), i.e.,

$$\|\tilde{Q}_i\|_{L^2(0,T;H^2(I)) \cap H^1(0,T;L^2(I)) \cap L^\infty(0,T;H^1(I))} \leq C(\|g_i\|_{H^{1/4}(0,T)} + \|h_i\|_{H^{3/4}(0,T)} + \|Q_i^0\|_{H^1(I)}). \quad (3.15)$$

We look for $Q = (Q_i)_{i=1}^3$ in the form $(Q_1, Q_2, Q_3) = (\tilde{Q}_1, \tilde{Q}_2, \tilde{Q}_3) + (\hat{Q}_1, \hat{Q}_2, \hat{Q}_3)$. Thus, $\hat{Q} = (\hat{Q}_1, \hat{Q}_2, \hat{Q}_3)$ satisfies

$$\begin{cases} L_A^i \frac{\partial \hat{Q}_i}{\partial t} - \frac{\partial}{\partial x} \left(N_A^i \frac{\partial \hat{Q}_i}{\partial x} \right) = \hat{f}_i & t \in (0, T), x \in I, i \in \{1, 2, 3\}, \\ \hat{Q}_1(1, t) = \hat{Q}_2(0, t) + \hat{Q}_3(0, t) & t \in (0, T), \\ \hat{Q}_1(0, t) = \hat{Q}_2(1, t) = \hat{Q}_3(1, t) = 0 & t \in (0, T), \\ -N_A^1 \frac{\partial \hat{Q}_1}{\partial x}(1, t) = -N_A^2 \frac{\partial \hat{Q}_2}{\partial x}(0, t) = -N_A^3 \frac{\partial \hat{Q}_3}{\partial x}(0, t) & t \in (0, T), \\ \hat{Q}_i(x, 0) = 0 & x \in I, i \in \{1, 2, 3\}, \end{cases} \quad (3.16)$$

where

$$\hat{f}_i = f_i - L_A^i \frac{\partial \tilde{Q}_i}{\partial t} + \frac{\partial}{\partial x} \left(N_A^i \frac{\partial \tilde{Q}_i}{\partial x} \right).$$

With Theorem 3.5, we have

$$\sum_{i=1}^3 \left[\|\hat{Q}_i\|_{L^\infty(0,T;H^1(I))} + \|\hat{Q}_i\|_{L^2(0,T;H^2(I))} + \|\hat{Q}_i\|_{H^1(0,T;L^2(I))} \right] \leq C \sum_{i=1}^3 \|\hat{f}_i\|_{L^2(0,T;L^2(I))},$$

The above estimate together with (3.15) give

$$\begin{aligned} \sum_{i=1}^3 \left[\|Q_i\|_{L^\infty(0,T;H^1(I))} + \|Q_i\|_{L^2(0,T;H^2(I))} + \|Q_i\|_{H^1(0,T;L^2(I))} \right] \\ \leq C \sum_{i=1}^3 (\|f_i\|_{L^2(0,T;L^2(I))} + \|g_i\|_{H^{1/4}(0,T)} + \|h_i\|_{H^{3/4}(0,T)} + \|Q_i^0\|_{H^1(I)}). \end{aligned}$$

The estimate for $(A_i)_{i=1}^3$ is obtained with the help of the first equation of system (3.1). ■

4. ESTIMATES OF NONLINEAR TERMS

Throughout this section, for $0 < T \leq \infty$, we set

$$I_T = I \times (0, T), \quad H^{2,1}(I_T) = L^2(0, T; H^2(I)) \cap H^1(0, T; L^2(I)).$$

4.1. Preliminary results. Let us recall some important lemmas which will be used later on.

Lemma 4.1. *Let $1/2 < s \leq 1$. There is a bounded extension operator from $\{f \in H^s(0, T) \mid f(0) = 0\}$ to $H^s(0, \infty)$, uniformly bounded with respect to $T > 0$.*

Lemma 4.2. *(i) Let us assume that $g \in H^{s_2}(0, T)$ with $g(0) = 0$ and $s_2 > s_1$, $s_1, s_2 \in (1/2, 1]$. Then there exists a constant $C > 0$ independent of T such that*

$$\|g\|_{H^{s_1}(0, T)} \leq C(\sqrt{T})^{1-s_1/s_2} \|g\|_{H^{s_2}(0, T)}. \quad (4.1)$$

(ii) Let us assume that $f \in H^{s_1}(0, T)$, $g \in H^{s_2}(0, T)$ with $s_1, s_2 \in (1/2, 1]$ and $f(0) = g(0) = 0$. Then there exists a constant $C > 0$, independent of T such that $fg \in H^s(0, T)$ for $s \in (1/2, 1)$ with $s_1 + s_2 - s > 1/2$ and

$$\|fg\|_{H^s(0, T)} \leq CT^\delta \|f\|_{H^{s_1}(0, T)} \|g\|_{H^{s_2}(0, T)}, \quad (4.2)$$

for some $\delta > 0$.

(iii) Let us assume that $f \in H^{s_1}(0, T)$, $g \in H^{s_2}(0, T)$ with $s_1, s_2 \in (1/2, 1]$ and $g(0) = 0$. Then there exists a constant $C > 0$, independent of T such that $fg \in H^s(0, T; U_3)$ for $s \in (1/2, 1)$ with $s_1 + s_2 - s > 1/2$ and

$$\|fg\|_{H^s(0, T)} \leq CT^\delta (\|f\|_{H^{s_1}(0, T)} + |f(0)| + \|g\|_{H^{s_2}(0, T)}), \quad (4.3)$$

for some $\delta > 0$.

Proof. *Step 1.* Let us first prove (4.1). Let \bar{g} be the extension of g to $(0, \infty)$ introduced in Lemma 4.1. Since $g(0) = 0$, we have from Lemma 4.1 that

$$\|g\|_{L^2(0, T)} \leq \sqrt{T} \|g\|_{L^\infty(0, T)} \leq \sqrt{T} \|\bar{g}\|_{L^\infty(0, \infty)} \leq \sqrt{T} \|\bar{g}\|_{H^{s_2}(0, \infty)} \leq C\sqrt{T} \|g\|_{H^{s_2}(0, T)}.$$

Let us now fix $1/2 < s_1 < 1$. By interpolation and the above inequality, we have

$$\begin{aligned} \|g\|_{H^{s_1}(0, T)} &\leq \|\bar{g}\|_{H^{s_1}(0, \infty)} \leq \|\bar{g}\|_{L^2(0, \infty)}^{1-s_1/s_2} \|\bar{g}\|_{H^{s_2}(0, \infty)}^{s_1/s_2} \leq C \|g\|_{L^2(0, T)}^{1-s_1/s_2} \|g\|_{H^{s_2}(0, T)}^{s_1/s_2} \\ &\leq C(\sqrt{T})^{1-s_1/s_2} \|g\|_{H^{s_2}(0, T)}, \end{aligned}$$

with C independent of T since $g(0) = 0$.

Step 2. We want to prove (4.2). We have \bar{f}, \bar{g} as extensions of f, g to $(0, \infty)$ respectively following Lemma 4.1. As $s_1 + s_2 - s > 1/2$, there exists $\varepsilon > 0$ such that $s_1 + s_2 - s - \varepsilon > 1/2$.

With [12, Proposition B1] and the estimate (4.1), we have

$$\begin{aligned} \|fg\|_{H^s(0, T)} &\leq \|\bar{f}\bar{g}\|_{H^s(0, \infty)} \leq C \|\bar{f}\|_{H^{s_1}(0, \infty)} \|\bar{g}\|_{H^{s_2-\varepsilon}(0, \infty)} \\ &\leq C \|f\|_{H^{s_1}(0, T)} \|g\|_{H^{s_2-\varepsilon}(0, T)} \leq T^\delta \|f\|_{H^{s_1}(0, T)} \|g\|_{H^{s_2}(0, T)}, \end{aligned}$$

where the constant C is independent of T as $f(0) = g(0) = 0$.

Step 3. To prove (4.3), we first write fg as

$$fg = (f - f(0))g + f(0)g.$$

Now we can use (4.2) to prove (4.3). ■

Lemma 4.3. *There exists a constant $C > 0$, independent of $T > 0$ and $0 < \gamma \leq 1$, such that, for all $f \in H^1(0, T; H^1(I))$ and $f(x, t) \geq \gamma$ in $I \times (0, T)$, the functions \sqrt{f} , $f^{1/4}$ and $1/f$ belong to $H^1(0, T; H^1(I))$ and the following estimates hold:*

$$\begin{aligned} \|\sqrt{f}\|_{H^1(0, T; H^1(I))} + \|\sqrt{f}\|_{L^\infty(0, T; H^1(I))} &\leq \frac{C}{\gamma^{3/2}} \left(\|f\|_{H^1(0, T; H^1(I))} + \|f\|_{L^\infty(0, T; H^1(I))} \right. \\ &\quad \left. + \|f\|_{H^1(0, T; H^1(I))} \|f\|_{L^\infty(0, T; H^1(I))} \right), \quad (4.4) \end{aligned}$$

$$\begin{aligned} \|f^{1/4}\|_{H^1(0, T; H^1(I))} + \|f^{1/4}\|_{L^\infty(0, T; H^1(I))} &\leq \frac{C}{\gamma^{7/4}} \left(\|f\|_{H^1(0, T; H^1(I))} + \|f\|_{L^\infty(0, T; H^1(I))} \right. \\ &\quad \left. + \|f\|_{H^1(0, T; H^1(I))} \|f\|_{L^\infty(0, T; H^1(I))} \right), \quad (4.5) \end{aligned}$$

and

$$\|1/f\|_{H^1(0,T;H^1(I))} + \|1/f\|_{L^\infty(0,T;H^1(I))} \leq \frac{C}{\gamma^3} \left(\|f\|_{H^1(0,T;H^1(I))} + \|f\|_{L^\infty(0,T;H^1(I))} + \|f\|_{H^1(0,T;H^1(I))} \|f\|_{L^\infty(0,T;H^1(I))} \right). \quad (4.6)$$

Proof. The proof is easy and left to the reader. \blacksquare

Lemma 4.4. *There exists a constant $C > 0$ such that, for all $T > 0$ and all f belonging to $H^{2,1}(I_T)$, the following estimates hold*

$$\left\| \frac{\partial f}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial f}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} \leq C (\|f(\cdot, 0)\|_{H^1(I)} + \|f\|_{H^{2,1}(I_T)}), \quad (4.7)$$

$$\|f(0, \cdot)\|_{H^{3/4}(0,T)} + \|f(1, \cdot)\|_{H^{3/4}(0,T)} \leq C (\|f(\cdot, 0)\|_{H^1(I)} + \|f\|_{H^{2,1}(I_T)}). \quad (4.8)$$

Proof. If $f \in H^{2,1}(I_T)$, then the fact that $\frac{\partial f}{\partial x}(0, \cdot)$ and $\frac{\partial f}{\partial x}(1, \cdot)$ belong to $H^{1/4}(0, T)$ can be deduced from [13, Chapter II, Lemma 3.4] or [11, Theorem 6.1], and the following estimate holds

$$\left\| \frac{\partial f}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial f}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} \leq C_T (\|f\|_{L^2(0,T;H^2(I)) \cap H^1(0,T;L^2(I))},$$

with a constant C_T which may depend on T . To obtain a constant independent of T , we proceed as follows. With [13, Chapter II, Lemma 3.4], we know that $f(\cdot, 0)$ belongs to $H^1(I)$. For some large $\lambda > 0$, we consider the following problem

$$\frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} + \lambda u = 0 \text{ in } I_\infty, \quad \frac{\partial u}{\partial x}(0, t) = \frac{\partial u}{\partial x}(1, t) = 0 \text{ for all } t \geq 0, \quad u(x, 0) = f(x, 0) \text{ for all } x \in I.$$

Due to [2, Part II, Chapter 1, Theorem 3.1], u belongs to $H^{2,1}(I_\infty)$ and

$$\|u\|_{H^{2,1}(I_T)} \leq \|u\|_{H^{2,1}(I_\infty)} \leq C \|f(\cdot, 0)\|_{H^1(I)}, \quad (4.9)$$

with $I_T = I \times (0, T)$. Let us set $\tilde{u} = u - f$. Then $\tilde{u} \in H^{2,1}(I_T)$ and $\tilde{u}(x, 0) = 0$ for all $x \in I$. Then, due to Lemma 4.1, there exists $u^* \in H^{2,1}(I_\infty)$ such that $u^* = \tilde{u}$ in $[0, T]$ and

$$\|u^*\|_{H^{2,1}(I_\infty)} \leq C \|\tilde{u}\|_{H^{2,1}(I_T)}.$$

Moreover, using [11, Theorem 6.1], we obtain

$$\begin{aligned} & \left\| \frac{\partial \tilde{u}}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial \tilde{u}}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} = \left\| \frac{\partial u^*}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial u^*}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} \\ & \leq \left\| \frac{\partial u^*}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,\infty)} + \left\| \frac{\partial u^*}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,\infty)} \leq C \|u^*\|_{H^{2,1}(I_\infty)} \leq C \|\tilde{u}\|_{H^{2,1}(I_T)}, \end{aligned}$$

where the constant C is independent of T . Finally, the above estimate and (4.9) yield

$$\begin{aligned} & \left\| \frac{\partial f}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial f}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} = \left\| \frac{\partial \tilde{u}}{\partial x}(0, \cdot) \right\|_{H^{1/4}(0,T)} + \left\| \frac{\partial \tilde{u}}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0,T)} \\ & \leq C \|\tilde{u}\|_{H^{2,1}(I_T)} \leq C (\|u\|_{H^{2,1}(I_T)} + \|f\|_{H^{2,1}(I_T)}) \leq C (\|f(\cdot, 0)\|_{H^1(I)} + \|f\|_{H^{2,1}(I_T)}). \end{aligned}$$

This completes the proof of estimate (4.7). Estimate (4.8) can be obtained in a similar manner. \blacksquare

4.2. **Analysis of nonlinear terms.** Let

$$\bar{A} \in [H^1(I)]^3 \quad \text{such that} \quad \gamma_{\bar{A}} > 0,$$

where $\gamma_{\bar{A}}$ is defined in (1.11). For $0 \leq t_0 < t_1$ and $\gamma > 0$, we introduce the space

$$B_{\bar{A}}(t_0, t_1; \gamma) = \left\{ (A, Q) \in \mathcal{E}(t_0, t_1) \mid A \in E_{\bar{A}}(t_0, t_1; \gamma) \right\},$$

where $E_{\bar{A}}(t_0, t_1; \gamma)$ is defined in (1.14). The ball $B_{\bar{A}}(t_0, t_1; \gamma, \mu)$ in $B_{\bar{A}}(t_0, t_1; \gamma)$ is defined as follows

$$B_{\bar{A}}(t_0, t_1; \gamma, \mu) = \left\{ (A, Q) \in B_{\bar{A}}(t_0, t_1; \gamma) \mid \|(A, Q)\|_{\mathcal{E}(t_0, t_1)} \leq \mu \right\}. \quad (4.10)$$

Our aim is to obtain different estimates for the nonlinear terms $F_{\bar{A}}^i, G_{\bar{A}}^i$ introduced in (2.2) - (2.4), in $I \times (t_0, t_1)$, for an arbitrary time interval (t_0, t_1) . For simplicity, we only treat the case when $(t_0, t_1) = (0, T)$, $\gamma \in (0, 1)$, and $\mu \geq 1$, but the results can be stated with obvious modifications for an arbitrary time interval (t_0, t_1) .

Now we prove the following proposition which will be used later on.

Proposition 4.5. *There exists a constant $C > 0$, independent of $\gamma \in (0, 1)$, of $\mu \geq 1$, and of $T \in (0, 1)$, such that, for all $(A, Q) \in B_{\bar{A}}(0, T; \gamma, \mu)$, we have*

$$\|A_i - \bar{A}_i\|_{L^\infty(0, T; H^1(I))} + \|A_i\|_{L^2(0, T; H^1(I))} \leq C\mu\sqrt{T} \quad (4.11)$$

$$\left\| \sqrt{A_i} - \sqrt{\bar{A}_i} \right\|_{L^\infty(0, T; H^1(I))} + \left\| \sqrt{A_i} \right\|_{L^2(0, T; H^1(I))} \leq C\mu \frac{\sqrt{T}}{\sqrt{\gamma}}, \quad (4.12)$$

$$\|Q_i\|_{L^2(0, T; H^{1+s}(I))} \leq C\mu T^{(1-s)/4}, \quad \text{for } s \in (0, 1), \quad (4.13)$$

$$\|Q_i\|_{L^\infty(I_T)} + \|A_i\|_{L^\infty(I_T)} + \left\| \sqrt{A_i} \right\|_{L^\infty(I_T)} \leq C\mu, \quad (4.14)$$

$$\left\| \sqrt{A_i} - \sqrt{\bar{A}_i} \right\|_{H^1(0, T; H^1(I))} \leq C\mu \frac{\sqrt{T}}{\sqrt{\gamma}}, \quad (4.15)$$

$$\|(A_i)^2 - (\bar{A}_i)^2\|_{H^1(0, T; H^1(I))} \leq C\mu^2. \quad (4.16)$$

Proof. We have

$$A_i(x, t) - \bar{A}_i = \int_0^t \partial_t A_i(x, s) \, ds \quad \text{and} \quad \sqrt{A_i}(x, t) - \sqrt{\bar{A}_i} = \int_0^t \partial_t \sqrt{A_i}(x, s) \, ds.$$

Therefore

$$\|A_i - \bar{A}_i\|_{L^\infty(0, T; H^1(0, 1))} \leq \sqrt{T} \|\partial_t A_i\|_{L^2(0, T; H^1(0, 1))} \leq \mu\sqrt{T}.$$

In a similar manner, we can easily obtain

$$\left\| \sqrt{A_i} - \sqrt{\bar{A}_i} \right\|_{L^\infty(0, T; H^1(0, 1))} \leq C\mu \frac{\sqrt{T}}{\sqrt{\gamma}}.$$

Using (4.12), we first obtain

$$\|A_i\|_{L^\infty(0, T; H^1(I))} \leq \|\bar{A}_i\|_{H^1(I)} + \|A_i - \bar{A}_i\|_{L^\infty(0, T; H^1(I))} \leq \mu + C\mu\sqrt{T},$$

where the constant C is independent of T . Thus

$$\|A_i\|_{L^2(0, T; H^1(0, 1))} \leq \sqrt{T} \|A_i\|_{L^\infty(0, T; H^1(I))} \leq C\mu(T + \sqrt{T}).$$

Similarly, we can show that $\left\| \sqrt{A_i} \right\|_{L^2(0, T; H^1(0, 1))} \leq \mu + C\mu \frac{T}{\sqrt{\gamma}}$. The estimate (4.13) can be proved by following the arguments of [16, Proposition 6.4]. ■

Estimate of $F_{\bar{A}}^i$.

Proposition 4.6. *For all $\mu \geq 1$ and $\gamma \in (0, 1)$, there exists a positive constant $C_F(\mu, \gamma)$, depending on μ and γ , but independent of $T \in (0, 1)$, such that, for all $(A, Q) \in B_{\bar{A}}(0, T; \gamma, \mu)$, all $(A^1, Q^1) \in B_{\bar{A}}(0, T; \gamma, \mu)$, and all $(A^2, Q^2) \in B_{\bar{A}}(0, T; \gamma, \mu)$, we have*

$$\left\| F_{\bar{A}}^i(A_i, Q_i) \right\|_{L^2(I_T)} \leq C_F(\mu, \gamma) T^\alpha, \quad (4.17)$$

$$\left\| F_{\bar{A}}^i(A_i^1, Q_i^1) - F_{\bar{A}}^i(A_i^2, Q_i^2) \right\|_{L^2(I_T)} \leq C_F(\mu, \gamma) T^\alpha \left\| (A_i^1, Q_i^1) - (A_i^2, Q_i^2) \right\|_{\mathcal{E}_T}, \quad (4.18)$$

for some $\alpha > 0$, independent of $T \in (0, 1)$, of $\mu \geq 1$, and of $\gamma \in (0, 1)$.

Proof. *Step 1. Proof of (4.17).* Let us recall that

$$\begin{aligned} F_{\bar{A}}^i(A_i, Q_i) &= -k_f \rho \frac{Q_i}{(A_i)^2} - \frac{2\rho Q_i}{(A_i)^2} \frac{\partial Q_i}{\partial x} + \frac{\rho(Q_i)^2}{(A_i)^3} \frac{\partial A_i}{\partial x} - \rho \left(\frac{1}{A_i} - \frac{1}{\bar{A}_i} \right) \frac{\partial Q_i}{\partial t} \\ &\quad + \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i}} - \frac{\nu}{2A_{i,0}\sqrt{\bar{A}_i}} \right) \frac{\partial Q_i}{\partial x} \right] - \frac{\beta}{2A_{i,0}\sqrt{A_i}} \frac{\partial A_i}{\partial x} - \frac{1}{4A_{i,0}A_i^{3/2}} \frac{\partial A_i}{\partial x} \frac{\partial Q_i}{\partial x}, \end{aligned}$$

- To estimate the first term of $F_{\bar{A}}^i$, we write

$$\left\| -k_f \rho \frac{Q_i}{(A_i)^2} \right\|_{L^2(I_T)} \leq \frac{k_f \rho}{\gamma^2} \|Q_i\|_{L^2(I_T)} \leq C \frac{\sqrt{T}}{\gamma^2} \|Q_i\|_{L^\infty(0,T;L^2(I))} \leq C \mu \frac{\sqrt{T}}{\gamma^2}.$$

- The estimate of the second term of $F_{\bar{A}}^i$ is obtained as follows

$$\left\| \frac{2\rho Q_i}{(A_i)^2} \frac{\partial Q_i}{\partial x} \right\|_{L^2(I_T)} \leq C \frac{\mu}{\gamma^2} \left\| \frac{\partial Q_i}{\partial x} \right\|_{L^2(I_T)} \leq C \mu \frac{\sqrt{T}}{\gamma^2} \left\| \frac{\partial Q_i}{\partial x} \right\|_{L^\infty(0,T;L^2(I))} \leq C \mu^2 \frac{\sqrt{T}}{\gamma^2}.$$

- To estimate the third term of $F_{\bar{A}}^i$ (the estimate of sixth term is similar), using (4.11) and (4.14), we have

$$\left\| \frac{\rho(Q_i)^2}{(A_i)^3} \frac{\partial A_i}{\partial x} \right\|_{L^2(I_T)} \leq C \frac{\mu^2}{\gamma^3} \left\| \frac{\partial A_i}{\partial x} \right\|_{L^2(I_T)} \leq C \mu^3 \frac{\sqrt{T}}{\gamma^3}.$$

- To estimate the fourth term of $F_{\bar{A}}^i$, with (4.11), we have

$$\left\| \rho \left(\frac{1}{A_i} - \frac{1}{\bar{A}_i} \right) \frac{\partial Q_i}{\partial t} \right\|_{L^2(I_T)} \leq C \frac{1}{\gamma^2} \|A_i - \bar{A}_i\|_{L^\infty(0,T;H^1(I))} \left\| \frac{\partial Q_i}{\partial t} \right\|_{L^2(I_T)} \leq C \mu^2 \frac{\sqrt{T}}{\gamma^2}.$$

- To estimate the fifth term of $F_{\bar{A}}^i$, with (4.12), we obtain

$$\begin{aligned} &\left\| \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i}} - \frac{\nu}{2A_{i,0}\sqrt{\bar{A}_i}} \right) \frac{\partial Q_i}{\partial x} \right] \right\|_{L^2(I_T)} \\ &\leq C \frac{1}{\gamma^2} \|\sqrt{A_i} - \sqrt{\bar{A}_i}\|_{L^\infty(0,T;H^1(I))} \left\| \frac{\partial Q_i}{\partial x} \right\|_{L^2(0,T;H^1(I))} \leq C \mu^2 \frac{\sqrt{T}}{\gamma^{5/2}}. \end{aligned}$$

- To estimate the last term of $F_{\bar{A}}^i$, using (4.13), we get

$$\left\| \frac{1}{4A_{i,0}A_i^{3/2}} \frac{\partial A_i}{\partial x} \frac{\partial Q_i}{\partial x} \right\|_{L^2(I_T)} \leq C \frac{1}{\gamma^{5/2}} \left\| \frac{\partial A_i}{\partial x} \right\|_{L^\infty(0,T;L^2(I))} \left\| \frac{\partial Q_i}{\partial x} \right\|_{L^2(0,T;H^s(I))} \leq C \mu^2 T^{(1-s)/4} \frac{1}{\gamma^{5/2}},$$

for all $\frac{1}{2} < s < 1$.

Step 2. Proof of (4.18). The Lipschitz estimate can be obtained as in Step 1. For clarity let us explain how we can prove it for the fifth term of $F_{\bar{A}}^i$:

$$\frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i}} - \frac{\nu}{2A_{i,0}\sqrt{\bar{A}_i}} \right) \frac{\partial Q_i}{\partial x} \right].$$

We have

$$\begin{aligned} & \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^1}} - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \right) \frac{\partial Q_i^1}{\partial x} \right] - \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^2}} - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \right) \frac{\partial Q_i^2}{\partial x} \right] \\ &= \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^1}} - \frac{\nu}{2A_{i,0}\sqrt{A_i^2}} \right) \frac{\partial Q_i^1}{\partial x} \right] + \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^2}} - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \right) \left(\frac{\partial Q_i^1}{\partial x} - \frac{\partial Q_i^2}{\partial x} \right) \right]. \end{aligned}$$

The first term can be estimated as in Step 1:

$$\begin{aligned} & \left\| \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^1}} - \frac{\nu}{2A_{i,0}\sqrt{A_i^2}} \right) \frac{\partial Q_i^1}{\partial x} \right] \right\|_{L^2(I_T)} \\ & \leq C \left\| \frac{1}{\sqrt{A_i^1}} - \frac{1}{\sqrt{A_i^2}} \right\|_{L^\infty(0,T;H^1(I))} \left\| \frac{\partial Q_i^1}{\partial x} \right\|_{L^2(0,T;H^1(I))} \\ & \leq C \frac{1}{\gamma^2} \left\| \sqrt{A_i^1} - \sqrt{A_i^2} \right\|_{L^\infty(0,T;H^1(I))} \left\| \frac{\partial Q_i^1}{\partial x} \right\|_{L^2(0,T;H^1(I))} \leq C\mu \frac{\sqrt{T}}{\gamma^2} \left\| (A_i^1, Q_i^1) - (A_i^2, Q_i^2) \right\|_{\mathcal{E}_T}. \end{aligned}$$

For the second term, as $(A^2, Q^2) \in B_{\bar{A}}(0, T; \gamma, \mu)$, we have $(\sqrt{A_i^2} - \sqrt{A_i})|_{t=0} = 0$. Thus, we can write

$$\begin{aligned} & \left\| \frac{\partial}{\partial x} \left[\left(\frac{\nu}{2A_{i,0}\sqrt{A_i^2}} - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \right) \left(\frac{\partial Q_i^1}{\partial x} - \frac{\partial Q_i^2}{\partial x} \right) \right] \right\|_{L^2(I_T)} \\ & \leq C \frac{1}{\gamma^2} \left\| \sqrt{A_i^2} - \sqrt{A_i} \right\|_{L^\infty(0,T;H^1(I))} \left\| \frac{\partial Q_i^1}{\partial x} - \frac{\partial Q_i^2}{\partial x} \right\|_{L^2(0,T;H^1(I))} \leq C\mu \frac{\sqrt{T}}{\gamma^{5/2}} \left\| (A_i^1, Q_i^1) - (A_i^2, Q_i^2) \right\|_{\mathcal{E}_T}. \end{aligned}$$

■

Estimate of $G_{\bar{A}}^i$.

Proposition 4.7. *For all $\mu \geq 1$ and $\gamma \in (0, 1)$, there exists a positive constant $C_G(\mu, \gamma)$, depending on μ and γ , but independent of $T \in (0, 1)$, such that, for all $(A, Q) \in B_{\bar{A}}(0, T; \gamma, \mu)$, all $(A^1, Q^1) \in B_{\bar{A}}(0, T; \gamma, \mu)$, and all $(A^2, Q^2) \in B_{\bar{A}}(0, T; \gamma, \mu)$, we have*

$$\left\| G_{\bar{A}}^i(A_i, Q_i) \right\|_{H^{1/4}(0,T)} \leq C_G(\mu, \gamma) T^\alpha, \quad (4.19)$$

$$\left\| G_{\bar{A}}^i(A_i^1, Q_i^1) - G_{\bar{A}}^i(A_i^2, Q_i^2) \right\|_{H^{1/4}(0,T)} \leq C_G(\mu, \gamma) T^\alpha \left\| (A_i^1, Q_i^1) - (A_i^2, Q_i^2) \right\|_{\mathcal{E}_T}, \quad (4.20)$$

for some $\alpha > 0$, independent of $T \in (0, 1)$, of $\mu \geq 1$, and of $\gamma \in (0, 1)$.

Proof. *Step 1. Proof of (4.19).* We only give estimate for $G_{\bar{A}}^1$. Let us recall

$$\begin{aligned} G_{\bar{A}}^1(A_1, Q_1) &= \frac{\beta}{A_{1,0}} \left(\sqrt{A_1}(1, t) - \sqrt{A_1}(1) \right) + \frac{\nu}{2A_{1,0}} \frac{\partial Q_1}{\partial x}(1, t) \left(\frac{1}{\sqrt{A_1}(1)} - \frac{1}{\sqrt{A_1}(1, t)} \right) \\ & \quad + \frac{1}{2} \rho \frac{(Q_1)^2(1, t)}{(A_1)^2(1, t)}. \end{aligned}$$

• Estimate of first term of $G_{\bar{A}}^1$. We apply Lemma 4.2 (i) with $s_1 = 1/4$, $s_2 = 1$ and $g = \sqrt{A_1}(1, \cdot) - \sqrt{A_1}(1)$ and relation (4.15), and we obtain

$$\begin{aligned} \left\| \sqrt{A_1}(1, \cdot) - \sqrt{A_1}(1) \right\|_{H^{1/4}(0,T)} & \leq CT^{3/8} \left\| \sqrt{A_1}(1, \cdot) - \sqrt{A_1}(1) \right\|_{H^1(0,T)} \\ & \leq CT^{3/8} \left\| \sqrt{A_1} - \sqrt{A_1} \right\|_{H^1(0,T;H^1(I))} \leq C\mu T^{7/8}. \end{aligned}$$

- Estimate of the second term of G_A^1 . We apply Lemma 4.2 (iii) with $s = 1/4$, $s_1 = 1/4$, $s_2 = 1$, $f = \frac{\partial Q_1}{\partial x}(1, \cdot)$, $g = \sqrt{A_1}(1, \cdot) - \sqrt{\bar{A}_1}(1)$ and Lemma 4.4, and we have

$$\begin{aligned} \left\| \frac{\partial Q_1}{\partial x}(1, \cdot) \left(\frac{1}{\sqrt{A_1}(1)} - \frac{1}{\sqrt{A_1}(1, \cdot)} \right) \right\|_{H^{1/4}(0, T)} &\leq C \left\| \frac{\partial Q_1}{\partial x}(1, \cdot) \left(\sqrt{A_1}(1) - \sqrt{A_1}(1, \cdot) \right) \right\|_{H^{1/4}(0, T)} \\ &\leq CT^\alpha \left(\left\| \frac{\partial Q_1}{\partial x}(1, 0) \right\| + \left\| \frac{\partial Q_1}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0, T)} \right) \left\| \sqrt{A_1}(1) - \sqrt{A_1}(1, \cdot) \right\|_{H^1(0, T)} \leq C\mu^2 T^\alpha. \end{aligned}$$

- Estimate of the last term in G_A^1 . We rewrite the last term of G_A^1 as follows

$$\frac{1}{2} \rho \frac{(Q_1)^2}{(A_1)^2}(1, t) = \frac{\rho}{2} \left((Q_1)^2 \left(\frac{1}{(A_1)^2} - \frac{1}{(\bar{A}_1)^2} \right) (1, t) + \frac{((Q_1)^2(1, t) - |Q_1^0|^2(1))}{(\bar{A}_1)^2(1)} + \frac{|Q_1^0|^2(1)}{(\bar{A}_1)^2(1)} \right). \quad (4.21)$$

First of all, using [12, Proposition B1], for any $s_0 \in (1/8, 1/4)$, we have

$$\|(Q_1)^2(1, \cdot)\|_{H^{1/4}(0, T)} \leq C \|Q_1(1, \cdot)\|_{H^{1/4+s_0}(0, T)}^2 \leq C \|Q_1(1, \cdot)\|_{H^{3/4}(0, T)}^2,$$

where the constant C can be chosen independent of T . Indeed, as $1/4 + s_0 < 1/2$, we can extend the function $Q_1(1, t)$ by zero to (T, ∞) and the estimate follows easily. Using Lemma 4.4 in the above estimate, we obtain

$$\|(Q_1)^2(1, \cdot)\|_{H^{1/4}(0, T)} \leq C\mu^2,$$

with the constant C is independent of T . Therefore, using Lemma 4.2 and relation (4.16) of Proposition 4.5, we have

$$\begin{aligned} \left\| (Q_1)^2 \left(\frac{1}{(A_1)^2} - \frac{1}{(\bar{A}_1)^2} \right) (1, \cdot) \right\|_{H^{1/4}(0, T)} &\leq C \|(Q_1)^2(1, \cdot)\|_{H^{1/4}(0, T)} \left\| \frac{1}{(A_1)^2(1, \cdot)} - \frac{1}{(\bar{A}_1)^2(1)} \right\|_{H^1(0, T)} \\ &\leq CT^\alpha \mu^2 \|(A_i)^2 - (\bar{A}_1)^2\|_{H^1(0, T; H^1(I))} \leq CT^\alpha \mu^4. \end{aligned}$$

To estimate second term of (4.21), we note that, by using Lemma 4.2, for any $s_0 \in (1/8, 1/4)$, we have

$$\begin{aligned} &\left\| \frac{(Q_1)^2(1, \cdot) - |Q_1^0|^2(1)}{(\bar{A}_1)^2(1)} \right\|_{H^{1/4}(0, T)} \\ &\leq C \|Q_1(1, \cdot) + Q_1^0(1)\|_{H^{1/4+s_0}(0, T)} \|Q_1(1, t) - Q_1^0(1)\|_{H^{1/4+s_0}(0, T)} \\ &\leq C\mu T^\alpha, \end{aligned}$$

for some C independent of T and $\alpha > 0$.

The last term of (4.21) can be estimated as follows

$$\left\| \frac{|Q_1^0|^2(1)}{(\bar{A}_1)^2(1)} \right\|_{H^{1/4}(0, T)} \leq \frac{|Q_1^0(1)|^2}{\gamma^2} \sqrt{T} \leq C\sqrt{T}.$$

Step 2. The Lipschitz estimate (4.20) can be obtained as in Step 1. ■

In Section 6, we are going to consider nonlinear Robin boundary conditions of the form

$$\begin{aligned} -\frac{\nu}{2A_{1,0}\sqrt{\bar{A}_1}} \frac{\partial Q_1}{\partial x}(0, t) + Q_1(0, t) &= H_1(Q_1, A_1) - h_1(t), \\ \text{and for } i = 1, 2, & \\ \frac{\nu}{2A_{i,0}\sqrt{\bar{A}_1}} \frac{\partial Q_i}{\partial x}(0, t) + Q_i(0, t) &= H_i(Q_i, A_i) - h_i(t), \end{aligned} \quad (4.22)$$

with

$$H_{\bar{A}}^1(Q_1, A_1) = \left[-\frac{\beta}{A_{1,0}} \left(\sqrt{A_1} - \sqrt{\bar{A}_1} \right) + \frac{\nu}{2A_{1,0}} \frac{\partial Q_1}{\partial x} \left(\frac{1}{\sqrt{A_1}} - \frac{1}{\sqrt{\bar{A}_1}} \right) - \frac{1}{2} \rho \frac{Q_1^2}{A_1^2} \right] \Big|_{x=1},$$

and for $i = 1, 2$,

$$H_{\bar{A}}^i(Q_i, A_i) = \left[\frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{\bar{A}_i} \right) - \frac{\nu}{2A_{i,0}} \frac{\partial Q_i}{\partial x} \left(\frac{1}{\sqrt{A_i}} - \frac{1}{\sqrt{\bar{A}_i}} \right) + \frac{1}{2} \rho \frac{Q_i^2}{A_i^2} \right] \Big|_{x=0}. \quad (4.23)$$

Estimate of $H_{\bar{A}}^i$. To study (1.8) in which the Dirichlet boundary condition (1.8)₅ are replaced by the nonlinear Robin boundary conditions stated in (4.22), we have to estimate the nonlinear terms defined in (4.23).

Proposition 4.8. *For all $\mu \geq 1$ and $\gamma \in (0, 1)$, there exists a positive constant $C_H(\mu, \gamma)$, depending on μ and γ , but independent of $T \in (0, 1)$, such that, for all $(A, Q) \in B_{\bar{A}}(0, T; \gamma, \mu)$, all $(A^1, Q^1) \in B_{\bar{A}}(0, T; \gamma, \mu)$, and all $(A^2, Q^2) \in B_{\bar{A}}(0, T; \gamma, \mu)$, we have*

$$\left\| H_{\bar{A}}^i(A_i, Q_i) \right\|_{H^{1/4}(0, T)} \leq C_H(\mu, \gamma) T^\alpha, \quad (4.24)$$

$$\left\| H_{\bar{A}}^i(A_i^1, Q_i^1) - H_{\bar{A}}^i(A_i^2, Q_i^2) \right\|_{H^{1/4}(0, T)} \leq C_H(\mu, \gamma) T^\alpha \left\| (A_i^1, Q_i^1) - (A_i^2, Q_i^2) \right\|_{\mathcal{E}_T}, \quad (4.25)$$

for some $\alpha > 0$, independent of $T \in (0, 1)$, of $\mu \geq 1$, and of $\gamma \in (0, 1)$.

Proof. *Step 1. Proof of (4.24).* We only give estimate for $H_{\bar{A}}^1$. Let us recall that

$$H_{\bar{A}}^1(Q_1, A_1) = \left[-\frac{\beta}{A_{1,0}} \left(\sqrt{A_1} - \sqrt{\bar{A}_1} \right) + \frac{\nu}{2A_{1,0}} \frac{\partial Q_1}{\partial x} \left(\frac{1}{\sqrt{A_1}} - \frac{1}{\sqrt{\bar{A}_1}} \right) - \frac{1}{2} \rho \frac{Q_1^2}{A_1^2} \right] \Big|_{x=1}.$$

Using the relation (4.12) of Proposition 4.5, we obtain

$$\left\| \sqrt{A_1}(1, \cdot) - \sqrt{\bar{A}_1}(1) \right\|_{H^{1/4}(0, T)} \leq \left\| \sqrt{A_1}(1, \cdot) - \sqrt{\bar{A}_1}(1) \right\|_{H^1(0, T)} \leq \frac{C\mu}{\sqrt{\gamma}} T^{1/2}$$

The term $\left[\frac{\partial Q_1}{\partial x} \left(\frac{1}{\sqrt{A_1}} - \frac{1}{\sqrt{\bar{A}_1}} \right) \right] \Big|_{x=1}$ can be estimated as follows

$$\begin{aligned} & \left\| \frac{\partial Q_1}{\partial x}(1, \cdot) \left(\frac{1}{\sqrt{A_1}(1, \cdot)} - \frac{1}{\sqrt{\bar{A}_1}(1)} \right) \right\|_{H^{1/4}(0, T)} \\ & \leq \left\| \frac{\partial Q_1}{\partial x}(1, \cdot) \right\|_{H^{1/4}(0, T)} \left\| \frac{1}{\sqrt{A_1}(1, \cdot)} - \frac{1}{\sqrt{\bar{A}_1}(1)} \right\|_{H^1(0, T)} \\ & \leq C \frac{\mu}{\gamma^{3/2}} \sqrt{T}. \end{aligned}$$

The estimate of the other term can be obtained similarly.

Step 2. The Lipschitz estimate (4.25) can be obtained as in Step 1. ■

5. EXISTENCE AND UNIQUENESS OF MAXIMAL SOLUTION

5.1. Proof of Theorem 1.2. Proof. *Step 1. Existence of a local-in-time strong solution.* We choose $M > 0$ and $\mu > 0$ such that

$$\sum_{i=1}^3 (\|A_i^0\|_{H^1(I)} + \|Q_i^0\|_{H^1(I)} + |g_{A^0}^i| + \|h_i\|_{H^{3/4}(0, 1)}) \leq M \quad \text{and} \quad \mu = 2C_{A^0} M, \quad (5.1)$$

where C_{A^0} is the continuity constant appearing in (3.14) corresponds to $\bar{A} = A^0$. Let us set

$$\gamma = \frac{1}{2} \gamma_{A^0} \quad \text{with} \quad \gamma_{A^0} = \min \left\{ A_1^0(x), A_2^0(x), A_3^0(x) \mid x \in I \right\} > 0.$$

Let $(\psi_i, \Phi_i)_{i=1}^3$ belong to $B_{A^0}(0, T; \gamma, \mu)$ (see (4.10)). We consider the following system:

$$\begin{aligned}
& \text{For } i \in \{1, 2, 3\}, (A_i, Q_i) \text{ satisfies} \\
& \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0, \quad t \in (0, T), \quad x \in I, \\
& L_{A^0}^i \frac{\partial Q_i}{\partial t} - \frac{\partial}{\partial x} \left(N_{A^0}^i \frac{\partial Q_i}{\partial x} \right) = F_{A^0}^i(\psi_i, \Phi_i), \quad t \in (0, T), \quad x \in I, \\
& Q_1(1, t) = Q_2(0, t) + Q_3(0, t), \quad t \in (0, T), \\
& Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \in (0, T), \\
& -N_{A^0}^1 \frac{\partial Q_1}{\partial x}(1, t) + g_{A^0}^1 + G_{A^0}^1(\psi_1, \Phi_1) = -N_{A^0}^2 \frac{\partial Q_2}{\partial x}(0, t) + g_{A^0}^2 + G_{A^0}^2(\psi_2, \Phi_2) \\
& \quad = -N_{A^0}^3 \frac{\partial Q_3}{\partial x}(0, t) + g_{A^0}^3 + G_{A^0}^3(\psi_3, \Phi_3), \quad t \in (0, T), \\
& A_i(x, 0) = A_i^0(x), \quad Q_i(x, 0) = Q_i^0(x), \quad x \in I.
\end{aligned} \tag{5.2}$$

We are going to show that, there exists $0 < T \leq 1$ such that the mapping

$$\mathcal{N} : (\psi_i, \Phi_i)_{i=1}^3 \mapsto (A_i, Q_i)_{i=1}^3,$$

where $(A_i, Q_i)_{i=1}^3$ is the solution to system (5.2), is a strict contraction in $B_{A^0}(0, T; \gamma, \mu)$.

Applying Theorem 3.7 to system (5.2), we obtain

$$\begin{aligned}
\| (A_i, Q_i)_{i=1}^3 \|_{\mathcal{E}(0, T)} \leq C_{A^0} \sum_{i=1}^3 \left(\| F_{A^0}^i(\psi_i, \Phi_i) \|_{L^2(0, T; L^2(I))} + \| G_{A^0}^i(\psi_i, \Phi_i) \|_{H^{1/4}(0, T)} + |g_{A^0}^i| \right. \\
\left. + \| h_i \|_{H^{3/4}(0, T)} + \| A_i^0 \|_{H^1(I)} + \| Q_i^0 \|_{H^1(I)} \right). \tag{5.3}
\end{aligned}$$

Since $(\psi_i, \Phi_i) \in B_{A^0}(0, T; \gamma, \mu)$, applying Proposition 4.6, Proposition 4.7 and relation (5.1), estimate (5.3) becomes

$$\| (A_i, Q_i)_{i=1}^3 \|_{\mathcal{E}(0, T)} \leq C_{A^0} M + C_{A^0} (C_F + C_G) T^\alpha \quad \text{for all } 0 < T \leq 1.$$

Therefore, with the choice of μ in relation (5.1), there exists $T > 0$ small enough such that

$$\| \mathcal{N}((\psi_i, \Phi_i)_{i=1}^3) \|_{\mathcal{E}(0, T)} \leq \mu.$$

Using the continuous embedding $L^\infty(0, T; H^1(I)) \hookrightarrow L^\infty(I_T)$, from (4.11) we obtain

$$\| A_i - A_i^0 \|_{L^\infty(I_T)} \leq C\sqrt{T},$$

with C independent of T since $A_i(0) = A_i^0$. By choosing T small enough, we get $A_i(x, t) \geq \gamma$ for all $(x, t) \in I_T$ because $\gamma = \gamma_{A^0}/2$. Therefore, \mathcal{N} maps $B_{A^0}(0, T; \gamma, \mu)$ into itself.

Now we will show that \mathcal{N} is a contraction. Let $(\psi_i^1, \Phi_i^1)_{i=1}^3$ and $(\psi_i^2, \Phi_i^2)_{i=1}^3$ belong to $B_{A^0}(0, T; \gamma, \mu)$. Using Theorem 3.7, Propositions 4.6 and 4.7, we obtain

$$\begin{aligned}
& \| \mathcal{N}((\psi_i^1, \Phi_i^1)_{i=1}^3) - \mathcal{N}((\psi_i^2, \Phi_i^2)_{i=1}^3) \|_{\mathcal{E}(0, T)} \\
& \leq C_{A^0} (C_F + C_G) T^\alpha \| (\psi_i^1, \Phi_i^1)_{i=1}^3 - (\psi_i^2, \Phi_i^2)_{i=1}^3 \|_{\mathcal{E}(0, T)}.
\end{aligned}$$

Thus \mathcal{N} is a contraction in $B_{A^0}(0, T; \gamma, \mu)$ for T small enough. The proof of the existence of $T \in (0, 1]$, for which the system (1.8) admits at least one strong solution over $[0, T]$, is complete.

Step 2. Existence of a maximal solution. Let us prove that any local-in-time strong solution may be extended to a maximal strong solution. Let (A, Q) be a local-in-time solution to (1.8) over $[0, T_1]$. We want to show that (A, Q) can be extended as a maximal strong solution over $[0, T_m)$, with $T_m > T_1$. We look for $(\widehat{A}, \widehat{Q}, T)$, with $T \geq T_1$ and $(\widehat{A}, \widehat{Q}) \in \mathcal{E}(0, T)$, such that

$$\begin{aligned}
& (\widehat{A}, \widehat{Q})(t) = (A, Q)(t) \quad \text{for all } t \in [0, T_1], \\
& (\widehat{A}, \widehat{Q}) \text{ is solution to (1.8) over } [0, T].
\end{aligned} \tag{5.4}$$

The set of triplets (\hat{A}, \hat{Q}, T) satisfying (5.4) is nonempty since (A, Q, T_1) satisfies (5.4). For a given triplet (\hat{A}, \hat{Q}, T) satisfying (5.4), we set

$$\hat{T} = \sup\{T \geq T_1 \mid (\hat{A}, \hat{Q}, T) \text{ satisfies (5.4)}\}.$$

If $\hat{T} = \infty$, then the proof is complete.

Let us assume that $\hat{T} < \infty$, and that (\hat{A}, \hat{Q}, T) satisfies (5.4) for all $T < \hat{T} < \infty$. We have to show that (1.15) holds for $(A, Q) = (\hat{A}, \hat{Q})$ and $T_m = \hat{T}$. We argue by contradiction. We assume that

$$\lim_{T \rightarrow \hat{T}} \left(\left\| (\hat{A}, \hat{Q}) \right\|_{\mathcal{E}(0, T)} + \max \left\{ |\hat{A}_i(x, T)|^{-1} \mid 1 \leq i \leq 3, x \in I \right\} \right) < \infty. \quad (5.5)$$

Let us set

$$\left\| (\hat{A}, \hat{Q}) \right\|_{\mathcal{E}(0, \hat{T})} = \mu_{\hat{T}} \text{ and } \sup \left\{ |\hat{A}_i(x, T)| \mid 1 \leq i \leq 3, x \in I, T \in [0, \hat{T}) \right\} = \gamma_{\hat{T}} > 0.$$

We are going to show that the solution (\hat{A}, \hat{Q}) can be extended to $[\hat{T}, \hat{T} + \varepsilon]$, for some $\varepsilon > 0$, so that (\hat{A}, \hat{Q}) is solution to system (1.8) over the time interval $[0, \hat{T} + \varepsilon]$.

We set $\bar{A} = A(x, \hat{T})$, $x \in I$. We consider the system over the time interval (\hat{T}, τ) with $\tau > \hat{T}$:

$$\begin{aligned} & \text{For } i \in \{1, 2, 3\}, (\hat{A}_i, \hat{Q}_i) \text{ satisfies} \\ & \frac{\partial \hat{A}_i}{\partial t} + \frac{\partial \hat{Q}_i}{\partial x} = 0, \quad t \in (\hat{T}, \tau), x \in I, \\ & L_A^i \frac{\partial \hat{Q}_i}{\partial t} - \frac{\partial}{\partial x} \left(N_A^i \frac{\partial \hat{Q}_i}{\partial x} \right) = F_A^i(\hat{A}_i, \hat{Q}_i), \quad t \in (\hat{T}, \tau), x \in I, \\ & \hat{Q}_1(1, t) = \hat{Q}_2(0, t) + \hat{Q}_3(0, t), \quad t \in (\hat{T}, \tau), \\ & \hat{Q}_1(0, t) = h_1(t), \quad \hat{Q}_2(1, t) = h_2(t), \quad \hat{Q}_3(1, t) = h_3(t), \quad t \in (\hat{T}, \tau), \\ & -N_A^1 \frac{\partial \hat{Q}_1}{\partial x}(1, t) + g_A^1 + G_A^1(\hat{A}_1, \hat{Q}_1) = -N_A^2 \frac{\partial \hat{Q}_2}{\partial x}(0, t) + g_A^2 + G_A^2(\hat{A}_2, \hat{Q}_2) \\ & \quad = -N_A^3 \frac{\partial \hat{Q}_3}{\partial x}(0, t) + g_A^3 + G_A^3(\hat{A}_3, \hat{Q}_3), \quad t \in (\hat{T}, \tau), \\ & \hat{A}_i(x, \hat{T}) = A_i(x, \hat{T}), \quad \hat{Q}_i(x, \hat{T}) = Q_i(x, \hat{T}), \quad x \in I, \end{aligned} \quad (5.6)$$

where the coefficients L_A^i , N_A^i and the nonlinear terms F_A^i , G_A^i , g_A^i are defined by (2.1)-(2.3). We consider the system

$$\begin{aligned} & \frac{\partial \hat{A}_i}{\partial t} + \frac{\partial \hat{Q}_i}{\partial x} = 0, \quad t \in (\hat{T}, \tau), x \in I, i \in \{1, 2, 3\}, \\ & L_A^i \frac{\partial \hat{Q}_i}{\partial t} - \frac{\partial}{\partial x} \left(N_A^i \frac{\partial \hat{Q}_i}{\partial x} \right) = F_A^i(\psi_i, \Phi_i), \quad t \in (\hat{T}, \tau), x \in I, i \in \{1, 2, 3\}, \\ & \hat{Q}_1(1, t) = \hat{Q}_2(0, t) + \hat{Q}_3(0, t), \quad t \in (\hat{T}, \tau), \\ & \hat{Q}_1(0, t) = h_1(t), \quad \hat{Q}_2(1, t) = h_2(t), \quad \hat{Q}_3(1, t) = h_3(t), \quad t \in (\hat{T}, \tau), \\ & -N_A^1 \frac{\partial \hat{Q}_1}{\partial x}(1, t) + g_A^1 + G_A^1(\psi_1, \Phi_1) = -N_A^2 \frac{\partial \hat{Q}_2}{\partial x}(0, t) + g_A^2 + G_A^2(\psi_2, \Phi_2) \\ & \quad = -N_A^3 \frac{\partial \hat{Q}_3}{\partial x}(0, t) + g_A^3 + G_A^3(\psi_3, \Phi_3), \quad t \in (\hat{T}, \tau), \\ & \hat{A}_i(x, \hat{T}) = A_i(x, \hat{T}), \quad \hat{Q}_i(x, \hat{T}) = Q_i(x, \hat{T}), \quad x \in I, i \in \{1, 2, 3\}. \end{aligned} \quad (5.7)$$

We set

$$\tilde{\gamma} = \gamma_{\hat{T}}/2 \text{ and } \tilde{\mu} = \hat{C}_{\bar{A}} \mu_{\hat{T}} + \mu_{\hat{T}}.$$

For $\tau > \widehat{T}$, we define the mapping $\widetilde{\mathcal{N}}_\tau$ in the ball $B_{\overline{A}}(\widehat{T}, \tau; \widetilde{\gamma}, \widetilde{\mu})$ by

$$\widetilde{\mathcal{N}}_\tau : (\psi_i, \Phi_i)_{i=1}^3 \mapsto (\widehat{A}_i, \widehat{Q}_i)_{i=1}^3,$$

where $(\widehat{A}_i, \widehat{Q}_i)_{i=1}^3$ is the solution to system (5.7) over $[\widehat{T}, \tau]$, and $B_{\overline{A}}(\widehat{T}, \tau; \widetilde{\gamma}, \widetilde{\mu})$ is defined in (4.10).

Applying Theorem 3.7 to system (5.7), as in Step 1, we can show that $\widetilde{\mathcal{N}}_\tau$ is a contraction in $B_{\overline{A}}(\widehat{T}, \tau; \widetilde{\gamma}, \widetilde{\mu})$ for $\tau - \widehat{T} > 0$ small enough. Thus the system (5.6) admits a solution over the time interval $[\widehat{T}, \tau]$ and the system (1.8) admits a solution over the time interval $[0, \tau]$. We have a contradiction with the definition of \widehat{T} . Thus (5.5) is false and (1.15) is proved. Hence, we have proved that any local-in-time strong solution may be extended to a maximal strong solution.

Step 3. Uniqueness of maximal solution. Let us prove that system (1.8) admits a unique maximal solution. Let (A, Q) be a maximal solution to system (1.8) over $[0, T_m)$, and let $(\widetilde{A}, \widetilde{Q})$ be another maximal solution to system (1.8) over $[0, \widetilde{T}_m)$. Let us assume that $T_m \leq \widetilde{T}_m$. Let us set

$$\widehat{T}_m = \sup \left\{ t \in [0, T_m) \mid (A, Q)(\tau) = (\widetilde{A}, \widetilde{Q})(\tau) \text{ for all } \tau \in [0, t] \right\}.$$

If $\widehat{T}_m = T_m = \widetilde{T}_m$, then the two maximal solutions are identical and the proof is complete.

If $\widehat{T}_m = T_m < \widetilde{T}_m$, then

$$\begin{aligned} & \lim_{T \rightarrow \widehat{T}_m} \left(\|(A, Q)\|_{\mathcal{E}(0, T)} + \max \left\{ |A_i(x, T)|^{-1} \mid 1 \leq i \leq 3, x \in [0, 1] \right\} \right) \\ &= \left(\|(\widetilde{A}, \widetilde{Q})\|_{\mathcal{E}(0, \widetilde{T}_m)} + \max \left\{ |\widetilde{A}_i(x, \widetilde{T}_m)|^{-1} \mid 1 \leq i \leq 3, x \in [0, 1] \right\} \right) < \infty, \end{aligned}$$

which is in contradiction with the fact that (A, Q) is a maximal solution to system (1.8) over $[0, T_m)$. Thus the proof is complete in that case too.

Let us examine the last case $\widehat{T}_m < T_m$. We have to treat separately the cases when $\widehat{T}_m > 0$ and the case when $\widehat{T}_m = 0$. The case when $\widehat{T}_m = 0$ can be treated with the same arguments as in Step 1. Let us treat the case when $\widehat{T}_m > 0$. We set

$$\begin{aligned} \mu &= \|(A, Q)\|_{\mathcal{E}(0, \widehat{T}_m)}, \quad \gamma = \min \{ A_i(x, t) \mid 1 \leq i \leq 3, x \in I, t \in [0, \widehat{T}_m] \}, \\ \widetilde{\mu} &= \|(\widetilde{A}, \widetilde{Q})\|_{\mathcal{E}(0, \widehat{T}_m)}, \quad \widetilde{\gamma} = \min \{ \widetilde{A}_i(x, t) \mid 1 \leq i \leq 3, x \in I, t \in [0, \widehat{T}_m] \}, \\ \widehat{\mu} &= 2\mu + 2\widetilde{\mu}, \quad \text{and} \quad \widehat{\gamma} = \min(\gamma/2, \widetilde{\gamma}/2). \end{aligned}$$

Now we set $\overline{A} = A(\widehat{T}_m)$. For $\widehat{T}_m < \tau < T_m$, we notice that the function $(\widehat{A}, \widehat{Q}) = (A, Q) - (\widetilde{A}, \widetilde{Q})$ satisfies the linear system (3.1) over $[\widehat{T}_m, \tau]$, with

$$\begin{aligned} f_i &= F_{\overline{A}}^i(A_i, Q_i) - F_{\overline{A}}^i(\widetilde{A}_i, \widetilde{Q}_i), \\ g_i &= G_{\overline{A}}^i(A_i, Q_i) - G_{\overline{A}}^i(\widetilde{A}_i, \widetilde{Q}_i), \\ h_i &= 0, \quad \widehat{A}_i(x, \widehat{T}_m) = 0, \quad \widehat{Q}_i(x, \widehat{T}_m) = 0. \end{aligned}$$

Using Theorem 3.7, Proposition 4.6 and Proposition 4.7, we obtain

$$\|(\widehat{A}, \widehat{Q})\|_{\mathcal{E}(\widehat{T}_m, \tau)} \leq C_{\overline{A}}(C_F + C_G)(\tau - \widehat{T}_m)^\alpha \|(\widehat{A}, \widehat{Q})\|_{\mathcal{E}(\widehat{T}_m, \tau)}.$$

We choose $\widehat{T}_m < \tau < T_m$ such that $\tau - \widehat{T}_m$ is small enough to have $C_{\overline{A}}(C_F + C_G)(\tau - \widehat{T}_m)^\alpha \leq 1/2$. From the above estimate we deduce that $\|(\widehat{A}, \widehat{Q})\|_{\mathcal{E}(\widehat{T}_m, \tau)} = 0$. Thus $(A, Q)(t) = (\widetilde{A}, \widetilde{Q})(t)$ for all $t \in [0, \tau]$. We have a contradiction with the definition of \widehat{T}_m , and the proof is complete. \blacksquare

5.2. Maximal solution for the (A, u) -system. We end this section by reformulating the system (1.4) – (1.7) in terms of the independent variables $A(x, t)$, the cross sectional area, and $u(x, t) := \frac{Q(x, t)}{A(x, t)}$, the average fluid velocity. Moreover, we prove an existence and uniqueness result for the system written in variables A and u .

The velocity $u_i(x, t)$ on the i -th vessel is

$$u_i(x, t) = \frac{Q_i(x, t)}{A_i(x, t)}, \quad x \in I, \quad t \in (0, T). \quad (5.8)$$

Then the system (1.4) – (1.7) can be written as:

$$\left\{ \begin{array}{l} \text{For } i \in \{1, 2, 3\}, (A_i, u_i) \text{ satisfies} \\ \frac{\partial A_i}{\partial t} + \frac{\partial}{\partial x}(A_i u_i) = 0, \quad t \in (0, T), \quad x \in I, \\ \frac{\partial}{\partial t}(A_i u_i) + \frac{\partial}{\partial x}(A_i u_i^2) + \frac{A_i}{\rho} \frac{\partial P_i}{\partial x} + k_f u_i = 0, \quad t \in (0, T), \quad x \in I, \\ P_i = P_{\text{ext}} + \frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) + \frac{\nu}{A_{i,0}} \frac{\partial}{\partial t}(\sqrt{A_i}), \\ A_i(x, 0) = A_i^0(x), \quad u_i(x, 0) = u_i^0(x), \quad x \in I, \end{array} \right. \quad (5.9)$$

with the following nonlinear boundary conditions at the branching points

$$\left\{ \begin{array}{l} A_1 u_1(1, t) = A_2 u_2(0, t) + A_3 u_3(0, t), \quad t \in (0, T), \\ P_1(1, t) + \frac{\rho}{2} u_1^2(1, t) = P_2(0, t) + \frac{\rho}{2} u_2^2(0, t) = P_3(0, t) + \frac{\rho}{2} u_3^2(0, t), \quad t \geq 0, \end{array} \right. \quad (5.10)$$

and with the following Dirichlet boundary conditions

$$u_1(0, t) = h_1(t), \quad u_2(1, t) = h_2(t), \quad u_3(1, t) = h_3(t), \quad t \in (0, T). \quad (5.11)$$

We look for solution to the system (5.9)–(5.11) in the space of functions $(A_i, u_i)_{i=1}^3$ satisfying

$$\left\{ \begin{array}{l} (A_i)_{i=1}^3 \in H^1(0, T; H^1(I))^3, \quad (u_i)_{i=1}^3 \in L^2(0, T; H^1(I))^3 \cap L^\infty(0, T; H^1(I))^3, \\ (A_i u_i)_{i=1}^3 \in L^2(0, T; H^2(I))^3 \cap H^1(0, T; L^2(I))^3, \end{array} \right. \quad (5.12)$$

equipped with the distance

$$\begin{aligned} d((A^1, u^1), (A^2, u^2)) &= \sum_{i=1}^3 \left(\|A_i^1 - A_i^2\|_{H^1(0, T; H^1(I)) \cap L^\infty(0, T; H^1(I))} \right. \\ &\quad \left. + \|u_i^1 - u_i^2\|_{L^2(0, T; H^1(I)) \cap L^\infty(0, T; H^1(I))} + \|A_i^1 u_i^1 - A_i^2 u_i^2\|_{L^2(0, T; H^2(I)) \cap H^1(0, T; L^2(I)) \cap L^\infty(0, T; H^1(I))} \right), \end{aligned} \quad (5.13)$$

where $A^j = (A_1^j, A_2^j, A_3^j)$ and $u^j = (u_1^j, u_2^j, u_3^j)$, $j = 1, 2$. As before, let us set

$$A = (A_i)_{i=1}^3, \quad u = (u_i)_{i=1}^3.$$

Let us recall that the constant $\gamma_A(0, T)$ is defined in (1.12).

Definition 5.1. *We say that a pair (A, u) is a strong solution to system (5.9) – (5.11) over the time interval $[0, T]$ when (A, u) satisfies the regularity assumptions in (5.12), $\gamma_A(0, T) > 0$, and when (A, u) satisfies (5.9) in the sense of distributions in $I \times (0, T)$ and (5.10)–(5.11) in the sense of traces.*

We say that (A, Q) is a maximal strong solution to system (5.9)–(5.10) over the time interval $[0, T_m)$ when either $T_m = \infty$, or $T_m < \infty$ and, for all $0 < T < T_m$, (A, u) is a strong solution to system (5.9)–(5.10) over the time interval $[0, T]$, and when

$$\lim_{T \rightarrow T_m} \left(d((A, u), (0, 0)) + \max \left\{ |A_i(x, T)|^{-1} \mid 1 \leq i \leq 3, \quad x \in [0, 1] \right\} \right) = \infty. \quad (5.14)$$

With the above definition it is quite easy to have an equivalence for the existence of strong solutions between the (A, Q) system ((1.4)–(1.7)) and the (A, u) system ((5.9) - (5.11)). More precisely, we have the following result:

Proposition 5.2. *Let (A, Q) be a maximal solution to the system (1.4)–(1.7) over $[0, T_m)$ with boundary conditions on the velocity*

$$u_1(0, t) = \frac{Q_1}{A_1}(0, t) = h_1(t), \quad u_2(1, t) = \frac{Q_2}{A_2}(1, t) = h_2(t), \quad u_3(1, t) = \frac{Q_3}{A_3}(1, t) = h_3(t), \quad (5.15)$$

(respectively with boundary conditions on the flow rate

$$Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t),) \quad (5.16)$$

if and only if, (A, u) is a maximal strong solution to the system (5.9)–(5.10), with Dirichlet boundary conditions on the velocity

$$u_1(0, t) = h_1(t), \quad u_2(1, t) = h_2(t), \quad u_3(1, t) = h_3(t) \quad (5.17)$$

(respectively with boundary conditions on the flow rate

$$u_1(0, t) = \frac{h_1(t)}{A_1(0, t)}, \quad u_2(1, t) = \frac{h_2(t)}{A_2(1, t)}, \quad u_3(1, t) = \frac{h_3(t)}{A_3(1, t)}). \quad (5.18)$$

Let us remark that, in Theorem 1.2, we have proved the existence of a unique maximal solution for the (A, Q) -system (1.4)–(1.6) with the classical Dirichlet boundary conditions (5.16). Similarly, we can show the existence of a unique maximal solution for the (A, Q) -system (1.4)–(1.6) with the nonlinear Dirichlet boundary conditions (5.15). More precisely, we have the following result.

Theorem 5.3. *Let us assume that, for $i = 1, 2, 3$, $A_i^0 > 0$, $A^0 \in [H^1(I)]^3$, $Q^0 \in [H^1(I)]^3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$ satisfying the compatibility conditions*

$$\begin{aligned} Q_1^0(1) &= Q_2^0(0) + Q_3^0(0), \\ Q_1^0(0) &= A_1^0(0)h_1(0), \quad Q_2^0(1) = A_2^0(1)h_2(0), \quad Q_3^0(1) = A_3^0(1)h_3(0). \end{aligned} \quad (5.19)$$

Then, the system (1.4)–(1.6) together with the nonlinear Dirichlet conditions (5.15) admits a unique maximal strong solution over $[0, T_m)$, for some $T_m > 0$. Both the solution and the maximal time of existence T_m are unique.

Proof. The proof is similar to that of Theorem 1.2. ■

We are now in a position to state the existence and uniqueness result for the (A, u) system (5.9) - (5.10). It is an immediate consequence of Theorems 1.2 and 5.3, and of Proposition 5.2.

Theorem 5.4. *Let us assume that, for $i = 1, 2, 3$, $A_i^0 > 0$, $A^0 \in [H^1(I)]^3$, $u^0 \in [H^1(I)]^3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$ satisfying the compatibility conditions*

$$\begin{aligned} A_1^0 u_1^0(1) &= A_2^0 u_2^0(0) + A_3^0 u_3^0(0), \\ u_1^0(0) &= h_1(0), \quad u_2^0(1) = h_2(0), \quad u_3^0(1) = h_3(0) \end{aligned} \quad (5.20)$$

(respectively the compatibility conditions

$$\begin{aligned} A_1^0 u_1^0(1) &= A_2^0 u_2^0(0) + A_3^0 u_3^0(0), \\ u_1^0(0) &= \frac{h_1(0)}{A_1^0(0)}, \quad u_2^0(1) = \frac{h_2(0)}{A_2^0(1)}, \quad u_3^0(1) = \frac{h_3(0)}{A_3^0(1)}. \end{aligned} \quad (5.21)$$

Then, the system (5.9) - (5.10) with the classical Dirichlet boundary conditions (5.17) (respectively the nonlinear Dirichlet boundary conditions (5.18)) admits a unique maximal strong solution over $[0, T_m)$, for some $T_m > 0$. Both the solution and the maximal time of existence T_m are unique.

6. ENERGY ESTIMATE FOR SYSTEM (1.8)

We are going to prove an energy identity for the maximal solution to system (1.8) over $[0, T_m)$. But this energy identity is also valid for solutions which do not satisfy the Dirichlet boundary conditions

$$Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad Q_3(1, t) = h_3(t), \quad t \in (0, T),$$

of system (1.8). This is why we consider a system corresponding to system 1.8, but in which the Dirichlet boundary conditions are not specified:

For $i \in \{1, 2, 3\}$, (A_i, Q_i) satisfies

$$\frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} = 0, \quad t \in (0, T_m), \quad x \in I,$$

$$\begin{aligned} \frac{\rho}{A_i} \frac{\partial Q_i}{\partial t} + \frac{\rho}{A_i} \left(\frac{2Q_i}{A_i} \frac{\partial Q_i}{\partial x} - \frac{Q_i^2}{A_i^2} \frac{\partial A_i}{\partial x} \right) + \frac{\beta}{2A_{i,0}\sqrt{A_i}} \frac{\partial A_i}{\partial x} + \frac{\nu}{4A_{i,0}A_i^{3/2}} \frac{\partial A_i}{\partial x} \frac{\partial Q_i}{\partial x} - \frac{\nu}{2A_{i,0}\sqrt{A_i}} \frac{\partial^2 Q_i}{\partial x^2} \\ = -k_f \rho \frac{Q_i}{A_i^2}, \quad t \in (0, T_m), \quad x \in I, \end{aligned}$$

$$Q_1(1, t) = Q_2(0, t) + Q_3(0, t), \quad t \in (0, T_m), \tag{6.1}$$

$$\begin{aligned} -\frac{\nu}{2A_{1,0}\sqrt{A_1}} \frac{\partial Q_1}{\partial x}(1, t) + \frac{\beta}{A_{1,0}} \left(\sqrt{A_1(1, t)} - \sqrt{A_{1,0}} \right) + \frac{1}{2} \rho \frac{Q_1^2(1, t)}{A_1^2(1, t)} \\ = -\frac{\nu}{2A_{2,0}\sqrt{A_2}} \frac{\partial Q_2}{\partial x}(0, t) + \frac{\beta}{A_{2,0}} \left(\sqrt{A_2(0, t)} - \sqrt{A_{2,0}} \right) + \frac{1}{2} \rho \frac{Q_2^2(0, t)}{A_2^2(0, t)} \\ = -\frac{\nu}{2A_{3,0}\sqrt{A_3}} \frac{\partial Q_3}{\partial x}(0, t) + \frac{\beta}{A_{3,0}} \left(\sqrt{A_3(0, t)} - \sqrt{A_{3,0}} \right) + \frac{1}{2} \rho \frac{Q_3^2(0, t)}{A_3^2(0, t)}, \quad t \in (0, T_m), \end{aligned}$$

$$A_i(x, 0) = A_i^0(x), \quad Q_i(x, 0) = Q_i^0(x), \quad x \in I.$$

We set

$$E(t) = \sum_{i=1}^3 \frac{1}{2} \int_0^1 \frac{Q_i^2}{A_i}(x, t) dx + \int_0^1 \frac{2\beta}{3\rho A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^3(x, t) dx + \int_0^1 \frac{\beta}{\rho \sqrt{A_{i,0}}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^2(x, t) dx.$$

In this section, at first we want to derive an energy estimate for the system (1.8). Let us notice that $E(t) \geq 0$ because

$$\frac{2\beta}{3\rho A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^3 + \frac{\beta}{\rho \sqrt{A_{i,0}}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^2 = \frac{2\beta}{3\rho A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^2 \left(\sqrt{A_i} + \frac{1}{2} \sqrt{A_{i,0}} \right). \tag{6.2}$$

Lemma 6.1. *If $(A_i, Q_i)_{i=1}^3$ is a maximal solution to system (6.1) over $[0, T_m)$, then we have*

$$\begin{aligned} E(t) + \int_0^t \int_0^1 \frac{\nu}{2A_{i,0}\rho\sqrt{A_i}} \left(\frac{\partial A_i}{\partial t} \right)^2(x, \tau) dx d\tau + k_f \int_0^t \int_0^1 \frac{Q_i^2}{A_i^2} dx d\tau \\ + \frac{1}{\rho} \int_0^t \left(Q_2(P_2(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_2^2(1, \tau)) d\tau + \frac{1}{\rho} \int_0^t \left(Q_3(P_3(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_3^2(1, \tau)) d\tau \right. \\ \left. - \frac{1}{\rho} \int_0^t \left(Q_1(P_1(0, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_1^2(0, \tau)) d\tau = E(0), \right. \end{aligned} \tag{6.3}$$

for all $t \in [0, T_m)$, where, for $i = 1, 2, 3$, P_i is defined by (1.4)₄.

Proof. We multiply equation (6.1)₃ by $u_i := \frac{Q_i}{A_i}$, we replace Q_i by $A_i u_i$, and we integrate over I . We end up with four terms. Let us analyse the four terms separately.

First term.

$$\int_0^1 \frac{\partial}{\partial t} (A_i u_i) \cdot u = \int_0^1 \left(\frac{\partial A_i}{\partial t} \cdot u_i^2 + \frac{A_i}{2} \frac{\partial}{\partial t} (u_i^2) \right) = \frac{1}{2} \int_0^1 \frac{\partial A_i}{\partial t} \cdot u_i^2 + \frac{1}{2} \left(\int_0^1 \frac{\partial A_i}{\partial t} \cdot u_i^2 + A \cdot \frac{\partial}{\partial t} (u_i^2) \right).$$

Thus,

$$\int_0^1 \frac{\partial}{\partial t} (A_i u_i) \cdot u_i = \frac{1}{2} \int_0^1 \frac{\partial A_i}{\partial t} \cdot u_i^2 + \frac{1}{2} \int_0^1 \frac{\partial}{\partial t} (A_i u_i^2). \quad (6.4)$$

Second term.

$$\int_0^1 \frac{\partial}{\partial x} (A_i u_i^2) \cdot u_i = \int_0^1 \left(\frac{\partial}{\partial x} (A_i u_i) \cdot u_i^2 + A u^2 \cdot \frac{\partial u}{\partial x} \right) = \int_0^1 \left(\frac{1}{2} \frac{\partial}{\partial x} (A u) \cdot u^2 + \frac{1}{2} \frac{\partial A_i}{\partial x} u_i^3 + \frac{3}{2} A_i u_i^2 \cdot \frac{\partial u_i}{\partial x} \right)$$

It can be written as

$$\int_0^1 \frac{\partial}{\partial x} (A_i u_i^2) \cdot u = \frac{1}{2} \int_0^1 \left(\frac{\partial Q_i}{\partial x} \cdot u_i^2 + \frac{\partial}{\partial x} (A_i u_i^3) \right) = -\frac{1}{2} \int_0^1 \frac{\partial A_i}{\partial t} \cdot u_i^2 + \frac{1}{2} (Q_i(1, t) u_i^2(1, t) - Q_i(0, t) u_i^2(0, t)). \quad (6.5)$$

Third term.

$$\begin{aligned} \frac{1}{\rho} \int_0^1 A_i \frac{\partial P_i}{\partial x} u_i &= \frac{1}{\rho} \left[- \int_0^1 \frac{\partial Q_i}{\partial x} (P_i - P_{\text{ext}}) + Q_i(1, t) (P_i(1, t) - P_{\text{ext}}) - Q_i(0, t) (P_i(0, t) - P_{\text{ext}}) \right] \\ &= \frac{\nu}{2A_{i,0}\rho} \int_0^1 \frac{1}{\sqrt{A_i}} \left(\frac{\partial A_i}{\partial t} \right)^2 + \frac{\beta}{A_{i,0}\rho} \int_0^1 \frac{\partial A_i}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}}) \\ &\quad + \frac{1}{\rho} Q_i(1, t) (P_i(1, t) - P_{\text{ext}}) - \frac{1}{\rho} Q_i(0, t) (P_i(0, t) - P_{\text{ext}}) \quad (6.6) \end{aligned}$$

We have

$$\begin{aligned} \int_0^1 \frac{\partial A_i}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}}) &= \int_0^1 \frac{\partial \sqrt{A_i}}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}}) 2\sqrt{A_i} = \int_0^1 \frac{\partial (\sqrt{A_i} - \sqrt{A_{i,0}})}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}}) 2\sqrt{A_i} \\ &= \frac{2}{3} \int_0^1 \frac{\partial}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}})^3 + \int_0^1 \sqrt{A_{i,0}} \frac{\partial}{\partial t} (\sqrt{A_i} - \sqrt{A_{i,0}})^2 \quad (6.7) \end{aligned}$$

This helps us to rewrite the third term.

Fourth term.

$$K_f \int_0^1 u_i^2 dx. \quad (6.8)$$

Thus, by adding (6.4), (6.5), (6.6) and (6.8), we obtain

$$\begin{aligned} \sum_{i=1}^3 \left(\frac{1}{2} \frac{d}{dt} \int_0^1 A_i u_i^2 dx + \int_0^1 \frac{\nu}{2A_{i,0} \rho \sqrt{A_i}} \left(\frac{\partial A_i}{\partial t} \right)^2 dx + \frac{\beta}{\rho \sqrt{A_{i,0}}} \frac{d}{dt} \int_0^1 (\sqrt{A_i} - \sqrt{A_{i,0}})^2 dx \right. \\ \left. + \frac{2\beta}{3\rho A_{i,0}} \frac{d}{dt} \int_0^1 (\sqrt{A_i} - \sqrt{A_{i,0}})^3 dx + k_f \int_0^1 u_i^2 dx \right) \\ + \frac{1}{\rho} \sum_{i=1}^3 \left(Q_i(P_i - P_{\text{ext}} + \frac{1}{2}u_i^2)(1, t) - Q_i(P_i - P_{\text{ext}} + \frac{1}{2}u_i^2)(0, t) \right) = 0. \end{aligned} \quad (6.9)$$

As $Q_1(1, t) = Q_2(0, t) + Q_3(0, t)$ and

$$(P_1 - P_{\text{ext}} + \frac{1}{2}u_1^2)(1, t) = (P_2 - P_{\text{ext}} + \frac{1}{2}u_2^2)(0, t) = (P_3 - P_{\text{ext}} + \frac{1}{2}u_3^2)(0, t),$$

we have

$$\begin{aligned} \sum_{i=1}^3 \left(Q_i(P_i - P_{\text{ext}} + \frac{1}{2}u_i^2)(1, t) - Q_i(P_i - P_{\text{ext}} + \frac{1}{2}u_i^2)(0, t) \right) \\ = Q_2(P_2 - P_{\text{ext}} + \frac{1}{2}\rho u_2^2)(1, t) + Q_3(P_3 - P_{\text{ext}} + \frac{1}{2}\rho u_3^2)(1, t) - Q_1(P_1 - P_{\text{ext}} + \frac{1}{2}\rho u_1^2)(0, t). \end{aligned}$$

The proof is complete. \blacksquare

If, in the energy identity (6.3), we substitute $Q_1(0, t)$, $Q_2(1, t)$, and $Q_3(2, t)$ by $h_1(t)$, $h_2(t)$ and $h_3(t)$ respectively, we do not obtain a stability estimate unless $h_1(t) = h_2(t) = h_3(t) = 0$. We are going to show that we can obtain a stability estimate if we replace the Dirichlet boundary conditions

$$Q_1(0, t) = h_1(t), \quad Q_2(1, t) = h_2(t), \quad \text{and} \quad Q_3(1, t) = h_3(t), \quad (6.10)$$

by the following nonlinear Robin boundary conditions

$$\begin{aligned} -\varepsilon \frac{\nu}{2A_{1,0} \sqrt{A_1^0}} \frac{\partial Q_1}{\partial x}(0, t) + Q_1(0, t) = \varepsilon H_1(Q_1, A_1) + h_1(t), \quad t \in (0, T_m), \\ \text{and for } i = 1, 2, \\ \varepsilon \frac{\nu}{2A_{i,0} \sqrt{A_i^0}} \frac{\partial Q_i}{\partial x}(1, t) + Q_i(1, t) = \varepsilon H_i(Q_i, A_i) + h_i(t), \quad t \in (0, T_m), \end{aligned} \quad (6.11)$$

with

$$\begin{aligned} H_1(Q_1, A_1) = \left[-\frac{\beta}{A_{1,0}} (\sqrt{A_1} - \sqrt{A_{1,0}}) + \frac{\nu}{2A_{1,0}} \frac{\partial Q_1}{\partial x} \left(\frac{1}{\sqrt{A_1}} - \frac{1}{\sqrt{A_{1,0}}} \right) - \frac{1}{2}\rho \frac{Q_1^2}{A_1^2} \right] \Big|_{x=0}, \\ \text{and for } i = 1, 2, \\ H_i(Q_i, A_i) = \left[\frac{\beta}{A_{i,0}} (\sqrt{A_i} - \sqrt{A_{i,0}}) - \frac{\nu}{2A_{i,0}} \frac{\partial Q_i}{\partial x} \left(\frac{1}{\sqrt{A_i}} - \frac{1}{\sqrt{A_{i,0}}} \right) + \frac{1}{2}\rho \frac{Q_i^2}{A_i^2} \right] \Big|_{x=1}. \end{aligned} \quad (6.12)$$

For the network represented in Figure 1, we consider the system (6.1) with the nonlinear Robin boundary conditions (6.11)-(6.12).

The notions of strong solution to system (6.1)-(6.11)-(6.12) over the time interval $[0, T]$, and of maximal strong solution over the time interval $[0, T_m)$, are similar to those in Definition 1.1.

Theorem 6.2. *Let us assume that, for $i = 1, 2, 3$, $A_i^0 > 0$, $A^0 \in [H^1(I)]^3$, $Q^0 \in [H^1(I)]^3$, $h_i \in H_{\text{loc}}^{3/4}([0, \infty))$ and*

$$Q_1^0(1) = Q_2^0(0) + Q_3^0(0). \quad (6.13)$$

Then, the system (6.1)-(6.11)-(6.12) admits a unique maximal strong solution over $[0, T_m)$, for some $T_m > 0$. Both the solution and the maximal time of existence T_m are unique.

Proof. The proof is similar to that of Theorem 1.2. \blacksquare

Proposition 6.3. *The maximal solution $(A_i, Q_i)_{i=1}^3$ to system (6.1)-(6.11)-(6.12) over $[0, T_m)$, whose existence is stated in Theorem 6.2, satisfies the following stability estimate*

$$\begin{aligned}
E(t) &+ \sum_{i=1}^3 \int_0^t \int_0^1 \frac{\nu}{2A_{i,0}\rho\sqrt{A_i}} \left(\frac{\partial A_i}{\partial t} \right)^2 (x, \tau) dx d\tau + k_f \sum_{i=1}^3 \int_0^t \int_0^1 \frac{Q_i^2}{A_i^2} dx d\tau \\
&+ \frac{1}{2\rho\varepsilon} \int_0^t Q_2^2(1, \tau) d\tau + \frac{1}{2\rho\varepsilon} \int_0^t Q_3^2(1, \tau) d\tau + \frac{1}{2\rho\varepsilon} \int_0^t Q_1^2(0, \tau) d\tau \\
&\leq E(0) + \frac{1}{2\rho\varepsilon} \int_0^t h_2^2(1, \tau) d\tau + \frac{1}{2\rho\varepsilon} \int_0^t h_3^2(1, \tau) d\tau + \frac{1}{2\rho\varepsilon} \int_0^t h_1^2(0, \tau) d\tau, \quad \text{for all } t \in [0, T_m).
\end{aligned} \tag{6.14}$$

Proof. Due to Lemma 6.1, to prove the proposition, it is sufficient to estimate the following boundary terms

$$\begin{aligned}
&\int_0^t \left(Q_2(1, \tau)(P_2(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_2^2(1, \tau)) \right) d\tau, \quad \int_0^t \left(Q_3(1, \tau)(P_3(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_3^2(1, \tau)) \right) d\tau \\
&\text{and} \quad \int_0^t \left(Q_1(0, \tau)(P_1(0, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_1^2(0, \tau)) \right) d\tau.
\end{aligned}$$

Let us only estimate the last term, the two others can be estimated similarly. The Robin boundary condition satisfied by Q_1 can be written in the form

$$\begin{aligned}
-Q_1(0, \tau) + h_1(t) &= \varepsilon \left(\frac{\beta}{A_{1,0}} \left(\sqrt{A_1} - \sqrt{A_{1,0}} \right) - \frac{\nu}{2A_{1,0}\sqrt{A_1}} \frac{\partial Q_1}{\partial x} + \frac{1}{2}\rho \frac{Q_1^2}{A_1^2}(0, \tau) \right) \\
&= \varepsilon \left(\frac{\beta}{A_{1,0}} \left(\sqrt{A_1} - \sqrt{A_{1,0}} \right) + \frac{\nu}{2A_{1,0}\sqrt{A_1}} \frac{\partial A_1}{\partial t} + \frac{1}{2}\rho \frac{Q_1^2}{A_1^2}(0, \tau) \right) \\
&= \varepsilon (P_1(0, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_1^2(0, \tau)).
\end{aligned}$$

Thus, we have

$$\varepsilon (P_1(0, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_1^2(0, \tau)) = -Q_1(0, \tau) + h_1(t).$$

and

$$\begin{aligned}
&-\int_0^t Q_1(0, \tau) \left(P_1(0, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_1^2(0, \tau) \right) d\tau \\
&= \frac{1}{\varepsilon} \int_0^t Q_1(0, \tau) \left(Q_1(0, \tau) - h_1(\tau) \right) d\tau \geq \frac{1}{2\varepsilon} \int_0^t Q_1^2(0, \tau) d\tau - \frac{1}{2\varepsilon} \int_0^t h_1^2(\tau) d\tau.
\end{aligned}$$

The nonlinear Robin boundary conditions at $x = 1$ satisfied by Q_i with $i = 2, 3$, correspond to

$$P_i(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_i^2(0, \tau) = Q_i(1, \tau) - h_i(t).$$

Thus, for $i = 2, 3$, we have

$$\int_0^t \left(Q_2(1, \tau)(P_2(1, \tau) - P_{\text{ext}} + \frac{1}{2}\rho u_2^2(1, \tau)) \right) d\tau \geq \frac{1}{2\varepsilon} \int_0^t Q_i^2(0, \tau) d\tau - \frac{1}{2\varepsilon} \int_0^t h_i^2(\tau) d\tau,$$

and the proof is complete. ■

7. A GENERAL NETWORK

We now consider a network constituted of N_s segments of length 1, numbered from $i = 1$ to $i = N_s$ and parametrized by $x \in I = (0, 1)$. The origins and extremities of the segments, corresponding to the points where $x = 0$ and $x = 1$ respectively, are the nodes of the network. The origins of segments are either inlet points or branching points, while the extremities are either outlet points or branching points. The set of branching points is $\{B_\ell \mid 1 \leq \ell \leq N_b\}$. At any branching point B_ℓ , the subsets of indices $J_0^\ell \subset \{1, \dots, N_s\}$ and $J_1^\ell \subset \{1, \dots, N_s\}$, corresponding to segments connected to B_ℓ by their origin and their extremity respectively, are nonempty. The subsets of indices $J_{\text{in}} \subset \{1, \dots, N_s\}$ and $J_{\text{out}} \subset \{1, \dots, N_s\}$ are nonempty, and they correspond to origins of segments which are inlet points and extremities of segments which are outlet points respectively. The boundary conditions at a branching point B_ℓ are

$$\begin{aligned} \sum_{i \in J_0^\ell} Q_i(0, t) &= \sum_{i \in J_1^\ell} Q_i(1, t), \\ P_{j_0^\ell}^i(0, t) + \frac{1}{2} \rho u_{j_0^\ell}^2(0, t) &= P_j(0, t) + \frac{1}{2} \rho u_j^2(0, t) \quad \text{for all } j \in J_0^\ell, \\ P_{j_1^\ell}^i(0, t) + \frac{1}{2} \rho u_{j_1^\ell}^2(0, t) &= P_j(1, t) + \frac{1}{2} \rho u_j^2(1, t) \quad \text{for all } j \in J_1^\ell, \end{aligned} \quad (7.1)$$

where $j_0^\ell = \min J_0^\ell$.

For this type of network, we consider the nonlinear system

$$\begin{aligned} \text{For } i \in \{1, \dots, N_s\}, (A_i, Q_i, u_i), \text{ with } u_i &= \frac{Q_i}{A_i}, \text{ satisfies} \\ \frac{\partial A_i}{\partial t} + \frac{\partial Q_i}{\partial x} &= 0, \quad t \in (0, T), x \in I, \\ \frac{\partial Q_i}{\partial t} + \frac{\partial}{\partial x} \left(\frac{Q_i^2}{A_i} \right) + \frac{A_i}{\rho} \frac{\partial P_i}{\partial x} &= -k_f \frac{Q_i}{A_i}, \quad t \in (0, T), x \in I \\ P_i &= P_{\text{ext}} + \frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) + \frac{\nu}{A_{i,0}} \frac{\partial}{\partial t} (\sqrt{A_i}), \\ A_i(x, 0) &= A_i^0, \quad Q_i(x, 0) = Q_i^0(x), \quad x \in I, \end{aligned} \quad (7.2)$$

with the following nonlinear boundary conditions at the branching points

$$\begin{aligned} \sum_{i \in J_0^\ell} Q_i(0, t) &= \sum_{i \in J_1^\ell} Q_i(1, t), \text{ for all } 1 \leq \ell \leq N_b \\ P_{j_0^\ell}^i(0, t) + \frac{1}{2} \rho u_{j_0^\ell}^2(0, t) &= P_j(0, t) + \frac{1}{2} \rho u_j^2(0, t) \quad \text{for all } 1 \leq \ell \leq N_b, \text{ and all } j \in J_0^\ell, \\ P_{j_1^\ell}^i(0, t) + \frac{1}{2} \rho u_{j_1^\ell}^2(0, t) &= P_j(1, t) + \frac{1}{2} \rho u_j^2(1, t) \quad \text{for all } 1 \leq \ell \leq N_b, \text{ and all } j \in J_1^\ell, \end{aligned}$$

with either the following Dirichlet boundary conditions

$$\begin{aligned} Q_i(0, t) &= h_i(t), \quad i \in J_{\text{in}}, \quad t \in (0, T), \\ Q_i(1, t) &= h_i(t), \quad i \in J_{\text{out}}, \quad t \geq 0, \end{aligned} \quad (7.3)$$

or the following nonlinear Robin boundary conditions

$$\begin{aligned} -\varepsilon \frac{\nu}{2A_{i,0}\sqrt{A_i^0}} \frac{\partial Q_i}{\partial x}(0, t) + Q_i(0, t) &= \varepsilon H_i^{\text{in}}(Q_1, A_1) + h_i(t), \quad i \in J_{\text{in}}, \quad t \in (0, T), \\ \varepsilon \frac{\nu}{2A_{i,0}\sqrt{A_i^0}} \frac{\partial Q_i}{\partial x}(1, t) + Q_i(1, t) &= \varepsilon H_i^{\text{out}}(Q_i, A_i) + h_i(t), \quad i \in J_{\text{out}}, \quad t \in (0, T), \end{aligned} \quad (7.4)$$

where

$$H_i^{\text{in}}(A_i, Q_i) = \left[-\frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) + \frac{\nu}{2A_{i,0}} \frac{\partial Q_i}{\partial x} \left(\frac{1}{\sqrt{A_i}} - \frac{1}{\sqrt{A_{i,0}}} \right) - \frac{1}{2} \rho \frac{Q_i^2}{A_i^2} \right] \Big|_{x=0}, \quad i \in J_{\text{in}},$$

and

$$H_i^{\text{out}}(A_i, Q_i) = \left[\frac{\beta}{A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right) - \frac{\nu}{2A_{i,0}} \frac{\partial Q_i}{\partial x} \left(\frac{1}{\sqrt{A_i}} - \frac{1}{\sqrt{A_{i,0}}} \right) + \frac{1}{2} \rho \frac{Q_i^2}{A_i^2} \right] \Big|_{x=1}, \quad i \in J_{\text{out}}.$$

We look for solutions to system (7.2)-(7.3), or system (7.2)-(7.4), in the space

$$\mathcal{E}_T = \left\{ (A_i, Q_i)_{i=1}^{N_s} \mid A_i \in H^1(0, T; H^1(0, 1)) \cap L^\infty(0, T; H^1(0, 1)), \right. \\ \left. Q_i \in L^2(0, T; H^2(0, 1)) \cap H^1(0, T; L^2(0, 1)) \cap L^\infty(0, T; H^1(0, 1)), A_i(\cdot, 0) = A_i^0 \text{ in } (0, 1) \right\}, \quad (7.5)$$

equipped with the norm

$$\left\| (A_i, Q_i)_{i=1}^{N_s} \right\|_{\mathcal{E}_T} = \sum_{i=1}^{N_s} \left(\|A_i\|_{H^1(0, T; H^1(0, 1))} + \|A_i\|_{L^\infty(0, T; H^1(0, 1))} + \|Q_i\|_{L^2(0, T; H^2(0, 1))} \right. \\ \left. + \|Q_i\|_{H^1(0, T; L^2(0, 1))} + \|Q_i\|_{L^\infty(0, T; H^1(0, 1))} \right). \quad (7.6)$$

Definition 7.1. We say that $(A_i, Q_i)_{i=1}^{N_s}$ is a strong solution to system (7.2)-(7.3) (or system (7.2)-(7.4)), over the time interval $[0, T]$, when $(A_i, Q_i)_{i=1}^{N_s} \in \mathcal{E}_T$,

$$A_i(x, t) > 0 \quad \text{for all } (x, t) \in [0, 1] \times [0, T], \quad (7.7)$$

and $(A_i, Q_i)_{i=1}^{N_s}$ satisfies the equations (7.2)₂₋₃ in the sense of distributions and the boundary and initial conditions in the sense of traces.

We say that $(A_i, Q_i)_{i=1}^{N_s}$ is a maximal strong solution to system (7.2)-(7.3) (or system (7.2)-(7.4)), over the time interval $[0, T_m)$, when either $T_m = \infty$, or $T_m < \infty$ and, for all $0 < T < T_m$, $(A_i, Q_i)_{i=1}^{N_s}$ is a strong solution to system (7.2) over the time interval $[0, T]$, and when

$$\lim_{T \rightarrow T_m} \left(\left\| (A_i, Q_i)_{i=1}^{N_s} \right\|_{\mathcal{E}_T} + \max \left\{ |A_i(x, T)|^{-1} \mid 1 \leq i \leq N_s, x \in [0, 1] \right\} \right) = \infty. \quad (7.8)$$

Theorem 7.2. Let $A_{i,0} > 0$ denote the sectional area of the i th vessel at equilibrium state, and let $Q_{i,0}$ belong to $H^1(I)$, and let assume that

$$\sum_{i \in J_0^\ell} Q_i(0, t) = \sum_{i \in J_\ell^t} Q_i(0, t) \quad \text{for all } \ell \in \{1, \dots, N_b\}. \quad (7.9)$$

Then, the system (7.2)-(7.3) (or system (7.2)-(7.4)) admits a unique maximal solution over a time interval $[0, T_m)$.

Proof. The proof is similar to those of Theorems 1.2 and 6.2. ■

We set

$$E(t) = \sum_{i=1}^{N_s} \frac{1}{2} \int_0^1 \frac{Q_i^2}{A_i}(x, t) dx + \int_0^1 \frac{2\beta}{3\rho A_{i,0}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^3(x, t) dx + \int_0^1 \frac{\beta}{\rho \sqrt{A_{i,0}}} \left(\sqrt{A_i} - \sqrt{A_{i,0}} \right)^2(x, t) dx.$$

Proposition 7.3. The maximal solution $(A_i, Q_i, P_i)_{i=1}^{N_s}$ to system (7.2)-(7.4), over $[0, T_m)$, whose existence is stated in Theorem 7.2, satisfies the following stability estimate

$$E(t) + \sum_{i=1}^{N_s} \int_0^t \int_0^1 \frac{\nu}{2A_{i,0}\rho\sqrt{A_i}} \left(\frac{\partial A_i}{\partial t} \right)^2(x, \tau) dx d\tau + k_f \sum_{i=1}^{N_s} \int_0^t \int_0^1 \frac{Q_i^2}{A_i^2} dx d\tau \\ \leq E(0) + \frac{1}{2\rho\varepsilon} \sum_{i \in J_{\text{in}} \cup J_{\text{out}}} \int_0^t h_i^2(\tau) d\tau, \quad \text{for all } t \in [0, T_m). \quad (7.10)$$

Proof. The proof is similar to that of Proposition 6.3. ■

REFERENCES

- [1] R. L. ARMENTANO, J. G. BARRA, J. LEVENSON, A. SIMON, AND R. H. PICHEL, *Arterial wall mechanics in conscious dogs. assessment of viscous, inertial, and elastic moduli to characterize aortic wall behavior*, *Circ. Res.*, 76 (1995), pp. 468–478.
- [2] A. BENSOUSSAN, G. DA PRATO, M. C. DELFOUR, AND S. K. MITTER, *Representation and control of infinite dimensional systems*, *Systems & Control: Foundations & Applications*, Birkhäuser Boston, Inc., Boston, MA, second ed., 2007.
- [3] P. J. BLANCO, S. M. WATANABE, E. A. DARI, M. A. R. F. PASSOS, AND R. A. FEIJÓO, *Blood flow distribution in an anatomically detailed arterial network model: criteria and algorithms*, *Biomechanics and Modeling in Mechanobiology*, 13 (2014), pp. 1303–1330.
- [4] P. J. BLANCO, S. M. WATANABE, M. A. R. F. PASSOS, P. A. LEMOS, AND R. A. FEIJÓO, *An anatomically detailed arterial network model for one-dimensional computational hemodynamics*, *IEEE Transactions on Biomedical Engineering*, 62 (2015), pp. 736–753.
- [5] S. ČANIĆ, C. J. HARTLEY, D. ROSENSTRAUCH, J. TAMBAČA, G. GUIDOBONI, AND A. MIKELIĆ, *Blood flow in compliant arteries: An effective viscoelastic reduced model, numerics, and experimental validation*, *Annals of Biomedical Engineering*, 34 (2006), pp. 575–592.
- [6] S. ČANIĆ AND E. H. KIM, *Mathematical analysis of the quasilinear effects in a hyperbolic model blood flow through compliant axi-symmetric vessels*, *Mathematical Methods in the Applied Sciences*, 26 (2003), pp. 1161–1186.
- [7] K. DEVAULT, P. A. GREMAUD, V. NOVAK, M. S. OLUFSEN, G. VERNIÈRES, AND P. ZHAO, *Blood flow in the circle of Willis: modeling and calibration*, *Multiscale Model. Simul.*, 7 (2008), pp. 888–909.
- [8] L. C. EVANS, *Partial differential equations*, vol. 19 of *Graduate Studies in Mathematics*, American Mathematical Society, Providence, RI, second ed., 2010.
- [9] M. A. FERNÁNDEZ, V. MILIŠIĆ, AND A. QUARTERONI, *Analysis of a geometrical multiscale blood flow model based on the coupling of ODEs and hyperbolic PDEs*, *Multiscale Model. Simul.*, 4 (2005), pp. 215–236.
- [10] L. FORMAGGIA, J. F. GERBEAU, F. NOBILE, AND A. QUARTERONI, *On the coupling of 3D and 1D Navier-Stokes equations for flow problems in compliant vessels*, *Comput. Methods Appl. Mech. Engrg.*, 191 (2001), pp. 561–582.
- [11] P. GRISVARD, *Caractérisation de quelques espaces d'interpolation*, *Arch. Rational Mech. Anal.*, 25 (1967), pp. 40–63.
- [12] G. GRUBB AND V. A. SOLONNIKOV, *Boundary value problems for the nonstationary Navier-Stokes equations treated by pseudo-differential methods*, *Math. Scand.*, 69 (1991), pp. 217–290 (1992).
- [13] O. A. LADYŽENSKAJA, V. A. SOLONNIKOV, AND N. N. URAL'CEVA, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. *Translations of Mathematical Monographs*, Vol. 23, American Mathematical Society, Providence, R.I., 1968.
- [14] R. LAL, B. MOHAMMADI, AND F. NICOU, *Data assimilation for identification of cardiovascular network characteristics*, *Int. J. Numer. Methods Biomed. Eng.*, 33 (2017), pp. e2824, 17.
- [15] J.-L. LIONS AND E. MAGENES, *Non-homogeneous boundary value problems and applications. Vol. II*, Springer-Verlag, New York-Heidelberg, 1972. Translated from the French by P. Kenneth, *Die Grundlehren der mathematischen Wissenschaften*, Band 182.
- [16] D. MAITY, J.-P. RAYMOND, AND A. ROY, *Maximal-in-time existence and uniqueness of strong solution of a 3d fluid-structure interaction model*, <https://hal.archives-ouvertes.fr/hal-02912001>, (2020).
- [17] A. C. I. MALOSSI, P. J. BLANCO, AND S. DEPARIS, *A two-level time step technique for the partitioned solution of one-dimensional arterial networks*, *Comput. Methods Appl. Mech. Engrg.*, 237/240 (2012), pp. 212–226.
- [18] G. I. MONTECINOS, L. O. MÜLLER, AND E. F. TORO, *Hyperbolic reformulation of a 1D viscoelastic blood flow model and ADER finite volume schemes*, *J. Comput. Phys.*, 266 (2014), pp. 101–123.
- [19] L. O. MÜLLER, G. LEUGERING, AND P. J. BLANCO, *Consistent treatment of viscoelastic effects at junctions in one-dimensional blood flow models*, *J. Comput. Phys.*, 314 (2016), pp. 167–193.
- [20] R. RAGHU AND C. A. TAYLOR, *Verification of a one-dimensional finite element method for modeling blood flow in the cardiovascular system incorporating a viscoelastic wall model*, *Finite Elem. Anal. Des.*, 47 (2011), pp. 586–592.
- [21] P. REYMOND, F. MERENDA, F. PERRIN, D. RÜFENACHT, AND N. STERGIOPOULOS, *Validation of a one-dimensional model of the systemic arterial tree*, *Am. J. Physiol. Heart Circ. Physiol.*, 297 (2009), pp. H208–H222.
- [22] B. N. STEELE, D. VALDEZ-JASSO, M. A. HAIDER, AND M. S. OLUFSEN, *Predicting arterial flow and pressure dynamics using a 1D fluid dynamics model with a viscoelastic wall*, *SIAM J. Appl. Math.*, 71 (2011), pp. 1123–1143.
- [23] M. STROCCHI, C. CONTARINO, Q. ZHANG, R. BONMASSARI, AND E. F. TORO, *A global mathematical model for the simulation of stenoses and bypass placement in the human arterial system*, *Appl. Math. Comput.*, 300 (2017), pp. 21–39.
- [24] X. WANG, J.-M. FULLANA, AND P.-Y. LAGRÉE, *Verification and comparison of four numerical schemes for a 1d viscoelastic blood flow model*, *Comput. Methods Biomech. Biomed. Engrg.*, 18 (2015), pp. 1704–1725.
- [25] X. WANG, S. NISHI, M. MATSUKAWA, A. GHIGO, P.-Y. LAGRÉE, AND J.-M. FULLANA, *Fluid friction and wall viscosity of the 1d blood flow model*, *J. Biomechanics*, 49 (2016), pp. 565–571.
- [26] M. ZAGZOULE AND J.-P. MARC-VERGNES, *A global mathematical model of the cerebral circulation in man*, *J. Biomechanics*, 19 (1986), pp. 1015–1022.
- [27] Y. ZOCALO, D. BIA, R. LLUBERAS, AND R. L. ARMENTANO, *Verification and comparison of four numerical schemes for a 1d viscoelastic blood flow model*, *Biological Research*, 41 (2008), pp. 227–233.

DEBAYAN MAITY
TIFR CENTRE FOR APPLICABLE MATHEMATICS,
560065 BANGALORE, KARNATAKA, INDIA.
Email address: debayan@tifrbng.res.in

JEAN-PIERRE RAYMOND
UNIVERSITÉ DE TOULOUSE & CNRS, UPS, INSTITUT DE MATHÉMATIQUES,
31062 TOULOUSE CEDEX 9, FRANCE.
Email address: raymond@math.univ-toulouse.fr

ARNAB ROY
INSTITUTE OF MATHEMATICS OF THE CZECH ACADEMY OF SCIENCES,
ŽITNÁ 25, 115 67 PRAHA 1, CZECH REPUBLIC.
Email address: royarnab244@gmail.com