

# Existence of Core Localized Toroidicity-induced Alfvén Eigenmode

G. Y. Fu

*Princeton Plasma Physics Laboratory, Princeton University*

*P. O. Box 451*

*Princeton, New Jersey 08543, U.S.A.*

(December 9, 1994)

## Abstract

The core-localized toroidicity-induced Alfvén eigenmode (TAE) is shown to exist at finite plasma pressure due to finite aspect ratio effects in tokamak plasma. The new critical beta for the existence of the TAE mode is given by  $\alpha \approx 3\epsilon + 2s^2$ , where  $\epsilon = r/R$  is the inverse aspect ratio,  $s$  is the magnetic shear and  $\alpha = -Rq^2 d\beta/dr$  is the normalized pressure gradient. In contrast, previous critical  $\alpha$  is given by  $\alpha \approx s^2$ . In the limit of  $s \ll \sqrt{r/R}$ , the new critical  $\alpha$  is greatly enhanced by the finite aspect ratio effects.

52.35.Bj, 52.55.Fa, 52.55.Pi

**MASTER**

Typeset using REVTeX

The toroidicity-induced Alfvén eigenmode [1,2] (TAE) exists in a tokamak plasma due to toroidal mode coupling and finite magnetic shear. The mode frequency is located inside a continuum gap induced by toroidicity. Recently, it has been shown [3–5] that there is a critical pressure gradient above which the TAE mode no longer exists. In the limit of small shear, the critical value is given by  $\alpha = \alpha_c = s^2/(1+s)$ , where  $s = rdq/qdr$  is the magnetic shear,  $\alpha = -Rq^2 d\beta/dr$  represents the product of magnetic field curvature and plasma pressure gradient with  $R$  being the major radius,  $r$  being the plasma radius,  $q$  being the safety factor and  $\beta$  being the plasma toroidal beta. This result implies that the TAE mode should not exist in the core of tokamak plasma where the values of  $\alpha$  and  $s$  are comparable and small for typical parameters. In particular,  $\alpha/s \gg 1$  asymptotically when  $r$  goes to zero. However, recent numerical results [6] indicate that a TAE mode, which is localized at a single gap, does exist in the core of the tokamak plasma, where  $\epsilon = r/R \sim 0.1$  and  $s \sim \alpha \sim \epsilon$ . We call this type of TAE mode the core-localized TAE mode. These core-localized modes are particularly susceptible to destabilization by fusion alpha particles in a tokamak plasma since the density profile of the alpha particles is sharply peaked at the center of plasma [6].

In this work, we will show that the effects of finite aspect ratio change the critical  $\alpha$  qualitatively using the high- $n$  ballooning mode equation. The previous results [3,4], based on the standard  $s - \alpha$  model ballooning mode equation [7], is only valid in the limit of  $\epsilon \ll s^2$ . We will show that our new critical  $\alpha$  reduces to  $\alpha_c = 3\epsilon$  in the limit of  $s \ll \sqrt{\epsilon} \ll 1$ . In the following, we will derive an analytic expression for critical  $\alpha$  by taking into account the effects of finite aspect ratio.

We start with the following ballooning mode equation:

$$\begin{aligned} \frac{\partial}{\partial \theta} [G_1(\theta) + G_2(\theta)h^2(\theta)] \frac{\partial}{\partial \theta} \Phi + \Omega^2 [G_1(\theta) + G_2(\theta)h^2(\theta)] (1 + 4\epsilon \cos \theta) \Phi \\ + \alpha [\cos \theta + h(\theta) \sin \theta] \Phi = 0 \end{aligned} \quad (1)$$

where  $h(\theta) = s\theta - \bar{\alpha} \sin \theta$ ,  $G_1(\theta) = 1 - 2(\epsilon + \Delta') \cos \theta$ ,  $G_2(\theta) = 1 + 2\Delta' \cos \theta$ ,  $\bar{\alpha} =$

## **DISCLAIMER**

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

$\alpha + 2\epsilon - 2(1 - s)\Delta'$ ,  $\Delta' = \epsilon(1/4 + \beta_p)$  is the radial derivative of the Shafranov shift, and  $\Omega$  is the mode frequency normalized to the Alfvén frequency. The poloidal beta is defined as  $\beta_p = 8\pi(\langle p \rangle - p)/B_p^2$  where  $\langle \rangle$  denotes a volume average.

Equation 1 is derived for a large aspect ratio, low-beta tokamak plasma by using the Shafranov shifted circle flux coordinates [8]. We note that the previous work [3,4] neglects the  $\epsilon$  term in  $G_1(\theta)$  and  $G_2(\theta)$  and assumes  $\alpha \gg \epsilon$ . In this limit, the Eq. (1) reduces to the standard  $s - \alpha$  model ballooning equation [7]:

$$\frac{\partial}{\partial \theta} [1 + h^2(\theta)] \frac{\partial}{\partial \theta} \Phi + \Omega^2 [1 + h^2(\theta)] (1 + 4\epsilon \cos \theta) \Phi + \alpha [\cos \theta + h(\theta) \sin \theta] \Phi = 0. \quad (2)$$

Comparing Eqs. (1) and (2), the new terms in Eq. (1), of order  $O(\epsilon)$ , come from the intrinsic toroidicity and pressure-induced Shafranov shift. We will show below that these  $\epsilon$  terms in  $G_1$  and  $G_2$  can not be neglected when  $\epsilon$  is comparable or larger than  $s^2$ .

To make analytic progress, we make the following transformation  $\Phi \rightarrow (1/\sqrt{F})\Phi$  with  $F = G_1(\theta) + G_2(\theta)h^2(\theta)$ , the Eq. (1) then becomes

$$\frac{\partial^2}{\partial \theta^2} \Phi + \left[ \Omega^2(1 + 4\epsilon \cos \theta) - \frac{F''}{2F} + \frac{(F')^2}{4F^2} + \frac{\alpha(\cos \theta + h(\theta) \sin \theta)}{F} \right] \Phi = 0 \quad (3)$$

where the prime denotes the derivative with respect to  $\theta$ . Assuming  $\epsilon \ll 1$  and  $s \sim \alpha \sim O(\epsilon)$ , we expand Eq. (3) to the second order of  $\epsilon$ . The resulting equation is given by

$$\frac{\partial^2}{\partial \theta^2} \Phi + \Omega^2(1 + 4\hat{\epsilon} \cos \theta) \Phi + \left[ \frac{H_1(\theta)}{1 + s^2 \theta^2} - \frac{H_2(\theta)}{(1 + s^2 \theta^2)^2} \right] \Phi = 0 \quad (4)$$

where the functions  $H_1$  and  $H_2$  are defined as

$$H_1(\theta) = (\alpha - \delta) \cos \theta - 2\alpha \Delta' \cos^2 \theta + (2\delta \Delta' + \alpha \bar{\alpha} - \bar{\alpha}^2) \sin^2 \theta \quad (5)$$

$$H_2(\theta) = (s - \bar{\alpha} \cos \theta)^2 + 2\delta(\delta - \alpha) \cos^2 \theta - (\delta^2 + 2\bar{\alpha}^2 - 2\alpha \bar{\alpha}) \sin^2 \theta \quad (6)$$

with  $\hat{\epsilon} \approx \epsilon + \Delta'$  and  $\delta = \epsilon + 2\Delta'$ .

Following Fu and Cheng [3], a dispersion relation for the TAE mode frequency can be derived by solving Eq. (4) asymptotically. The result is given by

$$\sqrt{\frac{\Omega^2 - \Omega_-^2}{\Omega_+^2 - \Omega^2}} = \frac{\pi}{4s}(1 + \epsilon + s)(\alpha_c - \alpha) \quad (7)$$

where  $\Omega_-$  and  $\Omega_+$  are the lower bound and the upper bound of the Alfvén continuum gap respectively. The critical  $\alpha$  for the existence of TAE mode,  $\alpha_c$ , is given by

$$\alpha_c = \frac{(1 + \epsilon)(\epsilon + 2\Delta') + s^2 - 2(\epsilon - \Delta')s}{1 + \epsilon + s}. \quad (8)$$

In the limit of  $\epsilon \rightarrow 0$  and  $\Delta' \rightarrow 0$ , the critical value given in Eq. (8) reduces to the previous results [3–5]. It should be noted that  $\Delta'$  is dependent on the poloidal beta which is proportional to  $\alpha$ . This implicit dependence can be made explicit in the core of plasma where we can expand the plasma pressure near  $r = 0$ . Thus, we have following relations near the center of plasma for typical pressure profile of form  $p \propto (1 - r^2)^b$  with  $b$  being a constant:  $\alpha = 4\epsilon\beta_p$  and  $\bar{\alpha} = \delta = 3\epsilon/2 + \alpha/2$ . The critical  $\alpha$  then reduces to

$$\alpha_c = \frac{3\epsilon(1 + \epsilon) + 2s^2 - 3\epsilon s}{1 + \epsilon + s} \quad (9)$$

It is instructive to note that in the limit of zero shear, the critical alpha is simply given by  $\alpha_c = 3\epsilon$  which corresponds to  $\beta_{pc} = 3/4$ . Thus, the finite aspect ratio effects greatly enhance the critical beta for small shear. It is also interesting to note that in the limit of  $\epsilon \rightarrow 0$ , the critical  $\alpha$  is a factor two larger than the previous results [3–5]. This factor of two comes from the pressure-induced Shafranov shift.

The critical  $\alpha$ , given by Eq. (9), has been confirmed by solving Eq.(1) numerically. Figure 1 shows that the analytic results for the critical  $\alpha$  (solid line) agree very well with the numerical results (solid dots). For comparison, we have shown in Fig. 2 both the previous results (solid line marked with  $r/R = 0$ ) and our new results (solid line marked with  $r/R = 0.1$ ). As a reference, the ballooning mode first stability boundary is also shown (dotted line). We observe that the effects of finite aspect ratio makes the critical  $\alpha$  much

closer to the first stability boundary for small value of shear. ( It was shown previously [3] that the critical  $\alpha$  for TAE mode approaches the first stability boundary asymptotically for large values of shear). Although our results are based on shifted-circle model equilibria, we conjecture that the critical  $\alpha$  for TAE mode is also close to the ballooning first stability limit in general equilibria, and that the TAE can generally exist as long as the tokamak plasma is ballooning stable.

Finally, the analytic critical  $\alpha$  in the limit of  $s^2 \ll \epsilon$ ,  $\alpha_c = 3\epsilon$ , has been confirmed in both magnitude and scaling by numerical calculations using a global stability code NOVA [6].

In conclusion, we have shown that the core-localized TAE mode exists at finite plasma pressure in tokamaks due to finite aspect ratio effects. The critical beta for the existence of TAE mode in the core region of plasma is given by  $\beta_p = 3/4$  in the limit of  $s \ll \sqrt{r/R} \ll 1$ .

The author gratefully acknowledges stimulating discussion with Drs. H. L. Berk, C. Z. Cheng and J. W. Van Dam. This work is supported by the U.S. Department of Energy under Contract No. DE-AC02-76-CHO-3073.

## REFERENCES

- [1] C. Z. Cheng and M. S. Chance, *Phys. Fluids* **29**, 3695 (1986).
- [2] C. Z. Cheng, L. Chen, and M. S. Chance, *Ann. Phys. NY* **161**, 21 (1984).
- [3] G. Y. Fu and C. Z. Cheng, *Phys. Fluids* **B2**, 985 (1990).
- [4] F. Zonca and L. Chen, *Bull. Am. Phys. Soc.* **37**, 1436 (1992).
- [5] J. W. Van Dam and H. L. Berk, *Bull. Am. Phys. Soc.* **38**, 1944 (1993).
- [6] G. Y. Fu et al., to be submitted to *Phys. Rev. Lett.*
- [7] J. W. Connor, R. J. Hastie and J. B. Taylor, *Phys. Rev. Lett.* **40**, 396 (1978).
- [8] J. D. Meiss and R. D. Hazeltine, *Phys. Fluids* **B2**, 2563 (1990).



## FIGURES

FIG. 1. The critical  $\alpha$  value,  $\alpha_c$ , as a function of shear  $s$  obtained with  $\epsilon = 0.1$  near the center of plasma.

FIG. 2. The critical  $\alpha$  for the TAE as a function of shear  $s$  from both the previous results (solid line marked with  $r/R = 0$ ) and the new results (solid line marked with  $r/R = 0.1$ ). The ballooning first stability boundary is also shown (dotted line).




Fig. 1




Fig. 2

# EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA  
 Prof. R.C. Cross, Univ. of Sydney, AUSTRALIA  
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA  
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA  
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA  
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA  
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM  
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM  
 Commission-European, DG. XII-Fusion Prog., BELGIUM  
 Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM  
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL  
 Prof. Dr. I.C. Nascimento, Instituto Fisica, Sao Paulo, BRAZIL  
 Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL  
 Documents Office, Atomic Energy of Canada Ltd., CANADA  
 Ms. M. Morin, CCFM/Tokamak de Varennes, CANADA  
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA  
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA  
 Prof. J. Teichmann, Univ. of Montreal, CANADA  
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA  
 Prof. T.W. Johnston, INRS-Energie, CANADA  
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA  
 Dr. C.R. James,, Univ. of Alberta, CANADA  
 Dr. P. Lukác, Komenského Univerzita, CZECHO-SLOVAKIA  
 The Librarian, Culham Laboratory, ENGLAND  
 Library, R61, Rutherford Appleton Laboratory, ENGLAND  
 Mrs. S.A. Hutchinson, JET Library, ENGLAND  
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS  
 P. Mähönen, Univ. of Helsinki, FINLAND  
 Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE  
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE  
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE  
 Prof. E. Economou, Univ. of Crete, GREECE  
 Ms. C. Rinni, Univ. of Ioannina, GREECE  
 Preprint Library, Hungarian Academy of Sci., HUNGARY  
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA  
 Dr. P. Kaw, Inst. for Plasma Research, INDIA  
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL  
 Librarian, International Center for Theo Physics, ITALY  
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY  
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY  
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY  
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN  
 Prof. I. Kawakami, Hiroshima Univ., JAPAN  
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN  
 Librarian, Naka Fusion Research Establishment, JAERI, JAPAN  
 Director, Japan Atomic Energy Research Inst., JAPAN  
 Prof. S. Itoh, Kyushu Univ., JAPAN  
 Research Info. Ctr., National Inst. for Fusion Science, JAPAN  
 Prof. S. Tanaka, Kyoto Univ., JAPAN  
 Library, Kyoto Univ., JAPAN  
 Prof. N. Inoue, Univ. of Tokyo, JAPAN  
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN  
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN  
 Dr. G.S. Lee, Korea Basic Sci. Ctr., KOREA  
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA  
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA  
 Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND  
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA  
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA  
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA  
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA  
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL  
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA  
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA  
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN  
 Dr. L. Stenflo, Univ. of UMEA, SWEDEN  
 Library, Royal Inst. of Technology, SWEDEN  
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN  
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND  
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS  
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY  
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. i Apparatus, USSR  
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR  
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR  
 Librarian, The Ukr.SSR Academy of Sciences, USSR  
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR  
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY  
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY  
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY  
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY  
 Librarian, Max-Planck-Institut, W. GERMANY