



## Advances in the Theory of Nonlinear Analysis and its Applications

ISSN: 2587-2648

Peer-Reviewed Scientific Journal

# Existence of Solutions for Nonlocal Boundary Value Problem of Hadamard Fractional Differential Equations

Subramanian Muthaiah<sup>a</sup>, Manigandan Murugesan<sup>a</sup>, Nandha Gopal Thangaraj<sup>a</sup>

<sup>a</sup>Department of Mathematics, Sri Ramakrishna Mission Vidyalaya College of Arts and Science, Coimbatore, Tamil Nadu, India.

---

### Abstract

We investigate the existence and uniqueness of solutions for Hadamard fractional differential equations with nonlocal integral boundary conditions, by using the Leray-Schauder nonlinear alternative, Leray Schauder degree theorem, Krasnoselskiis fixed point theorem, Schaefer's fixed point theorem, Banach fixed point theorem, Nonlinear Contractions. Two examples are also presented to illustrate our results.

*Keywords:* Hadamard fractional derivative; Hadamard fractional integral; Fractional differential equation; Integral boundary conditions; Existence; Fixed point.

*2010 MSC:* 26A33, 34A08, 34A12.

---

### 1. Introduction

Fractional differential equations have increased extensive consideration from both hypothetical and the applied perspectives as of late years. There are various applications in an assortment of fields, for example, chemical physics, permeable media, signal processing, viscoelasticity, aerodynamics, exhibiting anomalous diffusion, fluid flow, electrical systems, financial aspects, etc., rather than integer-order differential and integral operators, fractional-order differential operators are nonlocal and give the way to investigate the inherited properties of a few materials and procedures. The monographs [14, 15, 16, 18, 20] have ordinarily

---

*Email addresses:* [subramanianmcb@gmail.com](mailto:subramanianmcb@gmail.com) (Subramanian Muthaiah), [yogimani22@outlook.com](mailto:yogimani22@outlook.com) (Manigandan Murugesan), [nandhu792002@yahoo.co.in](mailto:nandhu792002@yahoo.co.in) (Nandha Gopal Thangaraj)

referred to the hypothesis of fractional derivatives and integrals and applications to differential equations of fractional order. For more points of interest and models, see [1, 4, 5, 6, 9, 10, 17, 24, 25, 26, 27, 28, 29, 21, 22, 23, 31], and the references in that.

In any case, it has seen that the greater part of the work on the point is concerned about Riemann-Liouville or Caputo type fractional differential equation. Other than these fractional derivatives, another sort of fractional derivatives established in the literature is the fractional derivative because of Hadamard made known to in 1892 [12], differs from the aforementioned derivatives in the sense that the kernel of the integral in the definition of Hadamard derivative contains a logarithmic function of an arbitrary exponent. A point by point depiction of Hadamard fractional derivatives and integral having been discovered in [2, 3, 7, 8, 11, 13, 30, 33]. Recently Qinghua et.al [19] discussed a Lyapunovtype inequality with the Hadamard fractional derivative. Similarly, Wang et.al [32] studied the nonlocal Hadamard fractional boundary value problem with Hadamard integral and discrete boundary conditions.

In this paper, we introduce nonlocal boundary value problem on the Hadamard fractional derivatives of the form:

$$\begin{aligned} {}^H\mathcal{D}^\varrho y(\tau) &= g(\tau, y(\tau)), \quad \tau \in [1, T], \\ y(1) &= 0, \quad y'(1) = 0, \quad {}^H\mathcal{D}^\varsigma y(T) = \omega {}^H\mathcal{J}^\gamma y(\varphi), \quad 1 < \varphi < T, \end{aligned} \tag{1}$$

where  ${}^H\mathcal{D}^\varrho, {}^H\mathcal{D}^\varsigma$  denotes the Hadamard fractional derivatives of order  $2 < \varrho \leq 3, 1 < \varsigma < 2$ , and  $g: [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is a given continuous function and  $\omega$  is positive real constant. The rest of the paper has organized as follows: section 2 has devoted to some fundamental concepts of fractional calculus with basic lemmas related to the given problem. The existence and uniqueness results based on, Leray-Schauder nonlinear alternative, Leray-Schauder degree theorem, Krasnoselskii’s fixed point theorem, Schaefer’s fixed point theorem, Banach fixed point theorem, and Nonlinear contractions have obtained in section 3. The validation of the results has done by providing examples in section 4.

## 2. Preliminaries

We begin with some basic definitions, properties, and lemmas with results [14, 18].

**Definition 2.1.** The Hadamard fractional integral of order  $\varrho \in \mathbb{R}^+$  of a function  $g \in L^p[b, c], 0 \leq b \leq \tau \leq c \leq \infty$ , is defined as

$$({}^H\mathcal{J}^\varrho g)(\tau) = \frac{1}{\Gamma(\varrho)} \int_b^\tau \left(\log \frac{\tau}{\sigma}\right)^{\varrho-1} g(\sigma) \frac{d\sigma}{\sigma}, \quad \varrho > 0.$$

**Definition 2.2.** Let  $0 < b < c < \infty, \delta = \left(\tau \frac{d}{d\tau}\right)$  and  $\mathcal{AC}_\delta^n[b, c] = \{g : [b, c] \rightarrow \mathbb{R} : \delta^{n-1}[g(\tau)] \in \mathcal{AC}[b, c]\}$ . The Hadamard fractional derivative of order  $\varrho > 0$  for a function  $g \in \mathcal{AC}_\delta^n[b, c]$  is defined as

$$({}^H\mathcal{D}^\varrho g)(\tau) = \frac{1}{\Gamma(n - \varrho)} \left(\tau \frac{d}{d\tau}\right)^n \int_1^\tau \left(\log \frac{\tau}{\sigma}\right)^{n-\varrho-1} g(\sigma) \frac{d\sigma}{\sigma},$$

where  $n - 1 < \varrho < n, n = [\varrho] + 1, [\varrho]$  denotes the integer part of the real number  $\varrho$  and  $\log(\cdot) = \log_e(\cdot)$ .

**Lemma 2.3.** Let  $b, \varrho, \varsigma > 0$ , then

$$\begin{aligned} ({}^H\mathcal{D}^\varsigma (\log \frac{\tau}{b})^{\varrho-1})(y) &= \frac{\Gamma(\varrho)}{\Gamma(\varrho - \varsigma)} \left(\log \frac{y}{b}\right)^{\varrho-\varsigma-1}, \\ ({}^H\mathcal{J}^\varsigma (\log \frac{\tau}{b})^{\varrho-1})(y) &= \frac{\Gamma(\varrho)}{\Gamma(\varrho + \varsigma)} \left(\log \frac{y}{b}\right)^{\varrho+\varsigma-1}. \end{aligned}$$

**Lemma 2.4.** Let  $\varrho > 0$  and  $y \in \mathcal{C}[1, \infty) \cap L^1[1, \infty)$ . Then the solution of Hadamard fractional differential equation  $({}^H\mathcal{D}^\varrho y)(\tau) = 0$  is given by

$$y(\tau) = \sum_{i=1}^n a_i (\log \tau)^{\varrho-i}.$$

and the following formula holds:

$${}^H\mathcal{J}^\varrho {}^H\mathcal{D}^\varrho y(\tau) = y(\tau) + \sum_{i=1}^n a_i (\log \tau)^{\varrho-i},$$

where  $a_i \in \mathbb{R}$ ,  $i = 1, 2, \dots, n$  and  $n - 1 < \varrho < n$ .

We define space  $\mathcal{P} = \mathcal{C}([1, T], \mathbb{R})$  the Banach space of all continuous functions from  $[1, T] \rightarrow \mathbb{R}$  endowed with a topology of uniform convergence with the norm defined by  $\|y\| = \sup\{|y(\tau)|, \tau \in [1, T]\}$ . Next, we present an auxiliary lemma which plays a key role in the sequel.

**Lemma 2.5.** For any  $\hat{g} \in \mathcal{C}([1, T], \mathbb{R})$ ,  $y \in \mathcal{C}([1, T], \mathbb{R})$ , the function  $y$  is the solution of the problem

$$\begin{aligned} {}^H\mathcal{D}^\varrho y(\tau) &= \hat{g}(\tau), \quad \tau \in [1, T], \\ y(1) &= 0, \quad y'(1) = 0, \quad {}^H\mathcal{D}^\varsigma y(T) = \omega {}^H\mathcal{J}^\gamma y(\varphi), \end{aligned} \tag{2}$$

if and only if

$$y(\tau) = {}^H\mathcal{J}^\varrho \hat{g}(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} \hat{g}(\varphi) - {}^H\mathcal{J}^{\varrho-\varsigma} \hat{g}(T) \right], \tag{3}$$

where

$$\Lambda = \frac{\Gamma(\varrho)}{\Gamma(\varrho - \varsigma)} (\log T)^{\varrho-\varsigma-1} - \omega \frac{\Gamma(\varrho)}{\Gamma(\varrho + \gamma)} (\log \varphi)^{\varrho+\gamma-1}. \tag{4}$$

*Proof.* Applying the operator  ${}^H\mathcal{J}^\varrho$  on the linear fractional differential equations in (2), we obtain

$$y(\tau) = {}^H\mathcal{J}^\varrho \hat{g}(\tau) + a_1 (\log \tau)^{\varrho-1} + a_2 (\log \tau)^{\varrho-2} + a_3 (\log \tau)^{\varrho-3}, \tag{5}$$

where  $a_1, a_2$  and  $a_3 \in \mathbb{R}$ , are arbitrary unknown constants. Using the boundary conditions (2) in (5), we get  $a_2, a_3 = 0$  and

$$a_1 = \frac{1}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} \hat{g}(\varphi) - {}^H\mathcal{J}^{\varrho-\varsigma} \hat{g}(T) \right].$$

Substituting the value of  $a_1, a_2$  in (5), we get the solution (3). This completes the proof. □

### 3. Existence and Uniqueness Results

In view of Lemma 2.5, we interpret an operator  $\mathcal{T} : \mathcal{P} \rightarrow \mathcal{P}$  as

$$\begin{aligned} \mathcal{T}(y)(\tau) &= {}^H\mathcal{J}^\varrho g(\sigma, y(\sigma))(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} g(\sigma, y(\sigma))(\varphi) \right. \\ &\quad \left. - {}^H\mathcal{J}^{\varrho-\varsigma} g(\sigma, y(\sigma))(T) \right]. \end{aligned} \tag{6}$$

In the sequel, we use the following expressions:

$$\begin{aligned} {}^H J^\varrho g(\sigma, y(\sigma))(\tau) &= \frac{1}{\Gamma(\varrho)} \int_1^\tau \left(\log \frac{\tau}{\sigma}\right)^{\varrho-1} g(\sigma, y(\sigma)) \frac{d\sigma}{\sigma}, \\ \omega {}^H J^{\varrho+\gamma} g(\sigma, y(\sigma))(\varphi) &= \omega \frac{1}{\Gamma(\varrho+\gamma)} \int_1^\varphi \left(\log \frac{\varphi}{\sigma}\right)^{\varrho+\gamma-1} g(\sigma, y(\sigma)) \frac{d\sigma}{\sigma}, \\ {}^H J^{\varrho-\varsigma} g(\sigma, y(\sigma))(T) &= \frac{1}{\Gamma(\varrho-\varsigma)} \int_1^T \left(\log \frac{T}{\sigma}\right)^{\varrho-\varsigma-1} g(\sigma, y(\sigma)) \frac{d\sigma}{\sigma}. \end{aligned}$$

Suitable for computation, we represent:

$$\Delta = \frac{(\log T)^\varrho}{\Gamma(\varrho+1)} + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \frac{\omega(\log \varphi)^{\varrho+\gamma}}{\Gamma(\varrho+\gamma+1)} + \frac{(\log T)^{\varrho-\varsigma}}{\Gamma(\varrho-\varsigma+1)} \right]. \tag{7}$$

Our first existence result is based on Leray-Schauder nonlinear alternative.

**Theorem 3.1.** *Let us speculate that  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function and the following conditions hold :*

- (E<sub>1</sub>) *There exists a function  $q \in \mathcal{C}([1, T], \mathbb{R}^+)$ , and  $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  nondecreasing such that  $|g(\tau, y)| \leq q(\tau)\psi(\|y\|)$  for each  $(\tau, y) \in [1, T] \times \mathbb{R}$ ;*
- (E<sub>2</sub>) *There exists a number  $\mathcal{L} > 0$  such that*

$$\begin{aligned} \frac{\mathcal{L}}{\|q\|\psi(\mathcal{L})} &> {}^H J^\varrho q(\sigma)(T) + \mathcal{G} \\ \mathcal{G} &= \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} q(\sigma)(\varphi) + {}^H J^{\varrho-\varsigma} q(\sigma)(T) \right]. \end{aligned}$$

Then there exists at least one solution for problem (1) on  $[1, T]$ .

*Proof.* To begin with, the operator  $\mathcal{T} : \mathcal{P} \rightarrow \mathcal{P}$  is described by (6). Next, we demonstrate that  $\mathcal{T}$  maps bounded sets into bounded sets in  $\mathcal{C}([1, T], \mathbb{R})$ . For a positive number  $\theta$ , let  $\mathcal{B}_\theta = \{y \in \mathcal{C}([1, T], \mathbb{R}) : \|y\| \leq \theta\}$  be a bounded set in  $\mathcal{C}([1, T], \mathbb{R})$ . Then, for each  $y \in \mathcal{B}_\theta$ , we have

$$\begin{aligned} |(\mathcal{T}y)(\tau)| &\leq {}^H J^\varrho |g(\sigma, y(\sigma))|(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \\ &\quad \left. + {}^H J^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(\tau) \right] \\ &\leq \psi(\|y\|) {}^H J^\varrho q(\sigma)(T) + \frac{\psi(\|y\|)(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} q(\sigma)(\varphi) \right. \\ &\quad \left. + {}^H J^{\varrho-\varsigma} q(\sigma)(T) \right], \end{aligned}$$

and consequently,

$$\|\mathcal{T}y\| \leq \psi(\|\theta\|) \left\{ {}^H J^\varrho q(\sigma)(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} q(\sigma)(\varphi) + {}^H J^{\varrho-\varsigma} q(\sigma)(T) \right] \right\}.$$

We shall proceed to prove that the operator  $\mathcal{J}$  maps bounded sets into equicontinuous sets of  $\mathcal{C}([1, T], \mathbb{R})$ . For  $\tau_1, \tau_2 \in [1, T]$  with  $\tau_1 < \tau_2$ , and  $y \in \mathcal{B}_\theta$  is a bounded set of  $\mathcal{C}([1, T], \mathbb{R})$ . Then we have

$$\begin{aligned} & |(\mathcal{J}y)(\tau_2) - (\mathcal{J}y)(\tau_1)| \\ & \leq |{}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma))|(\tau_2) - {}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma))|(\tau_1)| \\ & \quad + \frac{|(\log \tau_2)^{\varrho-1} - (\log \tau_1)^{\varrho-1}|}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \\ & \quad \left. + {}^H\mathcal{J}^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \\ & \leq \frac{\psi(\|\theta\|)}{\Gamma(\varrho)} \left| \int_0^{\tau_1} \left[ \left(\log \frac{\tau_2}{\sigma}\right)^{\varrho-1} - \left(\log \frac{\tau_1}{\sigma}\right)^{\varrho-1} \right] q(\sigma) \frac{d\sigma}{\sigma} \right. \\ & \quad \left. + \int_{\tau_1}^{\tau_2} \left(\log \frac{\tau_2}{\sigma}\right)^{\varrho-1} q(\sigma) \frac{d\sigma}{\sigma} \right| \\ & \quad + \frac{|(\log \tau_2)^{\varrho-1} - (\log \tau_1)^{\varrho-1}|}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} q(\sigma)(\varphi) + {}^H\mathcal{J}^{\varrho-\varsigma} q(\sigma)(T) \right]. \end{aligned}$$

Hence we have that right hand side of the above inequality tends to zero independent of  $y \in \mathcal{B}_\theta$  as  $\tau_2 - \tau_1 \rightarrow 0$ . Therefore, the operator  $\mathcal{J}(y)$  is equicontinuous and consequently, by Arzela-Ascoli theorem, it is completely continuous. Next, we demonstrate that the boundedness of the set of all solutions to equations  $y = \nu \mathcal{J}(y)$ ,  $0 < \nu < 1$ . Let  $y$  be a solution. Then, for  $\tau \in [1, T]$ , and using the computations in proving that  $\mathcal{J}$  is bounded, we have

$$\begin{aligned} |(\mathcal{J}y)(\tau)| & \leq \psi(\|y\|) \left\{ {}^H\mathcal{J}^\varrho q(\sigma)(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} q(\sigma)(\varphi) \right. \right. \\ & \quad \left. \left. + {}^H\mathcal{J}^{\varrho-\varsigma} q(\sigma)(T) \right] \right\} \\ & = \psi(\|y\|) {}^H\mathcal{J}^\varrho q(\sigma)(T) + \mathcal{G}. \end{aligned}$$

In view of  $(\mathcal{E}_2)$ , there exists  $\mathcal{L}$  such that  $\|y\| \neq \mathcal{L}$ . Let us set

$$\mathcal{M} = \{y \in \mathcal{C}([1, T], \mathbb{R}) : \|y\| < \mathcal{L}\}.$$

Bearing in mind that the operator  $\mathcal{J} : \overline{\mathcal{W}} \rightarrow \mathcal{C}([1, T], \mathbb{R})$  is continuous and completely continuous. From the of choice of  $\mathcal{W}$ , there is no  $y \in \partial \mathcal{W}$  such that  $y = \nu \mathcal{J}(y)$ ,  $0 < \nu < 1$ . Consequently, by the Leray-Schauder nonlinear alternative, we deduce that  $\mathcal{J}$  has a fixed point  $y \in \overline{\mathcal{W}}$  which is a solution of the problem (1).  $\square$

Our second existence result is based on Leray-Schauder degree theorem.

**Theorem 3.2.** Let  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function. Suppose that

$(\mathcal{E}_3)$  There exists constants  $0 \leq \phi < \frac{1}{\Delta}$ , and  $\mathcal{A} > 0$  such that

$$|g(\tau, y)| \leq \phi|y| + \mathcal{A} \quad \forall (\tau, y) \in [1, T] \times \mathbb{R},$$

where  $\Delta$  is described by (7). Then the boundary value problem (1) has at least one solution on  $[1, T]$ .

*Proof.* We interpret an operator  $\mathcal{J} : \mathcal{P} \rightarrow \mathcal{P}$  as in (6). In view of the fixed point problem

$$y = \mathcal{J}y, \tag{8}$$

we shall to prove the existence of at least one solution  $y \in \mathcal{C}([1, T])$  satisfying (8). Set a ball  $\mathcal{B}_\mathcal{R} \subset \mathcal{C}([1, T])$ , as

$$\mathcal{B}_\mathcal{R} = \{y \in \mathcal{P} : \max_{\tau \in \mathcal{C}([1, T])} |y(\tau)| < \mathcal{R}\},$$

with a constant radius  $\mathcal{R} > 0$ . Hence, we shall show that  $\mathcal{T} : \overline{\mathcal{B}}_{\mathcal{R}} \rightarrow \mathcal{C}([1, T])$  satisfies the following:

$$y \neq \zeta \mathcal{T}y, \quad \forall y \in \partial \mathcal{B}_{\mathcal{R}}, \quad \forall \zeta \in [1, T]. \tag{9}$$

We set

$$\mathcal{T}(\zeta, y) = \zeta \mathcal{I}y, \quad y \in \mathcal{P}, \quad \zeta \in [1, T].$$

As shown in Theorem 3.2 we have that the operator  $\mathcal{T}$  is continuous, uniformly bounded and equicontinuous. Then, by the Arzela-Ascoli theorem, a continuous map  $f_{\zeta}$  described by  $f_{\zeta}(y) = y - \mathcal{T}(\zeta, y) = y - \zeta \mathcal{I}y$  is completely continuous. If (9) holds, then the following Leray-Schauder degrees are well defined and by the homotopy invariance of topological degree, it follows that

$$\begin{aligned} \deg(f_{\zeta}, \mathcal{B}_{\mathcal{R}}, 0) &= \deg(\mathcal{J} - \zeta \mathcal{I}, \mathcal{B}_{\mathcal{R}}, 0) = \deg(f, \mathcal{B}_{\mathcal{R}}, 0) \\ &= \deg(f_0, \mathcal{B}_{\mathcal{R}}, 0) = \deg(\mathcal{J}, \mathcal{B}_{\mathcal{R}}, 0) = 1 \neq 0, \quad 0 \in \mathcal{B}_{\mathcal{R}}, \end{aligned}$$

where  $\mathcal{J}$  denotes the unit operator. By the nonzero property of Leray-Schauder degree,  $f_1(y) = y - \mathcal{J}y = 0$  for at least one  $y \in \mathcal{B}_{\mathcal{R}}$ . Let us assume that  $y = \zeta \mathcal{I}y$  for some  $\zeta \in [1, T]$  and for all  $\tau \in [1, T]$  so that

$$\begin{aligned} |(\mathcal{T}y)(\tau)| &\leq {}^H\mathcal{J}^{\varrho} |g(\sigma, y(\sigma))|(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \\ &\quad \left. + {}^H\mathcal{J}^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \\ &\leq \phi |y| + \mathcal{A} \left[ {}^H\mathcal{J}^{\varrho} q(\sigma)(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} q(\sigma)(\varphi) \right. \right. \\ &\quad \left. \left. + {}^H\mathcal{J}^{\varrho-\varsigma} q(\sigma)(T) \right] \right] \\ &= (\phi |y| + \mathcal{A}) \Delta \end{aligned}$$

which, on taking norm  $\sup_{\tau \in [1, T]} |y(\tau)| = \|y\|$  and solving for  $\|y\|$ , yields

$$\|y\| \leq \frac{\mathcal{A} \Delta}{1 - \phi \Delta}.$$

If  $\mathcal{R} = \frac{\Delta}{1 - \phi \Delta} + 1$ , inequality (9) holds. This completes the proof. □

Our third existence result is based on Krasnoselskii’s fixed point theorem.

**Theorem 3.3.** *Let  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function such that the following conditions hold:*

- ( $\mathcal{E}_4$ )  $|g(\tau, p_1) - g(\tau, p_2)| \leq \mathcal{S} |p_1 - p_2|, \forall \tau \in [1, T], p_1, p_2 \in \mathbb{R}, \mathcal{S} > 0.$
- ( $\mathcal{E}_5$ )  $|g(\tau, y(\tau))| \leq \xi(\tau)$  for  $(\tau, y) \in [1, T] \times \mathbb{R}$ , and  $\xi \in \mathcal{C}([1, T], \mathbb{R}^+)$  with  $\|\xi\| = \max_{\tau \in [1, T]} |\xi(\tau)|.$

If

$$\left\{ \frac{\mathcal{S}(\log T)^{\varrho-1}}{\Lambda} \left( \frac{\omega(\log \varphi)^{\varrho+\gamma}}{\Gamma(\varrho + \gamma + 1)} + \frac{(\log T)^{\varrho-\varsigma}}{\Gamma(\varrho - \varsigma + 1)} \right) \right\} < 1. \tag{10}$$

Then, there exists at least one solution for the problem (1) on  $[1, T]$ .

*Proof.* Let us interpret  $\mathcal{B}_{\theta} = \{y \in \mathcal{P} : \|y\| \leq \theta\}$ , where  $\theta \geq \|\xi\| \Delta$ . To prove the hypothesis of Krasnoselskii’s fixed point result, we split the operator  $\mathcal{T}$  given by (6) as  $\mathcal{T} = \mathcal{T}_1 + \mathcal{T}_2$  on  $\mathcal{B}_{\theta}$ , where

$$\begin{aligned} (\mathcal{T}_1 y)(\tau) &= {}^H\mathcal{J}^{\varrho} g(\sigma, y(\sigma))(\tau) \\ (\mathcal{T}_2 y)(\tau) &= \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} g(\sigma, y(\sigma))(\varphi) - {}^H\mathcal{J}^{\varrho-\varsigma} g(\sigma, y(\sigma))(T) \right]. \end{aligned}$$

For  $\hat{y}_1, \hat{y}_2 \in \mathcal{B}_\theta$ ,

$$\begin{aligned} & |(\mathcal{T}_1 \hat{y}_1)(\tau) + (\mathcal{T}_2 \hat{y}_2)(\tau)| \\ & \leq \sup_{\tau \in [1, T]} \left\{ {}^H J^\varrho |g(\sigma, y(\sigma))|(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \right. \\ & \quad \left. \left. + {}^H J^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \right\} \\ & \leq \|\xi\| \left\{ \frac{(\log T)^\varrho}{\Gamma(\varrho+1)} + \frac{(\log T)^{\varrho-1}}{\Lambda} \left( \frac{\omega (\log \varphi)^{\varrho+\gamma}}{\Gamma(\varrho+\gamma+1)} + \frac{(\log T)^{\varrho-\varsigma}}{\Gamma(\varrho-\varsigma+1)} \right) \right\} \\ & \leq \|\xi\| \Delta \leq \theta, \end{aligned}$$

which implies that  $\mathcal{T}_1 \hat{y}_1 + \mathcal{T}_2 \hat{y}_2 \in \mathcal{B}_\theta$ .

Now, we will show that  $\mathcal{T}_2$  is a contraction. Let  $p_1, p_2 \in \mathbb{R}$ ,  $\tau \in [1, T]$ . Then, using the assumption  $(\mathcal{E}_4)$  together with (10), we get

$$\|\mathcal{T}_2 p_1 - \mathcal{T}_2 p_2\| \leq \frac{S(\log T)^{\varrho-1}}{\Lambda} \left( \frac{\omega (\log \varphi)^{\varrho+\gamma}}{\Gamma(\varrho+\gamma+1)} + \frac{(\log T)^{\varrho-\varsigma}}{\Gamma(\varrho-\varsigma+1)} \right) \|p_1 - p_2\|.$$

By the assumption  $(\mathcal{E}_4)$ , it follows that the operator  $\mathcal{T}_2$  is a contraction. Next, we will show that  $\mathcal{T}_1$  is compact and continuous. Continuity of  $g$  implies that the operator  $\mathcal{T}_1$  is continuous. Also,  $\mathcal{T}_1$  is uniformly bounded on  $B_\theta$  as

$$\|\mathcal{T}_1 y\| \leq \frac{\|\xi\| (\log T)^\varrho}{\Gamma(\varrho+1)}.$$

Moreover, with  $\sup_{(\tau, y) \in [1, T] \times \mathcal{B}_\theta} |g(\tau, y)| = \hat{g} < \infty$  and  $\tau_1 < \tau_2$ ,  $\tau_1, \tau_2 \in [1, T]$ , we have

$$\begin{aligned} |(\mathcal{T}_1 y)(\tau_2) - (\mathcal{T}_1 y)(\tau_1)| &= |{}^H J^\varrho |g(\sigma, y(\sigma))|(\tau_2) - {}^H J^\varrho |g(\sigma, y(\sigma))|(\tau_1)| \\ &\leq \frac{\hat{g}}{\Gamma(\varrho)} \left| \int_0^{\tau_1} \left[ \left( \log \frac{\tau_2}{\sigma} \right)^{\varrho-1} - \left( \log \frac{\tau_1}{\sigma} \right)^{\varrho-1} \right] \frac{d\sigma}{\sigma} \right. \\ &\quad \left. + \int_{\tau_1}^{\tau_2} \left( \log \frac{\tau_2}{\sigma} \right)^{\varrho-1} \frac{d\sigma}{\sigma} \right|. \end{aligned} \tag{11}$$

Clearly, the right-hand sides of (11) tends to zero independent of  $y$  as  $\tau_2 - \tau_1 \rightarrow 0$ . Thus,  $\mathcal{T}_1$  is relatively compact on  $\mathcal{B}_\theta$ . Hence, by the Arzela-Ascoli Theorem,  $\mathcal{T}_1$  is compact on  $\mathcal{B}_\theta$ . Thus, all the assumptions of Krasnoselskii’s fixed point theorem are satisfied. Therefore, there exists at least one solution for problem (1) on  $[1, T]$ .  $\square$

Our next existence result is based on Schaefer’s fixed point theorem.

**Theorem 3.4.** *Let  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function. Assume that there exists a positive constant  $\hat{\mathcal{M}}$  such that  $|g(\tau, y)| \leq \hat{\mathcal{M}}$  for  $\tau \in [1, T]$ ,  $y \in \mathbb{R}$ . Then there exists at least one solution for problem (1) on  $[1, T]$ .*

*Proof.* To begin with, we demonstrate that the operator  $\mathcal{T} : \mathcal{P} \rightarrow \mathcal{P}$  is completely continuous. By continuity of the function  $g$ , it follows that the operator  $\mathcal{T}$  is continuous.

For a positive constant  $\theta$ , let  $\mathcal{B}_\theta = \{y \in \mathcal{P} : \|y\| \leq \theta\}$  be a bounded set in  $\mathcal{P}$ . Then, for  $\tau \in [1, T]$ , we obtain

$$\begin{aligned} |(\mathcal{J}y)(\tau)| &\leq {}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma))|(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \\ &\quad \left. + {}^H\mathcal{J}^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \\ &\leq \widehat{\mathcal{M}} \left\{ {}^H\mathcal{J}^\varrho(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma}(\varphi) + {}^H\mathcal{J}^{\varrho-\varsigma}(T) \right] \right\} = \widehat{\mathcal{M}}\Delta. \end{aligned}$$

Hence it follows that  $\mathcal{J}$  is uniformly bounded.

We shall proceed to prove that the operator  $\mathcal{J}$  is equicontinuous. For  $\tau_1, \tau_2 \in [1, T]$  with  $\tau_1 < \tau_2$ , we have  $|(\mathcal{J}y)(\tau_2) - (\mathcal{J}y)(\tau_1)|$

$$\begin{aligned} &\leq |{}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma))|(\tau_2) - {}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma))|(\tau_1)| \\ &\quad + \frac{|(\log \tau_2)^{\varrho-1} - (\log \tau_1)^{\varrho-1}|}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \\ &\quad \left. + {}^H\mathcal{J}^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \\ &\leq \frac{\widehat{\mathcal{M}}}{\Gamma(\varrho)} \left| \int_0^{\tau_1} \left[ \left(\log \frac{\tau_2}{\sigma}\right)^{\varrho-1} - \left(\log \frac{\tau_1}{\sigma}\right)^{\varrho-1} \right] \frac{d\sigma}{\sigma} + \int_{\tau_1}^{\tau_2} \left(\log \frac{\tau_2}{\sigma}\right)^{\varrho-1} \frac{d\sigma}{\sigma} \right| \\ &\quad + \frac{|(\log \tau_2)^{\varrho-1} - (\log \tau_1)^{\varrho-1}|}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma}(\varphi) + {}^H\mathcal{J}^{\varrho-\varsigma}(T) \right]. \end{aligned}$$

Hence we have that right hand side of the above inequality tends to zero independent of  $y \in \mathcal{B}_\theta$  as  $\tau_2 - \tau_1 \rightarrow 0$ . Therefore, the operator  $\mathcal{J}(y)$  is equicontinuous and consequently, by Arzela-Ascoli theorem, it is completely continuous. Next, we consider the set  $\mathcal{H} = \{y \in \mathcal{P} : y = v\mathcal{J}(y), 0 < v < 1\}$ . Then, we have to show that  $\mathcal{H}$  is bounded, let  $y \in \mathcal{H}$  and  $\tau \in [1, T]$ . Then

$$\|y\| \leq \widehat{\mathcal{M}} \left\{ \frac{(\log T)^\varrho}{\Gamma(\varrho+1)} + \frac{(\log T)^{\varrho-1}}{\Lambda} \left( \frac{\omega(\log \varphi)^{\varrho+\gamma}}{\Gamma(\varrho+\gamma+1)} + \frac{(\log T)^{\varrho-\varsigma}}{\Gamma(\varrho-\varsigma+1)} \right) \right\} = \widehat{\Theta}.$$

Thus,  $\mathcal{H}$  is bounded. Hence it follows by Schaefer’s fixed point result that the equation (1) has at least one solution on  $[1, T]$ . □

Next, we establish the uniqueness of solution using Banach fixed point theorem for the problem (1).

**Theorem 3.5.** *Let  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function satisfying the assumptions  $(\mathcal{E}_4)$ . In addition, it is assumed that  $S\Delta < 1$ , where  $\Delta$  is described by (7). Then there exists at most one solution for problem (1) on  $[1, T]$ .*

*Proof.* Let us interpret  $\sup_{\tau \in [1, T]} |g(\tau, 0)| = \Omega < \infty$ . Nominating  $\theta \geq \frac{\Omega\Delta}{1 - S\Delta}$ , we demonstrate that  $\mathcal{J}\mathcal{B}_\theta \subset \mathcal{B}_\theta$ ,



where  $\mathcal{B}_\theta = \{y \in \mathcal{P} : \|y\| \leq \theta\}$ . For  $y \in \mathcal{B}_\theta$ , we have

$$\begin{aligned}
 & |(\mathcal{J}y)(\tau)| \\
 & \leq \sup_{\tau \in [1, T]} \left\{ {}^H J^\varrho |g(\sigma, y(\sigma))|(\tau) + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} |g(\sigma, y(\sigma))|(\varphi) \right. \right. \\
 & \quad \left. \left. + {}^H J^{\varrho-\varsigma} |g(\sigma, y(\sigma))|(T) \right] \right\} \\
 & \leq (\mathcal{S}\theta + \mathcal{Q}) \sup_{\tau \in [1, T]} \left\{ {}^H J^\varrho(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma}(\varphi) + {}^H J^{\varrho-\varsigma}(T) \right] \right\} \\
 & \leq (\mathcal{S}\theta + \mathcal{Q})\Delta.
 \end{aligned} \tag{12}$$

Thus, it follows from (12) that  $\|(\mathcal{J}y)\| \leq \theta$ .

Now, for  $y, \hat{y} \in \mathcal{P}$ , we obtain

$$\begin{aligned}
 |\mathcal{J}y(\tau) - \mathcal{J}\hat{y}(\tau)| & \leq \sup_{\tau \in [1, T]} \left\{ {}^H J^\varrho |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(\tau) \right. \\
 & \quad \left. + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(\varphi) \right. \right. \\
 & \quad \left. \left. + {}^H J^{\varrho-\varsigma} |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(T) \right] \right\} \\
 & \leq \left[ \mathcal{G}\|y - \hat{y}\| {}^H J^\varrho(T) + \frac{\mathcal{G}\|y - \hat{y}\|(\log T)^{\varrho-1}}{\Lambda} \left( \omega {}^H J^{\varrho+\gamma}(\varphi) \right. \right. \\
 & \quad \left. \left. + {}^H J^{\varrho-\varsigma}(T) \right) \right] \\
 & = \mathcal{G}\Delta\|y - \hat{y}\|.
 \end{aligned}$$

Thus,

$$\|\mathcal{J}y - \mathcal{J}\hat{y}\| \leq \mathcal{S}\Delta\|y - \hat{y}\|.$$

Since  $\mathcal{S}\Delta < 1$  by the given assumption, therefore  $\mathcal{J}$  is a contraction. Hence it follows by Banach fixed point theorem that the equation (1) has at most one solution on  $[1, T]$ .  $\square$

Finally, we establish the uniqueness of solution using nonlinear contractions for the problem (1).

**Theorem 3.6.** *Let  $g : [1, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function satisfying the assumption:*

( $\mathcal{E}_6$ )  $|g(\tau, u) - g(\tau, v)| \leq f(\tau) \frac{|u-v|}{\vartheta + |u-v|}$ ,  $\forall \tau \in [1, T], u, v \geq 0$ , where  $f : [1, T] \rightarrow \mathbb{R}^+$  is continuous and  $\vartheta$  the constant described by

$$\vartheta = {}^H J^\varrho f(T) + \frac{(\log T)^{\varrho-1}}{\Lambda} \left[ \omega {}^H J^{\varrho+\gamma} f(\varphi) + {}^H J^{\varrho-\varsigma} f(T) \right].$$

Then there exists at most one solution for problem (1) on  $[1, T]$ .

*Proof.* Let us interpret the operator  $\mathcal{J} : \mathcal{P} \rightarrow \mathcal{P}$  as in (6) and the continuous nondecreasing function  $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  by

$$\Phi(\varpi) = \frac{\vartheta \varpi}{\vartheta + \varpi}, \quad \varpi > 0.$$

Now, for  $y, \hat{y} \in \mathcal{P}$  and for each  $\tau \in [1, T]$ , we obtain

$$\begin{aligned} |\mathcal{J}y(\tau) - \mathcal{J}\hat{y}(\tau)| &\leq \sup_{\tau \in [1, T]} \left\{ {}^H\mathcal{J}^\varrho |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(\tau) \right. \\ &\quad + \frac{(\log \tau)^{\varrho-1}}{\Lambda} \left[ \omega {}^H\mathcal{J}^{\varrho+\gamma} |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(\varphi) \right. \\ &\quad \left. \left. + {}^H\mathcal{J}^{\varrho-\varsigma} |g(\sigma, y(\sigma)) - g(\sigma, \hat{y}(\sigma))|(T) \right] \right\} \\ &\leq \left[ {}^H\mathcal{J}^\varrho \left( f(\sigma) \frac{|y - \hat{y}|}{\Phi + |y - \hat{y}|} \right)(T) \right. \\ &\quad + \frac{(\log T)^{\varrho-1}}{\Lambda} \left( \omega {}^H\mathcal{J}^{\varrho+\gamma} \left( f(\sigma) \frac{|y - \hat{y}|}{\Phi + |y - \hat{y}|} \right)(\varphi) \right. \\ &\quad \left. \left. + {}^H\mathcal{J}^{\varrho-\varsigma} \left( f(\sigma) \frac{|y - \hat{y}|}{\Phi + |y - \hat{y}|} \right)(T) \right) \right] \\ &\leq \frac{\Phi \|y - \hat{y}\|}{\vartheta} \left[ {}^H\mathcal{J}^\varrho f(T) + \frac{1}{\Lambda} \left( \omega {}^H\mathcal{J}^\varrho f(\varphi) + {}^H\mathcal{J}^{\varrho-\varsigma} f(T) \right) \right] \\ &= \Phi (\|y - \hat{y}\|). \end{aligned}$$

This implies that,  $\|\mathcal{J}y - \mathcal{J}\hat{y}\| \leq \Phi (\|y - \hat{y}\|)$ . Therefore  $\mathcal{J}$  is a nonlinear contraction. Hence it follows by nonlinear contractions that the equation (1) has at most one solution on  $[1, T]$ .  $\square$

#### 4. Examples

**Example 4.1.** Consider the following fractional-order boundary value problem

$${}^H\mathcal{D}^{\frac{9}{4}}y(\tau) = \frac{|y(\tau)|}{1 + 2|y(\tau)|} \cdot \frac{e^{(\log \tau)^{\varrho-1}}}{(1 + \tau)^2} + \frac{1}{1 + (\log \tau)^{\varrho-1}}, \quad \tau \in [1, T], \tag{13}$$

subject to the integral boundary conditions

$$y(1) = 0, \quad y'(1) = 0, \quad {}^H\mathcal{D}^{\frac{4}{3}}y(T) = \omega {}^H\mathcal{J}^{\frac{3}{2}}y(\varphi). \tag{14}$$

Here,  $\varrho = \frac{9}{4}$ ,  $\varsigma = \frac{4}{3}$ ,  $\gamma = \frac{3}{2}$ ,  $\omega = \frac{1}{40}$ ,  $\varphi = \frac{5}{4}$ ,  $T = 2$ .

In addition, we find that

$$\begin{aligned} |g(\tau, y(\tau))| &= \frac{1}{1 + (\log \tau)^{\varrho-1}} + \frac{|y|}{1 + 2|y|} \frac{1}{(1 + \tau)^2} \quad \text{as} \\ |g(\tau, p_1(\tau)) - g(\tau, p_2(\tau))| &\leq \frac{1}{4} \|p_1 - p_2\|. \end{aligned}$$

With the above specifics, we find that  $\Lambda \cong 0.5930036058127669$ ,  $\Delta \cong 1.1969781392352639$ . Thus, the presumptions of Theorem 3.3 are satisfied. Hence, by Theorem 3.3, the boundary value problem (13)-(14) has at least one solution on  $[1, T]$ .

**Example 4.2.** Consider the following fractional-order boundary value problem

$${}^H\mathcal{D}^{\frac{8}{3}}y(\tau) = \frac{\tau}{1 + \tau} + \frac{|y(\tau)|}{1 + |y(\tau)|} \cdot \frac{1}{(2 + \tau)^2}, \quad \tau \in [1, T], \tag{15}$$

subject to the integral boundary conditions

$$y(1) = 0, \quad y'(1) = 0, \quad {}^H\mathcal{D}^{\frac{5}{3}}y(T) = \omega {}^H\mathcal{J}^{\frac{6}{5}}y(\varphi). \quad (16)$$

Here,  $\varrho = \frac{8}{3}$ ,  $\varsigma = \frac{5}{3}$ ,  $\gamma = \frac{6}{5}$ ,  $\omega = \frac{1}{50}$ ,  $\varphi = \frac{6}{5}$ ,  $T = 2$ .

In addition, we find that

$$|g(\tau, y(\tau))| = \frac{\tau}{1+\tau} + \frac{|y|}{1+|y|} \frac{1}{(2+\tau)^2} \quad \text{as}$$

$$|g(\tau, p_1(\tau)) - g(\tau, p_2(\tau))| \leq \frac{1}{9} \|p_1 - p_2\|.$$

With the above specifics, we find that  $\Lambda = 1.3274143884549592$  and  $\Delta = 0.4217023815634139$ . Thus,  $\mathcal{S}\Delta \cong 0.046855820173712655 < 1$ , the presumptions of Theorem 3.5 are satisfied. Hence, by Theorem 3.5, the boundary value problem (15)-(16) has at most one solution on  $[1, T]$ .

## References

- [1] B. Ahmad, and A. Alsaedi, and Alaa. Alshariff, Existence results for fractional-order differential equations with nonlocal multi-point-strip conditions involving Caputo derivative, Adv. Difference Equ. 2015:1 (2015), 348.
- [2] B. Ahmad, SK. Ntouyas, On three-point Hadamard-type fractional boundary value problems, Int. Electron. J. Pure. Appl. Math. 8:4 2014 31-42.
- [3] B. Ahmad, SK. Ntouyas, RP. Agarwal, A. Alsaedi, New results for boundary value problems of Hadamard-type fractional differential inclusions and integral boundary conditions, Bound. Value Probl. 2013:1 (2013), 275.
- [4] B. Ahmad, SK. Ntouyas, RP. Agarwal, A. Alsaedi, Existence results for sequential fractional integro-differential equations with nonlocal multi-point and strip conditions, Fract. Calc. Appl. Anal. 18:1 (2015), 261-280.
- [5] S. Ali, T. Abdeljawad, K. Shah, F. Jarad, M. Arif, Computation of iterative solutions along with stability analysis to a coupled system of fractional order differential equations, Adv. Difference Equ. 2019 (2019), 215.
- [6] Z. Ali, A. Zada, K. Shah, On Ulam's Stability for a Coupled Systems of Nonlinear Implicit Fractional Differential Equations, Bull. Malays. Math. Sci. Soc. 42:5 2019 2681-2699.
- [7] A. Alsaedi, SK. Ntouyas, B. Ahmad, A. Hobiny, Nonlinear Hadamard fractional differential equations with Hadamard type nonlocal non-conserved conditions, Adv. Difference Equ. 2015:1 (2015), 285.
- [8] Y. Bai, H. Kong, Existence of solutions for nonlinear Caputo-Hadamard fractional differential equations via the method of upper and lower solutions, J. Nonlinear Sci. Appl. 10:1 (2017), 5744-5752.
- [9] P. Duraisamy, T. Nandha Gopal, Existence and uniqueness of solutions for a coupled system of higher order fractional differential equations with integral boundary conditions, Discontinuity, Nonlinearity, and Complexity. 7:1 (2018) 1-14.
- [10] P. Duraisamy, A.R. Vidhya Kumar, T. Nandha Gopal, M. Subramanian, Influence of nonlocal discrete and integral boundary conditions involving Caputo derivative in boundary value problem, J.Phys.Conf.Ser. 1139:1 (2018) 012014.
- [11] Y. Y. Gambo, F. Jarad, D. Baleanu, T. Abdeljawad, On Caputo modification of the Hadamard fractional derivatives, Adv. Difference Equ. 2014:1 (2014), 10.
- [12] J. Hadamard, Essai sur l'etude des fonctions donnees par leur developpement de Taylor, Journal de Mathematiques Pures et Appliquees 8 (1892) 101-186.
- [13] F. Jarad, T. Abdeljawad, D. Baleanu, Caputo-type modification of the Hadamard fractional derivatives, Adv. Difference Equ. 2012:1 (2012), 142.
- [14] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and applications of fractional differential equations, Amsterdam, Boston, Elsevier, 2006.
- [15] A. Kilbas, M. Saigo, R. K. Saxena, Generalized Mittag-Leffler function and generalized fractional calculus operators, Adv. Difference Equ. 15:1 (2004) 31-49.
- [16] J. Klafter, S. C. Lim, R. Metzler, Fractional dynamics: Recent advances, World Scientific, 2012.
- [17] SK. Ntouyas, Sina. Etemad, On the existence of solutions for fractional differential inclusions with sum and integral boundary conditions, Appl. Math. Comput. 266:1 (2016) 235-246.
- [18] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego-Boston-New York-London-Tokyo-Toronto, 1999.
- [19] M. Qinghua, M. Chao, J. Wang, A Lyapunov-type inequality for a fractional differential equation with Hadamard derivative, J. Math. Inequal. 11:1 (2017) 135-141.
- [20] J. Sabatier, O. P. Agrawal, J. A. Tenreiro Machado, Advances in fractional calculus: theoretical developments and applications in physics and engineering, Springer, Netherlands, 2007.
- [21] B. Samet, H. Aydi, Lyapunov-type inequalities for an anti-periodic fractional boundary value problem involving  $\psi$ -Caputo fractional derivative, J. Inequal. Appl. 2018 (2018) 286.
- [22] B. Samet, H. Aydi, On Some Inequalities Involving Liouville's Caputo Fractional Derivatives and Applications to Special Means of Real Numbers, Mathematics. 6:10 (2018) 1-9.

- [23] Samina, K. Shah, R.A. Khan, D. Baleanu, Study of implicit type coupled system of non-integer order differential equations with antiperiodic boundary conditions, *Math. Methods Appl. Sci.* 42:6 2019 1-10.
- [24] K. Shah, H. Khalil, R.A. Khan, Investigation of positive solution to a coupled system of impulsive boundary value problems for nonlinear fractional order differential equations, *Chaos Solitons Fractals.* 77 (2015) 240-246.
- [25] M. Subramanian, T. Nandha Gopal, Solvability of Liouville-Caputo Fractional Integro-Differential Equations With Non-Local Generalized Fractional Integral Boundary Conditions, *Global Journal of Engineering Science and Researches.* 6:4 (2019) 142-154.
- [26] M. Subramanian, A.R. Vidhya Kumar, T. Nandha Gopal, Analysis of fractional boundary value problem with non-local integral strip boundary conditions, *Nonlinear Stud.* 26:2 (2019) 445-454.
- [27] M. Subramanian, A.R. Vidhya Kumar, T. Nandha Gopal, A strategic view on the consequences of classical integral sub-strips and coupled nonlocal multi-point boundary conditions on a combined Caputo fractional differential equation, *Proc. Jangjeon Math. Soc.* 22:3 (2019) 437-453.
- [28] M. Subramanian, A.R. Vidhya Kumar, T. Nandha Gopal, A Fundamental Approach on Non-integer Order Differential Equation Using Nonlocal Fractional Sub-Strips Boundary Conditions, *Discontinuity, Nonlinearity, and Complexity.* 8:2 (2019) 189-199.
- [29] M. Subramanian, A.R. Vidhya Kumar, T. Nandha Gopal, Influence of coupled nonlocal slit-strip conditions involving Caputo derivative in fractional boundary value problem, *Discontinuity, Nonlinearity, and Complexity.* 8:4 (2019) 429-445.
- [30] P. Thiramanus, SK. Ntouyas, J. Tariboon, Positive solutions for Hadamard fractional differential equations on infinite domain, *Adv. Difference Equ.* 2016:1 (2016), 83.
- [31] A.R. Vidhya Kumar, P. Duraisamy, T. Nandha Gopal, M. Subramanian , Analysis of fractional differential equation involving Caputo derivative with nonlocal discrete and multi-strip type boundary conditions, *J.Phys.Conf.Ser.* 1139:1 (2018) 012020.
- [32] G. Wang, K. Pei, R. P. Agrawal, L. Zhang, B. Ahmad, Nonlocal Hadamard fractional boundary value problem with Hadamard integral and discrete boundary conditions on a half-line, *J. Comput. Appl. Math.* 343 (2018) 230-239.
- [33] W. Yukunthorn, B. Ahmad, SK. Ntouyas, J. Tariboon, On Caputo-Hadamard type fractional impulsive hybrid systems with nonlinear fractional integral conditions, *Nonlinear Anal. Hybrid Syst.* 19 (2019) 77-92.