

Existence of travelling wave solutions in delayed reaction–diffusion systems with applications to diffusion–competition systems

Wan-Tong Li¹, Guo Lin¹ and Shigui Ruan²

¹ School of Mathematics and Statistics, Lanzhou University, Lanzhou, Gansu 730000, People's Republic of China

² Department of Mathematics, University of Miami, PO Box 249085, Coral Gables, FL 33124-4250, USA

Received 1 June 2005, in final form 4 April 2006

Published 15 May 2006

Online at stacks.iop.org/Non/19/1253

Recommended by B Eckhardt

Abstract

This paper is concerned with the existence of travelling wave solutions in a class of delayed reaction–diffusion systems without monotonicity, which concludes two-species diffusion–competition models with delays. Previous methods do not apply in solving these problems because the reaction terms do not satisfy either the so-called quasimonotonicity condition or non-quasimonotonicity condition. By using Schauder's fixed point theorem, a new cross-iteration scheme is given to establish the existence of travelling wave solutions. More precisely, by using such a new cross-iteration, we reduce the existence of travelling wave solutions to the existence of an admissible pair of upper and lower solutions which are easy to construct in practice. To illustrate our main results, we study the existence of travelling wave solutions in two delayed two-species diffusion–competition systems.

Mathematics Subject Classification: 35K57, 35R20, 92D25

1. Introduction

In recent years, great attention has been paid to the study of the existence of travelling waves in reaction–diffusion systems with delays. In a pioneering work, Schaaf [27] systematically studied two scalar reaction–diffusion *equations* with a single discrete delay for the so-called Huxley nonlinearity as well as Fisher nonlinearity by using the phase space analysis, the maximum principle for parabolic functional differential equations and the general theory for ordinary functional differential equations. For reaction–diffusion *systems* with quasimonotonicity (QM) and a single discrete delay, Zou and Wu [36] established the existence

of travelling wave fronts by first truncating the unbounded domain and then passing to a limit. Wu and Zou [33] further considered more general reaction–diffusion systems with a single delay of the form

$$\frac{\partial}{\partial t}u(x, t) = D \frac{\partial^2}{\partial x^2}u(x, t) + f(u_t(x)), \quad (1.1)$$

where $t \in \mathbb{R}$, $x \in \mathbb{R}$, $D = \text{diag}(d_1, \dots, d_n)$, $d_i > 0$, $i = 1, \dots, n$, $f \in C([-\tau, 0], \mathbb{R}^n)$ is continuous and satisfies $f(\hat{\mathbf{0}}) = f(\hat{\mathbf{K}}) = \mathbf{0}$; here \hat{u} denotes the constant vector function on $[-\tau, 0]$ taking the value u , and for any fixed $x \in \mathbb{R}$, $u_t(x) \in C([-\tau, 0], \mathbb{R}^n)$ is defined by $u_t(x) = u(t + \theta, x)$, $\theta \in [-\tau, 0]$. If the reaction term f satisfies either the QM condition

(QM) there exists a matrix $\beta = \text{diag}(\beta_1, \beta_2, \dots, \beta_n)$ with $\beta_i \geq 0$ such that

$$f(\Phi) - f(\Psi) + \beta(\Phi(0) - \Psi(0)) \geq 0$$

for $\Phi, \Psi \in C([-\tau, 0], \mathbb{R}^n)$ with $\mathbf{0} \leq \Psi(s) \leq \Phi(s) \leq \mathbf{K}$, $s \in [-\tau, 0]$

or the non-quasimonotonicity (QM*) condition

(QM*) there exists a matrix $\beta = \text{diag}(\beta_1, \beta_2, \dots, \beta_n)$ with $\beta_i \geq 0$ such that

$$f(\Phi) - f(\Psi) + \beta(\Phi(0) - \Psi(0)) \geq 0$$

for $\Phi, \Psi \in C([-\tau, 0], \mathbb{R}^n)$ with (i) $\mathbf{0} \leq \Psi(s) \leq \Phi(s) \leq \mathbf{K}$, $s \in [-\tau, 0]$ and (ii) $e^{\beta s}(\Phi(s) - \Psi(s))$ is non-decreasing in $s \in [-\tau, 0]$;

then some existence results are established for travelling wave fronts connecting the trivial equilibrium $\mathbf{0}$ and the non-trivial equilibrium \mathbf{K} , where the well-known monotone iteration techniques for elliptic systems with advanced arguments are used [20, 23]. The results are applicable not only to delayed scalar equations (Lan and Wu [19]) but also to delayed systems, such as delayed diffusion–cooperation systems (Huang and Zou [14]) and the delayed Belousov–Zhabotinskii model (Huang and Zou [15]). Following Wu and Zou [33], Ma [22] employed the Schauder’s fixed point theorem to an operator used in Wu and Zou [33] in a properly chosen subset in the Banach space $C(\mathbb{R}, \mathbb{R}^n)$ equipped with the so-called exponential decay norm. The subset is constructed in terms of a pair of upper–lower solutions, which is less restrictive than the upper–lower solutions required in [33]. This makes the search for the pair of upper–lower solutions slightly easier. Since Ma [22] only considered delayed systems with quasimonotone reaction terms, Huang and Zou [15] extended the results of Ma [22] to a class of delayed systems with QM* reaction terms. For related results on reaction–diffusion equations with non-local delays, we refer to Ashwin *et al* [1], Al-Omari and Gourly [2], Billingham [3], Li, Ruan and Wang [21], Wang, Li and Ruan [31] and references cited therein.

However, it is quite common that the reaction term in a model system arising from a practical problem may not satisfy either the QM condition or the QM* condition. Two typical and important examples are the two species competition systems [26, 32]:

$$\begin{cases} \frac{\partial}{\partial t}u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2}u_1(x, t) + r_1u_1(x, t)[1 - a_1u_1(x, t) - b_1u_2(x, t - \tau_1)], \\ \frac{\partial}{\partial t}u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2}u_2(x, t) + r_2u_2(x, t)[1 - b_2u_1(x, t - \tau_2) - a_2u_2(x, t)] \end{cases} \quad (1.2)$$

and

$$\begin{cases} \frac{\partial}{\partial t}u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2}u_1(x, t) + r_1u_1(x, t)[1 - a_1u_1(x, t - \tau_1) - b_1u_2(x, t - \tau_2)], \\ \frac{\partial}{\partial t}u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2}u_2(x, t) + r_2u_2(x, t)[1 - b_2u_1(x, t - \tau_3) - a_2u_2(x, t - \tau_4)]. \end{cases} \quad (1.3)$$

Thus, it is worthwhile to further explore this topic for systems without either QM or QM*, and this constitutes the purpose of this paper.

In order to focus on the mathematical ideas and for the sake of simplicity, we consider a reaction–diffusion system of two equations with discrete delays, that is,

$$\begin{cases} \frac{\partial}{\partial t} u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2} u_1(x, t) + f_1(u_1(x, t - \tau_{11}), u_2(x, t - \tau_{12})), \\ \frac{\partial}{\partial t} u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2} u_2(x, t) + f_2(u_1(x, t - \tau_{21}), u_2(x, t - \tau_{22})), \end{cases} \tag{1.4}$$

where $d_i > 0, \tau_{ij} \geq 0, f_i : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function, $x \in (-\infty, \infty)$, and

(A1) $f_i(0, 0) = f_i(k_1, k_2) = 0$ for $i = 1, 2$.

(A2) There exist two positive constants $L_1 > 0$ and $L_2 > 0$ such that

$$\begin{aligned} |f_1(\phi_1, \psi_1) - f_1(\phi_2, \psi_2)| &\leq L_1 \|\Phi - \Psi\|, \\ |f_2(\phi_1, \psi_1) - f_2(\phi_2, \psi_2)| &\leq L_2 \|\Phi - \Psi\| \end{aligned}$$

for $\Phi = (\phi_1, \psi_1), \Psi = (\phi_2, \psi_2) \in C([-\tau, 0], \mathbb{R}^2)$ with $0 \leq \phi_i(s), \psi_i(s) \leq M_i, s \in [-\tau, 0], M_i > k_i$ is positive constant, $i = 1, 2$.

Since the results of Huang and Zou [15], Ma [22] and Wu and Zou [33] do not apply to delayed reaction–diffusion systems (1.2) and (1.3), we must search for new techniques that can be applied to our delayed reaction–diffusion system (1.4), at least for (1.2) and (1.3). To overcome the difficulty, we propose two new conditions on the reaction terms, which are to be called the **weak QM condition** (WQM) and the **weak QM* condition** (WQM*), respectively:

(WQM) Two positive numbers exist $\beta_1 > 0, \beta_2 > 0$ such that

$$\begin{aligned} f_1(\phi_1(s), \psi_1(s)) - f_1(\phi_2(s), \psi_1(s)) + \beta_1[\phi_1(0) - \phi_2(0)] &\geq 0, \\ f_1(\phi_1(s), \psi_1(s)) - f_1(\phi_1(s), \psi_2(s)) &\leq 0, \\ f_2(\phi_1(s), \psi_1(s)) - f_2(\phi_1(s), \psi_2(s)) + \beta_2[\psi_1(0) - \psi_2(0)] &\geq 0, \\ f_2(\phi_1(s), \psi_1(s)) - f_2(\phi_2(s), \psi_1(s)) &\leq 0 \end{aligned}$$

for $\phi_1(s), \phi_2(s), \psi_1(s), \psi_2(s) \in C([-\tau, 0], \mathbb{R})$ with

$$0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1, 0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2, s \in [-\tau, 0].$$

(WQM*) Two positive numbers exist $\beta_1 > 0, \beta_2 > 0$ such that

$$\begin{aligned} f_1(\phi_1(s), \psi_1(s)) - f_1(\phi_2(s), \psi_1(s)) + \beta_1[\phi_1(0) - \phi_2(0)] &\geq 0, \\ f_1(\phi_1(s), \psi_1(s)) - f_1(\phi_1(s), \psi_2(s)) &\leq 0, \\ f_2(\phi_1(s), \psi_1(s)) - f_2(\phi_1(s), \psi_2(s)) + \beta_2[\psi_1(0) - \psi_2(0)] &\geq 0, \\ f_2(\phi_1(s), \psi_1(s)) - f_2(\phi_2(s), \psi_1(s)) &\leq 0 \end{aligned}$$

for $\phi_1(s), \phi_2(s), \psi_1(s), \psi_2(s) \in C([-\tau, 0], \mathbb{R})$ with

$$\begin{cases} \text{(i)} & 0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1, 0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2, s \in [-\tau, 0]; \\ \text{(ii)} & e^{\beta_1 s} [\phi_1(s) - \phi_2(s)] \text{ and } e^{\beta_2 s} [\psi_1(s) - \psi_2(s)] \text{ are non-decreasing in } s \in [-\tau, 0]. \end{cases}$$

Since the nonlinear functions f_1 and f_2 in (1.4) have different monotonicity with respect to the first and second arguments in the first and second equations, respectively, following Pao [23] and Ye and Li [34], we introduce definitions of the upper and lower solutions, and a new cross-iteration scheme, which are different from those defined in Huang and Zou [15], Ma [22] and Wu and Zou [33]. By using such a scheme, we will construct a subset in the Banach space $C(\mathbb{R}, \mathbb{R}^2)$ equipped with the exponential decay norm and reduce the existence of travelling wave solutions to the existence of an admissible pair of upper and lower solutions which are easy to construct in practice. As applications, we shall show that system (1.2) satisfies the condition (WQM) while system (1.3) satisfies (WQM*) and establish the existence of travelling wave solutions in both models.

We remark that if $\tau_1 = \tau_2 = 0$, then (1.2) reduces to the following Lotka–Volterra diffusion–competition system:

$$\begin{cases} \frac{\partial}{\partial t} u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2} u_1(x, t) + r_1 u_1(x, t) [1 - a_1 u_1(x, t) - b_1 u_2(x, t)], \\ \frac{\partial}{\partial t} u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2} u_2(x, t) + r_2 u_2(x, t) [1 - b_2 u_1(x, t) - a_2 u_2(x, t)], \end{cases} \quad (1.5)$$

where u_1 and u_2 represent the densities of two competitive species in a one-dimensional habitat having infinite length and $d_1, d_2, r_1, r_2, a_1, a_2, b_1, b_2$ are some positive constants. The existence of travelling wave solutions of (1.5) has been extensively studied in the literature (Conley and Gardner [5], Gardner [8], Gourley and Ruan [9], Kanel and Zhou [16], Kan-on [17], Tang and Fife [29] and van Vuuren [30]). This model has a trivial (no species) equilibrium $E_0 = (0, 0)$, two semitrivial (one species only) spatially homogeneous equilibria [9]

$$E_1 = \left(\frac{1}{a_1}, 0 \right), E_2 = \left(0, \frac{1}{a_2} \right)$$

and a positive (two coexisting species) spatially homogeneous equilibrium

$$E^* = \left(\frac{a_2 - b_1}{a_1 a_2 - b_1 b_2}, \frac{a_1 - b_2}{a_1 a_2 - b_1 b_2} \right)$$

provided that $a_1 a_2 \neq b_1 b_2$ and either (i) $a_2 > b_1$ and $a_1 > b_2$ or (ii) $a_2 < b_1$ and $a_1 < b_2$. By using phase space analysis for the ordinary differential equations, Tang and Fife [29] and van Vuuren [30] showed that (1.5) has travelling front solutions connecting the equilibria E_0 and E^* . Kanel and Zhou [16] further proved that (1.5) has travelling front solutions connecting the equilibria E_1 and E^* . Conley and Gardner [5] and Gardner [8] showed that (1.5) has travelling front solutions connecting the equilibria E_1 and E_2 , where Conley index and degree theory methods have been developed. Other related results can be found in Gourley and Ruan [9], Hosono [11, 12], Kan-on [17], etc. We shall establish the existence of travelling waves in system (1.4), thus in systems (1.2) and (1.3), that connect the trivial equilibrium E_0 and the positive equilibrium E^* . Thus our results can be regarded as a generalization of the results of Tang and Fife [29] and van Vuuren [30] to the diffusion–competition models with delays.

This paper is organized as following. Section 2 is devoted to some preliminary discussions. In section 3, we establish a new cross-iteration scheme and apply it to obtain the existence of travelling wave solutions if the nonlinear reaction term satisfies the condition (WQM). In section 4, we use the non-standard ordering of the profile set and prove that similar results hold if the nonlinear reaction term satisfies the condition (WQM*). In section 5, we apply our main results to the diffusion–competition systems (1.2) and (1.3) and prove the existence of travelling wave solutions. The paper ends with a discussion in section 6.

2. Preliminaries

In this paper, we use the usual notations for the standard ordering in \mathbb{R}^2 . That is, for $u = (u_1, u_2)$ and $v = (v_1, v_2)$, we denote $u \leq v$ if $u_i \leq v_i$, $i = 1, 2$, and $u < v$ if $u \leq v$ but $u \neq v$. In particular, we denote $u \ll v$ if $u \leq v$ but $u_i \neq v_i$, $i = 1, 2$. If $u \leq v$, we also denote $(u, v) = \{w \in \mathbb{R}^2, u < w \leq v\}$, $[u, v) = \{w \in \mathbb{R}^2, u \leq w < v\}$, and $[u, v] = \{w \in \mathbb{R}^2, u \leq w \leq v\}$. Let $|\cdot|$ denote the Euclidean norm in \mathbb{R}^2 and $\|\cdot\|$ denote the supremum norm in $C([-\tau, 0], \mathbb{R}^2)$.

A *travelling wave solution* of (1.4) is a special translation invariant solution of the form $u_1(x, t) = \phi(x + ct)$, $u_2(x, t) = \psi(x + ct)$, where $(\phi, \psi) \in C^2(\mathbb{R}, \mathbb{R}^2)$ are the profiles of the wave that propagates through the one-dimensional spatial domain at a constant velocity

$c > 0$. Substituting $u_1(x, t) = \phi(x + ct)$, $u_2(x, t) = \psi(x + ct)$ into (1.4) and denoting $\phi_t(s) = \phi(t + s)$, $\psi_t(s) = \psi(t + s)$ and $x + ct$ by t , we find that (1.4) has a pair of travelling wave solutions if and only if the following wave equations

$$\begin{cases} d_1\phi'' - c\phi' + f_1^c(\phi_t, \psi_t) = 0, \\ d_2\psi'' - c\psi' + f_2^c(\phi_t, \psi_t) = 0 \end{cases} \tag{2.1}$$

with asymptotic boundary conditions

$$\lim_{t \rightarrow -\infty} \phi(t) = \phi_-, \quad \lim_{t \rightarrow +\infty} \phi(t) = \phi_+, \quad \lim_{t \rightarrow -\infty} \psi(t) = \psi_-, \quad \lim_{t \rightarrow +\infty} \psi(t) = \psi_+ \tag{2.2}$$

have a pair of solutions $(\phi(t), \psi(t))$ on \mathbb{R} , where $f_i^c(\phi, \psi) : C([-\tau, 0], \mathbb{R}) \rightarrow \mathbb{R}$, $i = 1, 2$, is given by

$$f_i^c(\phi, \psi) = f_i(\phi^c, \psi^c), \phi^c(s) = \phi(cs), \psi^c(s) = \psi(cs), s \in [-\tau, 0],$$

where $\tau = \max_{1 \leq i, j \leq 2} \{\tau_{ij}\}$, (ϕ_-, ψ_-) and (ϕ_+, ψ_+) are two equilibria of (2.1).

Without loss of generality, we let $\phi_- = 0$, $\phi_+ = k_1 > 0$, $\psi_- = 0$ and $\psi_+ = k_2 > 0$. Then boundary conditions (2.2) become

$$\lim_{t \rightarrow -\infty} \phi(t) = 0, \quad \lim_{t \rightarrow +\infty} \phi(t) = k_1, \quad \lim_{t \rightarrow -\infty} \psi(t) = 0, \quad \lim_{t \rightarrow +\infty} \psi(t) = k_2. \tag{2.3}$$

Let

$$C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) = \{(\phi, \psi) \in C(\mathbb{R}, \mathbb{R}^2) : 0 \leq \phi(s) \leq M_1, 0 \leq \psi(s) \leq M_2, s \in \mathbb{R}\}.$$

Define the operator $H = (H_1, H_2) : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ by

$$\begin{cases} H_1(\phi, \psi)(t) = f_1^c(\phi_t, \psi_t) + \beta_1\phi(t), \\ H_2(\phi, \psi)(t) = f_2^c(\phi_t, \psi_t) + \beta_2\psi(t). \end{cases} \tag{2.4}$$

Then (2.1) can be rewritten as following:

$$\begin{cases} d_1\phi''(t) - c\phi'(t) - \beta_1\phi(t) + H_1(\phi, \psi)(t) = 0, \\ d_2\psi''(t) - c\psi'(t) - \beta_2\psi(t) + H_2(\phi, \psi)(t) = 0. \end{cases} \tag{2.5}$$

Let

$$\begin{aligned} \lambda_1 &= \frac{c - \sqrt{c^2 + 4\beta_1 d_1}}{2d_1}, & \lambda_2 &= \frac{c + \sqrt{c^2 + 4\beta_1 d_1}}{2d_1}, \\ \lambda_3 &= \frac{c - \sqrt{c^2 + 4\beta_2 d_2}}{2d_2}, & \lambda_4 &= \frac{c + \sqrt{c^2 + 4\beta_2 d_2}}{2d_2}. \end{aligned}$$

Then

$$\lambda_1 < 0 < \lambda_2, \lambda_3 < 0 < \lambda_4,$$

$$d_1\lambda_i^2 - c\lambda_i - \beta_1 = 0, i = 1, 2 \text{ and } d_2\lambda_i^2 - c\lambda_i - \beta_2 = 0, i = 3, 4.$$

Define the operator $F = (F_1, F_2) : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ by

$$\begin{cases} F_1(\phi, \psi)(t) = \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} H_1(\phi, \psi)(s) ds + \int_t^{\infty} e^{\lambda_2(t-s)} H_1(\phi, \psi)(s) ds \right], \\ F_2(\phi, \psi)(t) = \frac{1}{d_2(\lambda_4 - \lambda_3)} \left[\int_{-\infty}^t e^{\lambda_3(t-s)} H_2(\phi, \psi)(s) ds + \int_t^{\infty} e^{\lambda_4(t-s)} H_2(\phi, \psi)(s) ds \right]. \end{cases} \tag{2.6}$$

We can see that the operator F is well defined and for any $(\phi, \psi) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$,

$$\begin{cases} d_1(F_1(\phi, \psi))''(t) - c(F_1(\phi, \psi))'(t) - \beta_1 F_1(\phi, \psi)(t) + H_1(\phi, \psi)(t) = 0, \\ d_2(F_2(\phi, \psi))''(t) - c(F_2(\phi, \psi))'(t) - \beta_2 F_2(\phi, \psi)(t) + H_2(\phi, \psi)(t) = 0. \end{cases} \tag{2.7}$$

Thus, a fixed point of F is a solution of (2.5), which is a travelling wave solution of (1.4) connecting $\mathbf{0} = (0, 0)$ and $\mathbf{K} = (k_1, k_2)$ if it satisfies (2.3).

In the following, we introduce the exponential decay norm. Let $\mu > 0$ such that $\mu < \min\{-\lambda_1, \lambda_2, -\lambda_3, \lambda_4\}$. Define

$$B_\mu(\mathbb{R}, \mathbb{R}^2) = \{\Phi \in C(\mathbb{R}, \mathbb{R}^2) : \sup_{t \in \mathbb{R}} |\Phi(t)| e^{-\mu|t|} < \infty\}$$

and

$$|\Phi|_\mu = \sup_{t \in \mathbb{R}} |\Phi(t)| e^{-\mu|t|}.$$

Then it is easy to check that $(B_\mu(\mathbb{R}, \mathbb{R}^2), |\cdot|_\mu)$ is a Banach space.

3. The case (WQM)

In this section, we consider the existence of travelling wave solutions of (2.1) when the delayed reaction terms f_1 and f_2 satisfy the condition (WQM).

We start our cross-iteration with a pair of upper and lower solutions of (2.1) defined as follows.

Definition 3.1. A pair of twice continuously differentiable functions $\bar{\Phi} = (\bar{\phi}, \bar{\psi})$ and $\underline{\Phi} = (\underline{\phi}, \underline{\psi}) \in C(\mathbb{R}, \mathbb{R}^2)$ are called an **upper** and a **lower solution** of (2.1), respectively, if $\bar{\Phi}$ and $\underline{\Phi}$ satisfy

$$\begin{cases} d_1 \bar{\phi}''(t) - c \bar{\phi}'(t) + f_1^c(\bar{\phi}_t, \bar{\psi}_t) \leq 0 \text{ on } \mathbb{R}, \\ d_2 \bar{\psi}''(t) - c \bar{\psi}'(t) + f_2^c(\bar{\phi}_t, \bar{\psi}_t) \leq 0 \text{ on } \mathbb{R}, \end{cases} \quad (3.1)$$

and

$$\begin{cases} d_1 \underline{\phi}''(t) - c \underline{\phi}'(t) + f_1^c(\underline{\phi}_t, \underline{\psi}_t) \geq 0 \text{ on } \mathbb{R}, \\ d_2 \underline{\psi}''(t) - c \underline{\psi}'(t) + f_2^c(\underline{\phi}_t, \underline{\psi}_t) \geq 0 \text{ on } \mathbb{R}. \end{cases} \quad (3.2)$$

In what follows, we assume that an upper solution $\bar{\Phi} = (\bar{\phi}, \bar{\psi})$ and a lower solution $\underline{\Phi} = (\underline{\phi}, \underline{\psi})$ of (2.1) are given so that

$$(P1) \quad \mathbf{0} \leq (\underline{\phi}, \underline{\psi}) \leq (\bar{\phi}, \bar{\psi}) \leq \mathbf{M} = (M_1, M_2);$$

$$(P2) \quad \lim_{t \rightarrow -\infty} (\bar{\phi}, \bar{\psi}) = \mathbf{0}, \quad \lim_{t \rightarrow +\infty} (\underline{\phi}, \underline{\psi}) = \lim_{t \rightarrow +\infty} (\bar{\phi}, \bar{\psi}) = \mathbf{K} = (k_1, k_2).$$

For the operator $H = (H_1, H_2)$ defined in section 2, we have the following result.

Lemma 3.2. Assume that (WQM) holds. Then

$$H_1(\phi_2, \psi_1)(t) \leq H_1(\phi_1, \psi_2)(t), \quad H_2(\phi_1, \psi_2)(t) \leq H_2(\phi_2, \psi_1)(t)$$

for $(\phi_i, \psi_i) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ with $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1$, $0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2$, $s \in \mathbb{R}$.

Proof. By (WQM), direct calculation shows that

$$\begin{aligned} & H_1(\phi_1, \psi_2)(t) - H_1(\phi_2, \psi_1)(t) \\ &= H_1(\phi_1, \psi_2)(t) - H_1(\phi_2, \psi_2)(t) + H_1(\phi_2, \psi_2)(t) - H_1(\phi_2, \psi_1)(t) \\ &\geq H_1(\phi_1, \psi_2)(t) - H_1(\phi_2, \psi_2)(t) \\ &= f_1^c(\phi_{1t}, \psi_{1t}) - f_1^c(\phi_{2t}, \psi_{1t}) + \beta_1(\phi_1(t) - \phi_2(t)) \\ &\geq 0, t \in \mathbb{R}. \end{aligned}$$

The inequality for H_2 can be established similarly. The proof is complete. \square

As a direct consequence of lemma 3.2, the following lemma is true.

Lemma 3.3. *Assume that (WQM) holds. Then*

$$F_1(\phi_2, \psi_1)(t) \leq F_1(\phi_1, \psi_2)(t), \quad F_2(\phi_1, \psi_2)(t) \leq F_2(\phi_2, \psi_1)(t)$$

for $(\phi_i, \psi_i) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ with $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1, 0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2, s \in \mathbb{R}$.

Now, we assume that there exists an upper solution $\bar{\Phi}(t) = (\bar{\phi}, \bar{\psi})$ and a lower solution $\underline{\Phi} = (\underline{\phi}, \underline{\psi})$ of (2.1) satisfying (P1) and (P2).

Define the following profile set.

$$\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) = \{(\phi, \psi) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2), \underline{\phi}(t) \leq \phi(t) \leq \bar{\phi}(t), \underline{\psi}(t) \leq \psi(t) \leq \bar{\psi}(t), t \in \mathbb{R}\}.$$

Obviously, $\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is non-empty.

Lemma 3.4. *Assume that (A2) holds. Then*

$$F = (F_1, F_2) : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$$

is continuous with respect to the norm $|\cdot|_\mu$ in $B_\mu(\mathbb{R}, \mathbb{R}^2)$.

Proof. We first prove that $H_1 : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_\mu$. If $\Phi = (\phi_1, \psi_1), \Psi = (\phi_2, \psi_2) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ satisfy

$$|\Phi - \Psi|_\mu = \sup_{t \in \mathbb{R}} |\Phi(t) - \Psi(t)|e^{-\mu|t|} < \delta,$$

then

$$\begin{aligned} & |H_1(\phi_1, \psi_1)(t) - H_1(\phi_2, \psi_2)(t)|e^{-\mu|t|} \\ &= |f_1^c(\phi_{1t}, \psi_{1t})(t) - f_1^c(\phi_{2t}, \psi_{2t})(t) + \beta_1(\phi_1(t) - \phi_2(t))|e^{-\mu|t|} \\ &\leq L_1 \|\Phi_t - \Psi_t\|_{C([-ct, 0], \mathbb{R}^2)} e^{-\mu|t|} + \beta_1 |\phi_1 - \phi_2|_\mu \\ &= L_1 \sup_{s \in [-ct, 0]} |\Phi(t+s) - \Psi(t+s)|e^{-\mu|t|} + \beta_1 |\phi_1 - \phi_2|_\mu \\ &\leq L_1 \sup_{\theta \in \mathbb{R}} |\Phi(\theta) - \Psi(\theta)|e^{-\mu|\theta|} e^{\mu c \tau} + \beta_1 |\Phi - \Psi|_\mu \\ &\leq (L_1 e^{\mu c \tau} + \beta_1) |\Phi - \Psi|_\mu. \end{aligned}$$

For any fixed $\varepsilon > 0$, let $\delta < \varepsilon / (L_1 e^{\mu c \tau} + \beta_1)$. If $\Phi = (\phi_1, \psi_1), \Psi = (\phi_2, \psi_2) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ satisfy $|\Phi - \Psi|_\mu < \delta$, then

$$\begin{aligned} & |H_1(\phi_1, \psi_1)(t) - H_1(\phi_2, \psi_2)(t)|e^{-\mu|t|} \\ &\leq (L_1 e^{\mu c \tau} + \beta_1) |\Phi - \Psi|_\mu < \varepsilon. \end{aligned}$$

Therefore, $|H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_\mu \leq \varepsilon$. That is, $H_1 : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_\mu$.

Now, we show that $F_1 : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_\mu$.

For $t \geq 0$, we find

$$\begin{aligned}
& |F_1(\phi_1, \psi_1)(t) - F_1(\phi_2, \psi_2)(t)| \\
& \leq \left| \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} |H_1(\phi_1, \psi_1)(s) - H_1(\phi_2, \psi_2)(s)| \, ds \right. \right. \\
& \quad \left. \left. + \int_t^{\infty} e^{\lambda_2(t-s)} |H_1(\phi_1, \psi_1)(s) - H_1(\phi_2, \psi_2)(s)| \, ds \right] \right| \\
& = \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)+\mu|s|} |H_1(\phi_1, \psi_1)(s) - H_1(\phi_2, \psi_2)(s)| e^{-\mu|s|} \, ds \right. \\
& \quad \left. + \int_t^{\infty} e^{\lambda_2(t-s)+\mu|s|} |H_1(\phi_1, \psi_1)(s) - H_1(\phi_2, \psi_2)(s)| e^{-\mu|s|} \, ds \right] \\
& \leq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_0^t e^{\lambda_1(t-s)+\mu s} \, ds + \int_{-\infty}^0 e^{\lambda_1(t-s)-\mu s} \, ds \right. \\
& \quad \left. + \int_t^{\infty} e^{\lambda_2(t-s)+\mu s} \, ds \right] |H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_{\mu} \\
& = \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_2 - \lambda_1}{(\mu - \lambda_1)(\lambda_2 - \mu)} e^{\mu t} + \frac{2\mu}{\lambda_1^2 - \mu^2} e^{\lambda_1 t} \right] \\
& \quad \times |H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_{\mu}.
\end{aligned}$$

Hence, we have

$$\begin{aligned}
& |F_1(\phi_1, \psi_1)(t) - F_1(\phi_2, \psi_2)(t)| e^{-\mu|t|} \\
& \leq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_2 - \lambda_1}{(\mu - \lambda_1)(\lambda_2 - \mu)} + \frac{2\mu}{\lambda_1^2 - \mu^2} e^{(\lambda_1 - \mu)t} \right] \\
& \quad \times |H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_{\mu} \\
& \leq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_2 - \lambda_1}{(\mu - \lambda_1)(\lambda_2 - \mu)} + \frac{2\mu}{\lambda_1^2 - \mu^2} \right] \\
& \quad \times |H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_{\mu}. \tag{3.3}
\end{aligned}$$

Similarly, for $t \leq 0$, we have

$$\begin{aligned}
& |F_1(\phi_1, \psi_1)(t) - F_1(\phi_2, \psi_2)(t)| e^{-\mu|t|} \\
& \leq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_2 - \lambda_1}{-(\mu + \lambda_1)(\lambda_2 + \mu)} + \frac{2\mu}{\lambda_2^2 - \mu^2} \right] |H_1(\phi_1, \psi_1) - H_1(\phi_2, \psi_2)|_{\mu}. \tag{3.4}
\end{aligned}$$

Thus, it follows from (3.3) and (3.4) that $F_1 : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_{\mu}$ in $B_{\mu}(\mathbb{R}, \mathbb{R}^2)$.

By using a similar argument as above, we can also prove that $F_2 : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_{\mu}$ in $B_{\mu}(\mathbb{R}, \mathbb{R}^2)$. This completes the proof. \square

Lemma 3.5. Assume that (WQM) holds. Then

$$F : \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \subset \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]).$$

Proof. For any $(\phi, \psi) \in \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$, by lemma 3.3, it is easy to see that

$$\begin{cases} F_1(\underline{\phi}, \bar{\psi}) \leq F_1(\phi, \psi) \leq F_1(\bar{\phi}, \underline{\psi}), \\ F_2(\bar{\phi}, \underline{\psi}) \leq F_2(\phi, \psi) \leq F_2(\underline{\phi}, \bar{\psi}). \end{cases}$$

Now, we only need to prove

$$\begin{cases} \underline{\phi} \leq F_1(\underline{\phi}, \bar{\psi}) \leq F_1(\bar{\phi}, \underline{\psi}) \leq \bar{\phi}, \\ \underline{\psi} \leq F_2(\bar{\phi}, \underline{\psi}) \leq F_2(\underline{\phi}, \bar{\psi}) \leq \bar{\psi}. \end{cases} \tag{3.5}$$

According to the definition of the operator F and the lower solution, we have

$$\begin{aligned} F_1(\underline{\phi}, \bar{\psi})(t) &= \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_{-\infty}^t e^{\lambda_2(t-s)} \right] H_1(\underline{\phi}, \bar{\psi})(s) ds \\ &\geq \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_{-\infty}^t e^{\lambda_2(t-s)} \right] (\beta_1 \underline{\phi}(s) + c \underline{\phi}'(s) - d_1 \underline{\phi}''(s)) ds \\ &= \underline{\phi}(t), t \in \mathbb{R}. \end{aligned}$$

In a similar way, we can prove that (3.5) holds for $t \in \mathbb{R}$. The proof is complete. □

Lemma 3.6. *Assume that (A2) and (WQM) hold. Then*

$$F : \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \rightarrow \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$$

is compact.

Proof. We first establish an estimate for F . For any $(\phi, \psi) \in \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$, we have

$$\begin{aligned} (F_1(\phi, \psi))'(t) &= \frac{\lambda_1 e^{\lambda_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_{-\infty}^t e^{-\lambda_1 s} H_1(\phi, \psi)(s) ds \\ &\quad + \frac{\lambda_2 e^{\lambda_2 t}}{d_1(\lambda_2 - \lambda_1)} \int_t^{\infty} e^{-\lambda_2 s} H_1(\phi, \psi)(s) ds. \end{aligned}$$

So

$$\begin{aligned} |(F_1(\phi, \psi))'|_{\mu} &\leq \sup_{t \in \mathbb{R}} \left[e^{-\mu|t|} \frac{|\lambda_1| e^{\lambda_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_{-\infty}^t e^{-\lambda_1 s} H_1(\phi, \psi)(s) ds \right. \\ &\quad \left. + e^{-\mu|t|} \frac{\lambda_2 e^{\lambda_2 t}}{d_1(\lambda_2 - \lambda_1)} \int_t^{\infty} e^{-\lambda_2 s} H_1(\phi, \psi)(s) ds \right] \\ &\leq \frac{|\lambda_1|}{d_1(\lambda_2 - \lambda_1)} \sup_{t \in \mathbb{R}} e^{\lambda_1 t - \mu|t|} \int_{-\infty}^t e^{-\lambda_1 s} e^{\mu|s|} e^{-\mu|s|} H_1(\phi, \psi)(s) ds \\ &\quad + \frac{\lambda_2}{d_1(\lambda_2 - \lambda_1)} \sup_{t \in \mathbb{R}} e^{\lambda_2 t - \mu|t|} \int_t^{\infty} e^{-\lambda_2 s} e^{\mu|s|} e^{-\mu|s|} H_1(\phi, \psi)(s) ds \\ &\leq \frac{|\lambda_1|}{d_1(\lambda_2 - \lambda_1)} |H_1(\phi, \psi)|_{\mu} \sup_{t \in \mathbb{R}} e^{\lambda_1 t - \mu|t|} \int_{-\infty}^t e^{-\lambda_1 s + \mu|s|} ds \\ &\quad + \frac{\lambda_2}{d_1(\lambda_2 - \lambda_1)} |H_1(\phi, \psi)|_{\mu} \sup_{t \in \mathbb{R}} e^{\lambda_2 t - \mu|t|} \int_t^{\infty} e^{-\lambda_2 s + \mu|s|} ds. \end{aligned}$$

Therefore, for $t > 0$, we have

$$\begin{aligned} |(F_1(\phi, \psi))'|_{\mu} &\leq \frac{|\lambda_1|}{d_1(\lambda_2 - \lambda_1)(-\lambda_1 - \mu)} |H_1(\phi, \psi)|_{\mu} \\ &\quad + \frac{\lambda_2}{d_1(\lambda_2 - \lambda_1)(\lambda_2 - \mu)} |H_1(\phi, \psi)|_{\mu} \\ &= \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_1}{(\lambda_1 + \mu)} + \frac{\lambda_2}{(\lambda_2 - \mu)} \right] |H_1(\phi, \psi)|_{\mu}. \end{aligned}$$

Similarly, for $t < 0$, we have

$$\begin{aligned} |(F_1(\phi, \psi))'|_\mu &\leq \frac{|\lambda_1|}{d_1(\lambda_2 - \lambda_1)(-\lambda_1 - \mu)} |H_1(\phi, \psi)|_\mu \\ &\quad + \frac{\lambda_2}{d_1(\lambda_2 - \lambda_1)} \left[\left| \frac{1}{(\lambda_2 - \mu)} - \frac{1}{(\lambda_2 + \mu)} \right| + \frac{1}{(\lambda_2 + \mu)} \right] |H_1(\phi, \psi)|_\mu \\ &\leq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\frac{\lambda_1}{(\lambda_1 + \mu)} + \frac{\lambda_2}{(\lambda_2 - \mu)} \right] |H_1(\phi, \psi)|_\mu. \end{aligned}$$

Since $H : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_\mu$ and the set $\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is uniformly bounded, there exists a constant C_1 such that $|(F_1(\phi, \psi))'|_\mu \leq C_1$. In a similar way, there exists a constant C_2 such that $|(F_2(\phi, \psi))'|_\mu \leq C_2$. Hence F is equicontinuous on $\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ and $F\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is uniformly bounded.

We next prove that $F : \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \rightarrow \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is compact.

Define $F^n(\phi, \psi)$ by

$$F^n(\phi, \psi)(t) = \begin{cases} F(\phi, \psi)(t), & t \in [-n, n]; \\ F(\phi, \psi)(n), & t \in (n, \infty); \\ F(\phi, \psi)(-n), & t \in (-\infty, -n). \end{cases}$$

Then, for any $n \geq 1$, F^n is equicontinuous and uniformly bounded. Ascoli–Arzela lemma implies that F^n is compact.

Since $\{F^n(\phi, \psi)\}_0^\infty$ is a compact series, and

$$\begin{aligned} &\sup_{t \in \mathbb{R}} |F^n(\phi, \psi)(t) - F(\phi, \psi)(t)|e^{-\mu|t|} \\ &= \sup_{t \in (-\infty, -n) \cup (n, \infty)} |F^n(\phi, \psi)(t) - F(\phi, \psi)(t)|e^{-\mu|t|} \\ &\leq 2C_0e^{-\mu n} \rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

where C_0 is a positive constant such that

$$|(\phi, \psi)| \leq C_0 \text{ for any } (\phi, \psi) \in \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]),$$

by proposition 2.1 in [35], we know that $\{F^n\}_0^\infty$ converges to F in $\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ with respect to the norm $|\cdot|_\mu$. Therefore F is compact. \square

Theorem 3.7. Assume that (A1), (A2) and (WQM) hold. If (2.1) has an upper solution $(\bar{\phi}, \bar{\psi}) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ and a lower solution $(\underline{\phi}, \underline{\psi}) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ such that (P1) and (P2) are satisfied, then (2.1) has a travelling wave solution satisfying (2.3).

Proof. From lemmas 3.4–3.6, we know that $F\Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \subset \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ and F is compact. By Schauder’s fixed point theorem there exists a fixed point $(\phi^*, \psi^*) \in \Gamma([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$, which is a solution of (2.1).

In order to prove that this solution is a travelling wave solution, we need to verify the asymptotic boundary condition (2.3).

By (P2) and the fact that

$$0 \leq (\underline{\phi}(t), \underline{\psi}(t)) \leq (\phi^*(t), \psi^*(t)) \leq (\bar{\phi}(t), \bar{\psi}(t)) \leq (M_1, M_2),$$

we see that

$$\lim_{t \rightarrow -\infty} (\phi^*(t), \psi^*(t)) = (0, 0) \text{ and } \lim_{t \rightarrow \infty} (\phi^*(t), \psi^*(t)) = (k_1, k_2).$$

Therefore, the fixed point $(\phi^*(t), \psi^*(t))$ satisfies the asymptotic boundary condition (2.3). The proof is complete. \square

From theorem 3.7 we can see that the existence of solutions of (2.1) and (2.3) is reduced to the existence of an admissible pair of upper and lower solutions. However, it is difficult to construct such a pair of upper and lower solutions in practice because they are required to be twice continuously differentiable on \mathbb{R} .

In order to overcome the difficulty, we introduce the following weaker definitions of upper and lower solutions of (2.1) than definition 3.1. More precisely, we do not require that the upper and lower solutions are twice continuously differentiable on \mathbb{R} and they are easy to construct in practice. See examples 5.1 and 5.5 in section 5.

Definition 3.8. A pair of continuous functions $\bar{\Phi} = (\bar{\phi}, \bar{\psi})$, $\underline{\Phi} = (\underline{\phi}, \underline{\psi}) \in C(\mathbb{R}, \mathbb{R}^2)$ is called a **weak upper solution** and a **weak lower solution** of (2.1), respectively, if constants $T_i, i = 1, \dots, m$ exist, such that $\bar{\Phi}$ and $\underline{\Phi}$ are twice continuously differentiable in $\mathbb{R} \setminus \{T_i : i = 1, \dots, m\}$ and satisfy

$$\begin{cases} d_1 \bar{\phi}''(t) - c \bar{\phi}'(t) + f_1^c(\bar{\phi}_t, \bar{\psi}_t) \leq 0, t \in \mathbb{R} \setminus \{T_i : i = 1, \dots, m\}, \\ d_2 \bar{\psi}''(t) - c \bar{\psi}'(t) + f_2^c(\bar{\phi}_t, \bar{\psi}_t) \leq 0, t \in \mathbb{R} \setminus \{T_i : i = 1, \dots, m\}, \end{cases} \tag{3.6}$$

and

$$\begin{cases} d_1 \underline{\phi}''(t) - c \underline{\phi}'(t) + f_1^c(\underline{\phi}_t, \underline{\psi}_t) \geq 0, t \in \mathbb{R} \setminus \{T_i : i = 1, \dots, m\}, \\ d_2 \underline{\psi}''(t) - c \underline{\psi}'(t) + f_2^c(\underline{\phi}_t, \underline{\psi}_t) \geq 0, t \in \mathbb{R} \setminus \{T_i : i = 1, \dots, m\}. \end{cases} \tag{3.7}$$

Lemma 3.9. Assume that (WQM) holds. If $\bar{\Phi}, \underline{\Phi} \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ are a weak upper solution and a weak lower solution of (2.1), respectively, satisfying (P1), (P2) and

$$\begin{cases} \bar{\phi}'(t+) \leq \bar{\phi}'(t-), \bar{\psi}'(t+) \leq \bar{\psi}'(t-), t \in \mathbb{R}, \\ \underline{\phi}'(t+) \geq \underline{\phi}'(t-), \underline{\psi}'(t+) \geq \underline{\psi}'(t-), t \in \mathbb{R}, \end{cases} \tag{3.8}$$

then

$$\underline{\Phi} \leq (F_1(\underline{\phi}, \underline{\psi}), F_2(\underline{\phi}, \underline{\psi})) \leq (F_1(\bar{\phi}, \bar{\psi}), F_2(\bar{\phi}, \bar{\psi})) \leq \bar{\Phi}, \tag{3.9}$$

and $(F_1(\underline{\phi}, \underline{\psi}), F_2(\underline{\phi}, \underline{\psi})), (F_1(\bar{\phi}, \bar{\psi}), F_2(\bar{\phi}, \bar{\psi})) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ are a lower and an upper solution of (2.1), respectively.

Proof. Without loss of generality, we assume that $\bar{\Phi}$ and $\underline{\Phi}$ are twice continuously differentiable in $\mathbb{R} \setminus \{T_i : i = 1, \dots, m\}$ with $-\infty < T_1 < T_2 < \dots < T_m < +\infty$. Denote $T_0 = -\infty$ and $T_{m+1} = +\infty$. It is easy to verify that $(F_1(\underline{\phi}, \underline{\psi}), F_2(\underline{\phi}, \underline{\psi}))$ is well defined and twice continuously differentiable. For any $t \in (T_k, T_{k+1}), 0 \leq k \leq m$, it follows from (2.6) and the definition of weak lower solution that

$$\begin{aligned} F_1(\underline{\phi}, \underline{\psi})(t) &= \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_t^{\infty} e^{\lambda_2(t-s)} \right] H_1(\underline{\phi}, \underline{\psi})(s) ds \\ &\geq \frac{1}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_t^{\infty} e^{\lambda_2(t-s)} \right] \\ &\quad \times \left(\beta_1 \underline{\phi}(s) + c \underline{\phi}'(s) - d_1 \underline{\phi}''(s) \right) ds \\ &= \underline{\phi}(t) + \frac{1}{\lambda_2 - \lambda_1} \left[\sum_{j=1}^k e^{\lambda_2(t-T_j)} \left(\underline{\phi}'(T_{j+}) - \underline{\phi}'(T_{j-}) \right) \right] \\ &\quad + \frac{1}{\lambda_2 - \lambda_1} \left[\sum_{j=k+1}^m e^{\lambda_1(t-T_j)} \left(\underline{\phi}'(T_{j+}) - \underline{\phi}'(T_{j-}) \right) \right] \\ &\geq \underline{\phi}(t). \end{aligned}$$

Using a similar argument, we can prove that (3.9) holds. Furthermore, (2.7) together with lemma 3.2 yields

$$\begin{aligned} 0 &= d_1(F_1(\underline{\phi}, \bar{\psi}))''(t) - c(F_1(\underline{\phi}, \bar{\psi}))'(t) - \beta_1 F_1(\underline{\phi}, \bar{\psi})(t) + H_1(\underline{\phi}, \bar{\psi})(t) \\ &\leq d_1(F_1(\underline{\phi}, \bar{\psi}))''(t) - c(F_1(\underline{\phi}, \bar{\psi}))'(t) - \beta_1 F_1(\underline{\phi}, \bar{\psi})(t) \\ &\quad + H_1(F_1(\underline{\phi}, \bar{\psi}), F_2(\underline{\phi}, \bar{\psi}))(t) \\ &= d_1(F_1(\underline{\phi}, \bar{\psi}))''(t) - c(F_1(\underline{\phi}, \bar{\psi}))'(t) + f_1^c(F_1(\underline{\phi}, \bar{\psi}), F_2(\underline{\phi}, \bar{\psi})). \end{aligned}$$

Similarly, we have

$$d_2(F_2(\bar{\phi}, \underline{\psi}))''(t) - c(F_2(\bar{\phi}, \underline{\psi}))'(t) + f_2^c(F_1(\bar{\phi}, \underline{\psi}), F_2(\bar{\phi}, \underline{\psi})) \geq 0.$$

Note that $(F_1(\underline{\phi}, \bar{\psi}), F_2(\bar{\phi}, \underline{\psi})) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \cap C^2(\mathbb{R}, \mathbb{R}^2)$; we conclude that it is a lower solution of (2.1).

In a similar way, we can prove that $(F_1(\bar{\phi}, \underline{\psi}), F_2(\underline{\phi}, \bar{\psi}))$ is an upper solution of (2.1). The proof is complete. \square

Theorem 3.10. Assume that (A1), (A2) and (WQM) hold. If (2.1) has a weak upper solution $(\bar{\phi}, \bar{\psi})$ and a weak lower solution $(\underline{\phi}, \underline{\psi})$ satisfying (P1), (P2) and (3.8), then (2.1) has a travelling wave solution satisfying (2.3).

4. The Case (WQM*)

In this section, we shall consider the existence of travelling wave solutions of (2.1) when the delayed reaction terms f_1 and f_2 satisfy the condition (WQM*).

In what follows, we assume that an upper solution $(\bar{\phi}(t), \bar{\psi}(t))$ and a lower solution $(\underline{\phi}(t), \underline{\psi}(t))$ satisfy (P1), (P2), and

(P3) $e^{\beta_1 s}[\bar{\phi}(s) - \underline{\phi}(s)]$ and $e^{\beta_2 s}[\bar{\psi}(s) - \underline{\psi}(s)]$ are non-decreasing for $s \in \mathbb{R}$.

To start with, we define the following profile set:

$$\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) = \left\{ (\phi, \psi) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) : \begin{array}{l} \text{(i) } (\underline{\phi}, \underline{\psi}) \leq (\phi, \psi) \leq (\bar{\phi}, \bar{\psi}), \\ \text{(ii) } e^{\beta_1 s}[\phi(s) - \underline{\phi}(s)], e^{\beta_1 s}[\bar{\phi}(s) - \phi(s)], \\ e^{\beta_2 s}[\psi(s) - \underline{\psi}(s)], e^{\beta_2 s}[\bar{\psi}(s) - \psi(s)] \\ \text{are non-decreasing for } s \in \mathbb{R} \end{array} \right\}.$$

It is easy to see that $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is non-empty. In fact, by (P3), we know that $e^{\beta_1 s}[\bar{\phi}(s) - \underline{\phi}(s)]$ and $e^{\beta_2 s}[\bar{\psi}(s) - \underline{\psi}(s)]$ are non-decreasing in $s \in \mathbb{R}$, $e^{\beta_1 s}[\bar{\phi}(s) - \bar{\phi}(s)] = 0$, and $e^{\beta_2 s}[\bar{\psi}(s) - \bar{\psi}(s)] = 0$. Thus $(\bar{\phi}, \bar{\psi})$ satisfies (ii) of $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$. Similarly, $(\underline{\phi}(t), \underline{\psi}(t))$ satisfies (ii) of $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$.

Lemma 4.1. Assume that (WQM*) holds. Then

$$H_1(\phi_2, \psi_1)(t) \leq H_1(\phi_1, \psi_2)(t), \quad H_2(\phi_1, \psi_2)(t) \leq H_2(\phi_2, \psi_1)(t),$$

where $\Phi = (\phi_1, \psi_1), \Psi = (\phi_2, \psi_2) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ with (i) $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1, 0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2$; (ii) $e^{\beta_1 s}[\phi_1(s) - \phi_2(s)]$ and $e^{\beta_2 s}[\psi_1(s) - \psi_2(s)]$ are non-decreasing for $s \in \mathbb{R}$.

Lemma 4.2. Assume that (WQM*) holds. Then

$$F_1(\phi_2, \psi_1)(t) \leq F_1(\phi_1, \psi_2)(t), \quad F_2(\phi_1, \psi_2)(t) \leq F_2(\phi_2, \psi_1)(t),$$

where $\Phi = (\phi_1, \psi_1), \Psi = (\phi_2, \psi_2) \in C_{[0, M]}(\mathbb{R}, \mathbb{R}^2)$ with (i) $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1, 0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2$; (ii) $e^{\beta_1 s}[\phi_1(s) - \phi_2(s)]$ and $e^{\beta_2 s}[\psi_1(s) - \psi_2(s)]$ are non-decreasing for $s \in \mathbb{R}$.

Lemma 4.3. Assume that (A2) and (WQM*) hold. Then $F : C_{[0, M]}(\mathbb{R}, \mathbb{R}^2) \rightarrow C(\mathbb{R}, \mathbb{R}^2)$ is continuous with respect to the norm $|\cdot|_\mu$ in $B_\mu(\mathbb{R}, \mathbb{R}^2)$.

The proofs of lemmas 4.1–4.3 are similar to those of lemmas 3.2–3.4 and are omitted here. By (P3), it is easy to see the following lemma.

Lemma 4.4. $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ is a closed, bounded and convex subset of $B_\mu(\mathbb{R}, \mathbb{R}^2)$.

Lemma 4.5. Assume that (WQM*) holds. If $c > 1 - \min\{\beta_1 d_1, \beta_2 d_2\}$, then

$$F\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \subset \Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]).$$

Proof. For any $(\phi, \psi) \in \Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$, repeating the argument in the proof of lemma 3.5, we have

$$\underline{\phi} \leq F_1(\phi, \psi) \leq \bar{\phi} \text{ and } \underline{\psi} \leq F_2(\phi, \psi) \leq \bar{\psi}.$$

This implies that (i) of $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$ holds. We now prove (ii) of $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$. Let $F_1(\phi, \psi) = \phi_1$ for $(\phi, \psi) \in \Gamma^*$, then

$$\begin{aligned} & e^{\beta_1 t} [\bar{\phi}(t) - \phi_1(t)] \\ &= \frac{e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_t^{+\infty} e^{\lambda_2(t-s)} \right] \\ & \quad \times \left\{ [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s)] - [\beta_1 \phi_1(s) + c\phi_1'(s) - d_1 \phi_1''(s)] \right\} ds \\ &= \frac{e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_t^{+\infty} e^{\lambda_2(t-s)} \right] \\ & \quad \times \left\{ [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\phi, \psi)(s)] \right. \\ & \quad \left. - [\beta_1 \phi_1(s) + c\phi_1'(s) - d_1 \phi_1''(s) - H_1(\phi, \psi)(s)] \right\} ds \\ &= \frac{e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \left[\int_{-\infty}^t e^{\lambda_1(t-s)} + \int_t^{+\infty} e^{\lambda_2(t-s)} \right] \\ & \quad \times [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\phi, \psi)(s)] ds. \end{aligned}$$

Hence, we have

$$\begin{aligned} & \frac{d}{dt} \{e^{\beta_1 t} [\bar{\phi}(t) - \phi_1(t)]\} \\ &= \frac{(\beta_1 + \lambda_1) e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_{-\infty}^t e^{\lambda_1(t-s)} [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\phi, \psi)(s)] ds \\ & \quad + \frac{(\beta_1 + \lambda_2) e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_t^{+\infty} e^{\lambda_2(t-s)} [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\phi, \psi)(s)] ds \\ &\geq \frac{(\beta_1 + \lambda_1) e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_{-\infty}^t e^{\lambda_1(t-s)} [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\bar{\phi}, \underline{\psi})(s)] ds \\ & \quad + \frac{(\beta_1 + \lambda_2) e^{\beta_1 t}}{d_1(\lambda_2 - \lambda_1)} \int_t^{+\infty} e^{\lambda_2(t-s)} [\beta_1 \bar{\phi}(s) + c\bar{\phi}'(s) - d_1 \bar{\phi}''(s) - H_1(\bar{\phi}, \underline{\psi})(s)] ds \\ &\geq 0, t \in \mathbb{R}. \end{aligned}$$

Similarly, we can prove that

$$e^{\beta_2 t} [\bar{\psi}(t) - F_2(\phi, \psi)(t)], e^{\beta_1 t} [F_1(\phi, \psi)(t) - \underline{\phi}(t)] \text{ and } e^{\beta_2 t} [F_2(\phi, \psi)(t) - \underline{\psi}(t)]$$

are non-decreasing in $t \in \mathbb{R}$. Thus, $F_1(\phi, \psi)$ satisfies (ii) of $\Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$. The proof is complete. \square

Similar to lemma 3.6 we have the following lemma.

Lemma 4.6. Assume that (A2) and (WQM*) hold. Then

$$F : \Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}]) \rightarrow \Gamma^*([\underline{\phi}, \underline{\psi}], [\bar{\phi}, \bar{\psi}])$$

is compact.

Now we state our main result in this section; its proof is similar to that of theorem 3.7.

Theorem 4.7. Assume that (A1), (A2) and (WQM*) hold. Assume further that (2.1) has an upper solution $(\bar{\phi}, \bar{\psi}) \in C_{[0, \mathbf{M}]}(\mathbb{R}, \mathbb{R}^2)$ and a lower $(\underline{\phi}, \underline{\psi}) \in C_{[0, \mathbf{M}]}(\mathbb{R}, \mathbb{R}^2)$ satisfying (P1)–(P3). Then, for any $c > 1 - \min\{\beta_1 d_1, \beta_2 d_2\}$, (2.1) has a travelling wave solution satisfying (2.3).

Similar to that in section 3, we have the following.

Theorem 4.8. Assume that (A1), (A2) and (WQM*) hold. If (2.1) has a weak upper solution $(\bar{\phi}, \bar{\psi}) \in C_{[0, \mathbf{M}]}(\mathbb{R}, \mathbb{R}^2)$ and a weak lower solution $(\underline{\phi}, \underline{\psi}) \in C_{[0, \mathbf{M}]}(\mathbb{R}, \mathbb{R}^2)$ such that (P1)–(P3) and (3.8) hold, then, for any $c > 1 - \min\{\beta_1 d_1, \beta_2 d_2\}$, (2.1) has a travelling wave solution satisfying (2.3).

Remark 4.9. If (WQM*) is satisfied, then we can always choose $\beta_i > 0$ sufficiently large such that $c > 1 - \min\{\beta_1 d_1, \beta_2 d_2\}$.

5. Applications

As mentioned in the introduction, in this section, we employ our conclusions in sections 3 and 4 to establish the existence of travelling wave solutions for systems (1.2) and (1.3).

Example 5.1. We consider the existence of the travelling wave solutions for the delayed diffusion–competition system (1.2), that is,

$$\begin{cases} \frac{\partial}{\partial t} u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2} u_1(x, t) + r_1 u_1(x, t)[1 - a_1 u_1(x, t) - b_1 u_2(x, t - \tau_1)], \\ \frac{\partial}{\partial t} u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2} u_2(x, t) + r_2 u_2(x, t)[1 - b_2 u_1(x, t - \tau_2) - a_2 u_2(x, t)]. \end{cases} \tag{5.1}$$

Assume that $c > 0$. Let $u_1(x, t) = u_1(x + ct) = \phi(s)$, $u_2(x, t) = u_2(x + ct) = \psi(s)$, $s \in \mathbb{R}$, and denote the coordinate s as t , then the corresponding wave system is

$$\begin{cases} d_1 \phi''(t) - c\phi'(t) + r_1 \phi(t)[1 - a_1 \phi(t) - b_1 \psi(t - c\tau_1)] = 0, \\ d_2 \psi''(t) - c\psi'(t) + r_2 \psi(t)[1 - b_2 \phi(t - c\tau_2) - a_2 \psi(t)] = 0. \end{cases} \tag{5.2}$$

We are interested in solutions of (5.2) satisfying

$$\lim_{t \rightarrow -\infty} \phi(t) = 0, \quad \lim_{t \rightarrow +\infty} \phi(t) = k_1, \quad \lim_{t \rightarrow -\infty} \psi(t) = 0, \quad \lim_{t \rightarrow +\infty} \psi(t) = k_2,$$

where

$$k_1 = \frac{a_2 - b_1}{a_1 a_2 - b_1 b_2} > 0, \quad k_2 = \frac{a_1 - b_2}{a_1 a_2 - b_1 b_2} > 0,$$

provided that

$$a_1 > b_2, a_2 > b_1. \tag{5.3}$$

For $\phi, \psi \in C([-\tau, 0], \mathbb{R})$, where $\tau = \max\{\tau_1, \tau_2\}$, denote

$$\begin{aligned} f_1(\phi, \psi) &= r_1\phi(0)[1 - a_1\phi(0) - b_1\psi(-\tau_1)], \\ f_2(\phi, \psi) &= r_2\psi(0)[1 - b_2\phi(-\tau_2) - a_2\psi(0)]. \end{aligned}$$

Obviously, (A1) and (A2) are satisfied. We now verify that $f = (f_1, f_2)$ satisfies (WQM).

Lemma 5.2. *The function f satisfies (WQM).*

Proof. For any $\Phi(s) = (\phi_1(s), \phi_2(s))$, $\Psi(s) = (\psi_1(s), \psi_2(s)) \in C([-\tau, 0], \mathbb{R}^2)$ with $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1$, $0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2$, in view of $M_1 > k_1$ and $M_2 > k_2$, we have $2a_1M_1 + b_1M_2 - 1 > 2a_1k_1 + b_1k_2 - 1 = a_1k_1 > 0$ and

$$\begin{aligned} &f_1(\phi_1, \psi_1) - f_1(\phi_2, \psi_1) \\ &= r_1\phi_1(0)[1 - a_1\phi_1(0) - b_1\psi_1(-\tau_1)] - r_1\phi_2(0)[1 - a_1\phi_2(0) - b_1\psi_1(-\tau_1)] \\ &= r_1[\phi_1(0) - \phi_2(0)] - r_1a_1\phi_1^2(0) - \phi_2^2(0) - r_1b_1\psi_1(-\tau_1)[\phi_1(0) - \phi_2(0)] \\ &= r_1[\phi_1(0) - \phi_2(0)][1 - a_1(\phi_1(0) + \phi_2(0)) - b_1\psi_1(-\tau_1)] \\ &\geq r_1[\phi_1(0) - \phi_2(0)][1 - 2a_1M_1 - b_1M_2] \\ &= -r_1(2a_1M_1 + b_1M_2 - 1)[\phi_1(0) - \phi_2(0)] \\ &= -\beta_1[\phi_1(0) - \phi_2(0)] \end{aligned}$$

and

$$\begin{aligned} &f_1(\phi_1, \psi_1) - f_1(\phi_1, \psi_2) \\ &= r_1\phi_1(0)[1 - a_1\phi_1(0) - b_1\psi_1(-\tau_1)] - r_1\phi_1(0)[1 - a_1\phi_1(0) - b_1\psi_2(-\tau_1)] \\ &= -r_1b_1\phi_1(0)(\psi_1(-\tau_1) - \psi_2(-\tau_1)) \\ &\leq 0. \end{aligned}$$

In a similar argument, we can prove that f_2 satisfies (WQM). The proof is complete. □

In order to apply theorem 3.10, we need to construct a weak upper solution and a weak lower solution for (5.2).

If $c > \max\{2\sqrt{d_1r_1}, 2\sqrt{d_2r_2}\}$, then there exist $0 < \lambda_1 < \lambda_2$ such that

$$d_1\lambda_i^2 - c\lambda_i + r_1 = 0, i = 1, 2,$$

and $0 < \lambda_3 < \lambda_4$ such that

$$d_2\lambda_i^2 - c\lambda_i + r_2 = 0, i = 3, 4.$$

For fixed

$$\eta \in \left(1, \min\left\{2, \frac{\lambda_2}{\lambda_1}, \frac{\lambda_4}{\lambda_3}\right\}\right) \tag{5.4}$$

and large constant $q > 0$, we consider the functions $l_1(t) = e^{\lambda_1 t} - qe^{\eta\lambda_1 t}$ and $l_2(t) = e^{\lambda_3 t} - qe^{\eta\lambda_3 t}$. It is easy to see that $l_1(t)$ and $l_2(t)$ have global maximum $m_1 > 0$ and $m_2 > 0$, respectively. Define

$$t_1 = \max\left\{t : l_1(t) = \frac{m_1}{2}\right\} \text{ and } t_3 = \max\left\{t : l_2(t) = \frac{m_2}{2}\right\}.$$

Then for any given $\lambda > 0$, there exists $\varepsilon_2 > 0$ and $\varepsilon_4 > 0$ such that

$$k_1 - \varepsilon_2 e^{-\lambda t_1} = l_1(t_1) = \frac{m_1}{2} \text{ and } k_2 - \varepsilon_4 e^{-\lambda t_3} = l_2(t_3) = \frac{m_2}{2}.$$

Note that (5.3) holds, we have $\varepsilon_0 > 0$, $\varepsilon_1 > 0$ and $\varepsilon_3 > 0$ such that

$$\begin{cases} a_1\varepsilon_1 - b_1\varepsilon_4 > \varepsilon_0, & a_2\varepsilon_3 - b_2\varepsilon_2 > \varepsilon_0, \\ a_1\varepsilon_2 - b_1\varepsilon_3 > \varepsilon_0, & a_2\varepsilon_4 - b_2\varepsilon_1 > \varepsilon_0. \end{cases} \quad (5.5)$$

For the above constants and suitable constants t_2, t_4 , we define the continuous functions as follows:

$$\bar{\phi}(t) = \begin{cases} e^{\lambda_1 t}, & t \leq t_2, \\ k_1 + \varepsilon_1 e^{-\lambda t}, & t \geq t_2, \end{cases} \quad \bar{\psi}(t) = \begin{cases} e^{\lambda_3 t}, & t \leq t_4 \\ k_2 + \varepsilon_3 e^{-\lambda t}, & t \geq t_4 \end{cases}$$

and

$$\underline{\phi}(t) = \begin{cases} e^{\lambda_1 t} - q e^{\eta \lambda_1 t}, & t \leq t_1, \\ k_1 - \varepsilon_2 e^{-\lambda t}, & t \geq t_1, \end{cases} \quad \underline{\psi}(t) = \begin{cases} e^{\lambda_3 t} - q e^{\eta \lambda_3 t}, & t \leq t_3, \\ k_2 - \varepsilon_4 e^{-\lambda t}, & t \geq t_3, \end{cases}$$

where $q > 0$ is large enough and $\lambda > 0$ is small enough. It is easy to see $M_1 = \sup_{t \in \mathbb{R}} \bar{\phi}(t) > k_1$, $M_2 = \sup_{t \in \mathbb{R}} \bar{\psi}(t) > k_2$, $\bar{\phi}(t)$, $\bar{\psi}(t)$, $\underline{\phi}(t)$ and $\underline{\psi}(t)$ satisfy (P1), (P2), (3.8) and

$$\min\{t_2, t_4\} - \max\{\tau_1, \tau_2\} \geq \max\{t_1, t_3\}$$

for sufficiently large $q > 0$ and sufficiently small $\lambda > 0$. We now prove that the continuous functions $(\bar{\phi}(t), \bar{\psi}(t))$ and $(\underline{\phi}(t), \underline{\psi}(t))$ are a weak upper solution and a weak lower solution of (5.2), respectively.

Lemma 5.3. Assume that (5.3) and (5.5) hold. Then $\bar{\Phi}(t) = (\bar{\phi}(t), \bar{\psi}(t))$ is a weak upper solution and $\underline{\Phi}(t) = (\underline{\phi}(t), \underline{\psi}(t))$ is a weak lower solution of (5.2).

Proof. For $t \leq t_2$, in view of $\underline{\psi}(t - c\tau_1) \geq 0$ for $t \in \mathbb{R}$ and $d_1\lambda_1^2 - c\lambda_1 + r_1 = 0$, we have

$$\begin{aligned} & d_1\bar{\phi}''(t) - c\bar{\phi}'(t) + r_1\bar{\phi}(t)[1 - a_1\bar{\phi}(t) - b_1\underline{\psi}(t - c\tau_1)] \\ & \leq d_1\bar{\phi}''(t) - c\bar{\phi}'(t) + r_1\bar{\phi}(t) \\ & = [d_1\lambda_1^2 - c\lambda_1 + r_1](k_1 + \varepsilon_1)e^{\lambda_1 t} = 0. \end{aligned}$$

For $t \geq t_2$, since $\underline{\psi}(t - c\tau_1) = k_2 - \varepsilon_4 e^{-\lambda(t - c\tau_1)}$, then

$$\begin{aligned} & d_1\bar{\phi}''(t) - c\bar{\phi}'(t) + r_1\bar{\phi}(t)[1 - a_1\bar{\phi}(t) - b_1\underline{\psi}(t - c\tau_1)] \\ & = e^{-\lambda t} \{d_1\varepsilon_1\lambda^2 + c\varepsilon_1\lambda + r_1(k_1 + \varepsilon_1)e^{-\lambda t}\}(b_1\varepsilon_4 e^{\lambda c\tau_1} - a_1\varepsilon_1). \end{aligned}$$

Let

$$I_1(\lambda) = d_1\varepsilon_1\lambda^2 + c\varepsilon_1\lambda + r_1(k_1 + \varepsilon_1)e^{-\lambda t}(b_1\varepsilon_4 e^{\lambda c\tau_1} - a_1\varepsilon_1).$$

Then, $a_1\varepsilon_1 - b_1\varepsilon_4 > \varepsilon_0$ implies that

$$I_1(0) = r_1(k_1 + \varepsilon_1)(b_1\varepsilon_4 - a_1\varepsilon_1) < 0,$$

and there exists a $\lambda_1^* > 0$ such that $I_1(\lambda) < 0$ for $\lambda \in (0, \lambda_1^*)$. Thus, we have

$$d_1\bar{\phi}''(t) - c\bar{\phi}'(t) + r_1\bar{\phi}(t)[1 - a_1\bar{\phi}(t) - b_1\underline{\psi}(t - c\tau_1)] \leq 0.$$

Similarly, there exists a $\lambda_2^* > 0$ such that for $\lambda \in (0, \lambda_2^*)$ we have

$$d_2\bar{\psi}''(t) - c\bar{\psi}'(t) + r_2\bar{\psi}(t)[1 - b_2\bar{\phi}(t - c\tau_2) - a_2\bar{\psi}(t)] \leq 0.$$

Taking $\lambda \in (0, \min(\lambda_1^*, \lambda_2^*))$, we see that our conclusion is true.

A similar argument applies to $\underline{\Phi}(t) = (\underline{\phi}(t), \underline{\psi}(t))$. The proof is complete. \square

By theorem 3.10, we have the following result.

Theorem 5.4. *Assume that (5.3) holds. Then for every $c > \max\{2\sqrt{d_1r_1}, 2\sqrt{d_2r_2}\}$, (5.1) has a travelling wave solution $(\phi(x + ct), \psi(x + ct))$ with wave speed c , which connects $(0, 0)$ and (k_1, k_2) . Furthermore, $\lim_{\xi \rightarrow -\infty} (\phi(\xi)e^{-\lambda_1\xi}, \psi(\xi)e^{-\lambda_3\xi}) = (1, 1)$, where $\xi = x + ct$.*

Example 5.5. We now consider the delayed diffusion–competition system (1.3), that is

$$\begin{cases} \frac{\partial}{\partial t} u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2} u_1(x, t) + r_1 u_1(x, t) [1 - a_1 u_1(x, t - \tau_1) - b_1 u_2(x, t - \tau_2)], \\ \frac{\partial}{\partial t} u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2} u_2(x, t) + r_2 u_2(x, t) [1 - b_2 u_1(x, t - \tau_3) - a_2 u_2(x, t - \tau_4)]. \end{cases} \quad (5.6)$$

The corresponding travelling wave system is

$$\begin{cases} d_1 \phi''(t) - c\phi'(t) + r_1 \phi(t) [1 - a_1 \phi(t - c\tau_1) - b_1 \psi(t - c\tau_2)] = 0, \\ d_2 \psi''(t) - c\psi'(t) + r_2 \psi(t) [1 - b_2 \phi(t - c\tau_3) - a_2 \psi(t - c\tau_4)] = 0. \end{cases} \quad (5.7)$$

For $\phi, \psi \in C([-\tau, 0], \mathbb{R})$ with $\tau = \max\{\tau_1, \tau_2, \tau_3, \tau_4\}$, we denote

$$\begin{aligned} f_1(\phi, \psi) &= r_1 \phi(0) [1 - a_1 \phi(-\tau_1) - b_1 \psi(-\tau_2)], \\ f_2(\phi, \psi) &= r_2 \psi(0) [1 - b_2 \phi(-\tau_3) - a_2 \psi(-\tau_4)]. \end{aligned}$$

Obviously, (A1) and (A2) are satisfied. We now verify that $f = (f_1, f_2)$ satisfies (WQM*).

Lemma 5.6. *Assume that τ_1 and τ_4 are small enough. Then the function f satisfies (WQM*).*

Proof. For any $\Phi(s) = (\phi_1(s), \phi_2(s))$, $\Psi(s) = (\psi_1(s), \psi_2(s)) \in C([-\tau, 0], \mathbb{R}^2)$ with (i) $0 \leq \phi_2(s) \leq \phi_1(s) \leq M_1$, $0 \leq \psi_2(s) \leq \psi_1(s) \leq M_2$ and (ii) $e^{\beta_1 s} [\phi_1(s) - \phi_2(s)]$ and $e^{\beta_2 s} [\psi_1(s) - \psi_2(s)]$ are non-decreasing in $s \in [-\tau, 0]$, we have

$$\begin{aligned} & f_1(\phi_1, \psi_1) - f_1(\phi_2, \psi_1) \\ &= r_1 \phi_1(0) [1 - a_1 \phi_1(-\tau_1) - b_1 \psi_1(-\tau_2)] - r_1 \phi_2(0) [1 - a_1 \phi_2(-\tau_1) - b_1 \psi_1(-\tau_2)] \\ &= r_1 [\phi_1(0) - \phi_2(0)] - r_1 a_1 [\phi_1(0)\phi_1(-\tau_1) - \phi_2(0)\phi_2(-\tau_1)] \\ &\quad - r_1 b_1 \psi_1(-\tau_2) [\phi_1(0) - \phi_2(0)] \\ &\geq (r_1 - r_1 b_1 M_2) [\phi_1(0) - \phi_2(0)] - r_1 a_1 \phi_1(0) [\phi_1(-\tau_1) - \phi_2(-\tau_1)] \\ &\quad - r_1 a_1 \phi_2(-\tau_1) [\phi_1(0) - \phi_2(0)] \\ &\geq r_1 (1 - b_1 M_2 - a_1 M_1) [\phi_1(0) - \phi_2(0)] \\ &\quad - r_1 a_1 \phi_1(0) e^{\beta_1 \tau_1} e^{-\beta_1 \tau_1} [\phi_1(-\tau_1) - \phi_2(-\tau_1)] \\ &\geq r_1 (1 - b_1 M_2 - a_1 M_1 - a_1 M_1 e^{\beta_1 \tau_1}) [\phi_1(0) - \phi_2(0)]. \end{aligned}$$

If $\tau_1 > 0$ is small enough, then we can choose $\beta_1 > 0$ such that

$$r_1 (1 - b_1 M_2 - a_1 M_1 - a_1 M_1 e^{\beta_1 \tau_1}) > -\beta_1.$$

We have

$$\begin{aligned} & f_1(\phi_1, \psi_1) - f_1(\phi_1, \psi_2) \\ &= r_1 \phi_1(0) [1 - a_1 \phi_1(-\tau_1) - b_1 \psi_1(-\tau_2)] - r_1 \phi_1(0) [1 - a_1 \phi_1(-\tau_1) - b_1 \psi_2(-\tau_2)] \\ &= -r_1 b_1 \phi_1(0) [\psi_1(-\tau_2) - \psi_2(-\tau_2)] \\ &\leq 0. \end{aligned}$$

Thus $f_1(\phi, \psi)$ satisfies (WQM*) if τ_1 is small enough.

In a similar way, we can prove that $f_2(\phi, \psi)$ satisfies (WQM*) if τ_4 is small enough. The proof is complete. \square

Remark 5.7. From the proof of lemma 5.6 we can see that if τ_1 and τ_4 are small enough, then we can always choose $\beta_i > 0$ sufficiently large such that $c > 1 - \min\{\beta_1 d_1, \beta_2 d_2\}$.

Now we define $\bar{\phi}(t)$, $\bar{\psi}(t)$, $\underline{\phi}(t)$ and $\underline{\psi}(t)$ as in example 5.1. It is easy to see that $\bar{\phi}(t)$, $\bar{\psi}(t)$, $\underline{\phi}(t)$ and $\underline{\psi}(t)$ satisfy (P1)–(P3) and (3.8).

Lemma 5.8. Assume that (5.3) and (5.5) hold. If τ_1 and τ_4 are small enough, then $\bar{\Phi}(t) = (\bar{\phi}(t), \bar{\psi}(t))$ is a weak upper solution and $\underline{\Phi}(t) = (\underline{\phi}(t), \underline{\psi}(t))$ is a weak lower solution of (5.7).

Proof. For $\bar{\phi}(t)$, we need to prove that

$$d_1 \bar{\phi}''(t) - c \bar{\phi}'(t) + r_1 \bar{\phi}(t) [1 - a_1 \bar{\phi}(t - c\tau_1) - b_1 \underline{\psi}(t - c\tau_2)] \leq 0. \quad (5.8)$$

For $t > t_2 + c\tau_1$ or $t < t_2$, the proof of (5.8) is similar to that of lemma 5.3: we omit it here.

For $t_2 < t < t_2 + c\tau_1$, we note that we can choose λ small enough such that

$$d_1 \bar{\phi}''(t) - c \bar{\phi}'(t) + r_1 \bar{\phi}(t) [1 - a_1 \bar{\phi}(t - c\tau_1) - b_1 \underline{\psi}(t - c\tau_2)] < 0,$$

for $t = t_2 + c\tau_1$. Since τ_1 is small enough and independent of $\bar{\phi}$, $\underline{\psi}$, and $\bar{\phi}''(t)$, $\bar{\phi}'(t)$, $\bar{\phi}(t)$ and $\underline{\psi}(t)$ are uniformly bounded and uniformly continuous for $t \in \mathbb{R} \setminus \{t_2, t_3\}$, it follows that (5.8) holds for $t_2 < t < t_2 + c\tau_1$.

Similarly, we can prove that $\bar{\psi}$ satisfies

$$d_2 \bar{\psi}''(t) - c \bar{\psi}'(t) + r_2 \bar{\psi}(t) [1 - b_2 \underline{\phi}(t - c\tau_3) - a_2 \bar{\psi}(t - c\tau_4)] \leq 0$$

and $\underline{\Phi}(t) = (\underline{\phi}(t), \underline{\psi}(t))$ is a weak lower solution. The proof is complete. \square

Theorem 5.9. Assume that (5.3) holds and τ_1 and τ_4 are sufficiently small. Then for every $c > \max\{2\sqrt{d_1 r_1}, 2\sqrt{d_2 r_2}\}$, (5.6) has a travelling wave solution $(\phi(x + ct), \psi(x + ct))$ with wave speed c , which connects $(0, 0)$ and (k_1, k_2) . Furthermore,

$$\lim_{\xi \rightarrow -\infty} (\phi(\xi) e^{-\lambda_1 \xi}, \psi(\xi) e^{-\lambda_3 \xi}) = (1, 1),$$

where $\xi = x + ct$.

Remark 5.10. We note that the delay of example 5.1 does not affect the existence of travelling wave solutions. However, the delays (τ_1 and τ_4) of example 5.5 do.

Remark 5.11. If $\tau = 0$, then (5.6) reduces to (1.5), that is,

$$\begin{cases} \frac{\partial}{\partial t} u_1(x, t) = d_1 \frac{\partial^2}{\partial x^2} u_1(x, t) + r_1 u_1(x, t) [1 - a_1 u_1(x, t) - b_1 u_2(x, t)], \\ \frac{\partial}{\partial t} u_2(x, t) = d_2 \frac{\partial^2}{\partial x^2} u_2(x, t) + r_2 u_2(x, t) [1 - b_2 u_1(x, t) - a_2 u_2(x, t)]. \end{cases} \quad (5.9)$$

Tang and Fife [29] and van Vuuren [30] proved that (5.9) has a bounded travelling wave front solution connecting $(0, 0)$ and (k_1, k_2) if and only if $c \geq \max\{2\sqrt{d_1 r_1}, 2\sqrt{d_2 r_2}\}$. Our results certainly include their results when $c > \max\{2\sqrt{d_1 r_1}, 2\sqrt{d_2 r_2}\}$. If $c = \max\{2\sqrt{d_1 r_1}, 2\sqrt{d_2 r_2}\}$, we can get the existence of travelling wave solution only by changing the definition of $\underline{\phi}$ and $\underline{\psi}$. However, our results cannot ensure that the bounded travelling wave solutions of systems (5.1) and (5.6) connecting $(0, 0)$ and (k_1, k_2) are monotonic. Numerical simulations indicate that the travelling wave solutions are monotonic. It would be interesting to study the monotonicity of the travelling wave solutions in the delayed diffusion–competition models.

6. Discussion

Coexistence of competing species is very common in natural systems (Durrett and Levin [6, 7], Silvertown *et al* [28]). In the classical competition models (described by ordinary differential equations and used quantities averaged over space) spatial heterogeneity is usually neglected, while in real systems species are distributed spatially; organisms experience different local environments, they consume resources locally and move around. It is known now that spatial heterogeneity is crucial to the dynamics of biological systems, in particular for the coexistence of some competing species (Durrett and Levin [6]).

Different types of mathematical models have been used to describe spatial heterogeneous environments. For example, in metapopulation models (Hanski and Gilpin [10]) space is represented as a set of patches with local interactions. Reaction–diffusion equation models (Cantrell and Cosner [4]) take account of space explicitly. Stochastic spatial models (Durrett and Levin [6, 7]) are a combination of these two approaches.

In this paper we considered a class of delayed reaction–diffusion systems without monotonicity. By using Schauder’s fixed point theorem, a new cross-iteration scheme was given to establish the existence of travelling wave solutions. More precisely, by using such a new cross-iteration, we reduced the existence of travelling wave solutions to the existence of an admissible pair of upper and lower solutions which are easy to construct in practice. The general results were then applied to study the existence of travelling wave solutions in delayed two-species diffusion–competition systems. It is interesting to note that the delays appearing in the interspecific competition terms do not affect the existence of travelling waves while the delays appearing in the intraspecific competition terms do. The existence of travelling wave solutions which connect the trivial equilibrium $(0, 0)$ and the positive equilibrium (k_1, k_2) indicates that there is a transition zone moving from the steady state with no species to the steady state with the coexistence of both species.

It is also known that the dynamics of reactive systems on supports of restricted geometry may deviate substantially from the predictions of mean-field descriptions (Provata *et al* [24]). To couple microscope level processes and the evolution of the macroscopic observables, nonlinear models on regular lattices have also been proposed (Kowalik *et al* [18], Provata *et al* [24], Rauch and Bar-Yam [25] and the references cited therein). The delayed lattice differential equations version of our model (1.4) takes the following form:

$$\begin{cases} \frac{du_n}{dt} = \sum_{j=1}^m a_j [u_{n+j}(t) - 2u_n(t) + u_{n-j}(t)] + f_1(u_n(t - \tau_{11}), v_n(t - \tau_{12})), \\ \frac{dv_n}{dt} = \sum_{j=1}^m b_j [v_{n+j}(t) - 2v_n(t) + v_{n-j}(t)] + f_2(u_n(t - \tau_{21}), v_n(t - \tau_{22})), \end{cases} \quad (6.1)$$

where $n \in \mathbb{Z}$, $m \geq 1$ is an integer, $a_j > 0$, $b_j > 0$, $1 \leq j \leq m$, $\tau \geq 0$. It would be interesting to combine the techniques and results in Huang *et al* [13] and the present paper to establish the existence of travelling waves for system (6.1) and apply the results to two species lattice competition models.

Acknowledgments

The authors are very grateful to an anonymous referee and the handling editor for their helpful comments and suggestions. W-T Li acknowledges support by the NNSF of China (10571078) and the Teaching and Research Award Program for Outstanding Young Teachers in Higher

Education Institutions of Ministry of Education of China. S Ruan's research was partially supported by NSF grant DMS-0412047 and the University of Miami.

References

- [1] Ashwin P B, Bartuccelli M V, Bridges T J and Gourly S A 2002 Travelling fronts for the KPP equation with spatio-temporal delay *Z. Angew. Math. Phys.* **53** 103–22
- [2] Al-Omari J and Gourly S A 2002 Monotone traveling fronts in age-structured reaction–diffusion model of a single species *J. Math. Biol.* **45** 294–312
- [3] Billingham J 2004 Dynamics of a strongly nonlocal reaction–diffusion population model *Nonlinearity* **17** 313–46
- [4] Cantrell R S and Cosner G C 2003 *Spatial Ecology via Reaction–Diffusion Equations* (Chichester, UK: Wiley)
- [5] Conley C and Gardner R 1984 An application of the generalized Mores index to traveling wave solutions of a competition reaction–diffusion model *Indiana Univ. Math. J.* **44** 319–43
- [6] Durrett R and Levin S 1994 The importance of being discrete (and spatial) *Theor. Pop. Biol.* **46** 363–94
- [7] Durrett R and Levin S 1998 Spatial aspects of interspecific competition *Theor. Pop. Biol.* **53** 30–43
- [8] Gardner R 1982 Existence and stability of traveling wave solutions of competition models: a degree theoretic approach *J. Diff. Eqns* **44** 343–64
- [9] Gourley S A and Ruan S 2003 Convergence and traveling fronts in functional differential equations with nonlocal terms: a competition model *SIAM J. Math. Anal.* **35** 806–22
- [10] Hanski I A and Gilpin M E 1996 *Metapopulation Biology: Ecology, Genetics, and Evolution* (New York: Academic)
- [11] Hosono Y 1995 Traveling waves for a diffusive Lotka–Volterra competition model, II: a geometric approach *Forma* **10** 235–57
- [12] Hosono Y 2003 Traveling waves for a diffusive Lotka–Volterra competition model, I: singular perturbation *Discrete Cont. Dyn. Syst. B* **3** 75–95
- [13] Huang J, Lu G and Ruan S 2005 Traveling wave solutions in delayed lattice differential equations with partial monotonicity *Nonlinear Anal. TMA* **60** 1331–50
- [14] Huang J and Zou X 2002 Traveling wavefronts in diffusive and cooperative Lotka–Volterra system with delays *J. Math. Anal. Appl.* **271** 455–66
- [15] Huang J and Zou X 2003 Existence of traveling wavefronts of delayed reaction–diffusion systems without monotonicity *Discrete Cont. Dyn. Syst.* **9** 925–36
- [16] Kanel J I and Zhou L 1996 Existence of wave front solutions and estimates of wave speed for a competition–diffusion system *Nonlinear Anal. TMA* **27** 579–87
- [17] Kan-on Y 1995 Parameter dependence of propagation speed of traveling waves for competition–diffusion equations *SIAM J. Math. Anal.* **26** 340–63
- [18] Kowalik M, Lipowski A and Ferreira A L 2002 Oscillations and dynamics in a two-dimensional prey–predator system *Phys. Rev. E* **66** 066107
- [19] Lan K and Wu J 2003 Traveling wavefronts of reaction–diffusion equations with and without delays *Nonlinear Anal. RWA* **4** 173–88
- [20] Leung A W 1989 *Systems of Nonlinear Partial Differential Equations with Applications to Biology and Engineering* (Dordrecht: Kluwer)
- [21] Li W T, Ruan S and Wang Z C 2005 On the diffusive Nicholson's Blowflies equation with nonlocal delays *J. Nonlinear Sci.* submitted
- [22] Ma S 2001 Traveling wavefronts for delayed reaction–diffusion systems via a fixed point theorem *J. Diff. Eqns* **171** 294–314
- [23] Pao C V 1992 *Nonlinear Parabolic and Elliptic Equations* (New York: Plenum)
- [24] Provata A, Nicolis G and Baras F 1999 Oscillatory dynamics in low-dimensional supports: a lattice Lotka–Volterra model *J. Chem. Phys.* **110** 8361–8
- [25] Rauch E M and Bar-Yam Y 2006 Long-range interactions and evolutionary stability in a predator–prey system *Phys. Rev. E* **73** 020903
- [26] Ruan S and Zhao X Q 1999 Persistence and extinction in two species reaction–diffusion systems with delays *J. Diff. Eqns* **156** 71–92
- [27] Schaaf K W 1987 Asymptotic behavior and traveling wave solutions for parabolic functional differential equations *Trans. Am. Math. Soc.* **302** 587–615
- [28] Silvertown J, Holtier S, Johnson J and Dale P 1992 Cellular automation models of interspecific competition for space—the effect of pattern on process *J. Ecol.* **80** 527–34

- [29] Tang M M and Fife P 1980 Propagating fronts for competing species equations with diffusion *Arch. Ration. Mech. Anal.* **73** 69–77
- [30] van Vuuren J H 1995 The existence of traveling plane waves in a general class of competition–diffusion systems *IMA J. Appl. Math.* **55** 135–48
- [31] Wang Z C, Li W T and Ruan S 2006 Traveling wave fronts in reaction–diffusion systems with spatio-temporal delays *J. Diff. Eqns* **222** 185–232
- [32] Wu J 1996 *Theory and Applications of Partial Functional Differential Equations* (New York: Springer)
- [33] Wu J and Zou X 2001 Traveling wave fronts of reaction–diffusion systems with delay *J. Dyn. Diff. Eqns* **13** 651–87
- [34] Ye Q X and Li Z Y 1994 *Introduction to Reaction Diffusion Equations* (Beijing: Science Press)
- [35] Zeidler E 1986 *Nonlinear Functional Analysis and its Applications: I, Fixed-Point Theorems* (New York: Springer)
- [36] Zou X and Wu J 1997 Existence of traveling wave fronts in delayed reaction–diffusion systems via the monotone iteration method *Proc. Am. Math. Soc.* **125** 2589–98