

RESEARCH

Open Access

Existence on positive solutions for boundary value problems of nonlinear fractional differential equations with p -Laplacian

Hongling Lu, Zhenlai Han*, Shurong Sun and Jian Liu

*Correspondence:
hanzhenlai@163.com
School of Mathematical Sciences,
University of Jinan, Jinan, Shandong
250022, P.R. China

Abstract

In this paper, we study the existence of positive solutions for the nonlinear fractional boundary value problem with a p -Laplacian operator

$$\begin{aligned} D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}u(t))) &= f(t, u(t)), \quad 0 < t < 1, \\ u(0) = u'(0) = u'(1) &= 0, \quad D_{0+}^{\alpha}u(0) = D_{0+}^{\alpha}u(1) = 0, \end{aligned}$$

where $2 < \alpha \leq 3$, $1 < \beta \leq 2$, D_{0+}^{α} , D_{0+}^{β} are the standard Riemann-Liouville fractional derivatives, $\phi_p(s) = |s|^{p-2}s$, $p > 1$, $\phi_p^{-1} = \phi_q$, $1/p + 1/q = 1$, and $f(t, u) \in C([0, 1] \times [0, +\infty), [0, +\infty))$. By the properties of Green's function, the Guo-Krasnosel'skii fixed-point theorem, the Leggett-Williams fixed-point theorem, and the upper and lower solutions method, some new results on the existence of positive solutions are obtained. As applications, examples are presented to illustrate the main results.

MSC: 34A08; 34B18; 35J05

Keywords: fractional boundary value problem; positive solution; upper and lower solutions; fixed-point theorems; p -Laplacian operator

1 Introduction

Recently, fractional differential equations have been of great interest. The motivation for those works stems from both the intensive development of the theory of fractional calculus itself and the applications such as economics, engineering and other fields [1–7]. Many people pay attention to the existence and multiplicity of solutions or positive solutions for boundary value problems of nonlinear fractional differential equations by means of some fixed-point theorems [8–24] (such as the Schauder fixed-point theorem, the Guo-Krasnosel'skii fixed-point theorem, the Leggett-Williams fixed-point theorem) and the upper and lower solutions method [25–27].

To the best of our knowledge, there are few papers devoted to the study of fractional differential equations with a p -Laplacian operator [22–24, 26–29]. Its theories and applications seem to be just being initiated.

Wang *et al.* [26] considered the following p -Laplacian fractional differential equations boundary value problems:

$$D_{0+}^{\gamma}(\phi_p(D_{0+}^{\alpha}u(t))) = f(t, u(t)), \quad 0 < t < 1,$$

$$u(0) = 0, \quad u(1) = au(\xi), \quad D_{0+}^{\alpha}u(0) = 0, \quad D_{0+}^{\alpha}u(1) = bD_{0+}^{\alpha}u(\eta),$$

where $1 < \alpha, \gamma \leq 2$, $0 \leq a, b \leq 1$, $0 < \xi, \eta < 1$, and D_{0+}^{α} is the standard Riemann-Liouville fractional derivative. $\phi_p(s) = |s|^{p-2}s$, $p > 1$, $\phi_p^{-1} = \phi_q$, $1/p + 1/q = 1$. They obtained the existence of at least one positive solution by means of the upper and lower solutions method.

Wang *et al.* [24] investigated the existence and multiplicity of concave positive solutions of a boundary value problem of a fractional differential equation with a p -Laplacian operator as follows:

$$D_{0+}^{\gamma}(\phi_p(D_{0+}^{\alpha}u(t))) + f(t, u(t), D_{0+}^{\rho}u(t)) = 0, \quad 0 < t < 1,$$

$$u(0) = u'(1) = 0, \quad u''(0) = 0, \quad D_{0+}^{\alpha}u(0) = 0,$$

where $2 < \alpha < 3$, $0 < \gamma < 1$, $0 < \rho \leq 1$, D_{0+}^{α} is the Caputo derivative. By using a fixed-point theorem, some results for multiplicity of concave positive solutions are obtained.

Chen *et al.* [28] considered the boundary value problem for a fractional differential equation with a p -Laplacian operator at resonance

$$D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}x(t))) = f(t, x(t), D_{0+}^{\alpha}x(t)), \quad t \in [0, 1],$$

$$D_{0+}^{\alpha}x(0) = D_{0+}^{\alpha}x(1) = 0,$$

where $0 < \alpha, \beta \leq 1$, $1 < \alpha + \beta \leq 2$, and D_{0+}^{α} is the Caputo fractional derivative. By using the coincidence degree theory, a new result on the existence of solutions is obtained.

Guoqing Chai [29] investigated the existence and multiplicity of positive solutions for a class of boundary value problems of fractional differential equations with a p -Laplacian operator

$$D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}u(t))) + f(t, u(t)) = 0, \quad 0 < t < 1,$$

$$u(0) = 0, \quad u(1) + \sigma D_{0+}^{\gamma}u(1) = 0, \quad D_{0+}^{\alpha}u(0) = 0,$$

where $1 < \alpha \leq 2$, $0 < \beta \leq 1$, $0 < \gamma \leq 1$, $0 \leq \alpha - \gamma - 1$, σ is a positive constant number, D_{0+}^{α} , D_{0+}^{β} , D_{0+}^{γ} are the standard Riemann-Liouville derivatives. By means of the fixed-point theorem on cones, some existence and multiplicity results of positive solutions are obtained.

Motivated by all the works above, in this paper, we deal with the following p -Laplacian fractional differential equation boundary value problem:

$$D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}u(t))) = f(t, u(t)), \quad 0 < t < 1, \tag{1.1}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^{\alpha}u(0) = D_{0+}^{\alpha}u(1) = 0, \tag{1.2}$$

where $2 < \alpha \leq 3$, $1 < \beta \leq 2$, D_{0+}^{α} , D_{0+}^{β} are the standard Riemann-Liouville fractional derivatives and $f(t, u) \in C([0, 1] \times [0, +\infty), [0, +\infty))$. By the properties of Green's function, the Guo-Krasnosel'skii fixed-point theorem, the Leggett-Williams fixed-point theorem, and

the upper and lower solutions method, some new results on the existence of positive solutions are obtained for the fractional differential equation boundary value problem (1.1) and (1.2).

The rest of this paper is organized as follows. In Section 2, we introduce some definitions and lemmas to prove our main results. In Section 3, we investigate the existence of a single positive solution for boundary value problems (1.1) and (1.2) by the upper and lower solutions method. In Section 4, we establish the existence of single and multiple positive solutions for boundary value problems (1.1) and (1.2) by fixed-point theorems. As applications, examples are presented to illustrate our main results in Section 3 and Section 4, respectively.

2 Preliminaries and lemmas

For the convenience of the reader, we give some background material from fractional calculus theory to facilitate the analysis of problem (1.1) and (1.2). These materials can be found in the recent literature, see [7, 8, 26, 27, 30–33].

Definition 2.1 [7] The fractional integral of order $\alpha > 0$ of a function $y : (0, +\infty) \rightarrow \mathbb{R}$ is given by

$$I_{0+}^{\alpha}y(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}y(s) ds$$

provided the right-hand side is pointwise defined on $(0, +\infty)$.

Definition 2.2 [7] The fractional derivative of order $\alpha > 0$ of a continuous function $y : (0, +\infty) \rightarrow \mathbb{R}$ is given by

$$D_{0+}^{\alpha}y(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt}\right)^n \int_0^t \frac{y(s)}{(t-s)^{\alpha-n+1}} ds,$$

where n is the smallest integer greater than or equal to α , provided that the right-hand side is pointwise defined on $(0, +\infty)$.

Lemma 2.1 [7] *Let $\alpha > 0$. If we assume $u \in D_{0+}^{\alpha}u \in L^1(0,1)$, then the fractional differential equation*

$$D_{0+}^{\alpha}u(t) = 0$$

has

$$u(t) = c_1t^{\alpha-1} + c_2t^{\alpha-2} + \dots + c_nt^{\alpha-n}, \quad c_i \in \mathbb{R}, i = 1, 2, \dots, n \text{ as a unique solution,}$$

where n is the smallest integer greater than or equal to α .

Lemma 2.2 [7] *Assume that $D_{0+}^{\alpha}u \in L^1(0,1)$ with a fractional derivative of order $\alpha > 0$. Then*

$$I_{0+}^{\alpha}D_{0+}^{\alpha}u(t) = u(t) + c_1t^{\alpha-1} + c_2t^{\alpha-2} + \dots + c_nt^{\alpha-n}$$

for some $c_i \in \mathbb{R}, i = 1, 2, \dots, n$, where n is the smallest integer greater than or equal to α .

Lemma 2.3 [27] *Let $y \in C[0,1]$ and $2 < \alpha \leq 3$. Then fractional differential equation boundary value problem*

$$D_{0+}^{\alpha} u(t) + y(t) = 0, \quad 0 < t < 1, \tag{2.1}$$

$$u(0) = u'(0) = u'(1) = 0 \tag{2.2}$$

has a unique solution

$$u(t) = \int_0^1 G(t,s)y(s) ds,$$

where

$$G(t,s) = \begin{cases} \frac{t^{\alpha-1}(1-s)^{\alpha-2}-(t-s)^{\alpha-1}}{\Gamma(\alpha)}, & s \leq t, \\ \frac{t^{\alpha-1}(1-s)^{\alpha-2}}{\Gamma(\alpha)}, & t \leq s. \end{cases} \tag{2.3}$$

Lemma 2.4 *Let $y \in C[0,1]$ and $2 < \alpha \leq 3, 1 < \beta \leq 2$. Then the fractional differential equation boundary value problem*

$$D_{0+}^{\beta} (\phi_p(D_{0+}^{\alpha} u(t))) = y(t), \quad 0 < t < 1, \tag{2.4}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^{\alpha} u(0) = D_{0+}^{\alpha} u(1) = 0 \tag{2.5}$$

has a unique solution

$$u(t) = \int_0^1 G(t,s)\phi_q\left(\int_0^1 H(s,\tau)y(\tau) d\tau\right) ds,$$

where

$$H(t,s) = \begin{cases} \frac{[t(1-s)]^{\beta-1}-(t-s)^{\beta-1}}{\Gamma(\beta)}, & s \leq t, \\ \frac{[t(1-s)]^{\beta-1}}{\Gamma(\beta)}, & t \leq s, \end{cases} \tag{2.6}$$

$G(t,s)$ is defined as (2.3).

Proof From Lemma 2.2 and $1 < \beta \leq 2$, we have

$$I_{0+}^{\beta} D_{0+}^{\beta} (\phi_p(D_{0+}^{\alpha} u(t))) = \phi_p(D_{0+}^{\alpha} u(t)) + c_1 t^{\beta-1} + c_2 t^{\beta-2} \quad \text{for some } c_1, c_2 \in \mathbb{R}.$$

In view of (2.4), we obtain

$$I_{0+}^{\beta} D_{0+}^{\beta} (\phi_p(D_{0+}^{\alpha} u(t))) = I_{0+}^{\beta} y(t).$$

Therefore,

$$\phi_p(D_{0+}^{\alpha} u(t)) = I_{0+}^{\beta} y(t) + C_1 t^{\beta-1} + C_2 t^{\beta-2} \quad \text{for some } C_1, C_2 \in \mathbb{R},$$

that is,

$$\phi_p(D_{0+}^\alpha u(t)) = \int_0^t \frac{(t-\tau)^{\beta-1}}{\Gamma(\beta)} y(\tau) d\tau + C_1 t^{\beta-1} + C_2 t^{\beta-2}.$$

By the boundary conditions $D_{0+}^\alpha u(0) = D_{0+}^\alpha u(1) = 0$, we have

$$C_2 = 0, \quad C_1 = - \int_0^1 \frac{(1-\tau)^{\beta-1}}{\Gamma(\beta)} y(\tau) d\tau.$$

Therefore, the solution $u(t)$ of fractional differential equation boundary value problem (2.4) and (2.5) satisfies

$$\begin{aligned} \phi_p(D_{0+}^\alpha u(t)) &= \int_0^t \frac{(t-\tau)^{\beta-1}}{\Gamma(\beta)} y(\tau) d\tau - \int_0^1 \frac{[t(1-\tau)]^{\beta-1}}{\Gamma(\beta)} y(\tau) d\tau \\ &= - \int_0^1 H(t, \tau) y(\tau) d\tau. \end{aligned}$$

Consequently, $D_{0+}^\alpha u(t) + \phi_q(\int_0^1 H(t, \tau) y(\tau) d\tau) = 0$. Thus, fractional differential equation boundary value problem (2.4) and (2.5) is equivalent to the following problem:

$$\begin{aligned} D_{0+}^\alpha u(t) + \phi_q\left(\int_0^1 H(t, \tau) y(\tau) d\tau\right) &= 0, \quad 0 < t < 1, \\ u(0) = u'(0) = u'(1) &= 0. \end{aligned}$$

Lemma 2.3 implies that fractional differential equation boundary value problem (2.4) and (2.5) has a unique solution

$$u(t) = \int_0^1 G(t, s) \phi_q\left(\int_0^1 H(s, \tau) y(\tau) d\tau\right) ds.$$

The proof is complete. □

Lemma 2.5 *Let $2 < \alpha \leq 3, 1 < \beta \leq 2$. The functions $G(t, s)$ and $H(t, s)$ defined by (2.3) and (2.6), respectively, are continuous on $[0, 1] \times [0, 1]$ and satisfy*

- (1) $G(t, s) \geq 0, H(t, s) \geq 0$ for $t, s \in [0, 1]$;
- (2) $G(t, s) \leq G(1, s), H(t, s) \leq H(s, s)$ for $t, s \in [0, 1]$;
- (3) $G(t, s) \geq t^{\alpha-1} G(1, s)$ for $t, s \in (0, 1)$;
- (4) *there exist two positive functions $\delta_1, \delta_2 \in C[0, 1]$ such that*

$$\min_{1/4 \leq t \leq 3/4} G(t, s) \geq \delta_1(s) \max_{0 \leq t \leq 1} G(t, s) = \delta_1(s) G(1, s) \quad \text{for } 0 < s < 1, \tag{2.7}$$

$$\min_{1/4 \leq t \leq 3/4} H(t, s) \geq \delta_2(s) \max_{0 \leq t \leq 1} H(t, s) = \delta_2(s) H(s, s) \quad \text{for } 0 < s < 1. \tag{2.8}$$

Proof Observing the expression of $G(t, s)$ and $H(t, s)$, it is easy to see that $G(t, s) \geq 0$ and $H(t, s) \geq 0$ for $s, t \in [0, 1]$.

From Lemma 3.1 in [27] and Lemma 2.4 in [8], we obtain (2) and (3).

In the following, we consider the existence of $\delta_1(s)$ and $\delta_2(s)$. Firstly, for given $s \in (0, 1)$, $G(t, s)$ is increasing with respect to t for $t \in (0, 1)$. Consequently, setting

$$g_1(t, s) = \frac{t^{\alpha-1}(1-s)^{\alpha-2} - (t-s)^{\alpha-1}}{\Gamma(\alpha)}, \quad g_2(t, s) = \frac{t^{\alpha-1}(1-s)^{\alpha-2}}{\Gamma(\alpha)},$$

we have

$$\begin{aligned} \min_{1/4 \leq t \leq 3/4} G(t, s) &= \begin{cases} g_1(\frac{1}{4}, s), & s \in (0, \frac{1}{4}], \\ g_2(\frac{1}{4}, s), & s \in [\frac{1}{4}, 1), \end{cases} \\ &= \begin{cases} \frac{1}{\Gamma(\alpha)} [(\frac{1}{4})^{\alpha-1}(1-s)^{\alpha-2} - (\frac{1}{4}-s)^{\alpha-1}], & s \in (0, \frac{1}{4}], \\ \frac{1}{\Gamma(\alpha)} (\frac{1}{4})^{\alpha-1}(1-s)^{\alpha-2}, & s \in [\frac{1}{4}, 1). \end{cases} \end{aligned}$$

Secondly, with the use of the monotonicity of $G(t, s)$, we have

$$\max_{0 \leq t \leq 1} G(t, s) = G(1, s) = \frac{1}{\Gamma(\alpha)} [(1-s)^{\alpha-2} - (1-s)^{\alpha-1}].$$

Thus, setting

$$\delta_1(s) = \begin{cases} \frac{(\frac{1}{4})^{\alpha-1}(1-s)^{\alpha-2} - (\frac{1}{4}-s)^{\alpha-1}}{(1-s)^{\alpha-2} - (1-s)^{\alpha-1}}, & s \in (0, \frac{1}{4}], \\ \frac{(\frac{1}{4})^{\alpha-1}(1-s)^{\alpha-2}}{(1-s)^{\alpha-2} - (1-s)^{\alpha-1}}, & s \in [\frac{1}{4}, 1), \end{cases}$$

then (2.7) holds.

Similar to Lemma 2.4 in [8], we choose

$$\delta_2(s) = \begin{cases} \frac{[\frac{3}{4}(1-s)]^{\beta-1} - (\frac{3}{4}-s)^{\beta-1}}{[s(1-s)]^{\beta-1}}, & s \in (0, r], \\ (\frac{1}{4s})^{\beta-1}, & s \in [r, 1). \end{cases}$$

The proof is complete. □

Lemma 2.6 *Let $2 < \alpha \leq 3$. If $y(t) \in C[0, 1]$ and $y(t) \geq 0$, then fractional differential equation boundary value problem (2.1) and (2.2) has a unique solution $u(t) \geq 0$, $t \in [0, 1]$.*

Proof From Lemma 2.3, the fractional differential equation boundary value problem (2.1) and (2.2) has a unique solution

$$u(t) = \int_0^1 G(t, s)y(s) ds.$$

In view of Lemma 2.5, we know $G(t, s)$ is continuous on $[0, 1] \times [0, 1]$ and $G(t, s) \geq 0$ for $t, s \in [0, 1]$. If $y(t) \in C[0, 1]$ and $y(t) \geq 0$, we obtain $u(t) \geq 0$. The proof is complete. □

Let $E_0 = \{u : u \in C^3[0, 1], \phi_p(D_{0+}^\alpha u) \in C^2[0, 1]\}$. Now, we introduce definitions about the upper and lower solutions of fractional differential equation boundary value problem (1.1) and (1.2).

Definition 2.3 [26] A function $\eta(t)$ is called an upper solution of fractional differential equation boundary value problem (1.1) and (1.2) if $\eta(t) \in E_0$ and $\eta(t)$ satisfies

$$\begin{aligned} D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}\eta(t))) &\geq f(t, \eta(t)), \quad 0 < t < 1, 2 < \alpha \leq 3, 1 < \beta \leq 2, \\ \eta(0) &\geq 0, \quad \eta'(0) \geq 0, \quad \eta'(1) \geq 0, \\ D_{0+}^{\alpha}\eta(0) &\leq 0, \quad D_{0+}^{\alpha}\eta(1) \leq 0. \end{aligned}$$

Definition 2.4 [26] A function $\xi(t)$ is called a lower solution of fractional differential equation boundary value problem (1.1) and (1.2) if $\xi(t) \in E_0$ and $\xi(t)$ satisfies

$$\begin{aligned} D_{0+}^{\beta}(\phi_p(D_{0+}^{\alpha}\xi(t))) &\leq f(t, \xi(t)), \quad 0 < t < 1, 2 < \alpha \leq 3, 1 < \beta \leq 2, \\ \xi(0) &\leq 0, \quad \xi'(0) \leq 0, \quad \xi'(1) \leq 0, \\ D_{0+}^{\alpha}\xi(0) &\geq 0, \quad D_{0+}^{\alpha}\xi(1) \geq 0. \end{aligned}$$

Definition 2.5 [8] The map θ is said to be a nonnegative continuous concave functional on a cone P of a real Banach space E provided that $\theta : P \rightarrow [0, +\infty)$ is continuous and

$$\theta(tx + (1-t)y) \geq t\theta(x) + (1-t)\theta(y)$$

for all $x, y \in P$ and $0 \leq t \leq 1$.

Lemma 2.7 [8] Let E be a Banach space, $P \subseteq E$ be a cone, and Ω_1, Ω_2 be two bounded open balls of E centered at the origin with $\bar{\Omega}_1 \subset \Omega_2$. Suppose that $\mathcal{A} : P \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow P$ is a completely continuous operator such that either

- (i) $\|\mathcal{A}x\| \leq \|x\|, x \in P \cap \partial\Omega_1$ and $\|\mathcal{A}x\| \geq \|x\|, x \in P \cap \partial\Omega_2$ or
- (ii) $\|\mathcal{A}x\| \geq \|x\|, x \in P \cap \partial\Omega_1$ and $\|\mathcal{A}x\| \leq \|x\|, x \in P \cap \partial\Omega_2$

holds. Then \mathcal{A} has a fixed point in $P \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

Let $a, b, c > 0$ be constants, $P_c = \{u \in P : \|u\| < c\}$, $P(\theta, b, d) = \{u \in P : b \leq \theta(u), \|u\| \leq d\}$.

Lemma 2.8 [8] Let P be a cone in a real Banach space E , $P_c = \{x \in P \mid \|x\| \leq c\}$, θ be a nonnegative continuous concave functional on P such that $\theta(x) \leq \|x\|$ for all $x \in \bar{P}_c$, and $P(\theta, b, d) = \{x \in P \mid b \leq \theta(x), \|x\| \leq d\}$. Suppose $\mathcal{B} : \bar{P}_c \rightarrow \bar{P}_c$ is completely continuous and there exist constants $0 < a < b < d \leq c$ such that

- (C1) $\{x \in P(\theta, b, d) \mid \theta(x) > b\} \neq \emptyset$ and $\theta(\mathcal{B}x) > b$ for $x \in P(\theta, b, d)$;
- (C2) $\|\mathcal{B}x\| < a$ for $x \leq a$;
- (C3) $\theta(\mathcal{B}x) > b$ for $x \in P(\theta, b, c)$ with $\|\mathcal{B}x\| > d$.

Then \mathcal{B} has at least three fixed points x_1, x_2 , and x_3 with

$$\|x_1\| < a, \quad b < \theta(x_2), \quad a < \|x_3\| \quad \text{with} \quad \theta(x_3) < b.$$

Let $E = C[0, 1]$ be endowed with $\|u\| = \max_{0 \leq t \leq 1} |u(t)|$. Define the cone $P \subset E$ by

$$P = \{u \in E \mid u(t) \geq 0\}.$$

Let the nonnegative continuous concave functional θ on the cone P be defined by

$$\theta(u) = \min_{1/4 \leq t \leq 3/4} |u(t)|.$$

Lemma 2.9 *Let $T : P \rightarrow E$ be the operator defined by*

$$Tu(t) := \int_0^1 G(t,s)\phi_q\left(\int_0^1 H(s,\tau)f(\tau,u(\tau))d\tau\right)ds.$$

Then $T : P \rightarrow P$ is completely continuous.

Proof Let $u \in P$, in view of the nonnegativeness and continuity of $G(t,s)$, $H(t,s)$, and $f(t,u(t))$, we have $T : P \rightarrow P$ is continuous.

Let $\Omega \subset P$ be bounded, i.e., there exists a positive constant $M > 0$ such that $\|u\| \leq M$ for all $u \in \Omega$. Let $L = \max_{0 \leq t \leq 1, 0 \leq u \leq M} |f(t,u)| + 1$, then for $u \in \Omega$, we have

$$\begin{aligned} |Tu(t)| &= \left| \int_0^1 G(t,s)\phi_q\left(\int_0^1 H(s,\tau)f(\tau,u(\tau))d\tau\right)ds \right| \\ &\leq L^{q-1} \int_0^1 G(t,s)\phi_q\left(\int_0^1 H(s,\tau)d\tau\right)ds \\ &\leq L^{q-1} \int_0^1 G(1,s)\phi_q\left(\int_0^1 H(\tau,\tau)d\tau\right)ds \\ &< +\infty. \end{aligned}$$

Hence, $T(\Omega)$ is uniformly bounded.

On the other hand, since $G(t,s)$ is continuous on $[0,1] \times [0,1]$, it is uniformly continuous on $[0,1] \times [0,1]$. Thus, for fixed $s \in [0,1]$ and for any $\varepsilon > 0$, there exists a constant $\delta > 0$ such that any $t_1, t_2 \in [0,1]$ and $|t_1 - t_2| < \delta$,

$$|G(t_1,s) - G(t_2,s)| < \frac{\varepsilon}{L^{q-1}\phi_q\left(\int_0^1 H(\tau,\tau)d\tau\right)}.$$

Then, for all $u \in \Omega$,

$$\begin{aligned} &|Tu(t_2) - Tu(t_1)| \\ &\leq \int_0^1 |G(t_2,s) - G(t_1,s)|\phi_q\left(\int_0^1 H(s,\tau)f(\tau,u(\tau))d\tau\right)ds \\ &\leq L^{q-1} \int_0^1 |G(t_2,s) - G(t_1,s)|\phi_q\left(\int_0^1 H(\tau,\tau)d\tau\right)ds \\ &= L^{q-1}\phi_q\left(\int_0^1 H(\tau,\tau)d\tau\right) \int_0^1 |G(t_2,s) - G(t_1,s)|ds \\ &< \varepsilon, \end{aligned}$$

that is to say, $T(\Omega)$ is equicontinuous. By the Arzela-Ascoli theorem, we have $T : P \rightarrow P$ is completely continuous. The proof is complete. \square

3 Existence of a single positive solution

In this section, for the sake of simplicity, we assume that

(H₁) $f(t, u)$ is nonincreasing to u ;

(H₂) There exists a continuous function $p(t) \geq 0, t \in [0, 1]$ such that

$$\int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, p(\tau)) d\tau\right) ds = q(t) \geq p(t), \tag{3.1}$$

$$\int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, q(\tau)) d\tau\right) ds \geq p(t). \tag{3.2}$$

Theorem 3.1 *Assume that (H₁) and (H₂) hold. Then the fractional differential equation boundary value problem (1.1) and (1.2) has at least one positive solution $\gamma(t)$.*

Proof From Lemma 2.9, we obtain $T(P) \subseteq P$. By direct computations, we have

$$D_{0+}^\beta(\phi_p(D_{0+}^\alpha(Tu)(t))) = f(t, u(t)), \quad 0 < t < 1, 2 < \alpha \leq 3, 1 < \beta \leq 2, \tag{3.3}$$

$$(Tu)(0) = (Tu)'(0) = (Tu)'(1) = 0, \quad D_{0+}^\alpha(Tu)(0) = D_{0+}^\alpha(Tu)(1) = 0. \tag{3.4}$$

Now, we prove that the functions $\eta(t) = Tp(t), \xi(t) = Tq(t)$ are upper and lower solutions of fractional differential equation boundary value problem (1.1) and (1.2), respectively.

From (H₁) and (H₂), we have

$$p(t) \leq q(t) = Tp(t), \quad Tq(t) \leq q(t) = Tp(t), \quad t \in [0, 1]. \tag{3.5}$$

Hence, $\xi(t) \leq \eta(t)$. By $T(P) \subseteq P$, we know $\xi(t), \eta(t) \in P$. From (3.3)-(3.5) we have

$$D_{0+}^\beta(\phi_p(D_{0+}^\alpha(\eta)(t))) - f(t, \eta(t)) \geq D_{0+}^\beta(\phi_p(D_{0+}^\alpha(Tp)(t))) - f(t, p(t)) = 0, \tag{3.6}$$

$$\eta(0) = \eta'(0) = \eta'(1) = 0, \quad D_{0+}^\alpha \eta(0) = D_{0+}^\alpha \eta(1) = 0,$$

$$D_{0+}^\beta(\phi_p(D_{0+}^\alpha(\xi)(t))) - f(t, \xi(t)) \leq D_{0+}^\beta(\phi_p(D_{0+}^\alpha(Tq)(t))) - f(t, q(t)) = 0, \tag{3.7}$$

$$\xi(0) = \xi'(0) = \xi'(1) = 0, \quad D_{0+}^\alpha \xi(0) = D_{0+}^\alpha \xi(1) = 0,$$

that is, $\eta(t)$ and $\xi(t)$ are upper and lower solutions of fractional differential equation boundary value problem (1.1) and (1.2), respectively.

Next, we show that the fractional differential equation boundary value problem

$$D_{0+}^\beta(\phi_p(D_{0+}^\alpha u(t))) = g(t, u(t)), \quad 0 < t < 1, 2 < \alpha \leq 3, 1 < \beta \leq 2, \tag{3.8}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^\alpha u(0) = D_{0+}^\alpha u(1) = 0 \tag{3.9}$$

has a positive solution, where

$$g(t, u(t)) = \begin{cases} f(t, \xi(t)), & \text{if } u(t) \leq \xi(t), \\ f(t, u(t)), & \text{if } \xi(t) \leq u(t) \leq \eta(t), \\ f(t, \eta(t)), & \text{if } \eta(t) \leq u(t). \end{cases} \tag{3.10}$$

Thus, we consider the operator $B : P \rightarrow E$ defined as follows:

$$Bu(t) := \int_0^1 G(t,s)\phi_q \left(\int_0^1 H(s,\tau)g(\tau,u(\tau)) d\tau \right) ds,$$

where $G(t,s)$ and $H(t,s)$ are defined as (2.3) and (2.6), respectively. It is clear that $Bu(t) \geq 0$ for all $u \in P$ and a fixed point of the operator B is a solution of the fractional differential equation boundary value problem (3.8) and (3.9).

Similar to Lemma 2.9, we know that B is a compact operator. By the Schauder fixed-point theorem, the operator B has a fixed point, that is, the fractional differential equation boundary value problem (3.8) and (3.9) has a positive solution.

Finally, we will prove that fractional differential equation boundary value problem (1.1) and (1.2) has at least one positive solution.

Suppose that $\gamma(t)$ is a solution of (3.8) and (3.9). Now, to complete the proof, it suffices to show that $\xi(t) \leq \gamma(t) \leq \eta(t)$, $t \in [0, 1]$.

Let $\gamma(t)$ be a solution of (3.8) and (3.9). We have

$$\gamma(0) = \gamma'(0) = \gamma'(1) = 0, \quad D_{0+}^\alpha \gamma(0) = D_{0+}^\alpha \gamma(1) = 0. \tag{3.11}$$

From (H_1) , we have

$$f(t, \eta(t)) \leq g(t, \gamma(t)) \leq f(t, \xi(t)), \quad t \in [0, 1]. \tag{3.12}$$

By (H_2) and (3.5), we obtain

$$f(t, q(t)) \leq g(t, \gamma(t)) \leq f(t, p(t)), \quad t \in [0, 1]. \tag{3.13}$$

By $p(t) \in P$ and (3.3), we can get

$$D_{0+}^\beta (\phi_p(D_{0+}^\alpha \eta(t))) = D_{0+}^\beta (\phi_p(D_{0+}^\alpha (Tp)(t))) = f(t, p(t)), \quad t \in [0, 1]. \tag{3.14}$$

Combining (3.4), (3.11)-(3.14), we have

$$D_{0+}^\beta (\phi_p(D_{0+}^\alpha \eta(t))) - D_{0+}^\beta (\phi_p(D_{0+}^\alpha \gamma(t))) = f(t, p(t)) - g(t, \gamma(t)) \geq 0, \quad t \in [0, 1], \tag{3.15}$$

$$(\eta - \gamma)(0) = (\eta - \gamma)'(0) = (\eta - \gamma)'(1) = 0, \tag{3.16}$$

$$D_{0+}^\alpha (\eta - \gamma)(0) = D_{0+}^\alpha (\eta - \gamma)(1) = 0.$$

Let $x(t) = \phi_p(D_{0+}^\alpha \eta(t)) - \phi_p(D_{0+}^\alpha \gamma(t))$. By (3.16), we obtain $x(0) = x(1) = 0$.

By Lemma 2.6, we know $x(t) \leq 0$, $t \in [0, 1]$, which implies that

$$\phi_p(D_{0+}^\alpha \eta(t)) \leq \phi_p(D_{0+}^\alpha \gamma(t)), \quad t \in [0, 1].$$

Since ϕ_p is monotone increasing, we obtain $D_{0+}^\alpha \eta(t) \leq D_{0+}^\alpha \gamma(t)$, that is, $D_{0+}^\alpha (\eta - \gamma)(t) \leq 0$. By Lemma 2.6, (3.15) and (3.16), we have $(\eta - \gamma)(t) \geq 0$. Therefore, $\eta(t) \geq \gamma(t)$, $t \in [0, 1]$.

In a similar way, we can prove that $\xi(t) \leq \gamma(t)$, $t \in [0, 1]$. Consequently, $\gamma(t)$ is a positive solution of fractional differential equation boundary value problem (1.1) and (1.2). This completes the proof. \square

Example 3.1 We consider the following fractional differential equation boundary value problem:

$$D_{0+}^{4/3}(\phi_p(D_{0+}^{5/2}u(t))) = t^2 + \frac{1}{\sqrt{u}}, \quad 0 < t < 1, \tag{3.17}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^{5/2}u(0) = D_{0+}^{5/2}u(1) = 0. \tag{3.18}$$

Clearly, $f(t, u) = t^2 + \frac{1}{\sqrt{u}}$ is nonincreasing relative to u . This shows that (H_1) holds.

Let $m(t) = t^{3/2}$. From Lemma 2.5, we have

$$\begin{aligned} n(t) &:= Tm(t) = \int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, m(\tau)) d\tau\right) ds \\ &= \int_0^1 \frac{G(t, s)}{G(1, s)}G(1, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, m(\tau)) d\tau\right) ds \\ &\geq t^{\alpha-1} \int_0^1 G(1, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, m(\tau)) d\tau\right) ds \end{aligned}$$

and $Tn(t) = T^2m(t) \in P$, there exist positive numbers d_1 and d_2 such that $Tm(t) \geq d_1m(t)$ and $T^2m(t) \geq d_2m(t)$.

Choosing a positive number $d_0 \leq \{1, d_1\}$ and combining the monotonicity of T , we have

$$T(d_0m(t)) \geq Tm(t) \geq d_1m(t) \geq d_0m(t), \quad T^2(d_0m(t)) \geq T(d_0m(t)) \geq d_0m(t).$$

Taking $p(t) = d_0t^{3/2}$, then we have

$$\begin{aligned} q(t) &= Tp(t) = \int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, d_0\tau^{3/2}) d\tau\right) ds \\ &\geq d_0t^{3/2} = p(t), \\ Tq(t) &= T^2p(t) = \int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, T(d_0\tau^{3/2})) d\tau\right) ds \\ &\geq d_0t^{3/2} = p(t). \end{aligned}$$

That is, the condition (H_2) holds. By Theorem 3.1, the fractional differential equation boundary value problem (3.17) and (3.18) has at least one positive solution.

4 Existence of single and multiple positive solutions

In this section, for convenience, we denote

$$\begin{aligned} M &= \left(\int_0^1 G(1, s)\phi_q\left(\int_0^1 H(\tau, \tau) d\tau\right) ds\right)^{-1}, \\ N &= \left(\int_{1/4}^{3/4} \delta_1(s)G(1, s)\phi_q\left(\int_{1/4}^{3/4} \delta_2(\tau)H(\tau, \tau) d\tau\right) ds\right)^{-1}. \end{aligned}$$

Theorem 4.1 Let $f(t, u)$ be continuous on $[0, 1] \times [0, +\infty)$. Assume that there exist two positive constants $a_2 > a_1 > 0$ such that

(A1) $f(t, u) \geq \phi_p(Na_1)$ for $(t, u) \in [0, 1] \times [0, a_1]$;

(A2) $f(t, u) \leq \phi_p(Ma_2)$ for $(t, u) \in [0, 1] \times [0, a_2]$.

Then the fractional differential equation boundary value problem (1.1) and (1.2) has at least one positive solution u such that $a_1 \leq \|u\| \leq a_2$.

Proof From Lemmas 2.3, 2.4, and 2.9, we get that $T : P \rightarrow P$ is completely continuous and fractional differential equation boundary value problem (1.1) and (1.2) has a solution $u = u(t)$ if and only if u solves the operator equation $u = Tu(t)$. In order to apply Lemma 2.7, we divide our proof into two steps.

Step 1. Let $\Omega_1 := \{u \in P \mid \|u\| < a_1\}$. For $u \in \partial\Omega_1$, we have $0 \leq u(t) \leq a_1$ for all $t \in [0, 1]$. It follows from (A1) that for $t \in [1/4, 3/4]$,

$$\begin{aligned} Tu(t) &= \int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, u(\tau)) d\tau\right) ds \\ &\geq \int_0^1 \delta_1(s)G(1, s)\phi_q\left(\int_0^1 \delta_2(\tau)H(\tau, \tau)f(\tau, u(\tau)) d\tau\right) ds \\ &\geq Na_1 \int_{1/4}^{3/4} \delta_1(s)G(1, s)\phi_q\left(\int_{1/4}^{3/4} \delta_2(\tau)H(\tau, \tau) d\tau\right) ds \\ &= a_1 = \|u\|. \end{aligned}$$

So,

$$\|Tu\| \geq \|u\| \quad \text{for } u \in \partial\Omega_1.$$

Step 2. Let $\Omega_2 := \{u \in P \mid \|u\| < a_2\}$. For $u \in \partial\Omega_2$, we have $0 \leq u(t) \leq a_2$ for all $t \in [0, 1]$. It follows from (A2) that for $t \in [0, 1]$,

$$\begin{aligned} \|Tu(t)\| &= \max_{0 \leq t \leq 1} \int_0^1 G(t, s)\phi_q\left(\int_0^1 H(s, \tau)f(\tau, u(\tau)) d\tau\right) ds \\ &\leq Ma_2 \int_0^1 G(1, s)\phi_q\left(\int_0^1 H(\tau, \tau) d\tau\right) ds \\ &= a_2 = \|u\|. \end{aligned}$$

Therefore,

$$\|Tu\| \leq \|u\| \quad \text{for } u \in \partial\Omega_2.$$

Then, by (ii) of Lemma 2.7, we complete the proof. □

Example 4.1 We consider the following fractional differential equation boundary value problem:

$$D_{0+}^{3/2}(\phi_p(D_{0+}^{5/2}u(t))) = \frac{5}{2} + \frac{\sqrt{u}}{10} + \frac{t^2}{4}, \quad 0 < t < 1, \tag{4.1}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^{5/2}u(0) = D_{0+}^{5/2}u(1) = 0. \tag{4.2}$$

Let $p = 2$. By a simple computation, we obtain $M = 11.25$, $N \approx 736.6099$. Choosing $a_1 = 0.003$, $a_2 = 0.25$, therefore

$$f(t, u) = \frac{5}{2} + \frac{\sqrt{u}}{10} + \frac{t^2}{4} \geq 2.5 > \phi_p(Na_1) \approx 2.2098 \quad \text{for } (t, u) \in [0, 1] \times [0, 0.003],$$

$$f(t, u) = \frac{5}{2} + \frac{\sqrt{u}}{10} + \frac{t^2}{4} \leq 2.8 < \phi_p(Ma_2) = 2.8125 \quad \text{for } (t, u) \in [0, 1] \times [0, 0.25].$$

With the use of Theorem 4.1, the fractional differential equation boundary value problem (4.1) and (4.2) has at least one solution u such that $0.003 \leq \|u\| \leq 0.25$.

Theorem 4.2 *Let $f(t, u)$ be continuous on $[0, 1] \times [0, +\infty)$. Assume that there exist constants $0 < a < b < c$ such that the following assumptions hold:*

- (B1) $f(t, u) < \phi_p(Ma)$ for $(t, u) \in [0, 1] \times [0, a]$;
- (B2) $f(t, u) \geq \phi_p(Nb)$ for $(t, u) \in [1/4, 3/4] \times [b, c]$;
- (B3) $f(t, u) \leq \phi_p(Mc)$ for $(t, u) \in [0, 1] \times [0, c]$.

Then the fractional differential equation boundary value problem (1.1) and (1.2) has at least three positive solutions u_1, u_2 , and u_3 with

$$\max_{0 \leq t \leq 1} |u_1(t)| < a, \quad b < \min_{1/4 \leq t \leq 3/4} |u_2(t)| < \max_{0 \leq t \leq 1} |u_2(t)| \leq c,$$

$$a < \max_{0 \leq t \leq 1} |u_3(t)| \leq c, \quad \min_{1/4 \leq t \leq 3/4} |u_3(t)| < b.$$

Proof From Lemmas 2.3, 2.4, and 2.9, we have $T : P \rightarrow P$ is completely continuous and fractional differential equation boundary value problem (1.1) and (1.2) has a solution $u = u(t)$ if and only if u satisfies the operator equation $u = Tu(t)$.

We show that all the conditions of Lemma 2.8 are satisfied. If $u \in \bar{P}_c$, then $\|u\| \leq c$. By (B3), we have

$$\begin{aligned} \|Tu(t)\| &= \max_{0 \leq t \leq 1} \left| \int_0^1 G(t, s) \phi_q \left(\int_0^1 H(s, \tau) f(\tau, u(\tau)) d\tau \right) ds \right| \\ &\leq Mc \int_0^1 G(1, s) \phi_q \left(\int_0^1 H(\tau, \tau) d\tau \right) ds \\ &\leq c. \end{aligned}$$

Hence, $T : \bar{P}_c \rightarrow \bar{P}_c$. In the same way, if $u \in \bar{P}_a$, then assumption (B1) yields $\|Tu\| < a$. Therefore, condition (C2) of Lemma 2.8 is satisfied.

To check condition (C1) of Lemma 2.6, we choose $u(t) = (b + c)/2$, $0 \leq t \leq 1$. It is easy to see that $u(t) = (b + c)/2 \in P(\theta, b, c)$, $\theta(u) = \theta((b + c)/2) > b$; consequently, $\{u \in P(\theta, b, c) \mid \theta(u) > b\} \neq \emptyset$. Hence, if $u \in P(\theta, b, c)$, then $b \leq u(t) \leq c$ for $1/4 \leq t \leq 3/4$. From assumption (B2), we have $f(t, u(t)) \geq \phi_p(Nb)$ for $1/4 \leq t \leq 3/4$. So,

$$\begin{aligned} \theta(Tu) &= \min_{1/4 \leq t \leq 3/4} |(Tu)(t)| \geq \min_{1/4 \leq t \leq 3/4} \int_0^1 G(t, s) \phi_q \left(\int_0^1 H(s, \tau) f(\tau, u(\tau)) d\tau \right) ds \\ &> \int_{1/4}^{3/4} \min_{1/4 \leq t \leq 3/4} G(t, s) \phi_q \left(\int_{1/4}^{3/4} \min_{1/4 \leq t \leq 3/4} H(s, \tau) f(\tau, u(\tau)) d\tau \right) ds \end{aligned}$$

$$\begin{aligned} &\geq Nb \int_{1/4}^{3/4} \delta_1(s)G(1,s)\phi_q\left(\int_{1/4}^{3/4} \delta_2(\tau)H(\tau,\tau) d\tau\right) ds \\ &= b, \end{aligned}$$

i.e., $\theta(Tu) > b$ for all $u \in P(\theta, b, c)$. Choosing $d = c$, this shows that condition (C1) of Lemma 2.8 is also satisfied.

In the same way, if $u \in P(\theta, b, c)$ and $\|Tu\| > c = d$, we also obtain $\theta(Tu) > b$. Then condition (C3) of Lemma 2.8 is also satisfied.

By Lemma 2.8, the fractional differential equation boundary value problem (1.1) and (1.2) has at least three positive solutions u_1, u_2 , and u_3 , satisfying

$$\begin{aligned} \max_{0 \leq t \leq 1} |u_1(t)| &< a, & b &< \min_{1/4 \leq t \leq 3/4} |u_2(t)|, \\ a &< \max_{0 \leq t \leq 1} |u_3(t)|, & \min_{1/4 \leq t \leq 3/4} |u_3(t)| &< b. \end{aligned}$$

The proof is complete. □

Example 4.2 We consider the following fractional differential equation boundary value problem:

$$D_{0+}^{3/2}(\phi_p(D_{0+}^{5/2}u(t))) = f(t, u), \quad 0 < t < 1, \tag{4.3}$$

$$u(0) = u'(0) = u'(1) = 0, \quad D_{0+}^{5/2}u(0) = D_{0+}^{5/2}u(1) = 0, \tag{4.4}$$

where

$$f(t, u) = \begin{cases} \frac{t}{100} + 737u^2 & \text{for } u \leq 1, \\ \frac{t}{100} + u + 736 & \text{for } u > 1. \end{cases}$$

Let $p = 2$. We obtain $M = 11.25, N \approx 736.6099$. Choosing $a = 0.01, b = 1, c = 72$, therefore

$$f(t, u) = \frac{t}{100} + 737u^2 \leq 0.0837 < \phi_2(Ma) = 0.1125 \quad \text{for } (t, u) \in [0, 1] \times [0, 0.01],$$

$$\begin{aligned} f(t, u) &= \frac{t}{100} + u + 736 \geq 737.0025 > \phi_2(Nb) \\ &\approx 736.6099 \quad \text{for } (t, u) \in [1/4, 3/4] \times [1, 72], \end{aligned}$$

$$f(t, u) = \frac{t}{100} + u + 736 \leq 809.001 < \phi_2(Mc) = 810 \quad \text{for } (t, u) \in [0, 1] \times [0, 72].$$

With the use of Theorem 4.2, the fractional differential equation boundary value problem (4.3) and (4.4) has at least three positive solutions u_1, u_2 , and u_3 with

$$\begin{aligned} \max_{0 \leq t \leq 1} |u_1(t)| &< 0.01, & 1 &< \min_{1/4 \leq t \leq 3/4} |u_2(t)| < \max_{0 \leq t \leq 1} |u_2(t)| \leq 72, \\ 0.01 &< \max_{0 \leq t \leq 1} |u_3(t)| \leq 72, & \min_{1/4 \leq t \leq 3/4} |u_3(t)| &< 1. \end{aligned}$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors declare that the study was realized in collaboration with the same responsibility. All authors read and approved the final manuscript.

Acknowledgements

The authors sincerely thank the reviewers for their valuable suggestions and useful comments that have led to the present improved version of the original manuscript. This research is supported by the Natural Science Foundation of China (11071143, 60904024, 61174217), Natural Science Outstanding Youth Foundation of Shandong Province (JQ201119) and supported by Shandong Provincial Natural Science Foundation (ZR2012AM009, ZR2010AL002, ZR2011AL007), also supported by Natural Science Foundation of Educational Department of Shandong Province (J11LA01).

Received: 31 October 2012 Accepted: 21 January 2013 Published: 11 February 2013

References

1. Oldham, KB, Spanier, J: *The Fractional Calculus*. Academic Press, New York (1974)
2. Podlubny, I: *Fractional Differential Equations*. Mathematics in Science and Engineering. Academic Press, New York (1999)
3. Agrawal, OP: Formulation of Euler-Lagrange equations for fractional variational problems. *J. Math. Anal. Appl.* **272**(1), 368-379 (2002)
4. Weitzner, H, Zaslavsky, GM: Some applications of fractional equations. *Commun. Nonlinear Sci. Numer. Simul.* **8**(3-4), 273-281 (2003)
5. Meral, FC, Royston, TJ, Magin, R: Fractional calculus in viscoelasticity: an experimental study. *Commun. Nonlinear Sci. Numer. Simul.* **15**(4), 939-945 (2010)
6. Machado, JT, Kiryakova, V, Mainardi, F: Recent history of fractional calculus. *Commun. Nonlinear Sci. Numer. Simul.* **16**(3), 1140-1153 (2011)
7. Kilbas, AA, Srivastava, HH, Trujillo, JJ: *Theory and Applications of Fractional Differential Equations*. Elsevier, Amsterdam (2006)
8. Bai, Z, Lü, H: Positive solutions for boundary value problem of nonlinear fractional differential equation. *J. Math. Anal. Appl.* **311**(2), 495-505 (2005)
9. Wang, G, Zhang, L, Ntouyas, SK: Existence of multiple positive solutions of a nonlinear arbitrary order boundary value problem with advanced arguments. *Electron. J. Qual. Theory Differ. Equ.* **2012**, 15 (2012)
10. Zhou, Y, Jiao, F, Li, J: Existence and uniqueness for p -type fractional neutral differential equations. *Nonlinear Anal.* **71**(7-8), 2724-2733 (2009)
11. Zhao, Y, Sun, S, Han, Z, Zhang, M: Positive solutions for boundary value problems of nonlinear fractional differential equations. *Appl. Math. Comput.* **217**(16), 6950-6958 (2011)
12. Zhao, Y, Sun, S, Han, Z, Li, Q: Positive solutions to boundary value problems of nonlinear fractional differential equations. *Abstr. Appl. Anal.* **2011**, Article ID 390543 (2011). doi:10.1155/2011/390543
13. Feng, W, Sun, S, Han, Z, Zhao, Y: Existence of solutions for a singular system of nonlinear fractional differential equations. *Comput. Math. Appl.* **62**(3), 1370-1378 (2011)
14. Zhao, Y, Sun, S, Han, Z, Li, Q: The existence of multiple positive solutions for boundary value problems of nonlinear fractional differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **16**(4), 2086-2097 (2011)
15. Yang, X, Wei, Z, Dong, W: Existence of positive solutions for the boundary value problem of nonlinear fractional differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **17**(1), 85-92 (2012)
16. Xu, X, Jiang, D, Yuan, C: Multiple positive solutions for the boundary value problem of a nonlinear fractional differential equation. *Nonlinear Anal.* **71**(10), 4676-4688 (2009)
17. Nyamoradi, N, Bashiri, T: Multiple positive solutions for nonlinear fractional differential systems. *Fract. Differ. Calc.* **2**(2), 119-128 (2012)
18. Ahmad, B, Nieto, JJ: Sequential fractional differential equations with three-point boundary conditions. *Comput. Math. Appl.* **64**(10), 3046-3052 (2012)
19. Nyamoradi, N, Bashiri, T: Existence of positive solutions for fractional differential systems with multi point boundary conditions. *Ann. Univ. Ferrara* (2012). doi:10.1007/s11565-012-0160-x
20. Nyamoradi, N, Javidi, M: Existence of multiple positive solutions for fractional differential inclusions with m -point boundary conditions and two fractional orders. *Electron. J. Differ. Equ.* **2012**, 187 (2012)
21. Ahmad, B, Nieto, JJ, Alsaedi, A, El-Shahed, M: A study of nonlinear Langevin equation involving two fractional orders in different intervals. *Nonlinear Anal., Real World Appl.* **13**(2), 599-606 (2012)
22. Han, Z, Lu, H, Sun, S, Yang, D: Positive solutions to boundary value problems of p -Laplacian fractional differential equations with a parameter in the boundary conditions. *Electron. J. Differ. Equ.* **2012**, 213 (2012)
23. Chen, T, Liu, W: An anti-periodic boundary value problem for fractional differential equation with p -Laplacian operator. *Appl. Math. Lett.* **25**(11), 1671-1675 (2012)
24. Wang, J, Xiang, H, Liu, Z: Existence of concave positive solutions for boundary value problem of nonlinear fractional differential equation with p -Laplacian operator. *Int. J. Math. Math. Sci.* **2010**, Article ID 495138 (2010). doi:10.1155/2010/495138
25. Liang, S, Zhang, J: Positive solutions for boundary value problems of nonlinear fractional differential equation. *Nonlinear Anal.* **71**(11), 5545-5550 (2009)
26. Wang, J, Xiang, H: Upper and lower solutions method for a class of singular fractional boundary value problems with p -Laplacian operator. *Abstr. Appl. Anal.* **2010**, Article ID 971824 (2010). doi:10.1155/2010/971824
27. Lu, H, Han, Z: Existence of positive solutions for boundary value problem of fractional differential equation with p -Laplacian operator. In: *Proceedings of the 2012 International Conference on Computer Science and Information Processing (CSIP)*, pp. 582-585 (2012)
28. Chen, T, Liu, W, Hu, Z: A boundary value problem for fractional differential equation with p -Laplacian operator at resonance. *Nonlinear Anal.* **75**(6), 3210-3217 (2012)

29. Chai, G: Positive solutions for boundary value problem of fractional differential equation with p -Laplacian operator. *Bound. Value Probl.* **2012**, 18 (2012)
30. El-Shahed, M: Positive solutions for boundary value problems of nonlinear fractional differential equation. *Abstr. Appl. Anal.* **2007**, Article ID 10368 (2007). doi:10.1155/2007/10368
31. Leggett, RW, Williams, LR: Multiple positive fixed points of nonlinear operators on ordered Banach spaces. *Indiana Univ. Math. J.* **28**(4), 673-688 (1979)
32. Krasnosel'skii, MA: *Positive Solution of Operator Equation*. Noordhoff, Groningen (1964)
33. Isac, G: *Leray-Schauder Type Alternatives, Complementarity Problems and Variational Inequalities*. Springer, New York (2006)

doi:10.1186/1687-1847-2013-30

Cite this article as: Lu et al.: Existence on positive solutions for boundary value problems of nonlinear fractional differential equations with p -Laplacian. *Advances in Difference Equations* 2013 2013:30.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
