


RESEARCH

Open Access



# Existence results for a fraction hybrid differential inclusion with Caputo–Hadamard type fractional derivative

Mohammad Esmael Samei<sup>1\*</sup> , Vahid Hedayati<sup>2,3</sup> and Shahram Rezapour<sup>3,4</sup>

\*Correspondence:  
[mesamei@gmail.com](mailto:mesamei@gmail.com);  
[mesamei@basu.ac.ir](mailto:mesamei@basu.ac.ir)

<sup>1</sup>Department of Mathematics,  
Faculty of Science, Bu-Ali Sina  
University, Hamedan, Iran  
Full list of author information is  
available at the end of the article

## Abstract

In this manuscript, we talk over the existence of solutions of a class of hybrid Caputo–Hadamard fractional differential inclusions with Dirichlet boundary conditions. Our results are based on the Arzelà–Ascoli theorem and some suitable theorems of fixed point theory. As well, to illustrate our results, we confront the exceptional case of the fractional differential inclusions with examples.

**MSC:** Primary 26A33; 34A08; secondary 34K37

**Keywords:** Caputo–Hadamard fractional derivative; Fractional differential inclusion; Hybrid boundary value problem

## 1 Introduction

During the last decade, the subject of fractional differential equations and inclusions has been developed intensively (for example, see [1–8] and the references therein). An excellent account on the study of fractional differential equations can be found in [9–12]. Hybrid fractional differential equations and inclusions, certain classes of equations and inclusions involve the fractional derivative of an unknown function hybrid with the non-linearity depending on it. Also, they have been examined by several researchers (for example, see [13–15]). Most of the workplace on the fractional differential equations and inclusions is based on Riemann–Liouville, Caputo, and Hadamard type fractional derivatives. In 2012, Jarad et al. modified the Hadamard fractional derivative into a more suitable one having physical interpretable initial conditions similar to the singles in the Caputo setting and called it Caputo–Hadamard fractional derivative. To determine the properties of the modified derivative, refer to [16]. In 2011, Zhao et al. investigated the existence results for the initial value problems of hybrid fractional integro-differential equation

$$D^q \left[ \frac{\theta(t)}{u(t, \theta(t))} \right] = v(t, \theta(t)),$$

with real-valued initial condition  $\theta(0) = \theta_0$ , where  $D^q$  denotes the Riemann–Liouville fractional derivative of order  $0 < q < 1$ ,  $t \in J = [0, T]$ ,  $u \in C(J \times \mathbb{R}, \mathbb{R} - \{0\})$ , and  $v \in C(J \times \mathbb{R}, \mathbb{R})$  (for more details, see [13]). In 2016, Ahmad et al. studied the existence of solutions for a

nonlocal boundary value problem of hybrid fractional integro-differential inclusions

$$\begin{cases} {}^C D^\alpha \left[ \frac{x(t) - \sum_{i=1}^m I^{\beta_i} h_i(t, x(t))}{f(t, x(t))} \right] \in G(t, x(t)), \\ x(0) = \mu(x), \quad x(1) = A, \end{cases}$$

where  ${}^C D^\alpha$  and  $I^\alpha$  denote the Caputo fractional derivative and Riemann–Liouville integral of order  $\alpha$ , respectively,  $t \in J = [0, 1]$ ,  $m \in \mathbb{N}$ ,  $A \in \mathbb{R}$ ,  $1 < \alpha \leq 2$ ,  $\beta_i > 0$  for  $i = 1, 2, \dots, m$ , the functions  $f : J \times \mathbb{R} \rightarrow \mathbb{R} - \{0\}$ ,  $h_i : J \times \mathbb{R} \rightarrow \mathbb{R}$  for  $i = 1, 2, \dots, m$ ,  $\mu : C(J, \mathbb{R}) \rightarrow \mathbb{R}$ , and the multifunction  $G : J \times \mathbb{R} \rightarrow P(\mathbb{R})$  satisfies certain conditions (for more details, see [14]).

Motivated by these articles, we look into the existence of solutions for the following hybrid Caputo–Hadamard fractional differential inclusion:

$$\begin{cases} {}^C_H D^\alpha \left[ \frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))} \right] \in K(t, x(t)), \\ x(1) = \mu(x), \quad x(e) = \eta(x), \end{cases} \quad (1)$$

where  ${}^C_H D^\alpha$  and  $I^\alpha$  denote the Caputo–Hadamard fractional derivative and Hadamard integral of order  $\alpha$ , respectively,  $t \in J = [1, e]$ ,  $n, m \in \mathbb{N}$ ,  $1 < \alpha \leq 2$ ,  $\beta_i > 0$  for  $i = 1, 2, \dots, n$ ,  $\gamma_i > 0$  for  $i = 1, 2, \dots, m$ , the functions  $f : J \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ ,  $g : J \times \mathbb{R}^{m+1} \rightarrow \mathbb{R} - \{0\}$ ,  $h_i : J \times \mathbb{R} \rightarrow \mathbb{R}$  for  $i = 1, 2, \dots, n$ , functions  $\mu, \eta$  map  $C(J, \mathbb{R})$  into  $\mathbb{R}$ , and the multifunction  $K : J \times \mathbb{R} \rightarrow P(\mathbb{R})$  satisfies certain conditions.

The Hadamard fractional integral of order  $\alpha > 0$  of a function  $y$  is defined by

$$I_a^\alpha y(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds,$$

where  $a > 0$ ,  $t > a$ . In particular,  $I_1^\alpha y(t) := I^\alpha y(t)$ .

**Definition 1** Let  $n \in \mathbb{N}$ ,  $n - 1 < \alpha \leq n$ ,  $y \in AC_\delta^n[a, b]$ , where  $0 < a < b < \infty$  and

$$AC_\delta^n[a, b] = \left\{ y : [a, b] \rightarrow \mathbb{R} : \delta^{n-1} y(t) \in AC[a, b], \delta = t \frac{d}{dt} \right\}.$$

(i) If  $\alpha \neq n$ , then the Caputo–Hadamard fractional derivative of order  $\alpha$  is defined by

$${}^C_H D_a^\alpha y(t) = \frac{1}{\Gamma(n - \alpha)} \int_a^t \left( \ln \frac{t}{s} \right)^{n-\alpha-1} \delta^n \frac{y(s)}{s} ds = I_a^{n-\alpha} \delta^n y(t).$$

(ii) If  $\alpha = n$ , then the Caputo–Hadamard fractional derivative of order  $n$  is defined by

$${}^C_H D_a^n y(t) = \delta^n y(t).$$

In particular,  ${}^C_H D_1^0 y(t) := y(t)$  and  ${}^C_H D_1^\alpha y(t) := {}^C_H D^\alpha y(t)$ .

**Lemma 2** ([16]) Let  $\alpha > 0$ ,  $n = [\alpha] + 1$ , and  $\beta > 0$ . Then

$${}^C_H D_a^\alpha \left( \ln \frac{t}{a} \right)^{\beta-1} = \frac{\Gamma(\beta)}{\Gamma(\beta - \alpha)} \left( \ln \frac{t}{a} \right)^{\beta-\alpha-1} \quad \text{and} \quad {}^C_H D_a^\alpha \left( \ln \frac{t}{a} \right)^k = 0,$$

where  $\beta > n$ ,  $k = 0, 1, \dots, n - 1$ , and  ${}^C_H D_a^\alpha c = 0$  for all  $c \in \mathbb{R}$ .

**Lemma 3** ([16]) *Let  $n \in \mathbb{N}$ ,  $n - 1 < \alpha \leq n$ , and  $x \in AC^n_\delta[a, b]$ . Then one has*

$$I_a^\alpha ({}_H^C D_a^\alpha)x(t) = x(t) + \sum_{i=0}^{n-1} c_i \left( \ln \frac{t}{a} \right)^i,$$

where  $c_0, c_1, \dots, c_{n-1} \in \mathbb{R}$ .

**Lemma 4** ([16]) *Let  $n \in \mathbb{N}$ ,  $n - 1 < \alpha \leq n$ , and  $x \in C[a, b]$ . Then one has  ${}_H^C D_a^\alpha (I_a^\alpha)x(t) = x(t)$ .*

Let  $(X, \|\cdot\|)$  be a normed space. Denote the set of compact and convex subsets of  $X$ , the set of closed subsets of  $X$ , and the set of bounded subsets of  $X$  by  $P_{cp,cv}(X)$ ,  $P_{cl}(X)$ , and  $P_{bd}(X)$ , respectively. An element  $x \in X$  is called a fixed point of  $T : X \rightarrow 2^X$  whenever  $x \in T(x)$ . A multifunction  $T : J \rightarrow 2^{\mathbb{R}}$  is said to be measurable whenever the function  $t \mapsto d(x, T(t))$  is measurable for all  $x \in \mathbb{R}$ . A multifunction  $T : J \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a Caratheodory multifunction whenever the map  $t \mapsto T(t, x)$  is measurable for all  $x \in \mathbb{R}$  and the map  $x \mapsto T(t, x)$  is upper semi-continuous for almost all  $t \in J$ . Also, a Caratheodory multifunction  $T : J \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is called  $L^1$ -Caratheodory whenever, for each  $\rho > 0$ , there exists  $\phi_\rho \in L^1(J, \mathbb{R}^+)$  such that

$$\|T(t, x)\| = \sup\{|v| : v \in T(t, x)\} \leq \phi_\rho(t)$$

for all  $x \in \mathbb{R}$  with  $|x| \leq \rho$  and almost all  $t \in [1, e]$ .

Suppose that  $x$  belongs to  $C(J, \mathbb{R})$ . We define the set of the selections of a multifunction  $T : J \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  at  $x$  by

$$S_{T,x} = \{v \in L^1[1, e] | v(t) \in T(t, x(t)) : \forall t \in J\}.$$

**Lemma 5** ([17]) *The selections  $S_{T,x} \neq \emptyset$  for all  $x$  belong to  $C(J, \mathbb{R})$  whenever a multifunction  $T : J \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is an  $L^1$ -Caratheodory multifunction.*

**Lemma 6** ([17]) *Let  $T : J \times X \rightarrow P_{cp,cv}(X)$  be an  $L^1$ -Caratheodory multifunction and linear mapping  $\Theta : L^1(J, X) \rightarrow C(J, X)$  be continuous. Then the operator*

$$\begin{cases} \Theta \circ S_T : C(J, X) \rightarrow P_{cp,cv}(C(J), X), \\ (\Theta \circ S_T)(x) = \Theta(S_{T,x}), \end{cases}$$

*is a closed graph operator in  $C(J, X) \times C(J, X)$ .*

**Lemma 7** ([17]) *If a multifunction  $T : J \times X \rightarrow P_{cl}(X)$  is compact and has a closed graph, then it is upper semi-continuous.*

Let  $B_r(0)$  and  $\overline{B_r(0)}$  be the open and closed balls centered at the origin 0 of radius  $r$  in a Banach algebra  $X$ , respectively.

**Theorem 8** ([18]) *Suppose that  $A, B, C$  from  $X$  to  $X$  are three operators such that*

- $\frac{1}{A}$  is well-defined and one-to-one,
- $A$  and  $C$  are Lipschitzians with the Lipschitz constants  $q_1$  and  $q_2$ , respectively,

- (c)  $B$  is completely continuous,  
 (d)  $Mq_1 + q_2 < \frac{1}{2}$ , where

$$M = \|B(\overline{B_r(0)})\| = \sup\{|Bx| : x \in \overline{B_r(0)}\}. \quad (2)$$

Then either

- (i) the operator equation  $AxBx + Cx = x$  has a solution in  $\overline{B_r(0)}$ , or  
 (ii) there exists  $u \in X$  with  $\|u\| = r$  such that  $\mu u = AuBu + Cu$  for some  $\mu > 1$ .

**Theorem 9** ([19]) Let  $A, C : \overline{B_r(0)} \rightarrow X$  and  $B : \overline{B_r(0)} \rightarrow P_{cp,cv}(X)$  be three operators such that

- (a)  $A$  and  $C$  are single-valued Lipschitz with the Lipschitz constants  $q_1$  and  $q_2$ , respectively;  
 (b)  $B$  is upper-semi continuous and compact;  
 (c)  $Mq_1 + q_2 < \frac{1}{2}$ , where  $M$  is defined by (2).

Then either

- (i) the operator inclusion  $x \in AxBx + Cx$  has a solution, or  
 (ii) there exists  $u \in X$  with  $\|u\| = r$  such that  $\mu u \in AuBu + Cu$  for some  $\mu > 1$ .

## 2 Main results

In this section, by using fixed point theorems, we look into the existence of solution for the boundary value problem (1). In the first step, by applying the following lemma, we break down the solution kind of hybrid fractional differential inclusion (1).

**Lemma 10** Let  $y \in C(J, \mathbb{R})$  and

$$\frac{x(\cdot) - f(\cdot, x(\cdot), I^{\beta_1} h_1(\cdot, x(\cdot)), I^{\beta_2} h_2(\cdot, x(\cdot)), \dots, I^{\beta_n} h_n(\cdot, x(\cdot)))}{g(\cdot, x(\cdot), I^{\gamma_1} x(\cdot), I^{\gamma_2} x(\cdot), \dots, I^{\gamma_m} x(\cdot))} \in AC_{\delta}^2[1, e].$$

Then the unique solution of the hybrid fractional differential equation

$$\begin{aligned} {}^C_H D^{\alpha} \left[ \frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))} \right] \\ = y(t), \end{aligned} \quad (3)$$

via the boundary conditions  $x(1) = \mu(x)$  and  $x(e) = \eta(x)$ , is

$$\begin{aligned} x(t) = & g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t)) \\ & \times \left[ \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \right. \\ & \left. + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t \right] \\ & + f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t))) \end{aligned} \quad (4)$$

for all  $t \in J$ , where

$$a_0 = \frac{\mu(x) - f(1, \underbrace{\mu(x), 0, \dots, 0}_m)}{g(1, \underbrace{\mu(x), 0, \dots, 0}_m)},$$

$$a_1 = \frac{\eta(x) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))|_{t=e}}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))|_{t=e}}.$$

*Proof* Let  $x(t)$  be a solution of equation (3). By using the operator  $I^\alpha$  on both sides of equation (3) and applying Lemma 3, we have

$$\frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))}$$

$$= I^\alpha y(t) + c_0 + c_1 \ln t,$$

where  $c_0, c_1 \in \mathbb{R}$  denote arbitrary constants. Thus

$$x(t) = g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))$$

$$\times \left[ \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + c_0 + c_1 \ln t \right]$$

$$+ f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t))).$$

Now, by utilizing the boundary conditions, we conclude  $c_0 = a_0$  and

$$c_1 = a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0.$$

Conversely, it is easy to check that equation (4) satisfies the boundary conditions  $x(1) = \mu(x)$  and  $x(e) = \eta(x)$ . On the other hand, by deformation equation (4) to

$$\frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))}$$

$$= I^\alpha y(t) + a_0 + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t,$$

and by applying Lemmas 2 and 4, we have

$${}_H^C D^\alpha \left[ \frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))} \right] = y(t),$$

where  $a_0, a_1$  are defined in Lemma 10. This finishes our proof.  $\square$

A function  $x$  belonging to  $C(J, \mathbb{R})$  is a solution of problem (1) whenever it satisfies the boundary conditions, and there exists a function  $y \in S_{K,x}$  such that

$$\begin{aligned} x(t) = & g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t)) \\ & \times \left[ \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \right. \\ & \left. + \left( a_1 \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t \right] \\ & + f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t))) \end{aligned}$$

for  $t \in J$ , where

$$\begin{aligned} a_0 &= \frac{\mu(x) - f(1, \mu(x), \overbrace{0, \dots, 0}^n)}{g(1, \mu(x), \underbrace{0, \dots, 0}_m)}, \\ a_1 &= \frac{\eta(x) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))|_{t=e}}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))|_{t=e}}. \end{aligned}$$

Let  $\mathcal{X} = C(J, \mathbb{R})$  denote the Banach algebra of all continuous functions from  $J$  to  $\mathbb{R}$  endowed with the norm defined by  $\|x\| = \sup\{|x(t)| : t \in J\}$ .

**Theorem 11** Suppose that

(H<sub>1</sub>) The multifunction  $K : J \times \mathbb{R} \rightarrow P_{cp,cv}(\mathbb{R})$  is an  $L^1$ -Caratheodory such that

$$\|K(t, x)\| = \sup\{|y| : y \in K(t, x)\} \leq p(t)\varphi(|x|)$$

for  $(t, x) \in J \times \mathbb{R}$ , where  $p \in C(J, \mathbb{R}^+)$  and  $\varphi : [0, \infty) \rightarrow (0, \infty)$  is a continuous and increasing function;

(H<sub>2</sub>) Function  $f : J \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is continuous, and there exists  $\phi \in C(J, \mathbb{R}^+)$  such that

$$|f(t, x_1, x_2, \dots, x_{n+1}) - f(t, x'_1, x'_2, \dots, x'_{n+1})| \leq \phi(t) \sum_{i=1}^{n+1} |x_i - x'_i|,$$

where  $x_i, x'_i \in \mathbb{R}$  for  $i = 1, 2, \dots, n+1$ ,  $t \in J$ , and  $n \in \mathbb{N}$ ;

(H<sub>3</sub>) Function  $g : J \times \mathbb{R}^{m+1} \rightarrow \mathbb{R} - \{0\}$  is continuous, and there exists  $\psi \in C(J, \mathbb{R}^+)$  such that

$$|g(t, x_1, x_2, \dots, x_{m+1}) - f(t, x'_1, x'_2, \dots, x'_{m+1})| \leq \psi(t) \sum_{i=1}^{m+1} |x_i - x'_i|,$$

where  $x_i, x'_i \in \mathbb{R}$  for  $i = 1, 2, \dots, m+1$ ,  $t \in J$ , and  $m \in \mathbb{N}$ ;

(H<sub>4</sub>) Functions  $h_i : J \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous for  $i = 1, 2, \dots, n$ , and there exists  $p_i \in C(J, \mathbb{R}^+)$  for  $i = 1, 2, \dots, n$  such that  $|h_i(t, x) - h_i(t, x')| \leq p_i(t)|x - x'|$  for  $i = 1, 2, \dots, n$ , where  $x, x' \in \mathbb{R}$ ,  $n \in \mathbb{N}$ ,  $t \in J$ ;

(H<sub>5</sub>) There exist constants  $M_0, M_1 > 0$  such that

$$\left| \frac{\eta(x) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))|_{t=e}}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))|_{t=e}} \right| < M_1$$

and

$$\left| \frac{\mu(x) - f(1, \mu(x), \underbrace{0, \dots, 0}_n)}{g(1, \mu(x), \underbrace{0, \dots, 0}_m)} \right| < M_0$$

for all  $x \in C(J, \mathbb{R})$ .

The fractional differential inclusion (1) admits a solution whenever, for  $i = 1, 2, \dots, n$ ,

$$\begin{aligned} & \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \\ & + \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right) < \frac{1}{2}, \end{aligned}$$

where  $G_0 = \sup_{t \in J} |g(t, \underbrace{0, \dots, 0}_{m+1})|$ ,  $F_0 = \sup_{t \in J} |f(t, \underbrace{0, \dots, 0}_{n+1})|$ ,  $H_{i,0} = \sup_{t \in J} |h_i(t, 0)|$ , and

$$r > \frac{G_0 \left[ \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] + \|\phi\| \sum_{i=1}^n \frac{H_{i,0}}{\Gamma(\beta_i+1)} + F_0}{1 - \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) - \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right)}.$$

*Proof* Let  $\overline{B_r(0)} = \{x \in \mathcal{X} : \|x\| \leq r\}$ . Define operators  $A : \overline{B_r(0)} \rightarrow \mathcal{X}$ ,  $B : \overline{B_r(0)} \rightarrow P(\mathcal{X})$  and  $C : \overline{B_r(0)} \rightarrow \mathcal{X}$  by

$$\begin{aligned} (Ax)(t) &= g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t)), \\ (Bx)(t) &= \left\{ v(t) : v(t) = \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \right. \\ &\quad \left. + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t, y \in S_{K,x} \right\}, \\ (Cx)(t) &= f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t))), \end{aligned}$$

respectively. We establish that the operators  $A$ ,  $B$ , and  $C$  satisfy all the conditions of Theorem 9. Since the operator  $K$  is  $L^1$ -Caratheodory,  $S_{K,x} \neq \emptyset$  for all  $x \in \mathcal{X}$ . First, we show that  $Bx$  is a convex subset of  $\mathcal{X}$  for all  $x \in \overline{B_r(0)}$ . Consider  $x \in \overline{B_r(0)}$  and  $v_1, v_2 \in Bx$ . Choose  $y_1, y_2 \in S_{K,x}$  such that

$$\begin{aligned} v_1(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y_1(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_1(s)}{s} ds - a_0 \right) \ln t, \end{aligned}$$

$$\begin{aligned} v_2(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y_2(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_2(s)}{s} ds - a_0 \right) \ln t. \end{aligned}$$

At present, for any  $0 \leq \lambda \leq 1$ , we have

$$\begin{aligned} \lambda v_1(t) + (1-\lambda)v_2(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{\lambda y_1(s) + (1-\lambda)y_2(s)}{s} ds + a_0 \\ &\quad + \left[ a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{\lambda y_1(s) + (1-\lambda)y_2(s)}{s} ds - a_0 \right] \ln t. \end{aligned}$$

Since  $K$  has convex values, hence  $S_{K,x}$  is convex and  $\lambda y_1 + (1-\lambda)y_2 \in S_{K,x}$ . So  $\lambda v_1 + (1-\lambda)v_2 \in Bx$ , which results in that  $Bx$  is a convex subset of  $\mathcal{X}$ . Now, we show that  $Bx$  is a compact subset of  $\mathcal{X}$  for all  $x \in \overline{B_r(0)}$ . Let  $x \in \overline{B_r(0)}$  and  $\{v_n\}$  be a sequence in  $Bx$ . For each  $n$ , choose  $y_n \in S_{K,x}$  such that

$$\begin{aligned} v_n(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds - a_0 \right) \ln t. \end{aligned}$$

Since  $K$  is compact-valued,  $\{y_n\}$  has a subsequence which pointwise converges to  $y$ . We denote this subsequence again by  $\{y_n\}$ . Since  $y(t) \in K(t, x(t))$  and  $\|K(t, x(t))\| \leq p(t)\varphi(|x(t)|)$ , by using Lebesgue dominated convergence theorem  $y \in S_{K,x}$ , we have

$$\begin{aligned} v_n(t) \rightarrow v(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t \end{aligned}$$

pointwise on  $J$ . Let  $t_1 < t_2 \in J$ . Then we have

$$\begin{aligned} |v_n(t_2) - v_n(t_1)| &\leq \frac{1}{\Gamma(\alpha)} \left| \int_1^{t_2} \left( \ln \frac{t_2}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds \right. \\ &\quad \left. - \int_1^{t_1} \left( \ln \frac{t_1}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds \right| \\ &\quad + \left| a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds - a_0 \right| \\ &\quad \times (\ln t_2 - \ln t_1) \\ &\leq \frac{1}{\Gamma(\alpha)} \int_1^{t_1} \left| \left( \ln \frac{t_2}{s} \right)^{\alpha-1} - \left( \ln \frac{t_1}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} \left| \left( \ln \frac{t_2}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds \end{aligned}$$



$$+ \left( |a_1| + \frac{1}{\Gamma(\alpha)} \int_1^e \left| \left( \ln \frac{1}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds + |a_0| \right) \\ \times (\ln t_2 - \ln t_1).$$

As  $t_1 \rightarrow t_2$ , the right-hand side of the above inequality tends to zero. Hence  $\{v_n\}$  is an equicontinuous sequence. Therefore  $\{v_n\}$  has a subsequence that uniformly converges to  $v \in Bx$ , which proves that  $Bx$  is a compact subset of  $\mathcal{X}$ . In this section, we shall show that the operator  $B$  is compact on  $\overline{B_r(0)}$ . Let  $S \subset \overline{B_r(0)}$ . We prove that  $\overline{B(S)}$  is a compact set in  $\mathcal{X}$ . To do this, we apply Arzelà–Ascoli theorem and show that  $B(S)$  is a uniformly bounded and equicontinuous set. Let  $x \in S$ ,  $v \in Bx$ , and  $t_1 < t_2 \in J$ . Choose  $y \in S_{K,x}$  such that

$$v(t) = \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \\ + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t.$$

Therefore, we have

$$|v(t)| \leq \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{|y(s)|}{s} ds + |a_0| \\ + \left( |a_1| + \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{|y(s)|}{s} ds + |a_0| \right) \ln t \\ \leq \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{\|p\|\varphi(r)}{s} ds \\ + M_0 + \left( M_1 + \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{\|p\|\varphi(r)}{s} ds + M_0 \right) \\ \leq \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1.$$

Thus,  $\|v\| \leq \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1$ . Also, we have

$$|v(t_2) - v(t_1)| \leq \frac{1}{\Gamma(\alpha)} \left| \int_1^{t_2} \left( \ln \frac{t_2}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - \int_1^{t_1} \left( \ln \frac{t_1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds \right| \\ + \left| a_0 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right| (\ln t_2 - \ln t_1) \\ \leq \frac{1}{\Gamma(\alpha)} \int_1^{t_1} \left| \left( \ln \frac{t_2}{s} \right)^{\alpha-1} - \left( \ln \frac{t_1}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds \\ + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} \left| \left( \ln \frac{t_2}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds \\ + \left( |a_1| + \frac{1}{\Gamma(\alpha)} \int_1^e \left| \left( \ln \frac{1}{s} \right)^{\alpha-1} \right| \frac{\|p\|\varphi(r)}{s} ds + |a_0| \right) \\ \times (\ln t_2 - \ln t_1).$$

As  $t_1 \rightarrow t_2$ , the right-hand side of the above inequality tends to zero. Thus,  $B(S)$  is a uniformly bounded and equicontinuous set in  $\mathcal{X}$  and  $B$  is a compact operator. In the following,

we shall show that the operator  $B$  is upper semi-continuous on  $\overline{B_r(0)}$ . To do this, we apply Lemma 7 and show that  $B$  has a closed graph. Let  $x_n \rightarrow x_0$  and  $v_n \in Bx_n$  ( $n \geq 1$ ) with  $v_n \rightarrow v_0$ . We prove that  $v_0 \in Bx_0$ . For each natural number  $n$ , choose  $y_n \in S_{K,x_n}$  such that

$$\begin{aligned} v_n(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_n(s)}{s} ds - a_0 \right) \ln t. \end{aligned}$$

Define the continuous linear operator  $\theta : L^1(J, \mathbb{R}) \rightarrow \mathcal{X}$  by

$$\begin{aligned} \theta(y)(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t. \end{aligned}$$

By using Lemma 6,  $\theta \circ S_K$  is a closed graph operator. Since  $v_n \in \theta(S_{K,x_n})$  for all  $n$  and  $x_n \rightarrow x_0$ , there exists  $y_0 \in S_{K,x_0}$  such that

$$\begin{aligned} v_0(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y_0(s)}{s} ds + a_0 \\ &\quad + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y_0(s)}{s} ds - a_0 \right) \ln t. \end{aligned}$$

Hence,  $v_0 \in Bx_0$ . This implies that  $B$  has a closed graph, and as a result,  $B$  is upper semi-continuous. Now, we show that the operators  $A, C$  are Lipschitz on  $\overline{B_r(0)}$ . Let  $x, y \in \overline{B_r(0)}$ . Then we have

$$\begin{aligned} |Ax(t) - Ay(t)| &= |g(t, x(t), I^{\gamma_1}x(t), I^{\gamma_2}x(t), \dots, I^{\gamma_m}x(t)) \\ &\quad - g(t, y(t), I^{\gamma_1}y(t), I^{\gamma_2}y(t), \dots, I^{\gamma_m}y(t))| \\ &\leq \psi(t) \left[ |x(t) - y(t)| + \sum_{i=1}^m |I^{\gamma_i}x(t) - I^{\gamma_i}y(t)| \right] \\ &\leq \psi(t) \left[ |x(t) - y(t)| \right. \\ &\quad \left. + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i)} \int_1^t \left( \ln \frac{t}{s} \right)^{\gamma_i-1} \frac{|x(s) - y(s)|}{s} ds \right] \\ &\leq \|\psi\| \left[ \|x - y\| + \|x - y\| \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i + 1)} \right], \end{aligned}$$

which implies that

$$\|Ax - Ay\| \leq \underbrace{\|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i + 1)} \right)}_{q_1} \|x - y\| = q_1 \|x - y\|.$$

Also,

$$\begin{aligned}
 |Cx(t) - Cy(t)| &= |f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t))) \\
 &\quad - f(t, y(t), I^{\beta_1} h_1(t, y(t)), I^{\beta_2} h_2(t, y(t)), \dots, I^{\beta_n} h_n(t, y(t)))| \\
 &\leq \phi(t) \left[ |x(t) - y(t)| \right. \\
 &\quad \left. + \sum_{i=1}^n |I^{\beta_i} h_i(t, x(t)) - I^{\beta_i} h_i(t, y(t))| \right] \\
 &\leq \phi(t) \left[ |x(t) - y(t)| \right. \\
 &\quad \left. + \sum_{i=1}^n \frac{1}{\Gamma(\beta_i)} \int_1^t \left( \ln \frac{t}{s} \right)^{\beta_i-1} \frac{|h_i(s, x(s)) - h_i(s, y(s))|}{s} ds \right] \\
 &\leq \phi(t) \left[ |x(t) - y(t)| \right. \\
 &\quad \left. + \sum_{i=1}^n \frac{1}{\Gamma(\beta_i)} \int_1^t \left( \ln \frac{t}{s} \right)^{\beta_i-1} \frac{p_i(s) |x(s) - y(s)|}{s} ds \right] \\
 &\leq \|\phi\| \left( \|x - y\| + \|x - y\| \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i + 1)} \right) \\
 &= \underbrace{\|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i + 1)} \right)}_{q_2} \|x - y\| = q_2 \|x - y\|.
 \end{aligned}$$

Since

$$M = \|B(\overline{B_r(0)})\| = \sup_{x \in B_r(0)} \{\sup Bx(t), t \in J\} \leq \frac{2\|p\|\varphi(r)}{\Gamma(\alpha + 1)} + 2M_0 + M_1,$$

we have

$$\begin{aligned}
 Mq_1 + q_2 &\leq \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha + 1)} + 2M_0 + M_1 \right) \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i + 1)} \right) \\
 &\quad + \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i + 1)} \right) < \frac{1}{2}.
 \end{aligned}$$

Hence, all the conditions of Theorem 9 hold. Let  $u \in \mathcal{X}$ ,  $\|u\| = r$ ,  $\mu > 1$ , and  $\mu u \in AuBu + Cu$ . Then we have

$$\begin{aligned}
 |u(t)| &\leq |g(t, u(t), I^{\gamma_1} u(t), I^{\gamma_2} u(t), \dots, I^{\gamma_m} u(t))| \\
 &\quad \times \left| \frac{1}{\Gamma(\alpha)} \int_1^t \left( \ln \frac{t}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds + a_0 \right|
 \end{aligned}$$

$$\begin{aligned}
& + \left( a_1 - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \ln \frac{1}{s} \right)^{\alpha-1} \frac{y(s)}{s} ds - a_0 \right) \ln t \Big| \\
& + |f(t, u(t), I^{\beta_1} h_1(t, u(t)), I^{\beta_2} h_2(t, u(t)), \dots, I^{\beta_n} h_n(t, u(t)))| \\
& \leq \left[ |g(t, u(t), I^{\gamma_1} u(t), I^{\gamma_2} u(t), \dots, I^{\gamma_m} u(t)) - g(t, \underbrace{0, 0, \dots, 0}_{m+1})| \right. \\
& \quad + |g(t, \underbrace{0, 0, \dots, 0}_{m+1})| \Big] \left( \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) \\
& \quad + |f(t, u(t), I^{\beta_1} h_1(t, u(t)), I^{\beta_2} h_2(t, u(t)), \dots, I^{\beta_n} h_n(t, u(t))) \\
& \quad - f(t, \underbrace{0, 0, \dots, 0}_{n+1})| + |f(t, \underbrace{0, 0, \dots, 0}_{n+1})| \\
& \leq \left[ \psi(t) \left( |u(t)| + \sum_{i=1}^m |I^{\gamma_i} u(t)| \right) + G_0 \right] \left[ \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] \\
& \quad + \phi(t) \left( |u(t)| + \sum_{i=1}^n |I^{\beta_i} h_i(t, u(t))| \right) + F_0 \\
& \leq \left[ \|\psi\| \|u\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) + G_0 \right] \left[ \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] \\
& \quad + \|\phi\| \left( \|u\| + \sum_{i=1}^n \frac{\|p_i\| \|u\| + H_{i,0}}{\Gamma(\beta_i+1)} \right) + F_0
\end{aligned}$$

for some  $y \in S_{K,u}$ . Therefore

$$\begin{aligned}
\|u\| & \leq \left[ \|\psi\| \|u\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) + G_0 \right] \left[ \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] \\
& \quad + \|\phi\| \left( \|u\| + \sum_{i=1}^n \frac{\|p_i\| \|u\| + H_{i,0}}{\Gamma(\beta_i+1)} \right) + F_0.
\end{aligned}$$

As a result, we have

$$\|u\| \leq \frac{G_0 \left[ \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] + \|\phi\| \sum_{i=1}^n \frac{H_{i,0}}{\Gamma(\beta_i+1)} + F_0}{1 - \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \left( \frac{2\|p\|\varphi(\|u\|)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) - \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right)}.$$

Since  $\|u\| = r$ , we have

$$r \leq \frac{G_0 \left[ \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] + \|\phi\| \sum_{i=1}^n \frac{H_{i,0}}{\Gamma(\beta_i+1)} + F_0}{1 - \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) - \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right)} < r.$$

This is an apparent contradiction, and conclusion (ii) of Theorem 9 does not hold. Hence, there exists  $x \in \overline{B_r(0)}$  such that  $x \in Ax \cup Bx + Cx$ , and it is a solution of hybrid fractional differential inclusion (1).  $\square$

In the following, we review the existence of solution for the hybrid fractional differential equation

$$\begin{cases} {}^C_H D^\alpha \left[ \frac{x(t) - f(t, x(t), I^{\beta_1} h_1(t, x(t)), I^{\beta_2} h_2(t, x(t)), \dots, I^{\beta_n} h_n(t, x(t)))}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t), \dots, I^{\gamma_m} x(t))} \right] = k(t, x(t)), \\ x(1) = \mu(x), \quad x(e) = \eta(x), \end{cases} \quad (5)$$

where  ${}^C_H D^\alpha$  and  $I^\alpha$  denote the Caputo–Hadamard fractional derivative and Hadamard integral of order  $\alpha$ , respectively,  $t \in J = [1, e]$ ,  $n, m \in \mathbb{N}$ ,  $1 < \alpha \leq 2$ ,  $\beta_i > 0$  ( $i = 1, 2, \dots, n$ ),  $\gamma_i > 0$  ( $i = 1, 2, \dots, m$ ), the functions  $f : J \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ ,  $g : J \times \mathbb{R}^{m+1} \rightarrow \mathbb{R} - \{0\}$ ,  $h_i : J \times \mathbb{R} \rightarrow \mathbb{R}$  ( $i = 1, 2, \dots, n$ ),  $k : J \times \mathbb{R} \rightarrow \mathbb{R}$ , and  $\mu, \eta : C(J, \mathbb{R}) \rightarrow \mathbb{R}$  satisfy certain conditions. It is clear that equation (5) is a special case of inclusion (1), where  $K(t, x(t)) = \{k(t, x(t))\}$ . With argument similar to the proof of Theorem 11, and applying Theorem 8, we can prove the following theorem.

**Theorem 12** *Suppose that*

(H<sub>1</sub>') *The function  $k : J \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous map such that  $|k(t, x)| \leq p(t)\varphi(|x|)$  for all  $(t, x) \in J \times \mathbb{R}$ , where  $p \in C(J, \mathbb{R}^+)$  and  $\varphi : [0, \infty) \rightarrow (0, \infty)$  is a continuous and increasing function;*

(H<sub>2</sub>') *If  $\frac{(x_1, x_2, \dots, x_{m+1})}{g(t, x_1, x_2, \dots, x_{m+1})} = \frac{(y_1, y_2, \dots, y_{m+1})}{g(t, y_1, y_2, \dots, y_{m+1})}$ , then*

$$(x_1, x_2, \dots, x_{m+1}) = (y_1, y_2, \dots, y_{m+1})$$

*for all  $t \in J$  and  $(x_1, x_2, \dots, x_{m+1}) \in \mathbb{R}^{m+1}$ .*

*Also, assume that conditions (H<sub>2</sub>), (H<sub>3</sub>), (H<sub>4</sub>), and (H<sub>5</sub>) in Theorem 11 hold. If*

$$\begin{aligned} & \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha + 1)} + 2M_0 + M_1 \right) \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i + 1)} \right) \\ & + \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i + 1)} \right) < \frac{1}{2}, \end{aligned}$$

*where  $r$  is the constant that was introduced in Theorem 11, then the fractional differential equation (5) has at least one solution on  $J$ .*

**Example 1** Consider the following hybrid Caputo–Hadamard fractional inclusion:

$$\begin{cases} {}^C_H D^{\frac{3}{2}} \left[ \frac{x(t) - 3 - \frac{t^2}{100(1+t^2)} (\sin x(t) + \cos(I^{\frac{1}{3}} (\sin t \tan^{-1} x(t))) + \frac{|\frac{1}{2}(\frac{t^2|x(t)|}{5(1+|x(t)|)})|}{1 + |\frac{1}{2}(\frac{t^2|x(t)|}{5(1+|x(t)|)})|}}{1 + \frac{e^{-\pi t}}{1000(1+e^{-\pi t})} (\frac{|x(t)| + |\frac{2}{3}x(t)| + |\frac{3}{4}x(t)|}{1 + |x(t)| + |\frac{2}{3}x(t)| + |\frac{3}{4}x(t)|})} \right] \in [0, e^t \sin t + 2], \\ x(1) = \sin(x(\frac{3}{2})), \quad x(e) = \cos(x(\frac{5}{2})). \end{cases} \quad (6)$$

Here  $\alpha = \frac{3}{2}$ ,  $m = n = 2$ ,  $\beta_1 = \frac{1}{3}$ ,  $\beta_2 = \frac{1}{2}$ ,  $\gamma_1 = \frac{2}{3}$ ,  $\gamma_2 = \frac{3}{4}$ ,

$$g(t, x_1, x_2, x_3) = 1 + \frac{(|x_1| + |x_2| + |x_3|)e^{-\pi t}}{1000(1 + (|x_1| + |x_2| + |x_3|)(1 + e^{-\pi t}))},$$

$$f(t, x_1, x_2, x_3) = 3 + \frac{t^2}{100(1+t^2)} \left( \sin x_1 + \cos x_2 + \frac{|x_3|}{1+|x_3|} \right),$$

$K(t, x) = [0, e^t \sin x + 2]$ ,  $h_1(t, x) = \sin t \tan^{-1} x$ ,  $h_2(t, x) = \frac{t^2|x|}{5(1+|x|)}$ ,  $\mu(y) = \sin(y(\frac{3}{2}))$ , and  $\eta(y) = \cos(y(\frac{5}{2}))$  for all  $x, x_1, x_2$ , and  $x_3 \in \mathbb{R}$ ,  $t \in J$  and  $y \in C(J, \mathbb{R})$ . Obviously,

$$|g(t, x_1, x_2, x_3) - g(t, x'_1, x'_2, x'_3)| \leq \frac{e^{-\pi t}}{1000(1+e^{-\pi t})} \sum_{i=1}^3 |x_i - x'_i|,$$

$$|f(t, x_1, x_2, x_3) - f(t, x'_1, x'_2, x'_3)| \leq \frac{t^2}{100(1+t^2)} \sum_{i=1}^3 |x_i - x'_i|,$$

$$|h_1(t, x_1) - h_1(t, x'_1)| \leq \sin t |x_1 - x'_1|, |h_2(t, x_1) - h_2(t, x'_1)| \leq \frac{t^2}{5} |x_1 - x'_1|,$$

$$\left| \frac{\eta(y) - f(t, y(t), I^{\beta_1} h_1(t, y(t)), I^{\beta_2} h_2(t, y(t)))|_{t=e}}{g(t, x(t), I^{\gamma_1} x(t), I^{\gamma_2} x(t))|_{t=e}} \right| < 4.03,$$

$$\left| \frac{\mu(y) - f(1, \mu(y), 0, 0)}{g(1, \mu(y), 0, 0)} \right| < 4.03,$$

$\|K(t, x)\| \leq e^t + 2$ ,  $F_0 \leq 3.01$ ,  $G_0 = 1$ , and  $H_{1,0} = H_{2,0} = 0$  for all  $x_1, x_2, x_3, x'_1, x'_2, x'_3$ , and  $x \in \mathbb{R}$ ,  $t \in J$ , and  $y \in C(J, \mathbb{R})$ . Now, by setting  $p(t) = e^t + 2$ ,  $\varphi(x) = 1$ ,  $\psi(t) = \frac{e^{-\pi t}}{1000(1+e^{-\pi t})}$ ,  $\phi(t) = \frac{t^2}{100(1+t^2)}$ ,  $p_1(t) = \sin t$ ,  $p_2(t) = \frac{t^2}{5}$ ,  $M_0 = 4.03$ , and  $M_1 = 4.03$  for all  $t \in J$  and  $x \in \mathbb{R}^+$ , we have  $\|p\| = e^e + 2$ ,  $\|\phi\| \leq \frac{1}{100}$ ,  $\|\psi\| \leq \frac{1}{1000}$ ,  $\|p_1\| = 1$ , and  $\|p_2\| = \frac{e^2}{5}$ . Hence

$$\begin{aligned} & \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \\ & + \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right) \leq 0.14606 < \frac{1}{2}. \end{aligned}$$

Therefore, all the conditions in Theorem 11 are satisfied, and problem (6) has at least one solution on  $\overline{B_r(0)}$ , where

$$r > \frac{G_0 \left[ \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right] + \|\phi\| \sum_{i=1}^n \frac{H_{i,0}}{\Gamma(\beta_i+1)} + F_0}{1 - \|\psi\| \left( 1 + \sum_{i=1}^m \frac{1}{\Gamma(\gamma_i+1)} \right) \left( \frac{2\|p\|\varphi(r)}{\Gamma(\alpha+1)} + 2M_0 + M_1 \right) - \|\phi\| \left( 1 + \sum_{i=1}^n \frac{\|p_i\|}{\Gamma(\beta_i+1)} \right)} \geq 47.68.$$

#### Acknowledgements

The authors would like to thank the editors and the anonymous reviewers for their constructive comments and suggestions that have helped to improve the present paper. Also, the authors are very grateful to the submission editor of Springer Nature, Dhanshi Nagargoje, for helpful guidance. We thank Dear Vanesa Mae Ledesma for good guidance and following the paper with patience.

#### Funding

Not applicable.

#### Availability of data and materials

Not applicable.

#### Ethics approval and consent to participate

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

**Consent for publication**

Not applicable.

**Authors' contributions**

All authors contributed equally and significantly in this manuscript, and they read and approved the final manuscript.

**Author details**

<sup>1</sup>Department of Mathematics, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran. <sup>2</sup>Department of Mathematics, Shahid Madani Educational Institution, Hamedan, Iran. <sup>3</sup>Department of Mathematics, Azarbaijan Shahid Madani University, Tabriz, Iran. <sup>4</sup>Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 15 January 2019 Accepted: 10 April 2019 Published online: 30 April 2019

**References**

1. Hedayati, V., Rezapour, S.: The existence of solution for a  $k$ -dimensional system of fractional differential inclusions with anti-periodic boundary value conditions. *Filomat* **30**(6), 1601–1613 (2016). <https://doi.org/10.2298/FIL1606601H>
2. Ahmad, B., Ntouyas, S.K.: Boundary value problem for fractional differential inclusions with four-point integral boundary conditions. *Surv. Math. Appl.* **6**, 175–193 (2011)
3. Agarwal, R.P., Ahmad, B.: Existence theory for anti-periodic boundary value problems of fractional differential equations and inclusions. *Comput. Math. Appl.* **62**, 1200–1214 (2011). <https://doi.org/10.1016/j.camwa.2011.03.001>
4. Baleanu, D., Nazemi, S.Z., Rezapour, S.: Existence and uniqueness of solutions for multi-term nonlinear fractional integro-differential equations. *Adv. Differ. Equ.* **2013**, 368 (2013). <https://doi.org/10.1186/1687-1847-2013-368>
5. Agarwal, R.P., Baleanu, D., Hedayati, V., Rezapour, S.: Two fractional derivative inclusion problems via integral boundary condition. *Appl. Math. Comput.* **257**, 205–212 (2015). <https://doi.org/10.1016/j.amc.2014.10.082>
6. Rezapour, S., Hedayati, V.: On a Caputo fractional differential inclusion with integral boundary condition for convex-compact and nonconvex-compact valued multifunctions. *Kragujev. J. Math.* **41**(1), 143–158 (2017). <https://doi.org/10.5937/KgJMath1701143R>
7. Ghorbanian, R., Hedayati, V., Postolache, M., Rezapour, S.: On a fractional differential inclusion via a new integral boundary condition. *J. Inequal. Appl.* **2014**, 319 (2014). <https://doi.org/10.1186/1029-242X-2014-319>
8. Agarwal, R.P., Baleanu, D., Hedayati, V., Rezapour, S.: Two fractional derivative inclusion problems via integral boundary condition. *Appl. Math. Comput.* **257**, 205–212 (2015). <https://doi.org/10.1016/j.amc.2014.10.082>
9. Hilfer, R.: *Applications of Fractional Calculus in Physics*. World Scientific, Singapore (2000). <https://doi.org/10.1142/3779>
10. Kilbas, A.A., Srivastava, H.M., Trujillo, J.J.: *Theory and Applications of Fractional Differential Equations*. North-Holland Mathematics Studies. Elsevier, Amsterdam (2006)
11. Podlubny, I.: *Fractional Differential Equations*. Academic Press, Slovak Republic (1999)
12. Kac, V., Cheung, P.: *Quantum Calculus*. Universitext. Springer, New York (2002)
13. Zhao, Y., Sun, S., Han, Z., Li, Q.: Theory of fractional hybrid differential equations. *Comput. Math. Appl.* **62**(3), 1312–1324 (2011). <https://doi.org/10.1016/j.camwa.2011.03.041>
14. Ahmad, B., Ntouyas, S., Tariboon, J.: On hybrid Caputo fractional integro-differential inclusions with nonlocal conditions. *J. Nonlinear Sci. Appl.* **9**, 4235–4246 (2016). <https://doi.org/10.22436/jnsa.009.06.65>
15. Baleanu, D., Hedayati, V., Rezapour, S., Al Qurashi, M.M.: On two fractional differential inclusions. *SpringerPlus* **5**(1), 882 (2016). <https://doi.org/10.1186/s40064-016-2564-z>
16. Jarad, F., Abdeljawad, T., Baleanu, D.: Caputo-type modification of the Hadamard fractional derivatives. *Adv. Differ. Equ.* **2012**, 142 (2012). <https://doi.org/10.1186/1687-1847-2012-142>
17. Djebali, S., Górniewicz, L., Ouahab, A.: *Solution Sets for Differential Equations and Inclusions*. De Gruyter Series in Nonlinear Analysis and Applications. Germany (2012). <https://doi.org/10.1515/9783110293562>
18. Dhage, B.: On some nonlinear alternatives of Leray-Schauder type and functional integral equations. *Arch. Math.* **42**, 11–23 (2006)
19. Dhage, B.: On solvability of operator inclusions  $x \in axb + cx$  in Banach algebras and differential inclusions. *Commun. Appl. Anal.* **14**(4), 567–596 (2010)